

1 Factors Associated with Lumbar Intervertebral Disc Degeneration in the Elderly

2

3 Mika Hangai, MD<sup>a</sup>, Koji Kaneoka, MD, PhD<sup>b\*</sup>, Shinya Kuno, PhD<sup>c</sup>, Shiro Hinotsu, MD,  
4 PhD<sup>d</sup>, Masataka Sakane, MD, PhD<sup>d</sup>, Naotaka Mamizuka, MD, PhD<sup>a</sup>,  
5 Shinsuke Sakai, MD<sup>a</sup>, Naoyuki Ochiai, MD, PhD<sup>d</sup>

6

7 <sup>a</sup> Department of Orthopaedic Surgery, Institute of Clinical Medicine, Graduate School of  
8 Comprehensive Human Sciences, University of Tsukuba, Japan

9 <sup>b</sup> Faculty of Sports Science, Waseda University, Japan

10 <sup>c</sup> Institute of Health and Sports Sciences, Graduate School of Comprehensive Human Sciences,  
11 University of Tsukuba, Japan

12 <sup>d</sup> Critical-Path Research and Education Integrated Leading Center, University of Tsukuba,  
13 Japan

14 \* Corresponding author

15

16 Abstract

17 BACKGROUND CONTEXT: Lumbar intervertebral disc degeneration (DD) precedes  
18 degenerative diseases of the lumbar spine. Various factors in addition to normal aging are  
19 reported to be associated with DD, and recently atherosclerosis and risk factors for  
20 cardiovascular diseases (cardiovascular risk factors) have received much attention; however, the  
21 links between these risk factors and DD are unclear.

22 PURPOSE: By correlating magnetic resonance images with suspected degenerative disc risk  
23 factors such as obesity, cardiovascular risk factors and atherosclerosis we hope to clarify the

1 factors associated with DD.

2 STUDY DESIGN/SETTING: An observational study.

3 PATIENT SAMPLE: Two hundred seventy adults (51 - 86 years old) who participated in a health  
4 promotion program.

5 OUTCOME MEASURES: DD evaluated based on the signal intensity of MR T2-weighted  
6 mid-sagittal images of the lumbar spine.

7 METHODS: Age, gender, body mass index (BMI), low-density lipoprotein cholesterol (LDLc),  
8 triglyceride, glycosylated hemoglobin, brachial-ankle pulse wave velocity as an index of  
9 atherosclerosis, osteo-sono-assessment index calculated from quantitative ultrasound assessment  
10 of the calcaneus as an index of bone mineral density, history of low back pain, smoking and  
11 drinking habits, and physical loading related to occupations and sports were assessed. The  
12 univariate relationships between DD and the variables were evaluated, and finally, odds ratios  
13 (OR) and 95% confidence intervals for the associations of each factor with DD were calculated  
14 using logistic regression at each disc level.

15 RESULTS: Aging correlated significantly with DD of L1/2 (OR, 2.14), L2/3 (OR, 3.56), L3/4  
16 (OR, 2.84) and L4/5 (OR, 3.05); high BMI, with L2/3 (OR, 2.98), L3/4 (OR, 3.58), L4/5 (OR,  
17 2.32) and L5/S1 (OR, 3.34); high LDLc, with L4/5 (OR, 2.65); occupational lifting, with L1/2  
18 (OR, 4.25); and sports activities, with L5/S1 (OR, 3.36).

19 CONCLUSIONS: Aging, high BMI, high LDLc, occupational lifting and sports activities are  
20 associated with DD. The results of this study raise our index of suspicion that cardiovascular risk  
21 factors and particular physical loading may contribute to DD; however additional studies are  
22 required to further investigate associations between DD and these factors.

23

1 Introduction

2 Lumbar intervertebral discs degenerate with normal aging [1-4]. One study using magnetic  
3 resonance imaging (MRI) on healthy women reported about 30% of the subjects in their twenties  
4 had lumbar intervertebral disc degeneration (DD), while about 90% of the subjects in their  
5 seventies had DD [4]. Nevertheless, DD also precedes lumbar degenerative diseases, such as  
6 lumbar disc herniation and lumbar spinal canal stenosis; therefore, various studies aiming to  
7 identify risk factors for DD have been completed. The importance of heredity has become  
8 increasingly clear following the twin studies of Battie et al.[1,5] and Sambrook et al.[6]. It is  
9 reported that individuals with specific gene polymorphisms might develop DD earlier than  
10 individuals without these genetic variants [7-9]. In addition, acquired factors, including obesity  
11 [3,10,11], diabetes mellitus [12,13], smoking [12,14,15], physical loading related to occupations  
12 and sports [5,16-19], and bone mineral density (BMD) [3,20,21], have been associated with DD.  
13 In industrialized nations, the human life span has been extended because of medical advances  
14 and improvements in eating habits; therefore, we are concerned that the number of patients who  
15 have lumbar degenerative diseases will increase, and thereby medical expenses also. Accordingly,  
16 we need to clarify which acquired factors promote DD so as to establish preventive measures  
17 against DD.

18 Recently, suspicions have arisen that atherosclerosis and occlusion of the abdominal aorta and  
19 the lumbosacral arteries result in insufficient blood flow and nutrition for the lumbar  
20 intervertebral discs and, consequently, might contribute to DD [2,22-24]. Kauppila et al.  
21 examined and confirmed this association between DD and atherosclerosis using autopsies,  
22 radiographs and MR aortographs [2,22,23]. It is well known atherosclerosis is accelerated by  
23 some factors which are known as cardiovascular risk factors. Interestingly, obesity, diabetes

1 mellitus and smoking are not only suspected risk factors for DD, but are also cardiovascular risk  
2 factors. Furthermore, obesity and smoking is highly correlated with lifestyle, and type 2 diabetes  
3 can be triggered by overeating or lack of exercise in people predisposed to type 2 diabetes by  
4 specific hereditary factors. Jhavar et al. noted this, and prospectively and longitudinally have  
5 examined the associations between lumbar disc herniation and cardiovascular risk factors in  
6 nurses [12]. They found that lumbar disc herniation was associated with diabetes, high  
7 cholesterol, hypertension, smoking and a myocardial infarction before age 60. If an association  
8 between atherosclerosis, cardiovascular risk factors and DD is conclusively shown, this suggests  
9 that behaviors that help prevent atherosclerosis and cardiovascular diseases, which have only a  
10 weak association with orthopedic diseases, by early improvements in lifestyle, would reduce the  
11 frequency not only of cardiovascular diseases but also of DD and lumbar degenerative diseases  
12 as well. This could lead to the establishment of preventive measures against DD and lumbar  
13 degenerative diseases.

14 Consequently, we constructed this observational study to clarify the associations between DD,  
15 evaluated by MRI, and suspected risk factors for DD including atherosclerosis and  
16 cardiovascular risk factors.

17

## 18 Materials and Methods

### 19 Participants

20 In 2002 and 2003, we publicly recruited adults over 50 years old to participate in a health  
21 promotion program by advertising in local newspapers in three municipalities (Tsukuba City  
22 with 190,000 inhabitants, Ogano Town with 12,500 and Taiyo Village with 11,000). The program  
23 included light weight- and endurance-training for about one year and physical examinations

1 before and after the exercise program. The main purposes of this project were the development  
2 of an index of older people's health and a total exercise program to maintain and improve their  
3 living function and the establishment of a health promotion system. When we recruited  
4 participants, we did not ask whether they had suffered any low back disorders.

5 Figure 1 shows the study profile. Three hundred forty-eight individuals received written and oral  
6 information about the study. Forty-five participants who did not provide written  
7 informed-consent or had any serious illness or abnormality according to screening by physicians  
8 involved in this project were excluded. Three hundred three individuals were examined with  
9 MRI. The imaging was performed on 255 participants before the exercise program and after  
10 12-18 months on 48 participants who could not be examined before the program for various  
11 reasons. We excluded 33 participants: 16 because of poor images and 17 because of uncertainty  
12 about their disc levels. The study's subjects consisted of 270 adults (96 males and 174 females)  
13 with a mean age of 68.4 years (range: 51-86 years). This study was approved by the Human  
14 Subjects Institutional Review Board of the Institute of Health and Sport Sciences, University of  
15 \* \* \* \* \* (Notification Number 29).

16  
17 Assessment of disc degeneration

18 We examined MR T2 density-weighted fast spin-echo saggital images (TR 4000 ms/TE 125 ms)  
19 of the subjects' lumbar spine using a 0.2-Tesla imager (AIRIS Mate, Hitachi Medical AG, Tokyo,  
20 Japan) with a surface coil. Slice thickness was 6.0 mm with no interslice gap, matrix 256 × 256  
21 and field of view 300 × 300 mm. The imager was a low field system. However, Merl et al.  
22 reported no statistically relevant difference in diagnostic accuracy between high-field and  
23 low-field (0.2-Tesla) MRI as measured by clinical or surgical gold standard in the spine, and a

1 high degree of concordance between high- and low-field MRI has also been confirmed [25]. In  
2 our study, a total of 1,347 intervertebral discs from L1/2 to L5/S1 in the mid-sagittal images  
3 were assessed. The degree of degeneration was classified into five grades by a modified  
4 Pfirrmann's classification (Table 1, Figure 2), with Grades IV and V considered degenerated.  
5 Pfirrmann et al. used both signal intensity and disc height categories for classification [26].  
6 However, it was difficult to assess the images of adults over 50 years old by Pfirrmann's  
7 classification because the grades from signal intensity and disc height disagreed. It has been  
8 reported that signal intensity loss is significantly correlated with the morphological level of DD  
9 and also with both the water and proteoglycan content of a disc [27]. Therefore, we used a  
10 grading based on signal intensity only. The images were assessed independently by two  
11 orthopedic surgeons who were unaware of the subject's status. Finally, a consensus decision was  
12 reached in conference and was used for the statistical analysis.

13 We assessed interrater and intrarater agreement by calculating kappa statistics (dichotomous  
14 variables). The kappa coefficient for interrater reliability was 0.81. Repeated measurements were  
15 done at 2-month intervals for 40 images (200 discs), chosen at random, to estimate intrarater  
16 reliability. The intrarater agreement coefficient was 0.89.

17  
18 MRI for evaluating DD was performed on 222 participants (Group A) before the exercise  
19 program, and after 12-18 months on 48 participants (Group B) who could not be examined  
20 before the program. In addition, Group A involved 88 participants (Group A') of whom a pair of  
21 MR images were taken – one before the program and the other 18 months later. To verify that  
22 there was no significant progression of DD due to aging of the participants over 18 months and  
23 thereby to confirm that the 48 images for group B were appropriate for analysis, we chose 40

1 pairs of images at random from Group A' and compared images within each pair. The agreement  
2 coefficient from kappa statistics for the pair of images was 0.90, which was higher than the  
3 interrater and intrarater agreement coefficients. Therefore, no progression due to aging over 18  
4 months could be detected by our grading method.

#### 5 6 Physical findings

7 Each subject's height and weight were recorded and their body mass index (BMI) was  
8 calculated.

#### 9 10 Laboratory measurements

11 Blood samples were collected and serum low-density lipoprotein cholesterol (LDLc),  
12 triglyceride (TG) and glycosylated hemoglobin (HbA<sub>1c</sub>) were measured. All blood samples were  
13 obtained in the morning following an overnight fast.

#### 14 15 An index of atherosclerosis

16 Pulse wave velocity (PWV) is known to be an indicator of arterial stiffness, and has been  
17 regarded as a marker reflecting vascular damage [28,29]. PWV is the speed at which a pulse  
18 travels between two points and is related to the square root of the elastic modulus according to  
19 the Moens-Korteweg equation [30]. Therefore, the stiffer the artery, the faster the PWV. To  
20 assess atherosclerosis we used a simple and noninvasive method of measuring brachial-ankle  
21 PWV (baPWV) that has recently been made available. The subjects were examined in a supine  
22 resting position, using an automated device with an oscillometric method (form PWV/ABI, Colin  
23 AG, Ltd, Aichi, Japan) [28,29]. The average of the left and right baPWVs in each participant was

1 used for the analysis.

2

3 A total index of bone mineral density

4 Quantitative ultrasound assessment of the calcaneus was performed using an Acoustic  
5 Osteo-Screener (AOS-100, Aloka AG, Tokyo, Japan) [31]. The speed of sound (SOS) and the  
6 transmission index (TI) were measured, and from them the osteo sono-assessment index (OSI),  
7 which is given by  $OSI = TI \times SOS^2$ , was calculated as a total index of BMD.

8

9 Questionnaire and telephone interview

10 A questionnaire containing demographic data, physical and mental health status and lifestyle  
11 factors was given to the participants to complete. The following three questions concerning  
12 experience of low back pain (LBP), current smoking habits and current drinking habits were  
13 used for the analysis: Have you experienced LBP during your lifetime? Do you smoke  
14 cigarettes? Do you drink alcoholic beverages?

15 We conducted a telephone interview about occupational and sports histories to clarify the  
16 associations between DD and physical loading. We asked the following two questions about the  
17 occupation that the participants had been engaged in for the longest time: Have you ever engaged  
18 in an occupation that involved lifting weights of more than 10 kg for more than one third of your  
19 working hours? Have you worked as an occupational driver? The question concerning sports  
20 history was as follows: Have you ever participated in a sport more than 3 times per week for  
21 more than 5 years, and, if so, what kind of sport?

22

23 Statistical analysis

1 The values measured as continuous data were divided into binary categorical variables to make  
2 the prediction rule clear. The cutoff points between low and high values were set at 70 years for  
3 age and 25 kg/m<sup>2</sup> for BMI according to the criteria of the Japan Society for the Study of Obesity  
4 [32]. The cutoff points were set at 140 mg/dl (3.6 mmol/L) for LDLc and 150 mg/dl (1.7  
5 mmol/L) for TG according to those of the Japan Atherosclerosis Society [33], 5.8% for HbA<sub>1c</sub>  
6 from the standard range (4.3-5.8) of the Japan Diabetes Society [34]; and 1400 cm/sec for  
7 baPWV according to a report suggesting baPWV as a new marker for cardiovascular risk [35].  
8 There are no generally accepted criteria of OSI, so the cutoff point was chosen to be 2.4  
9 arbitrarily to provide equal-size groups for statistical analysis. We categorized experience of LBP  
10 during lifetime, current smoking and drinking habits as “with” or “without”. We used only two  
11 categories for the smoking and drinking habits because we did not ask our subjects details of  
12 their histories of smoking and drinking in our questionnaire. The participants who answered  
13 “yes” to the respective questions about occupational and sports histories were categorized into  
14 the occupational lifting group, the occupational driving group and the sports group. Table 2  
15 shows the characteristics of the participants. The univariate relationships between DD and the  
16 variables were evaluated using chi-square tests at each disc level, and the odds ratios (ORs) were  
17 calculated. The variables that showed substantial correlation ( $P < 0.10$ ) with DD at one or more  
18 disc levels and the gender, for adjusting gender differences, were entered into logistic regression  
19 analysis at each disc level. The ORs and their 95% confidence intervals (CI) were calculated for  
20 DD, and P values less than 0.05 were considered significant. Co linearity among the independent  
21 variables was considered for inclusion in the regression model. The Pearson’s correlation  
22 coefficient between age and baPWV was 0.48, and the correlation coefficients among other  
23 variables were less than 0.21.

1 We analyzed each disc level separately because it has been suggested that the effects of  
2 mechanical factors such as occupational and sports histories could vary with disc level.  
3 Statistical analyses were performed using the computer software JMP version 5.1 (SAS Institute  
4 Inc., Cary, NC) [36].

## 5 6 Results

7 Disc degeneration (Grades IV, V) was observed in 140 discs (52%) at L1/2, 159 discs (59%) at  
8 L2/3, 159 discs (59%) at L3/4, 181 discs (67%) at L4/5 and 156 discs (58%) at L5/S1. The  
9 proportions of DD were 57.3% (274 discs) in males and 60.0% (521 discs) in females but these  
10 were not significant.

## 11 12 Univariate Results

13 Results of the univariate analysis are shown in Table 3. Aging was substantially correlated ( $P <$   
14  $0.10$ ) with DD from L1/2 to L4/5; high BMI, with L2/3, L3/4 and L5/S1; high LDLc, with L4/5  
15 and L5/S1; high baPWV, with L1/2 and L2/3; low OSI, with L5/S1; occupational lifting, with  
16 L1/2 and L4/5; and sports, with L5/S1. Gender, TG, HbA<sub>1c</sub>, history of LBP, current smoking and  
17 drinking habits and occupational driving had no correlation with DD.

## 18 19 Multivariate Results

20 Considering clinical and statistical significance, our logistic regression model was constructed  
21 using eight variables as shown in Table 4. Aging was significantly correlated with L1/2 (OR  
22 2.14; CI, 1.15-3.99), L2/3 (OR, 3.56; CI, 1.86-7.03), L3/4 (OR, 2.84; CI, 1.50-5.50) and L4/5  
23 (OR, 3.05; CI, 1.55-6.16); high BMI, with L2/3 (OR, 2.98; CI, 1.52-6.05), L3/4 (OR, 3.58; CI,

1 1.85-7.21), L4/5 (OR, 2.32; CI, 1.18-4.72) and L5/S1 (OR, 3.34; CI, 1.70-6.81); high LDLc, with  
2 L4/5 (OR, 2.65; CI, 1.33-5.52); occupational lifting, L1/2 (OR, 4.25; CI, 1.60-13.48); and sports,  
3 with L5/S1 (OR, 3.36; CI, 1.31-9.90). Gender, baPWV and OSI were not correlated with DD at  
4 any level.

5

## 6 Discussion

7 We have clarified the factors associated with MRI evaluated DD. Aging, obesity, high LDLc,  
8 occupational lifting and sports activities are associated with DD at one or more disc levels.  
9 Atherosclerosis had previously been associated with DD [2,22,23]. Cardiovascular risk factors  
10 that accelerate atherosclerosis have also been suspected of being associated with DD [3,10-15].  
11 We had the opportunity to examine baPWV by a simple and noninvasive method, and so we used  
12 baPWV as an index of atherosclerosis and investigated the association between DD and baPWV.  
13 However, we did not find a significant association. It is reported that baPWV may provide  
14 qualitatively similar information to aortic PWV; however, it also reflects muscular arterial  
15 stiffness of the leg [37]. In addition, PWV is thought to be a marker of the early stages of  
16 atherosclerosis [38]. We surmise that the blood flow to the lumbar spine was maintained even in  
17 our high baPWV group participants and that baPWV may not be a good enough indicator of the  
18 blood flow to discs.

19 Among the cardiovascular risk factors that we investigated, high BMI was significantly  
20 associated with DD at four disc levels. The association between DD and obesity has been  
21 previously reported [3,11] and Liuke et al. found that past obesity is also associated with DD  
22 more than present obesity [10]. Therefore, it seems that obesity has some influence on the  
23 process of DD. Further research to elucidate the exact mechanism is needed because both direct

1 mechanical stress on the intervertebral discs and indirect effects of atherosclerosis as a  
2 cardiovascular risk factor are suspected to be mechanisms through which obesity affects DD.  
3 Our study showed that high LDLc was also significantly associated with DD at L4/5 with a high  
4 OR of 2.65. In a cross-sectional study on patients with long-term nonspecific LBP, Kauppila et al.  
5 studied the association between LDLc and DD but no correlation was found [22]. On the other  
6 hand, Jhawar et al. reported a significant association between self-reported lumbar disc  
7 herniation and high cholesterol in the prospective Nurses' Health Study, and indicated that high  
8 cholesterol might affect DD [12]. However, this study had the limitation that identification of  
9 lumbar disc herniation was judged only from questionnaires which asked whether participants  
10 had a physician-diagnosis of herniated lumbar disc and confirmation by computed tomography  
11 or MRI—the researchers did not perform their own examinations. Although our results do not  
12 show the cause because our study is an observational study, we directly evaluated DD by MRI,  
13 and found that DD was significantly associated with high LDLc. However, in our study DD was  
14 not associated with baPWV measured as an index of atherosclerosis, and even though LDLc is a  
15 systemic factor, only one disc level was significant. Accordingly, our results also indicate the  
16 possibility that high LDLc influences DD in other ways than through atherosclerosis, and that  
17 additional factors that we did not investigate affect DD. A future prospective study should clarify  
18 the associations between DD, atherosclerosis and LDLc.

19 It has been reported that diabetic sand rats had more dehydrated discs than those of a control  
20 group [13], and diabetes increased the risk of lumbar disc herniation in the Nurses' Health Study  
21 [12]. However, high HbA<sub>1c</sub> was not significantly correlated with DD in our results. Our high  
22 HbA<sub>1c</sub> group involved a subgroup that had abnormal levels of HbA<sub>1c</sub>, but were not diagnosed  
23 with diabetes. If DD has only a weak association with hyperglycemia as previously reported [39],

1 it would not be possible for a study such as ours to clarify the association.

2 Similarly to diabetes, smoking has a reportedly weak association with DD [14], and there were a  
3 small proportion of current smokers among our subjects. Therefore, our inability to demonstrate  
4 an association between smoking and DD may be more indicative of an inadequate sample size  
5 than the lack of an actual relationship.

6 By investigating whether physical loading is associated with DD, level by level, we found that  
7 occupational lifting was significantly associated with L1/2 and sports activities were  
8 significantly associated with L5/S1. It has been reported that heavier lifetime occupational  
9 physical loading was associated with DD in the upper lumbar region [5], and weightlifting led to  
10 slightly more degeneration in the lower thoracic spine [18]. Although we did not assess the  
11 thoracic spine, we suspect that physical loading during lifting of heavy objects, regardless of  
12 occupation or sport, is concentrated throughout the thoraco-lumbar junction and causes DD.

13 Several investigators previously reported that gymnasts [17] and soccer players [19] had a higher  
14 incidence of DD and that DD in elite athletes with LBP at the Sydney Olympic Games was most  
15 common at L5/S1 [16]. Our result supports these studies. The degree and the disc level of DD  
16 might vary with the type of sports; however, we were unable to identify the type of sports  
17 activities and the disc level that were most likely to result in DD because our sports group was  
18 too small and it was necessary to group together all sports.

19 Our major study limitation is that our study is an observational study. Accordingly, the  
20 correlations between DD and several factors in our study do not indicate their causes. In addition,  
21 because of the number of independent variables compared and the methods available, it is  
22 possible that some of the significant associations identified may be artifactual. However, this  
23 work lays the foundation necessary to further elucidate the causes and mechanisms of DD and

1 we hope that more detailed and prospective studies based on our results would be able to clarify  
2 them. And, the accumulated influence of cardiovascular risk factors and physical loading on DD  
3 also has to be estimated for the establishment of preventive measures against DD and lumbar  
4 degenerative diseases. Furthermore, our results should be interpreted in view of a potential  
5 selection bias that our subjects were Japanese in a health promotion program. Our participants  
6 might therefore have been more interested in health-care and lifestyle than the average person,  
7 and people who had severe LBP and osteoporosis might have excluded themselves from our  
8 study.

9

10 Conclusions

11 In our observational study, aging, high BMI, high LDLc, occupational lifting and sports activities  
12 were independently associated with DD at one or more disc levels. The results of this study raise  
13 our index of suspicion that cardiovascular risk factors and particular physical loading may  
14 promote DD, but additional studies are required to investigate this further. We hope that our  
15 findings will be useful in establishing preventive measures against DD and lumbar degenerative  
16 diseases.

17

18 Acknowledgements

19 This research is supported by the Special Coordination Funds for Promoting Science and  
20 Technology of the Ministry of Education, Culture, Sports, Science and Technology in Japan.

21 The authors thank Dr. M. Matsuda and his colleagues for providing baPWV data, and Dr. T.  
22 Fukunaga and his colleagues for providing OSI data.

23

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

References

[1] Battie MC, Videman T, Parent E. Lumbar disc degeneration: epidemiology and genetic influences. *Spine*. 2004;29:2679-90.

[2] Kauppila LI, McAlindon T, Evans S, Wilson PW, Kiel D, Felson DT. Disc degeneration/back pain and calcification of the abdominal aorta. A 25-year follow-up study in Framingham. *Spine*. 1997;22:1642-7; discussion 1648-9.

[3] Nanjo Y, Morio Y, Nagashima H, Hagino H, Teshima R. Correlation between bone mineral density and intervertebral disk degeneration in pre- and postmenopausal women. *J Bone Miner Metab*. 2003;21:22-7.

[4] Powell MC, Wilson M, Szypryt P, Symonds EM, Worthington BS. Prevalence of lumbar disc degeneration observed by magnetic resonance in symptomless women. *Lancet*. 1986;2:1366-7.

[5] Battie MC, Videman T, Gibbons LE, Fisher LD, Manninen H, Gill K. 1995 Volvo Award in clinical sciences. Determinants of lumbar disc degeneration. A study relating lifetime exposures and magnetic resonance imaging findings in identical twins. *Spine*. 1995;20:2601-12.

[6] Sambrook PN, MacGregor AJ, Spector TD. Genetic influences on cervical and lumbar disc degeneration: a magnetic resonance imaging study in twins. *Arthritis Rheum*. 1999;42:366-72.

[7] Kawaguchi Y, Kanamori M, Ishihara H, Ohmori K, Matsui H, Kimura T. The association of lumbar disc disease with vitamin-D receptor gene polymorphism. *J Bone Joint Surg Am*. 2002;84-A:2022-8.

- 1 [8] Seki S, Kawaguchi Y, Chiba K, Mikami Y, Kizawa H, Oya T, Mio F, Mori M, Miyamoto  
2 Y, Masuda I, Tsunoda T, Kamata M, Kubo T, Toyama Y, Kimura T, Nakamura Y, Ikegawa  
3 S. A functional SNP in CILP, encoding cartilage intermediate layer protein, is associated  
4 with susceptibility to lumbar disc disease. *Nat Genet.* 2005;37:607-12.
- 5 [9] Solovieva S, Lohiniva J, Leino-Arjas P, Raininko R, Luoma K, Ala-Kokko L, Riihimaki  
6 H. COL9A3 gene polymorphism and obesity in intervertebral disc degeneration of the  
7 lumbar spine: evidence of gene-environment interaction. *Spine.* 2002;27:2691-6.
- 8 [10] Liuke M, Solovieva S, Lamminen A, Luoma K, Leino-Arjas P, Luukkonen R, Riihimaki  
9 H. Disc degeneration of the lumbar spine in relation to overweight. *Int J Obes (Lond).*  
10 2005;29:903-8.
- 11 [11] Yoshimura N, Dennison E, Wilman C, Hashimoto T, Cooper C. Epidemiology of chronic  
12 disc degeneration and osteoarthritis of the lumbar spine in Britain and Japan: a  
13 comparative study. *J Rheumatol.* 2000;27:429-33.
- 14 [12] Jhawar BS, Fuchs CS, Colditz GA, Stampfer MJ. Cardiovascular risk factors for  
15 physician-diagnosed lumbar disc herniation. *Spine J.* 2006;6:684-91.
- 16 [13] Ziv I, Moskowitz RW, Krause I, Adler JH, Maroudas A. Physicochemical properties of the  
17 aging and diabetic sand rat intervertebral disc. *J Orthop Res.* 1992;10:205-10.
- 18 [14] Battie MC, Videman T, Gill K, Moneta GB, Nyman R, Kaprio J, Koskenvuo M. 1991  
19 Volvo Award in clinical sciences. Smoking and lumbar intervertebral disc degeneration:  
20 an MRI study of identical twins. *Spine.* 1991;16:1015-21.
- 21 [15] Oda H, Matsuzaki H, Tokuhashi Y, Wakabayashi K, Uematsu Y, Iwahashi M.  
22 Degeneration of intervertebral discs due to smoking: experimental assessment in a  
23 rat-smoking model. *J Orthop Sci.* 2004;9:135-41.

- 1 [16] Ong A, Anderson J, Roche J. A pilot study of the prevalence of lumbar disc degeneration  
2 in elite athletes with lower back pain at the Sydney 2000 Olympic Games. *Br J Sports*  
3 *Med.* 2003;37:263-6.
- 4 [17] Sward L, Hellstrom M, Jacobsson B, Nyman R, Peterson L. Disc degeneration and  
5 associated abnormalities of the spine in elite gymnasts. A magnetic resonance imaging  
6 study. *Spine.* 1991;16:437-43.
- 7 [18] Videman T, Battie MC, Gibbons LE, Manninen H, Gill K, Fisher LD, Koskenvuo M.  
8 Lifetime exercise and disk degeneration: an MRI study of monozygotic twins. *Med Sci*  
9 *Sports Exerc.* 1997;29:1350-6.
- 10 [19] Videman T, Sarna S, Battie MC, Koskinen S, Gill K, Paananen H, Gibbons L. The  
11 long-term effects of physical loading and exercise lifestyles on back-related symptoms,  
12 disability, and spinal pathology among men. *Spine.* 1995;20:699-709.
- 13 [20] Harada A, Okuizumi H, Miyagi N, Genda E. Correlation between bone mineral density  
14 and intervertebral disc degeneration. *Spine.* 1998;23:857-61; discussion 862.
- 15 [21] Miyakoshi N, Itoi E, Murai H, Wakabayashi I, Ito H, Minato T. Inverse relation between  
16 osteoporosis and spondylosis in postmenopausal women as evaluated by bone mineral  
17 density and semiquantitative scoring of spinal degeneration. *Spine.* 2003;28:492-5.
- 18 [22] Kauppila LI, Mikkonen R, Mankinen P, Peltto-Vasenius K, Maenpaa I. MR aortography  
19 and serum cholesterol levels in patients with long-term nonspecific lower back pain.  
20 *Spine.* 2004;29:2147-52.
- 21 [23] Kauppila LI, Penttila A, Karhunen PJ, Lalu K, Hannikainen P. Lumbar disc degeneration  
22 and atherosclerosis of the abdominal aorta. *Spine.* 1994;19:923-9.
- 23 [24] Kurunlahti M, Kerttula L, Jauhiainen J, Karppinen J, Tervonen O. Correlation of

- 1 diffusion in lumbar intervertebral disks with occlusion of lumbar arteries: a study in adult  
2 volunteers. *Radiology*. 2001;221:779-86.
- 3 [25] Merl T, Scholz M, Gerhardt P, Langer M, Laubenberger J, Weiss HD, Gehl HB, Wolf KJ,  
4 Ohnesorge I. Results of a prospective multicenter study for evaluation of the diagnostic  
5 quality of an open whole-body low-field MRI unit. A comparison with high-field MRI  
6 measured by the applicable gold standard. *Eur J Radiol*. 1999;30:43-53.
- 7 [26] Pfirrmann CW, Metzdorf A, Zanetti M, Hodler J, Boos N. Magnetic resonance  
8 classification of lumbar intervertebral disc degeneration. *Spine*. 2001;26:1873-8.
- 9 [27] Benneker LM, Heini PF, Anderson SE, Alini M, Ito K. Correlation of radiographic and  
10 MRI parameters to morphological and biochemical assessment of intervertebral disc  
11 degeneration. *Eur Spine J*. 2005;14:27-35.
- 12 [28] Tomiyama H, Yamashina A, Arai T, Hirose K, Koji Y, Chikamori T, Hori S, Yamamoto Y,  
13 Doba N, Hinohara S. Influences of age and gender on results of noninvasive  
14 brachial-ankle pulse wave velocity measurement--a survey of 12517 subjects.  
15 *Atherosclerosis*. 2003;166:303-9.
- 16 [29] Yamashina A, Tomiyama H, Takeda K, Tsuda H, Arai T, Hirose K, Koji Y, Hori S,  
17 Yamamoto Y. Validity, reproducibility, and clinical significance of noninvasive  
18 brachial-ankle pulse wave velocity measurement. *Hypertens Res*. 2002;25:359-64.
- 19 [30] O'Rourke MF. Wave travel and reflection in the arterial system. *J Hypertens Suppl*.  
20 1999;17:S45-7.
- 21 [31] Naka H. Aos-100. *Nippon Rinsho*. 2004;62 Suppl 2:299-304.
- 22 [32] The. New criteria for obesity and 'obesity disease'. *J Jpn Soc Obes*. 2000;6:18-28.
- 23 [33] Japan ASi. Japan Atherosclerosis Society (JAS) guidelines for diagnosis and treatment of

- 1 atherosclerotic cardiovascular disease. Tokyo: JAS; 2002.
- 2 [34] Shima K, Kuwajima M. Standardization for glycohemoglobin (HbA1c) measurements  
3 and its evaluation. *Nippon Rinsho*. 1998;56 Suppl 3:59-67.
- 4 [35] Yamashina A, Tomiyama H, Arai T, Hirose K, Koji Y, Hirayama Y, Yamamoto Y, Hori S.  
5 Brachial-ankle pulse wave velocity as a marker of atherosclerotic vascular damage and  
6 cardiovascular risk. *Hypertens Res*. 2003;26:615-22.
- 7 [36] Sall J, Creighton L, Lehman A. *JMP Start Statistics Third Edition*. Third Edition. ed. Cary,  
8 NC: SAS Institute Inc; 2004.
- 9 [37] Sugawara J, Hayashi K, Yokoi T, Cortez-Cooper MY, DeVan AE, Anton MA, Tanaka H.  
10 Brachial-ankle pulse wave velocity: an index of central arterial stiffness? *J Hum*  
11 *Hypertens*. 2005;19:401-6.
- 12 [38] Wilkinson IB, Cockcroft JR, Webb DJ. Pulse wave analysis and arterial stiffness. *J*  
13 *Cardiovasc Pharmacol*. 1998;32 Suppl 3:S33-7.
- 14 [39] Videman T, Battie MC, Gibbons LE, Kaprio J, Koskenvuo M, Kannus P, Raininko R,  
15 Manninen H. Disc degeneration and bone density in monozygotic twins discordant for  
16 insulin-dependent diabetes mellitus. *J Orthop Res*. 2000;18:768-72.

17  
18  
19  
20  
21