

Tetsu Shimizu, Naoki Takaya and Akira Nakamura

From the Faculty of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan.

*Running title: *An L-glucose catabolic pathway*

To whom correspondence should be addressed: Akira Nakamura, Faculty of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan. Tel; +81-29-853-6637, E-mail; nakamura.akira.fm@u.tsukuba.ac.jp

Keywords: L-glucose; homochirality; *Paracoccus*

Background: L-glucose, the enantiomer of D-glucose, was believed not to be utilized by any organisms.

Results: An L-glucose-utilizing bacterium was isolated and its L-glucose catabolic pathway was identified genetically and enzymatically.

Conclusion: L-glucose was utilized via a novel pathway to pyruvate and D-glyceraldehyde-3-phosphate

Significance: This might lead to an understanding of homochirality in sugar metabolism.

SUMMARY

An L-glucose-utilizing bacterium, *Paracoccus* sp. 43P, was isolated from soil by enrichment cultivation in a minimal medium containing L-glucose as the sole carbon source. In cell free extracts from this bacterium, NAD⁺-dependent L-glucose dehydrogenase was detected as having sole activity toward L-glucose. This enzyme, LgdA, was purified and the *lgdA* gene was found to be located in a cluster of putative inositol catabolic genes. LgdA showed similar dehydrogenase activity toward *scyllo*- and *myo*-inositols. L-gluconate dehydrogenase activity was also detected in cell free extracts, which represents the reaction product of LgdA activity toward L-glucose. Enzyme purification and gene cloning revealed that the corresponding gene resides in a nine gene cluster, the *lgn* cluster, which may participate in aldonate incorporation and assimilation. Kinetic and reaction product analysis of each

gene product in the cluster indicated that they sequentially metabolize L-gluconate to glycolytic intermediates, D-glyceraldehyde-3-phosphate, and pyruvate through reactions of C-5 epimerization by dehydrogenase/reductase, dehydration, phosphorylation, and aldolase reaction, using a pathway similar to L-galactonate catabolism in *Escherichia coli*. Gene disruption studies indicated that the identified genes are responsible for L-glucose catabolism.

All living organisms show homochirality toward organic compounds, such as amino acids, with L- rather than D- amino acids utilized for protein synthesis. The same is true for sugars, where D-sugars are more abundant than L-sugars in natural environments. Accordingly, most D-sugars can be utilized by many organisms, but L-sugar usage is less common, indicating that homochirality also occurs in sugar metabolism. A good example of this is glucose, where D-glucose is the most abundant sugar in nature, and most organisms use D-glucose as an energy source via well-known pathways such as the Embden-Meyerhof-Parnas, the pentose phosphate, and the Entner-Doudoroff pathways. Meanwhile, L-glucose is not present in nature, and few instances of L-glucose usage have appeared since Rudney reported that mammalian cells, yeasts, and some bacteria could not utilize this epimer (1) with several reports also confirming that L-glucose was not utilized by other organisms (2, 3). As for enzymes with

specificity toward L-glucose, Sasajima and Sinskey reported L-glucose dehydrogenase activity in *Pseudomonas caryophylli* (4), but no further reports concerning this finding are currently available.

To determine whether there are organism(s) capable of utilizing L-glucose and to obtain further insights into homochirality in sugar metabolism, isolation of L-glucose-utilizing organisms and analysis of their catabolic pathways is necessary. Here we describe the isolation of an L-glucose-utilizing bacterium and its catabolic pathway for L-glucose metabolism is characterized.

EXPERIMENTAL PROCEDURES

Bacterial Strains, Plasmids and Media-- *Escherichia coli* DH10B was used as a host for plasmid construction. pGEM-T Easy (Promega) and pUC19 (Takara Bio) were used for gene cloning, and BL21(DE3) and plasmids pET21a(+) and pET28a(+) (Novagen) for enzyme production. pBBR1MCS2 (5) was used for constructing suicide plasmids. *E. coli* S17-1 λ pir acted as a donor for biparental mating of cells. LB medium supplemented with appropriate antibiotics was used for *E. coli* cultivation.

For screening of L-glucose-utilizing microorganisms, L-glucose minimal medium (L-Glc MM¹; 10 mM NH₄Cl, 10 mM potassium phosphate (pH 7.0), 10 mM MgSO₄, 10 mM KCl, 0.2 % (v/v) Hutner's trace element solution [per liter, 2.2 g ZnSO₄·7H₂O, 1.1 g H₃BO₃, 0.5 g MnCl₂·4H₂O, 0.5 g FeSO₄·7H₂O, 0.16 g CoCl₂·6H₂O, 0.16 g CuSO₄·5H₂O, and 0.11 g (NH₄)₆Mo₇O₂₄·4H₂O] containing 14 mM L-glucose; L-glucose was separately sterilized by filtration and added to the medium) was used, and for L-glucose dehydrogenase (L-GDH) purification, D-galactose minimal medium (20 mM NH₄Cl, 20 mM potassium phosphate (pH 7.0), 10 mM MgSO₄, 10 mM KCl, 0.2 % (v/v) Hutner's trace element solution, 0.1 % yeast extract containing 28 mM D-galactose; D-galactose was sterilized separately by filtration) was used.

PCR Amplification and Sequencing-- PCR primers used in this study are listed in supplemental Table 1, and amplification was

conducted under standard conditions. When appropriate, fragments were cloned into pUC19, pET21a or pET28a vectors after digestion with restriction enzymes whose sites corresponded to the primers. Nucleotide sequencing was conducted with a CEQ8000XL sequencer (Beckman Coulter).

Enzyme Assay Substrates-- L-glucose and inositol derivatives were purchased from Sigma-Aldrich and Hokko Chemical Industry, respectively. L-gluconate and D-galactonate were prepared as potassium salts by hypiodite-in-methanol oxidization (6) of L-glucose and D-galactose, respectively. D-idonic acid was obtained by base hydrolysis of D-idono-1,4-lactone from Carbosynth. D-2-keto-3-deoxygalactonate (KDGal) was enzymatically synthesized from D-galactonate by the recombinant D-galactonate dehydratase (DgoD) from *Escherichia coli* DH10B (7). DgoD was produced and purified as described below. D-2-keto-3-deoxy-6-phosphogalactonate (KDPGal) was enzymatically synthesized from KDGal using recombinant LgnF cloned in this study.

L-Glucose-Utilizing Microorganism Screening-- Soil samples were inoculated in 10 ml L-Glc MM and cultivated at 30 °C for 48 h with shaking at 300 rpm. Enrichment cultivation was conducted by inoculating one ml of the cultures to fresh L-Glc MM and cultivating under the same conditions, which was repeated ten times. The cultures were then serially diluted and plated onto L-Glc MM agar plates to isolate L-glucose-utilizing microorganisms. L-glucose consumption was monitored using the 3,5-dinitrosalicylic acid (DNS) method to determine the amount of reducing sugars in the culture supernatants (8).

Detection of Enzyme Activities toward L-Glucose in Strain 43P Cell-Free Extracts-- Strain 43P was cultured in L-Glc MM at 30 °C for 20 h and the cells were collected by centrifugation at 17,400 x g for 10 min. The cells were then resuspended in 100 mM Tris-HCl, pH 7.0 containing 10 % (v/v) glycerol and 1 mM DTT (Buffer A) and disrupted by sonication with a SONIFIER 250 (Branson). After centrifugation at 17,400 x g for 10 min., the cell-free extracts were used to detect enzyme

activities toward L-glucose. For isomerase activity, extracts were incubated with 10 mM L-glucose in 100 mM Tris-HCl, pH 7.5 for 2 h at 30 °C, and the reaction was terminated by addition of trichloroacetic acid. Formation of ketose was analyzed by TLC according to the method described by Lobanok *et al.* (9). For kinase activity, the reaction was performed as described above with the buffer containing 1 mM MgCl₂ and 10 mM ATP, and the reaction mixture was analyzed by TLC with silica gel plates (Merck, Darmstadt) as described by Hemker *et al.* (10). Dehydrogenase activity was assayed as described below.

Purification of Native L-glucose dehydrogenase (L-GDH) and L-Gluconate Dehydrogenase (L-GnDH)-- Strain 43P was pre-cultured in 200 ml LB medium at 30 °C for 20 h. After collecting the cells by centrifugation at 5,800 x g for 15 min, they were transferred to 1 liters of D-galactose minimal medium, and cultivation was continued for 24 h. The cells were then harvested by centrifugation, washed, and resuspended with Buffer A. Cell-free extracts were prepared by sonication and centrifugation at 27,000 x g for 30 min at 4 °C, and applied to a DEAE-cellulose column (DE52, Whatman International; ϕ 2.5 cm x d 15 cm) equilibrated with Buffer A. After washing the column with Buffer A, proteins were eluted with a linear gradient of 0-500 mM NaCl in Buffer A. The active fractions were eluted at 80-120 mM NaCl. To the fractions containing L-GDH activity, (NH₄)₂SO₄ was added to the solution to 20 % saturation. This solution was then applied to a Butyl-Toyopearl 650M column (Tosoh Corporation; ϕ 2.5 cm x d 15 cm) equilibrated with Buffer A containing 1.5 M (NH₄)₂SO₄ and washed with the same buffer. Proteins were eluted with a linear gradient of 1.5-0 M (NH₄)₂SO₄ in Buffer A and the active fractions eluted at 0.6-0.5 M (NH₄)₂SO₄ were dialyzed against Buffer A. The solution was loaded onto a Mono Q 5/50 GL column (GE Healthcare) equilibrated with Buffer A and washed with the same buffer. Proteins were eluted with a linear gradient of 0-250 mM NaCl in Buffer A. The active fractions eluted at 150 mM NaCl were concentrated by osmotic ultrafiltration (11) against dialysis membrane, size 20 (Wako

chemical) with PEG 20,000 at 4 °C for 3 h, and applied to a Superdex 200 10/300 GL column (GE Healthcare) pre-equilibrated with Buffer A containing 150 mM NaCl and eluted at a flow rate of 0.5 ml min⁻¹. Fractions containing L-GDH activity were dialyzed against 100 mM Tris-HCl, pH 8.0, containing 50 % (v/v) glycerol and 1 mM DTT.

For purification of L-GnDH, strain 43P was cultivated similar to that described above, except the second cultivation was conducted with 200 ml L-Glc MM containing 0.1 % yeast extract for 6 h. Cell-free extracts were prepared with Buffer B (100 mM Tris-HCl, pH 8.5, containing 20 % (v/v) glycerol and 2.5 mM DTT), as described above, and applied to a DEAE-cellulose column (DE52; ϕ 1 cm x d 15 cm) pre-equilibrated with Buffer B. After washing the column with Buffer B containing 50 mM NaCl, proteins were eluted with a linear gradient of 50-500 mM NaCl in Buffer B. The fractions containing L-GnDH activity eluted at 170-200 mM NaCl were dialyzed against Buffer B, and applied again to the same column equilibrated with Buffer B. After washing the column with Buffer B containing 120 mM NaCl, proteins were eluted with a linear gradient of 120-250 mM NaCl in Buffer B. The active fractions were directly loaded onto a Mono Q 5/50 GL column equilibrated with Buffer B containing 200 mM NaCl and washed with the same buffer. Proteins were eluted with a linear gradient of 200-350 mM NaCl in Buffer B. The active fractions were eluted at 280 mM NaCl, which were concentrated by Amicon Ultra-0.5 mL 10 K (Milipore) at 12000 x g for 7 min at 4 °C. The concentrated proteins were applied to a Superdex 200 10/300 GL column pre-equilibrated with Buffer B containing 150 mM NaCl, and eluted at a flow rate of 0.5 ml min⁻¹. Fractions containing L-GnDH activity were dialyzed against 100 mM Tris-HCl, pH 8.5, containing 50 % (v/v) glycerol and 1 mM DTT.

Purification steps were monitored by SDS-PAGE as described by Laemmli (12).

Determination of N-terminal Amino Acid Sequences-- Purified L-GDH and L-GnDH were resolved by 10% SDS-PAGE and electroblotted onto a polyvinylidene difluoride membrane. The bands corresponding to the enzymes were

excised and used for Edman degradation and sequencing that was carried out by Hokkaido System Science.

Sequencing Analyses of Genes Encoding L-GDH and L-GnDH and their Regions-- Based on the N-terminal amino acid sequence of L-GDH and an internal conserved amino acid sequence derived from Pden_1680, two degenerate primers, L-GDH_Nter and L-GDH_midR, were synthesized and used for PCR amplification from strain 43P genomic DNA. The amplified 500 bp fragment was cloned into the pGEM-T Easy vector and sequenced. Based on the obtained sequence, two primers, 43P-LGDH-inverseF and 43P-L-GDH-inverseR, were designed and used for inverse-PCR (13) with a mixture of *EcoRI*-digested and self-ligated strain 43P genomic DNA. The amplified 3.5 kb fragment was cloned into the pGEM-T Easy vector, and sequenced.

Cloning and sequencing of the L-GnDH gene was conducted as described above except for the primers used: L-GnDH_Nter and L-GnDH_midR for the first PCR, and 43P-LGnDH-inverseF and 43P-LGnDH-inverseR for the inverse-PCR. For inverse-PCR, *PstI*-digested and self-ligated genomic DNA was used as a template.

To obtain sequence information for the regions around the L-GDH and L-GnDH genes, inverse-PCR and primer walking were conducted using the primers listed in supplemental Table 1. The nucleotide sequences of the resulting fragments were then determined.

Recombinant Enzyme Production and Purification-- The ORFs of *lgdA*, *lgnE*, *lgnF*, *lgnG*, *lgnH* and *lgnI* of strain 43P, and *E. coli dgoD* were PCR-amplified with the appropriate primers from genomic DNA of strain 43P and *E. coli* DH10B, respectively (supplemental Table 1). After digestion with restriction enzymes whose sites were located in the primers, *lgdA* was cloned into pET21a(+) and the remaining genes were cloned in pET28a(+). The His₆-tags in the vectors were attached at the LgdA C-terminus and at the N-termini for all other genes.

E. coli BL21(DE3) harboring these plasmids was cultured in LB medium containing

appropriate antibiotics at 37 °C until the optical densities of the cultures reached 0.5. Isopropyl-β-D-thiogalactopyranoside (IPTG, final concentration 0.1 mM) was then added and culturing was continued for 3 h. For *lgnE*, culturing was carried out at 28 °C and the cultivation time after IPTG addition was prolonged to 12 h. Cells were harvested by centrifugation and disrupted by sonication in 50 mM Tris-HCl, pH 8.0, 10% (v/v) glycerol, 0.5 M NaCl, and 10 mM imidazole. The resulting cell-free extracts were loaded onto a HisTrap HP column (GE Healthcare) and His₆-tagged proteins were eluted with 0.5 M imidazole. The purified proteins were dialyzed against 100 mM Tris-HCl, pH 8.0 containing 50 % (v/v) glycerol and 1 mM DTT.

Enzyme Assays-- L-GDH, L-GnDH, and L-5-keto-gluconate reductase activities were assayed using a DU 800 spectrophotometer (Beckman Coulter) to measure reduction/oxidation of NAD⁺ and NADPH at 340 nm and 25 °C. The reactions were conducted in 28 mM L-glucose and 1 mM NAD⁺ in 100 mM Tris-HCl, pH 8.0, 2 mM potassium L-gluconate and 1 mM NAD⁺ in 100 mM Tris-HCl, pH 9.0 and 0.5 mM L-5-keto-gluconate and 0.2 mM NADPH in 100 mM HEPES-KOH, pH 7.0, respectively.

D-idonate dehydratase activity was assayed by measuring the reducing activity of the product using DNS assay as described by Kuorelahti *et al.* (14). The enzyme was mixed with 10 mM D-idonate, 0.5 mM FeCl₂, 20 mM DTT, in 100 mM Tris-HCl, pH 7.5, in a total volume of 100 μl, and the mixture was incubated at 30 °C for 15 min. The reaction was terminated by addition of 200 μl DNS reagent and heated for 5 min in boiling water. After dilution with 900 μl water, absorbance at 535 nm was measured. The molar absorption coefficient ($\epsilon = 91 \text{ M}^{-1} \text{ cm}^{-1}$) was obtained from a standard curve with KDGal that was produced by recombinant DgoD.

KDGal kinase activity was assayed at 25 °C by measuring the amount of ADP produced using a coupled assay with pyruvate kinase (PK) and lactate dehydrogenase (LDH). The enzyme was mixed with 2 mM KDGal, 1 mM ATP, 10 mM MgCl₂, 0.2 mM NADH, 1 mM

phosphoenolpyruvate containing 1.5 U LDH and 2 U PK (Sigma-Aldrich), in 100 mM HEPES-KOH, pH 7.5. The ADP produced by the kinase was used for production of pyruvate by PK, which was converted to lactate by LDH with concomitant oxidation of NADH.

KDPGal aldolase activity was assayed by measuring the reaction products pyruvate and D-glyceraldehyde-3-phosphate (Gap) by LDH and GapDH reactions, respectively, based on the method described by Wong *et al* (15). The enzyme was mixed with 1 mM KDPGal in 100 mM potassium phosphate, pH 7.5, containing 0.2 mM NADH and 1.5 U LDH or 2 mM NAD⁺ and 1.5 U GapDH (Sigma-Aldrich). The mixture was incubated at 25 °C and the pyruvate or Gap produced was quantified by concomitant oxidation/reduction of NADH or NAD⁺, respectively.

One unit of each enzyme activity was defined as the amount of the enzyme which reduce/oxidize 1 μmol of NAD(P)⁺/NADPH for LgdA, LgnH, LgnI, LgnF and LgnG, and that which produce 1 μmol of KDGal for LgnE per min, under the assay condition.

Kinetic parameters K_m and k_{cat} were calculated using the fitting tool of Origin ver. 6.0 (OriginLab) with the basic Michaelis-Menten equation. Protein concentrations were determined by the method of Lowry *et al.* (16) with bovine serum albumin as a standard.

TLC and HPLC Analyses of Enzyme Reaction Products-- Enzyme reaction products were routinely analyzed by TLC with silica gel plates (Merck) using ethyl acetate-methanol-acetate-water (6:1.5:1.5:1) as the mobile phase. Spots were detected by diphenylamine-aniline-phosphoric acid reagent (17). HPLC analysis was performed with a LC-2000 plus system (JASCO). Samples were applied to an IC Sep COREGEL 87H column (7.8 x 300 mm, Tokyo Chemical Industry) and eluted with 4 mM H₂SO₄ at a flow rate of 0.6 ml/min and 30 °C. Detection was conducted using a refractive index detector (RI-2031 plus, JASCO) and a chiral detector (OR-2090 plus, JASCO).

Purification of Reaction Products-- The LgdA reaction product was produced in a 10 ml reaction mixture containing 2 U of LgdA, 100

mM Tris-HCl, pH 8.0, 50 mM L-glucose, 2 mM NAD⁺, with 100 mM pyruvate and 2 U D-lactate dehydrogenase (Oriental Yeast) to regenerate NAD⁺. The reaction mixture was incubated at 30 °C for 8 h and treated with activated charcoal to remove NAD⁺ and NADH. The supernatant was then applied to a Dowex 1X8 column (formate form, 200-400 mesh, Wako Pure Chemical). After washing with deionized water, the product was eluted with 100 mM formate. The eluted sample was concentrated by lyophilization and further purified by preparative TLC with a preparative silica gel plate (Merck) as above.

The LgnH reaction product was produced in a 10 ml reaction mixture containing 1 U LgnH, 100 mM Tris-HCl, pH 8.0, 20 mM potassium L-gluconate, 2 mM NAD⁺, and 40 mM *scyllo*-inosose (Hokko Chemical Industry) and 2 U *myo*-inositol dehydrogenase (Sigma-Aldrich) for regeneration of NAD⁺. After 8 h incubation at 30 °C, the reaction mixture was treated with activated charcoal and applied to a DOWEX 1X8 column (Cl⁻ form). The column was washed with water and the reaction product was eluted with 200 mM HCl. Then a 10 % (w/v) slurry of calcium carbonate was added to the eluted fraction until the solution pH was neutral. After excess amounts of calcium carbonate were removed, the solution was lyophilized and the resulting pellet was washed twice with 70% ethanol to remove CaCl₂. The precipitate was resuspended in water and an appropriate amount of Dowex 50W (H⁺ form, 200-400 mesh) was added to the solution and incubated for 30 min with shaking at room temperature so that it was dissolved in water as a free acid. Finally, the resin was removed and the sample was lyophilized.

For production of LgnI reaction products, a reaction mixture (20 ml) containing 1 U LgnI, 100 mM potassium acetate, pH 5.0, 20 mM LgnH reaction product prepared as above, and 30 mM NADPH was incubated at 30 °C for 2 h with periodic pH adjustment to <6.0 by acetic acid addition. The reaction mixture was then treated with activated charcoal and loaded onto a Dowex 50W column. The column was washed with deionized water, and the flow-through fraction was concentrated with a rotary

evaporator to remove the acetic acid. The dried sample was dissolved in water and loaded onto a Dowex 1x8 column (formate form, 200-400 mesh). After the column was washed with deionized water, the product was eluted with 100 mM formate and lyophilized.

Likewise, the LgnE reaction product was produced in a 10 ml reaction mixture containing 2 U LgnE, 100 mM Tris-acetate, pH 7.5, 50 mM D-iodonate, 20 mM DTT and 0.5 mM FeCl₂. After 5 h incubation at 30 °C, the reaction mixture was filtered through a 0.22 µm filter to remove iron precipitates and passed over a AG 501-X8 column (Bio-rad, 20-50 mesh) to remove proteins and negatively-charged iron complexes. The flow-through fraction was loaded onto a Dowex 1X8 column (acetate form, 200-400 mesh) and the column was washed with 80 mM acetate. The product was eluted with 200 mM potassium acetate, pH 7.0, and concentrated to ca. 300 µl by lyophilization. One ml ethanol was added to the solution and the precipitates were washed twice with ethanol to remove potassium acetate (18).

NMR Analyses and Assignment of Reaction Products-- NMR analyses were carried out at 25 °C with an Avance-500 apparatus (500 MHz, Bruker) at the University of Tsukuba Research Facility Center of Science and Technology with the purified reaction products dissolved in deuterium oxide. For samples of free acids and aldonolactones, the pH was adjusted to 8.0-9.0 by sodium deuteroxide. All chemical shifts were referenced to a 4,4-dimethyl-4-silapentane-1-sulfonic acid internal standard (0 ppm). LgdA and LgnE reaction product spectra were assigned by comparison with those available on the SDBS web of the National Institute of Advanced Industrial Science and Technology, Japan (<http://riodb01.ibase.aist.go.jp/sdbs/>, Feb 15, 2012.) and that reported by Kuorelahti *et al.* (14), respectively. Spectra for other compounds were assigned by HH-COSY and Hetero-nuclear Multiple Quantum Coherence (HMQC) analyses.

Construction of Gene Disruption Mutants of strain 43P-- Gene disruption was carried out by inserting a kanamycin resistant gene (*Km^r*) into the target genes by homologous recombination.

For construction of a mobile suicide vector, the *mob* region of pBBR1MCS2 was amplified with a primer set of Mob_F and Mob_R and cloned into the pGEM-T Easy vector to generate pGEM-mob, which was subsequently digested with *SphI* and *SalI*. The resultant fragment containing *mob* was cloned into pUC19 to generate pUC19-mob. Homologous regions for target genes were amplified by PCR with corresponding primer sets as follows: *lgdA*, LgdA1700_EcoF and LgdA1700_BamR; *lgnE*, LgnE2000_HinF and LgnE2000_XbaR; *lgnH*, LgnH1500_EcoF and LgnH1500_BamR; and *lgnI*, LgnI1500_XbaF and LgnI1500_SacIR. Each amplified fragment was digested with appropriate restriction enzymes and cloned into pUC19 to generate pUC19-*lgdA*, -*lgnE*, -*lgnH* and -*lgnI*, respectively. The *Km^r* cassette amplified by PCR from pBBR1MCS2 with primers PkanF and KanR was introduced into the *EcoRV* and the *SmaI* sites of pUC19-*lgnE* and -*lgnI*, respectively. Alternatively, primers Pkan_SacIIF and Kan_SacIIR were used for amplification of the *Km^r* cassette, which was digested with *SacII* and cloned into the *SacII* sites of pUC19-*lgdA* and pUC19-*lgnH*. Finally, the *mob* cassette from pUC19-mob was introduced into the *EcoRI* site of pUC19-*lgdA*::kan, -*lgnE*::kan, *lgnH*::kan and the *HindIII-XbaI* site of pUC19-*lgnI*::kan. Introduction of the suicide plasmids into strain 43P cells was achieved by biparental mating with *E. coli* S17-1 *λpir* as a donor (19). Mating was carried out at a ratio of 1:10 (donor:recipient) for 8 h at 30 °C on a cellulose membrane filter placed on an LB agar plate without antibiotics. Transconjugants were selected by LB medium containing 100 µg/ml kanamycin and 15 µg/ml nalidixic acid, and proper integration was confirmed by PCR amplification with primers listed in Table S1.

Phylogenetic Analyses-- The amino acid sequences of the L-glucose-catabolic genes and related sequences from the KEGG genome database (<http://www.genome.jp/kegg/>; April 23, 2012), together with those of bacterial genes in the same protein families having known functions (Gfo/Idh/MocA, Zn-containing ADH, Short-chain dehydrogenase, UxaA, and Aldolase_KDPG_KHG_2 for LgdA, LgnH,

LgnI, LgnE and LgnG, respectively) that were selected from the "reviewed" entries from the UniProt KB database (<http://www.uniprot.org/>), were aligned with clustal X2 software (20) and unrooted phylogenetic trees were visualized using TreeView software (21).

RESULTS

Isolation of L-Glucose-Utilizing Bacteria-- Several L-glucose-utilizing bacteria were isolated from soil samples by enrichment cultivation in L-Glc MM, on which strain 43P showed a maximum growth rate with concomitant decrease of reducing sugars in the medium (Fig. 1). Phylogenetic analysis based on 16S rRNA gene sequences showed that strain 43P is included in the genus *Paracoccus* cluster, with the closest-related species being *P. denitrificans* (96.8%, supplemental Fig. 1), although it should be noted that the *P. denitrificans* strain NBRC 102528^T did not grow on L-Glc MM (Fig. 1). These results indicate that strain 43P assimilates L-glucose, and this strain should be classified in the genus *Paracoccus*. Therefore, we sought to further analyze L-glucose catabolism in strain 43P.

L-GDH, the First Enzyme in the L-Glucose Catabolic Pathway-- To determine the L-glucose catabolic pathway in strain 43P, we used cell-free extracts prepared from cells grown in L-Glc MM as an enzyme source and L-glucose as a substrate to analyze enzyme activities known to occur in the first reactions in sugar catabolism, such as kinase, isomerase, and dehydrogenase activity. No activities beyond NAD⁺-dependent L-GDH activity were detected, suggesting that the first reaction in this pathway is L-glucose oxidation. Since this activity was also detected when the strain was cultivated with D-galactose as the sole carbon source, we purified L-GDH from cells grown in a minimal medium containing D-galactose (supplemental Table 2). The purified enzyme showed a single band of ca. 40 kDa on SDS-PAGE (supplemental Fig. 2A), and its N-terminal amino acid sequence MSNAEKALGVALIGTGFMGK exhibited 80% identity with that of *P. denitrificans* PD1222 Pden_1680, a putative oxidoreductase that is a member of the Gfo/Idh/MocA family. Therefore,

we assumed that the gene encoding L-GDH in strain 43P is an ortholog of Pden_1680, and then identified this gene by PCR and DNA sequencing as described in the EXPERIMENTAL PROCEDURES.

As expected, this gene, termed *lgdA*, showed 84% amino acid sequence identity with Pden_1680 and was located in a cluster of putative inositol-catabolic genes (Fig. 2A), the organization of which was quite similar to that of strain PD1222 with 76-89% identities, except that Pden_1673 encodes a hypothetical protein of 61 amino acids that was not present in strain 43P.

Recombinant LgdA was produced with an *Escherichia coli* pET system and purified (supplemental Fig. 2C). Purified LgdA showed NAD⁺-dependent L-GDH activity (Table 1), which is comparable to the native enzyme ($k_{cat} = 710 \pm 24 \text{ min}^{-1}$, $K_m = 44.4 \pm 6.3 \text{ mM}$) and also had dehydrogenase activity toward *myo*-inositol, *scyllo*-inositol and D-glucose (Table 1), with the highest activity observed for *scyllo*-inositol. The reaction products of LgdA from an L-glucose source were analyzed by NMR and HPLC, which showed ¹H and ¹³C chemical shifts that were identical to potassium D-gluconate (supplemental Fig. 3A) and optical rotation that was identical to potassium L-gluconate (Fig. 3A). These data indicate that LgdA catalyzes dehydrogenation of the L-glucose C1-hydroxyl group to form L-glucono-1,5-lactone, which would in turn be spontaneously hydrolyzed to L-gluconate.

The Pathway downstream of L-Gluconate-- Next, we evaluated enzyme activity toward L-gluconate with cell-free extracts of strain 43P cultivated in L-Glc MM, and again an NAD⁺ dependent L-GnDH activity was detected. The enzyme catalyzing the reaction was partially purified by about 100-fold (supplemental Table 3) and showed a 35 kDa band on SDS-PAGE that corresponded to the activity (supplemental Fig. 2B). This protein had an N-terminal amino acid sequence of MKALIEKEGHTVIGEISEP, which has 100% identity with strain PD1222 Pden_4931, a putative oxidoreductase that is a member of the Zn-containing alcohol dehydrogenase family. As such, the same PCR-based strategy used to identify *lgdA* was

adopted to obtain sequence information for the gene for L-GnDH and its surrounding region.

The gene for L-GnDH was found in a nine-gene cluster corresponding to Pden_4924 to Pden_4932 in strain PD1222 with amino acid sequence identities of 71-91% (Fig. 2B), and was designated *lgnA-lgnI*. At the cluster's upstream end, we also found a gene corresponding to Pden_4923, termed *lgnR*, which may function as a transcriptional regulator of the cluster. Based on the Pden gene annotations, the first four genes of the cluster, *lgnA-lgnD*, presumably encode subunits of an ABC transporter for polar amino acids, while *lgnE-lgnI* encode, respectively, a galactarate dehydratase, a KDGal kinase, a KDPGal aldolase, an alcohol dehydrogenase (L-GnDH), and a short-chain dehydrogenase/reductase. This information led us to consider the possibility that L-gluconate can be assimilated by enzymes encoded in this cluster, and this was assessed by analyzing each enzyme activity using purified recombinant enzymes (supplemental Fig. 2C).

i. L-Gluconate Dehydrogenation by LgnH. As expected, LgnH showed L-GnDH activity (Table 1). The activity was solely dependent on NAD⁺, not NADP⁺. The enzyme also showed dehydrogenase activity toward L-galactonate, the C-4 epimer of L-gluconate, while no activity was observed toward other epimers, such as D-idonate (C-5) and L-mannonate (C-2) and uronic acids such as D-glucuronate and D-galacturonate. NMR analysis of the reaction product from L-gluconate showed chemical shifts identical to that of commercial D-5-keto-gluconate (supplemental Fig. 3B), while HPLC analysis showed opposite optical rotation to D-5-keto-gluconate (Fig. 3B). These data clearly indicate that LgnH catalyzes dehydrogenation of the L-gluconate C-5 hydroxyl group to form L-5-keto-gluconate.

ii. LgnI-Mediated L-5-Keto-gluconate Reduction. LgnI is a member of the short chain dehydrogenase/reductase family and is therefore predicted to catalyze NAD(P)⁺/NAD(P)H dependent oxidoreductase activity. Consequently, reduction of the purified reaction product of LgnH, L-5-keto-gluconate, was observed with

LgnI in an NADPH-dependent manner (Table 1). The reaction product showed the same chemical shifts and peak in NMR and HPLC analyses, respectively, as an authentic D-idonate, the C-5 epimer of L-gluconate (supplemental Fig. 3C and Fig. 3C). Therefore, we concluded that LgnI is responsible for the reduction of L-5-keto-gluconate to form D-idonate.

iii. LgnE-Mediated Dehydration of D-Idonate. When LgnE was incubated with D-idonate in the presence of FeCl₂ and DTT, the formation of reducing sugars was detected (Table 1). We did not detect any activity toward meso-galactarate, although LgnE showed sequence similarity to galactarate dehydratase (GarD) from *E. coli* K-12 (38%). NMR and HPLC analyses of the reaction products showed, respectively, the same chemical shifts and peak as KDGal, which was prepared from D-galactonate using recombinant *E. coli* D-galactonate dehydratase (DgoD) (supplemental Fig. 3D and Fig. 3D). These data indicate that LgnE catalyzes dehydration of D-idonate to produce KDGal.

iv. Phosphorylation and Aldol Cleavage of KDGal by LgnF and LgnG. Since LgnF and LgnG showed amino acid sequence similarity to *E. coli* DgoK (33%) and DgoA (43%), which catalyze ATP-dependent phosphorylation of KDGal to form KDPGal, and aldol-cleavage of KDPGal to form pyruvate and Gap, respectively, we tested their corresponding activities.

As expected, LgnF produced ADP from ATP only in the presence of KDGal, indicating that it phosphorylates KDGal (Table 1). Also, pyruvate and Gap formation was enzymatically detected when LgnG was incubated with KDPGal. These data clearly show that LgnF catalyzes phosphorylation of KDGal to form KDPGal, which is then converted to pyruvate and Gap by LgnG.

Involvement of the Identified Genes in the L-Glucose Catabolic Pathway-- To determine how the identified genes participate in the L-glucose catabolic pathway, we constructed gene disruption mutants of *lgnA*, *lgnE*, *lgnH*, and *lgnI*, based on strain 43P, by inserting a *Km^r* cassette into the respective genes (supplemental Fig. 4). Growth of cells harboring these

disrupted genes in minimal medium containing L-glucose or related compounds was observed. As expected, the *Algda*, *AlgnH* and *AlgnI* mutants did not grow in the presence of "upstream" compounds of the pathway, such as L-glucose for *Algda*, and L-glucose and L-gluconate for *AlgnH* and *AlgnI* mutants, whereas they could grow using "downstream" compounds (Fig. 4). These results strongly indicate that these genes are responsible for L-glucose catabolism.

In contrast, the *AlgnE* mutant showed rather weak but distinct growth in the presence of each compound tested. It may be possible that strain 43P possesses a paralog to *lgnE*, as the related strain, PD1222, possesses such a paralog, Pden_4671, which exhibits 53% identity.

Though *LgnH* utilized L-galactonate as a substrate as well as L-gluconate, none of the gene disrupted mutants including the *AlgnH* mutant showed growth defects with L-galactonate (Fig. 4D), suggesting that these genes are not involved in L-galactonate catabolism (see below). Also, none of the mutants showed growth defects with *scyllo*- and *myo*-inositols, except the *Algda* mutation affected the growth with *scyllo*-inositol severely (Fig. 4E and F). These results indicate that *lgdA* is required for both L-glucose and *scyllo*-inositol utilization, whereas the genes in the *lgn* cluster are not involved in inositol catabolism.

DISCUSSION

This study sought to describe the L-glucose-catabolic pathway in *Paracoccus* sp. 43P, a model for which is shown in Fig. 5. Based on the gene organization, this pathway could be divided into two parts, oxidation of L-glucose by *LgdA* and L-gluconate catabolism by enzymes encoded by the *lgn* cluster.

LgdA showed sequence similarity to Gfo/IDH/MocA family proteins, and its gene was located in a putative inositol utilization gene cluster of strain 43P. Kinetic analysis of purified *LgdA* showed efficient oxidation of *scyllo*-inositol rather than L-glucose, and the *Algda* mutation affected the growth with *scyllo*-inositol. Together, these results strongly indicate that the physiological role of *LgdA* involves *scyllo*-inositol catabolism. Among

inositol dehydrogenases (IDHs), *myo*-IDH of *Bacillus subtilis* (IolG) is the best-characterized enzyme, and was reported to catalyze oxidation of the axial hydroxyl group of *myo*-inositol and D-*chiro*-inositol (22). IolG also catalyzes oxidation of α -D-glucose and α -D-xylose (23), which possess the same stereo-configuration of *myo*-inositol hydroxyl groups. A β -anomer of L-glucose possesses the same stereo-configuration as *scyllo*-inositol, *i.e.* hydroxyl groups occupy all equatorial positions, so it is possible that *LgdA* utilizes both L-glucose and *scyllo*-inositol as substrates. There are two *scyllo*-IDHs that have been characterized at an enzyme level, IolW and IolX from *B. subtilis* (24), and these enzymes showed low sequence identity with *LgdA* (22.1% and 26.3%, respectively). IolX showed no activity toward L-glucose, whereas IolW had only low activity ($k_{cat}/K_m = 0.37 \text{ min}^{-1}\text{mM}^{-1}$; data from Drs. Dong-min Kang and Ken-ichi Yoshida)² that was about 1/50 that of *LgdA*. These findings may indicate that *LgdA* has a unique substrate recognition mechanism.

On the other hand, results for the *Algda* mutant showed a clear involvement of this gene in L-glucose catabolism. Although L-glucose is an unnatural sugar, these results indicate that, at least for strain 43P, bacterial inositol dehydrogenases may be involved in both inositol and sugar catabolism.

The reactions downstream of the *LgdA* reaction product, L-gluconate, are analogous to those of the *E. coli* L-galactonate catabolic pathway, which is the C-4 epimer of L-gluconate (25, 26). Both the L-aldonates are epimerized at the C-5 position by dehydrogenase/reductase reactions (D/L conversion) to produce D-aldonates, converted to D-2-keto-3-deoxyaldonates by dehydratases, phosphorylated at C-6 position by kinases, and then converted by aldolases to the glycolysis substrates pyruvate and Gap. Comparison of the amino acid sequences of the *lgn* enzymes, *LgnH*, *LgnI*, *LgnE*, *LgnF* and *LgnG*, with those of corresponding enzymes in the *E. coli* L-galactonate pathway, YjjN, UxaB, UxaA, KdgK and Eda, respectively, showed limited sequence identities of about 30%, with *LgnI* and *LgnF* in particular showing no similarities to

their corresponding proteins. Moreover, the mutant deficient in LgnH or LgnI could not grow on L-gluconate, but did not show any growth defects on L-galactonate (Fig. 4D). Taken together, we conclude that the *lgn* cluster enzymes are specific to L-gluconate rather than L-galactonate.

The D/L conversion by the dehydrogenase/reductase reactions are also seen in the catabolism of other six-carbon L-aldonates, such as L-gulonate and L-idonate, in *E. coli* (27, 28). Like the pathway described in this study, the final products of these catabolic pathways are pyruvate and Gap, where Gap is used as a key-intermediate for substrate-level phosphorylation in the downstream portion of the Embden-Meyerhof-Parnas pathway. Since L-glyceraldehyde-3-phosphate is not utilized as a substrate by GapDH, but rather shows a bactericidal effect (29), the D/L conversion may be important for obtaining the Gap D-form following cleavage reactions that are catalyzed by aldolases. In contrast, the fungal L-galactonate catabolic pathway does not involve this process, with L-2-keto-3-deoxygalactonate, formed by L-galactonate dehydratase, being directly cleaved to form pyruvate and L-glyceraldehyde. However, the latter product is not utilized directly, but is reduced to an achiral compound, glycerol (30, 31). Therefore, a homochirality issue is likely present in the catabolism of glyceraldehyde, the minimum chiral compound, not in six-carbon sugars. This notion agrees with the recent findings that D-glyceraldehyde might have been selectively synthesized in the prebiotic era, via a formose reaction in the presence of L-amino acids (32).

Phylogenetic analyses based on the amino acid sequences of the enzymes in this pathway and their related sequences showed that they are located in deeply-branched lineages that are distinct from those of enzymes in the corresponding protein families whose functions are known, and from KEGG genome database sequences, except for those from *P. denitrificans* (supplemental Fig. 5). The lineages containing LgdA, LgnF and LgnG were included in the clusters of *Alphaproteobacteria*, whereas those of LgnH, LgnI and LgnE diverged at branch

points with clusters including sequences from other bacterial affiliations, *i. e.*, clusters containing sequences mainly from *Bacteroidetes* for LgnH and LgnI, and *Betaproteobacteria* for LgnE. This divergence may indicate a unique evolutionary origin of this catabolic pathway.

There are several microorganisms in the class *Alpha-*, *Beta-*, and *Gammaproteobacteria* that possess sets of all six orthologs, or five orthologs corresponding to the enzymes for L-gluconate assimilation, which can be assumed as "potential" L-glucose or L-gluconate utilizing organisms from a metabolic pathway standpoint. These orthologs were widely distributed within the clusters of those from the genome sequences, and there are no organisms other than *P. denitrificans*, which possesses all such orthologs found in strain 43P in the same or nearby clusters. Most of these organisms likely cannot assimilate L-glucose or L-gluconate, because there are previous no reports on the assimilation of such compounds. Rather, it may be that either no other organisms can utilize such compounds or different catabolic pathways for L-glucose or L-gluconate exist if any of these organisms can indeed utilize them.

P. denitrificans utilized L-gluconate (Fig. 4B) but not L-glucose (Fig. 1). In our preliminary experiments, recombinant Pden_1680 from strain NBRC 102528^T showed L-GDH activity similar to LgdA ($k_{cat} = 836 \pm 12 \text{ min}^{-1}$, $K_m = 78.0 \pm 8.2 \text{ mM}$), indicating that this strain possesses the ability to utilize L-glucose. The reason why *P. denitrificans* cannot assimilate L-glucose is not known, but could be because the gene cluster including Pden_1680 is not transcribed or L-glucose cannot be incorporated into the cells, at least under our culture conditions.

L-glucose is not found in natural environment, and therefore the physiological role of this catabolic pathway is unknown. However, our data clearly shows that the *lgn* gene cluster is specifically responsible for L-gluconate utilization. This finding may indicate the presence of L-gluconate in nature, which is believed to be not existed as well as L-glucose.

We are now attempting to analyze the L-glucose catabolic pathway of other isolates we obtained, which will lead to a better understanding of L-glucose catabolism.

REFERENCES

1. Rudney, H. (1940) The utilization of L-glucose by mammalian tissue and bacteria. *Science* **92**, 112-113
2. Livesey, G., and Brown, J.C. (1995) Whole body metabolism is not restricted to D-sugars because energy metabolism of L-sugars fits a computational model in rats. *J. Nutr.* **125**, 3020-3029
3. Sun, H.J., Saccomanno, V., Hedlund, B., and McKay, C.P. (2009) Stereo-specific Glucose consumption may be used to distinguish between chemical and biological reactivity on Mars: A preliminary test on Earth. *Astrobiol.* **9**, 443-446
4. Sasajima, K., and Shinsky, A. (1979) Oxidation of L-glucose by a *Pseudomonad*. *Biochim. Biophys. Acta.* **571**, 120-126
5. Kovach, M.E., Elzer, P.H., Hill, D.S., Robertson, G.T., Farris, M.T., Roop, R.M., and Peterson, K.M. (1995) Four new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene* **166**, 175-176
6. Moore, S., and Link, K.P. (1940) Carbohydrate characterization. *J. Biol. Chem.* **133**, 293-311
7. Babbit, P.C., Mrachko, G.T., Hasson, M.S., Huisman, G.W., Kolter, R., Ringe, D., Petsko, G.A., Kenyou, G.L., and Gerlt, J.A. (1995) A functionally diverse enzyme superfamily that abstracts the alpha protons of carboxylic acid. *Science* **267**, 1159-1161
8. Miller, G.L. (1959) Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal. Chem.* **31**, 426-428
9. Lobanok, A.G., Sapunova, L.I., Dikhtievski, Y.O., and Kazakevich, I.O. (1998) Screening of glucose isomerase-producing microorganisms. *World. J. Microbiol. Biotechnol.* **14**, 259-262
10. Hemker, M., Stratmann, A., Goeke, A., Schröder, W., Lenz, J., Piepersberg, W., and Pape, H. (2001) Identification, cloning, expression, and characterization of the extracellular acarbose-modifying glycosyltransferase, AcbD, from *Actinoplanes* sp. strain SE50. *J. Bacteriol.* **183**, 4484-4492
11. Howe, A.F., Groom, T, and Carter, R.G. (1964) Use of polyethylene glycol in the concentration of protein solution. *Anal. Biochem.* **9**, 443-453
12. Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685
13. Ochman, H., Gerber, A.S., and Hartl, D.L. (1988) Genetic application of an inverse polymerase chain reaction. *Genetics* **120**, 621-623
14. Kuorelahti, S., Jouhten, P., Maaheimo, H., Penttilä, M., and Richard, P. (2006) L-galactonate dehydratase is a part of the fungal path for D-galacturonic acid catabolism. *Mol. Microbiol.* **61**, 1060-1068
15. Wong, T.Y., and Yao, X.T. (1994) The Dely-Doudoroff pathway of galactose metabolism in *Azotobacter vinelandii*. *Appl. Environ. Microbiol.* **60**, 2065-2068
16. Lowry, O.H. (1951) Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265-275
17. Anderson, K., Li, S.C., and Li, Y.T. (2000) Diphenylamine-aniline-phosphoric acid reagent, a versatile spray reagent for revealing glycoconjugates on thin-layer chromatography plates. *Anal. Biochem.* **287**, 337-339
18. Groeve, M.M., Depreitere, V., Desmet, T., Soetaert, W. (2009) Enzymatic production of α -D-galactose 1-phosphate by lactose phosphorolysis. *Biotechnol. Lett.* **31**, 1873-1877.
19. Simon, R., Prier, U., and Puhler, A. (1983) A broad host range mobilization system for in vivo genetic engineering: transposon mutagenesis in gram negative bacteria. *Nat. Biotechnol.* **1**, 784-791
20. Larkinn, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H.,

- Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J., and Higgins, D.G. (2007) CLUSTAL W and CLUSTAL_X version 2.0. *Bioinformatics* **23**, 2947-2948
21. Page, R.D. (1996) TreeView: an application to display phylogenetic tree on personal computers. *Comput. Appl. Biosci.* **12**, 357-358
 22. Yoshida, K., Yamaguchi, M., Morinaga, T., Kinehara, M., Ikeuchi, M., Ashida, H., and Fujita, Y. (2008) *myo*-inositol catabolism in *Bacillus subtilis*. *J. Biol. Chem.* **283**, 10415-10424
 23. Ramaley, R., Fujita, Y., and Freese, E. (1979) Purification and properties of *Bacillus subtilis* inositol dehydrogenase. *J. Biol. Chem.* **254**, 7684-7690
 24. Morinaga, T., Ashida, H., and Yoshida, K. (2010) Identification of two *scyllo*-inositol dehydrogenase from *Bacillus subtilis*. *Microbiology* **156**, 1538-1546
 25. Cooper, R.A. (1979) The pathway for L-galactonate catabolism in *Escherichia coli* K12. *FEBS. Lett.* **103**, 216-220
 26. Reed, J.L., Patel, T.R., Chen, K.H., Joyce, A.R., Applebee, M.K., Herring, C.D., Bui, O.T., Knight, E.M., Fong, S.S., and Palsson, B.O. (2006) System approach to refining genome annotation. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 17480-17484
 27. Cooper, R.A. (1980) The pathway for L-gulonate catabolism in *Escherichia coli* K-12 and *Salmonella typhimurium* LT-2. *FEBS. Lett.* **115**, 63-67.
 28. Bausch, C., Peekhaus, N., Utz, U., Blais, T., Murray, E., Lowary, T., and Conway, T. (1998) Sequence analysis of the GntII (subsidiary) system for gluconate metabolism reveals novel pathway for L-idonic acid catabolism in *Escherichia coli*. *J. Bacteriol.* **180**, 3701-3710
 29. Kalyananda, M.J., Engel, R., and Tropp, B.E. (1987) Metabolism of L-glyceraldehyde 3-phosphate in *Escherichia coli*. *J. Bacteriol.* **169**, 2488-2493
 30. Zhang, L., Thiewes, H., and van Kan, J.A. (2011) D-galacturonic acid catabolic pathway in *Botrytis cinerea*. *Fungal. Genet. Biol.* **48**, 990-997
 31. Hilditch, S., Berghäll, S., Kalkkinen, N., Penttilä, M., and Richard, P. (2007) The missing link in the fungal D-galacturonate pathway: identification of the L-threo-3-deoxy-hexulose aldolase. *J. Biol. Chem.* **282**, 26195-26201
 32. Breslow, R., and Cheng, Z.L. (2010) L-amino acid catalyze the formation of an excess of D-glyceraldehyde, and thus of other D-sugars, under credible prebiotic conditions. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 5723-5725

FOOTNOTES

* This work was supported by the NISR Research Grant from the Noda Institute for Scientific Research, Japan and Grants-in-Aid for Scientific Research (B) from Japan Society for Promotion of Science.

¹The abbreviations used are: L-Glc MM, L-glucose minimal medium; L-GDH, L-glucose dehydrogenase; DNS, 3,5-dinitrosalicylic acid; L-GnDH, L-gluconate dehydrogenase; KDGal, D-2-keto-3-deoxygalactonate; KDPGal, D-2-keto-3-deoxy-6-phosphogalactonate; IPTG, Isopropyl- β -D-thiogalactopyranoside; HMQC, Hetero-nuclear Multiple Quantum Coherence; Gap, D-glyceraldehyde-3-phosphate; LDH, L-lactate dehydrogenase; PK, pyruvate kinase; RI, refractive index; OR, optical rotation; IDH, inositol dehydrogenase.

²To whom correspondence may be addressed: Department of Biofunctional Chemistry, University of Kobe, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan. Tel.: +81-90-4279-6263; E-mail: ken-yoshi@kobe-u.ac.jp

Data deposition: The nucleotide sequences of the 16S rRNA gene, the gene clusters containing *lgdA* and *lgn* genes of *Paracoccus* sp. 43P were deposited to Genbank/EMBL/DDBJ databases under accession numbers AB727354, AB727355, and AB727356, respectively.

FIGURE LEGENDS

Figure 1. Growth (open symbols) and L-glucose consumption (closed symbols) of strains 43P (circles) and NBRC 102528^T (triangles) in L-Glc MM. Growth was monitored by measuring the absorbance at 600 nm. The DNS method was used to determine the L-glucose concentration by measuring the reducing sugars in the culture medium of strains 43P and NBRC 102528^T. Cultivation was conducted in three independent cultures and average values \pm S. D. are shown.

Figure 2. Gene organization of the clusters containing *lgdA* (A) and *lgn* (B). Genes encoding the enzymes for L-glucose catabolism are shaded, and sequence identities to the corresponding *P. denitrificans* PD1222 genes are shown below.

Figure 3. HPLC analysis of LgdA (A), LgnH (B), LgnI (C), and LgnE (D) reaction products detected by a refractive index (RI) and chiral detector (OR). Black and grey lines indicate chromatograms of the reaction products and authentic compounds, L-gluconate (A), D-5-keto-gluconate (B), D-idonate (C), and KDGal (D), respectively.

Figure 4. Growth of strain 43P and gene disruption mutants in minimal media containing L-glucose (A), L-gluconate (B), D-idonate (C), L-galactonate (D), *scyllo*-inositol (E), and *myo*-inositol (F). Open circles, strain 43P; closed circles, Δ *lgdA* strain; open triangles, Δ *lgnE* strain; closed triangles, Δ *lgnH* strain; open squares, Δ *lgnI* strain. In panel (B), the growth of strain NBRC 102528^T is also shown (closed squares). Cultivation was conducted in three independent cultures and average values \pm S. D. are shown.

Figure 5. A model of the L-glucose catabolic pathway in strain 43P based on the results of this study. Altered moieties in each enzyme reaction are shown in grey.

Table 1. Kinetic parameters of the L-glucose catabolic enzymes.

Enzyme	Substrate	K_m (mM)	k_{cat} (min^{-1})	k_{cat}/K_m ($\text{min}^{-1} \text{mM}^{-1}$)
LgdA	L-glucose ^a	59.7 ± 5.7	1040 ± 28	17.4
	D-glucose ^a	88.7 ± 8.3	215 ± 36	2.42
	<i>scyllo</i> -inositol ^a	3.70 ± 0.4	705 ± 12	190
	<i>myo</i> -inositol ^a	53.3 ± 8.6	572 ± 30	10.7
	NAD ^{+b}	2.24 ± 0.4	742 ± 20	309
	NADP ^{+b}	-	n.d. ^c	-
LgnH	L-gluconate ^d	0.659 ± 0.023	573 ± 5	869
	L-galactonate ^d	0.224 ± 0.011	178 ± 2	828
	NAD ^{+e}	0.155 ± 0.006	543 ± 4	3500
	NADP ^{+e}	-	n.d. ^c	-
LgnI	L-5-keto-gluconate ^f	0.180 ± 0.020	1200 ± 33	6640
	NADPH ^g	0.085 ± 0.004	1910 ± 90	22500
	NADH ^g	-	n.d. ^h	-
LgnE	D-idonate	0.454 ± 0.072	376 ± 12	828
LgnF	KDGal	0.629 ± 0.054	1710 ± 34	2720
	KDPGal ⁱ	0.587 ± 0.023	4290 ± 270	7310
LgnG	KDPGal ^j	0.466 ± 0.012	3900 ± 31	8370

The assays were conducted in triplicate, and average values \pm S. D. are shown.

^a The activity was measured with a fixed concentration of 10 mM NAD⁺.

^b The activity was measured with a fixed concentration of 20 mM *scyllo*-inositol.

^c Activity was not detected with 10 mM NADP⁺.

^d The activity was measured with a fixed concentration of 2 mM NAD⁺.

^e The activity was measured with a fixed concentration of 10 mM L-gluconate.

^f The activity was measured with a fixed concentration of 0.2 mM NADPH.

^g The activity was measured with a fixed concentration of 2 mM L-5-keto-gluconate.

^h Activity was not detected with 0.2 mM NADH.

ⁱ The activity was measured by pyruvate production.

^j The activity was measured by Gap production.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5