

DA
3081
2002
03

**ALKYLPHENOLS AND PHYTOESTROGENS,
THEIR INTERACTION WITH FOOD FACTORS
AND MECHANISMS OF ACTION *IN VITRO***

**Division of Agricultural and Forest Engineering
Doctoral Degree Program in Agricultural Sciences
University of Tsukuba**

TERENCE P.N. TALORETE

寄贈
タロレンス P. N.
テロレテ氏

03302700

CONTENTS

CHAPTER 1	1
General Introduction	
CHAPTER 2	8
Alkylphenolic compounds and their effect on the injury rate, survival and acetylcholinesterase activity of the rat neuronal cell line PC12	
2.1. Introduction	9
2.2. Materials and Methods	13
2.3. Results and Discussion	16
2.4. Summary	21
CHAPTER 3	22
Phytoestrogens genistein and daidzein enhance the acetylcholinesterase activity of the rat pheochromocytoma cell line PC12 by binding to the estrogen receptor	
3.1. Introduction	23
3.2. Materials and Methods	25
3.3. Results and Discussion	28
3.4. Summary	31
CHAPTER 4	32
<i>Agaricus blazei</i> (Class Basidiomycotina) aqueous extract enhances the expression of c-Jun protein in MCF7 cells	
3.2. Introduction	33
3.3. Materials and Methods	36
3.4. Results	42
3.5. Discussion	44
3.1. Summary	48
CHAPTER 5	49
General Discussion and Conclusion	
REFERENCES	55
LIST OF PAPERS	80

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
1	Chemical structure of <i>p</i> -alkylphenols. n = 5 (pentyl), 6 (hexyl), 7 (heptyl), 8 (octyl), 9 (nonyl).	69
2	Lactate dehydrogenase activity (expressed as cell injury rate) of PC12 cells after 15 min incubation with varying concentrations of different alkylphenols. Results based from three independent replicates are expressed as percent cell injury \pm SEM.	70
3	Ethidium-bromide-stained agarose gel showing genomic DNA from the rat pheochromocytoma cell line PC 12. Results show no fragmentation of DNA after 4 h treatment with the indicated concentrations of <i>p</i> -nonylphenol (NP) and <i>p</i> -octylphenol (OP). PC12 DNA was extracted using the Sodium Iodide Method and electrophoresed on a 1.5% agarose gel. C, control; MW, molecular weight marker. Results represent three independent replicates.	71
4	Acetylcholinesterase (AChE) activity of PC12 cells after 24 h incubation with 40 ng ml ⁻¹ nerve growth factor (NGF), which stimulates AChE release, and varying concentrations of different alkylphenols. Results based from at least three independent replicates are expressed as relative acetylcholinesterase activity, which is obtained by dividing the mean AChE activity of cells incubated with alkylphenolic compounds and NGF by the mean AChE activity of cells incubated with NGF alone (positive control), \pm SEM. Relative AChE activity of positive control is therefore 1. Results show significant decreases compared to control ($p < 0.05$, <i>t</i> -test), except for those with asterisk (*).	72
5	Acetylcholinesterase (AChE) activity of PC12 cells after 24 h incubation with 40 ng ml ⁻¹ nerve growth factor (NGF), which stimulates AChE release, and varying concentrations of the phytoestrogens genistein and daidzein, with or without 1.5 μ M tamoxifen, a known estrogen receptor (ER) antagonist. Results based from three independent replicates are expressed as relative acetylcholinesterase activity, which is obtained by dividing the mean AChE activity of cells incubated with the test compounds and NGF by the mean AChE activity of cells incubated with NGF alone (positive control), \pm SEM. Relative AChE activity of positive control is	73

therefore 1. Results show significant differences ($p < 0.05$, t-test) in the AChE activities of cells in the presence or absence of tamoxifen at phytoestrogen concentrations of up to $0.325 \mu\text{M}$, indicating involvement of the ER in AChE induction.

- 6 Proliferative activity of MCF7 cells after 6 days incubation with the phytoestrogens genistein and daidzein in the absence (*Fig. 6a, top*) and presence (*Fig. 6b, bottom*) of $1.5 \mu\text{M}$ tamoxifen, a known estrogen receptor antagonist. Results based from three independent replicates are expressed as relative estrogenic (MTT) activity, which is obtained by dividing the mean proliferative activity of cells incubated with the test compounds with or without tamoxifen by the mean proliferative activity of the control (no additions), \pm SD. Relative estrogenic activity of control is therefore 1. Results show increased proliferation of MCF7 cells with increasing phytoestrogen concentration (*Fig. 6a*), which was effectively blocked by tamoxifen (*Fig. 6b*). Data analyzed using one-way ANOVA showed significant differences ($p < 0.05$) starting at $0.02 \mu\text{M}$ isoflavone concentration. 74
- 7 Acetylcholinesterase (AChE) activity of PC12 cells after 24 h incubation with 40 ng ml^{-1} nerve growth factor (NGF), which stimulates AChE release, and varying concentrations of 17β -estradiol (E2) and E2 plus $0.08 \mu\text{M}$ genistein (G) and $0.08 \mu\text{M}$ daidzein (D) or $1.5 \mu\text{M}$ tamoxifen (TAM). Results based from three independent replicates are expressed as relative acetylcholinesterase activity, which is obtained by dividing the mean AChE activity of cells incubated with the test compounds and NGF by the mean AChE activity of cells incubated with NGF alone (positive control), \pm SEM. Relative AChE activity of positive control is therefore 1. Results show no significant differences ($p > 0.05$, t-test) in the AChE activities of cells incubated with E2 in the presence or absence D, G, or TAM, suggesting that the presence of an estrogenic compound alone, in this case, E2, does not necessarily lead to AChE induction. 75
- 8 MTT activity of MCF-7 cells in the presence or absence of $1 \mu\text{M}$ *p*-nonylphenol (NP), 1 nM 17β -estradiol (E2) and 5% (v/v) *Agaricus* extract (AE). Results based from at least four independent replicates are expressed as relative MTT activity, which is obtained by dividing the mean MTT activity of the cells incubated with the test compounds and/or AE by the mean activity of the control (no additions), \pm SD. Relative MTT activity of control is therefore 1. The statistical analyses of this figure are shown in Table 3 (page 42). 76
- 9 MTT Activity of MCF-7 cells in the presence of varying concentrations of *p*-nonylphenol (NP), plus or minus 1 nM 17β -estradiol (E2) and 5% (v/v) aqueous *Agaricus* extract (AE). Results based from four independent replicates are expressed as relative MTT activity, which is 77

obtained by dividing the mean MTT activity of cells incubated with the test compounds and/or AE by the mean MTT activity of the control (no additions), \pm SD. Relative MTT activity of control is therefore 1. Two-way ANOVA showed significant differences between the <plus NP/E2/AE> and <plus NP/AE> treatments compared to control ($p < 0.0001$) while the differences in the MTT activity of cells incubated in different NP concentrations within the same treatments are not statistically significant ($P > 0.05$).

- 10 Western blotting results showing the expression of the c-Jun/API protein in MCF7 cells after 6 days incubation in the presence or absence of 1 μ M *p*-nonylphenol (NP), 1 nM 17 β -estradiol (E2), and 5% (v/v) aqueous *Agaricus* extract (AE). Lane 1, plus NP only; 2, plus NP, E2 & AE; 3, plus NP & E2; 4, plus NP & AE; 5, plus E2 only; 6, plus E2 & AE; 7, plus AE only; 8, no addition (control). Results show enhanced expression of the c-Jun protein in the presence of 5% (v/v) aqueous *Agaricus* extract (Lane 7). 78
- 11 Protein kinase C (PKC) assay results showing PKC activity in MCF7 cells after 24 h incubation in the presence or absence of 1 μ M *p*-nonylphenol (NP), 1 nM 17 β -estradiol (E2), and 5% (v/v) aqueous *Agaricus* extract (AE). Lane 1, no addition (control); 2, plus E2 only; 3, plus NP only; 4, plus AE only; 5, plus E2 & AE; 6, plus NP & AE; +PKC, positive control; -PKC, negative control. This means that the PKC enzyme is not a factor in the observed results. 79

ACKNOWLEDGEMENTS

This thesis would not have been possible without the kindness and support of very important people:

- **Dr. Hiroko Isoda**, my direct supervisor in the Cell Biology Research Group, for her excellent guidance and unparalleled kindness. Dr. Isoda is not only an outstanding researcher and a highly knowledgeable expert in her field, but she is also an excellent human being. Our weekly discussions and highly interactive seminars have contributed largely not only to my intellectual growth but also to that of other students. My thanks will never be enough.
- **Professor Takaaki Maekawa**, my direct professor, for his trust and kindness in accepting me in his laboratory, for the support that I have always felt ever since I joined his *kenkyushitsu*, and for his excellent suggestions in the improvement of this thesis. My *gratitude also extends to Prof. Maekawa's hardworking secretary Ms. Nobuko Akamatsu* for her support and kindness.
- **Professor Suguira, Professor Satake and Professor Tanaka**, my dissertation committee members, for their helpful suggestions in the improvement of this thesis, and also for their support.
- The **Cell Group Members** (past and present), for their kind assistance, friendship, company and encouragement.
- My former mentors and advisers, **NAST Academician Professor Lourdes J. Cruz** and **Professor Wilfredo L. Barraquio** of the University of the Philippines (Diliman), and **Dr. Takanori Higashira**, formerly with the National Institute of Bioscience and Human-Technology (AIST-Tsukuba), for leading me through the right path in science.
- My dear parents, **Exequiel Talorete** and **Cosette Nemenzo Talorete**, and my dear aunt, **Natalie Talorete Mackey**, for their unending support for my education and loving encouragement.

Finally, I am deeply indebted to the **Japanese government** and to the **Japanese people** for the Japanese Government (Monbusho) scholarship, and for this rare opportunity that I have gratefully obtained during my stay in Tsukuba. (My master's thesis was also undertaken in Tsukuba many years ago through the auspices of the Japan International Cooperation Agency.)

THANK YOU SO MUCH FOR EVERYTHING.