

# A novel preoperative fusion analysis using three-dimensional MDCT combined with three-dimensional MRI for patients with hilar cholangiocarcinoma

著者別名	大城 幸雄, 佐々木 亮孝, 那須 克宏, 大河内 信弘
journal or publication title	Clinical imaging
volume	37
number	4
page range	772-774
year	2013-07
権利	(C) 2013 Elsevier Inc. NOTICE: this is the author's version of a work that was accepted for publication in Clinical imaging. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Clinical imaging, VOL37, ISSUE4, 2013, <a href="http://dx.doi.org/10.1016/j.clinimag.2013.02.002">http://dx.doi.org/10.1016/j.clinimag.2013.02.002</a>
URL	<a href="http://hdl.handle.net/2241/119605">http://hdl.handle.net/2241/119605</a>

doi: 10.1016/j.clinimag.2013.02.002

Clinical Imaging

Case report

**A novel preoperative fusion analysis using 3-dimensional MDCT  
combined with 3-dimensional MRI for patients with hilar  
cholangiocarcinoma**

Yukio Oshiro, M.D.<sup>a</sup>, Ryoko Sasaki, M.D., Ph.D., F.I.C.S.<sup>a,\*</sup>,  
Katsuhiko Nasu, M.D., Ph.D.<sup>b</sup>, Nobuhiro Ohkohchi, M.D., Ph.D.<sup>a</sup>

*<sup>a</sup>Department of Organ Transplantation, Gastroenterology and  
Hepato-biliary surgery, University of Tsukuba, Graduate School of  
Comprehensive Human Sciences, 1-1-1 Tennodai, Tsukuba 305-8575,  
Japan*

*<sup>b</sup>Department of radiology, University of Tsukuba, Graduate School of  
Comprehensive Human Sciences, 1-1-1 Tennodai, Tsukuba 305-8575,  
Japan*

*\*Corresponding address, Reprint request:*

Ryoko Sasaki, M.D., Ph.D.

Department of Organ Transplantation, Gastroenterology and Hepato-biliary surgery,  
University of Tsukuba, Graduate School of Comprehensive Human Sciences, 1-1-1  
Tennodai, Tsukuba 305-8575, Japan

TEL: +81-29-853-3221

FAX: +81-29-853-3222

Email: rsasaki.dr@hotmail.co.jp

***An abbreviated title:***

3-dimensional fusion MDCT combined with MRCP

1    **Abstract**

2    The purpose of the present study was to evaluate the anatomical relationship between  
3    the tumor, portal veins, hepatic arteries, and hilar hepatic ducts at the hepatic hilum  
4    using a novel preoperative fusion analysis for patients with hilar cholangiocarcinoma.  
5    This involved combining 3-dimensional multidetector-row computed tomography  
6    (3D-MDCT) with 3-D magnetic resonance imaging (MRI). This novel fusion imaging  
7    technique can play an important clinical role for patients undergoing surgery for hilar  
8    cholangiocarcinoma.

9

10    ***Key words:***

11    hilar cholangiocarcinoma, 3-dimensional imaging, fusion imaging, MDCT, MRI

12

## 1 **1. Introduction**

2 To determine the optimal surgical procedure for patients with hilar  
3 cholangiocarcinoma, it is essential to know the anatomical relationship between the  
4 tumor and the vessels around the hepatic hilum in individual patients. Several  
5 institutions have recently developed 3-dimensional multidetector-row computed  
6 tomography (3D-MDCT) cholangiography or drip infusion 3D-MDCT for use with  
7 patients with hilar cholangiocarcinoma [1-4]. However, it has been said that these  
8 radiological examinations have several problems with side effects such as cholangitis or  
9 anaphylactic shock due to infusion of the contrast material into the vein or bile duct.  
10 Therefore, we were looking forward to a new modality and method. To our knowledge  
11 there have been no published reports concerning the preoperative visualization of  
12 3D-MDCT fusion images combined with 3D-magnetic resonance  
13 cholangiopancreatography (MRCP) images. The purpose of the present study was to  
14 preoperatively evaluate portal veins and hilar hepatic ducts at the hepatic hilum in  
15 patients with hilar cholangiocarcinoma.

## 16 **2. Case**

17 The patient was a 52-year-old man. He consulted the Tsukuba University Hospital  
18 complaining of jaundice. Laboratory blood samples showed elevated total bilirubin and  
19 carcinogenic antigen (CA) 19-9, and MDCT and MRI were subsequently performed  
20 (Fig. 1).

21 MDCT images were acquired using an IDT-16 imager (Philips, Eindhoven, The  
22 Netherlands). Scan settings included a pitch of 17, a 0.75-second scan time per rotation,  
23 a table speed of 12 mm/rotation, and a detector configuration of  $0.75 \times 16$  mm. A power  
24 injector was used to administer 100 mL iopamidol (370 mgI/mL) at 4 mL/s through a  
25 20-G high pressure intravenous catheter. The MDCT protocol obtained 3 sets of hepatic  
26 images in succession, including images of arterial, and those of portal venous, and  
27 hepatic venous phases. The arterial phase images were obtained 5 seconds after peak  
28 aortic enhancement, portal venous phase images were taken 70 seconds after starting the  
29 injection, and the hepatic venous phase began 180 seconds after starting the injection.  
30 The MRI was performed on a 1.5 Tesla MRI imager (Achieva Nova Dual, Philips  
31 Medical Systems, Eindhoven, The Netherlands)

32 The preoperative fusion examination consisted of 4 steps. We first combined heavily  
33 T2-weighted MRCP images with axial thin slice 3D-T1-turbo field echo (TFE) images.  
34 Second, we combined the thin TFE images with thin MDCT axial images. Third, we  
35 superimposed MRCP data onto the axial CT images, focusing on the area around the  
36

1 hepatic hilum. Forth, we checked the following three registration landmarks: 1) the right  
2 border along the umbilical portion, 2) the bifurcation of the right and left hepatic arteries,  
3 and 3) the bifurcation of the right and left portal veins. Although we were able to  
4 delineate the hilar cholangiocarcinoma on the reconstructed 3D image by tracing the  
5 enhanced tumor on the 2D MDCT, we were not able to detect the tumor on the MRCP  
6 as expected. However, we confirmed that the resultant 3D fusion image was correct by  
7 checking the three registration landmarks (**Fig. 2**).

8 We used the Synapse Vincent medical imaging system (Fujifilm Medical, Tokyo),  
9 which was developed specifically for 3D visualization and virtual resection of the liver.  
10 This software offers standardized computation of liver anatomy functions and  
11 volumetric risk analysis based on 2D-CT imaging. The structures of the liver, portal  
12 vein, hepatic artery, hepatic vein, and tumor were extracted from MDCT scan data. One  
13 of the most important points of this novel fusion imaging is to examine MDCT during  
14 the patient's expiratory phase, which is the same condition of MRI. The user can  
15 visualize 3D images of the structures mentioned above as well as those of the hepatic  
16 arterial, portal, and hepatic venous phases. These images can also be rotated separately  
17 or simultaneously (**Fig. 2**).

18 We diagnosed the patient's disease as locally advanced hilar cholangiocarcinoma, T2,  
19 N1, M0, Stage IIB according to the UICC-TNM classification [5], and Type IIIb based  
20 on the Bithmuth- Corlette classification [6]. The patient and his family agreed to this  
21 procedure and provided informed consent. The patient tolerated MDCT and MRI  
22 without serious complications such as skin rash or other allergic adverse effects.

23 A chief surgeon (R.S.) performed the surgical resection. Based on preoperative  
24 imaging, we determined that the most appropriate surgical procedure was left hepatic  
25 lobectomy combined with total caudate lobectomy and extrahepatic bile duct resection  
26 (**Fig. 3**). No concomitant vascular resection was performed. Operative time was 581  
27 minutes, and estimated blood loss was 1215 ml. The patient's postoperative course was  
28 stable and uneventful, and he was discharged on postoperative day 14. Postoperative  
29 clinical pathology revealed that there was no residual carcinoma (R0). The patient is  
30 now alive 1 year and 1 month after surgery with no evidence of recurrence.

## 31 **Discussion**

32 To determine the optimal surgical approach to hilar cholangiocarcinoma for  
33 individual patients, it is essential to perform a detailed evaluation of the anatomical  
34 structures at the hepatic hilum. By using newly developed software and a novel method,  
35 we were able to easily analyze the hepatic arteries, portal veins, hepatic veins, and hilar  
36

1 bile ducts without performing 3D-MDCT cholangiography or drip infusion 3D-MDCT.

2 A more recent advance is the development of image integration, which refers to the  
3 process of registering previously acquired MRI and CT scans of the heart with the  
4 mapping space during an ablation procedure [7]. Further, a urology study has reported  
5 the usefulness of CT-MRI fusion imaging in radiotherapy planning for localized prostate  
6 cancer [8].

7 Several limitations of this novel modality should be addressed. First, at present this  
8 approach requires a considerable amount of time (about 5–6 hours) to create detailed  
9 images. Second, as described in the Patients and Methods section, obtaining images  
10 during the expiratory phase is one of the most important aspects of this method.  
11 Obtaining satisfactory images will be easiest with patients who are able to cooperate in  
12 this aspect. Third, this study involved only one patient, and our results will therefore  
13 need to be confirmed with larger populations.

14 In conclusion, this novel fusion imaging technique should play an important role in  
15 preoperative planning for patients undergoing surgery for hilar cholangiocarcinoma.

#### 16 **Acknowledgements**

17 The authors thank Mr. K. Kotera and Ms. K. Suzuki of Fujifilm Medical Co., Ltd, for  
18 their kind support.  
19  
20

## 1   **References**

- 2
- 3   1.   Sasaki R, Kondo T, Oda T, et al. Impact of three-dimensional analysis of  
4       multidetector row computed tomography cholangiography in operative planning  
5       for hilar cholangiocarcinoma. *Am J Surg* 2011;202:441-448.  
6
- 7   2.   Takahashi K, Sasaki R, Kondo T, et al. Preoperative 3D volumetric analysis for  
8       liver congestion applied in a patient with hilar cholangiocarcinoma. *Langenbecks*  
9       *Arch Surg* 2010;395:761-765.  
10
- 11  3.   Endo I, Shimada H, Sugioka M, et al. Role of three-dimensional imaging in  
12       operative planning for hilar cholangiocarcinoma. *Surgery* 2007;142:666-675.  
13
- 14  4.   Hansen C, Wieferich J, Ritter F, et al. Illustrative visualization of 3D planning  
15       models for augmented reality in liver surgery. *Int J comput Assist Radiol Surg*  
16       2010;2:133-141.  
17
- 18  5.   Greene FL, Page DL, Fleming ID, et al. *AJCC Cancer Staging Manual*, 6th edn.  
19       New York: Springer-Verlag, 2002.  
20
- 21  6.   Bithmuth H, CorletteMB. Intrahepatic cholangioenteric anastomosis in carcinoma  
22       of the hilus of the liver. *Surg Gynecol Obstet* 1975;140:170-177.  
23
- 24  7.   Govil A, Calkins H, Spragg DD, et al. Fusion of imaging technologies: how, when,  
25       and for whom? *J Interv Card Electrophysiol* 2011;32:195-203.  
26
- 27  8.   Tanaka H, Hayashi S, Ohtakara K, et al. Usefulness of CT-MRI fusion in  
28       radiotherapy planning for localized prostate cancer. *J Radiat Res*  
29       2011;52:782-788.  
30

## Figure legends


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17

**Fig. 1** Axial MDCT and MRCP. (A) Axial MDCT demonstrates bilateral intrahepatic bile duct dilatation. (B) MRCP demonstrates that the bilateral intrahepatic bile ducts were dilated, and that the confluence of the right and left hepatic ducts is tapered and obstructed at the hepatic hilum.


**Fig. 2** Preoperative fusion analysis using 3D-MDCT combined with 3D MRI imaging (A) and a virtual dissection line when performing left hepatic lobectomy (B). The red line indicates the planned dissection line. Novel modality was successfully described and the surgeon and surgical staff could determine the appropriate dissection line preoperatively.

**Fig. 3** The actual liver cut surface shows the stump of the left hepatic vein, the middle hepatic vein, and an orifice of the right hepatic duct. This ductal margin was confirmed as histologically negative. The relationship between the right hepatic duct and right portal vein was the same as anticipated preoperatively.






A




B

Figure 1.



- : portal vein
- : bile duct
- : hepatic vein
- : tumor

A



B

Figure 2.




Figure 3.