

Vertical distribution and temporal changes of ¹³⁷Cs in soil profiles under various land uses after the Fukushima Dai-ichi Nuclear Power Plant accident

著者別名	高橋 純子, 田村 憲司, 恩田 裕一
journal or publication title	Journal of environmental radioactivity
volume	139
page range	351-361
year	2015-01
権利	(C) 2014 Elsevier Ltd. NOTICE: this is the author's version of a work that was accepted for publication in Journal of environmental radioactivity. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Journal of environmental radioactivity, 139, 2015. DOI:10.1016/j.jenvrad.2014.07.004
URL	http://hdl.handle.net/2241/00124265

doi: 10.1016/j.jenvrad.2014.07.004

Total (without litter)	6192	3901	4596	3718
---------------------------	------	------	------	------

- a: weight per unit volume (225 or 450 cm³) of soil samples collected using scraper plate after oven heating.
- b: calculated from depth (cm) and bulk density (g cm⁻³).
- c: one sample in U8 container was measured for each layer.
- d: standard deviation from counting statistics of sample measurement.
- e: calculated from mass per unit area and concentration for each layer.