

The fixed point set of a holomorphic isometry,
the intersection of two real forms in a
Hermitian symmetric space of compact type and
symmetric triads

著者別名	田崎 博之
journal or publication title	International journal of mathematics
volume	26
number	6
page range	1541005
year	2015-06
権利	(C) World Scientific Publishing Company. Electronic version of an article published as International Journal of Mathematics, 26, 6, 2015, 1541005. DOI: 10.1142/S0129167X15410050, http://www.worldscientific.com/worldscinet/ijm
URL	http://hdl.handle.net/2241/00125752

doi: 10.1142/S0129167X15410050

THE FIXED POINT SET OF A HOLOMORPHIC ISOMETRY, THE INTERSECTION OF TWO REAL FORMS IN A HERMITIAN SYMMETRIC SPACE OF COMPACT TYPE AND SYMMETRIC TRIADS

OSAMU IKAWA, MAKIKO SUMI TANAKA, AND HIROYUKI TASAKI

ABSTRACT. We show a necessary and sufficient condition that the fixed point set of a holomorphic isometry and the intersection of two real forms of a Hermitian symmetric space of compact type are discrete and prove that they are antipodal sets in the cases. We also consider some relations between the intersection of two real forms and the fixed point set of a certain holomorphic isometry.

1. INTRODUCTION

In [15], [17] and [18] the second and third authors showed that the intersection of two real forms in a Hermitian symmetric space of compact type is an antipodal set if the intersection is discrete. The notion of an antipodal set of a Riemannian symmetric space was introduced by Chen-Nagano [3]. We showed the main results of this paper in a special case in [8]. In this paper we show a necessary and sufficient condition that the fixed point set of a holomorphic isometry of a Hermitian symmetric space of compact type is discrete and prove that the discrete fixed point set is an antipodal set. We also show a necessary and sufficient condition that the intersection of two real forms in a Hermitian symmetric space of compact type is discrete and consider some relations between the intersection of two real forms and the fixed point set of a certain holomorphic isometry by the use of the symmetric triads introduced by the first author in [6].

We roughly explain how to use symmetric triads in order to obtain a necessary and sufficient condition that the intersection of two real forms is discrete. In an irreducible Hermitian symmetric space $M = G/K$ of

The first author was partly supported by the Grant-in-Aid for Science Research (C) 2013 (No. 25400070), Japan Society for the Promotion of Science.

The second author was partly supported by the Grant-in-Aid for Science Research (C) 2013 (No. 23540108), Japan Society for the Promotion of Science.

The third author was partly supported by the Grant-in-Aid for Science Research (C) 2013 (No. 24540064), Japan Society for the Promotion of Science.

compact type two real forms L_1 and L_2 are determined by two involutive anti-holomorphic isometries τ_1 and τ_2 . These involutive isometries τ_1 and τ_2 determine two symmetric pairs (G, F_1) and (G, F_2) . The triad (G, F_1, F_2) defines a symmetric triad, by which we can describe the intersection $L_1 \cap L_2$ and obtain a necessary and sufficient condition that $L_1 \cap L_2$ is discrete.

The organization of this paper is as follows. In Section 2 we briefly review some fundamental results on Hermitian symmetric spaces of compact type, their antipodal sets and real forms.

In Section 3 we describe the fixed point set of a holomorphic isometry of a Hermitian symmetric space of compact type and obtain a necessary and sufficient condition that the fixed point set is discrete. If a holomorphic isometry is contained in the identity component of the group of holomorphic isometries, we can describe its fixed point set by the root system of the Lie algebra of the group of holomorphic isometries. There are two sequences of irreducible Hermitian symmetric spaces of compact type whose groups of holomorphic isometries are not connected. In these cases we describe the fixed point set of a holomorphic isometry which is not contained in the identity component in another way and obtain a necessary and sufficient condition that the fixed point set is discrete. In the cases where the fixed point sets are discrete, we describe them as orbits of certain Weyl groups.

In Section 4 we first describe a great antipodal set of each irreducible Hermitian symmetric space M of compact type as an orbit of the Weyl group. We second investigate two real forms in M and their intersection from a viewpoint of symmetric triads.

In Section 5 we also investigate a relation between the intersection of two real forms in M and the fixed point set of a certain holomorphic isometry on M from a viewpoint of symmetric triads.

The authors would like to thank Katsuya Mashimo and Kurando Baba for useful conversations. They are also indebted to the referee, whose comments improved the manuscript.

2. HERMITIAN SYMMETRIC SPACES OF COMPACT TYPE

In this section we review some fundamental results on Hermitian symmetric spaces of compact type. We also review their antipodal sets and real forms which we need later.

We construct a Hermitian symmetric space of compact type as an adjoint orbit in a compact semisimple Lie algebra. Let G be a connected compact semisimple Lie group and \mathfrak{g} its Lie algebra, which is

a compact semisimple Lie algebra. We take an $\text{Ad}(G)$ -invariant inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{g} . We take a nonzero element $J \in \mathfrak{g}$ satisfying $(\text{ad}J)^3 = -\text{ad}J$. The adjoint orbit $M = \text{Ad}(G)J \subset \mathfrak{g}$ is a Hermitian symmetric space of compact type with respect to the induced metric from $\langle \cdot, \cdot \rangle$. Define a closed subgroup K of G by

$$K = \{k \in G \mid \text{Ad}(k)J = J\}.$$

Its Lie algebra \mathfrak{k} is given by

$$\mathfrak{k} = \{X \in \mathfrak{g} \mid [J, X] = 0\}.$$

The subspace

$$\mathfrak{m} = \{[J, X] \mid X \in \mathfrak{g}\}$$

is the orthogonal complement of \mathfrak{k} , thus we have an orthogonal direct sum decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{m}$. The automorphism $e^{\pi \text{ad}J}$ is involutive. The subalgebra \mathfrak{k} is the $(+1)$ -eigenspace and the subspace \mathfrak{m} is the (-1) -eigenspace of $e^{\pi \text{ad}J}$. The operator $\text{ad}J$ defines an $\text{Ad}(K)$ -invariant complex structure on \mathfrak{m} which can be identified with the tangent space of M at J , hence it defines an $\text{Ad}(G)$ -invariant complex structure on M . It is known that any Hermitian symmetric space of compact type is constructed in this manner.

In a Riemannian symmetric space M we denote by s_x the geodesic symmetry at $x \in M$. A subset S of M is an *antipodal set* if $s_x(y) = y$ for any $x, y \in S$. The 2-number $\#_2 M$ of M is the maximum of the cardinality of antipodal sets of M . We call an antipodal set of M *great* if it attains $\#_2 M$. These were introduced by Chen-Nagano [3].

A great antipodal set of a Hermitian symmetric space of compact type is described in the following way.

Theorem 2.1 ([16]). *Let $M = \text{Ad}(G)J \subset \mathfrak{g}$ be a Hermitian symmetric space of compact type. A great antipodal set of M is represented as $M \cap \mathfrak{t}$ for a maximal abelian subalgebra \mathfrak{t} of \mathfrak{g} containing J . In particular, a great antipodal set of M is an orbit of the Weyl group of \mathfrak{g} with respect to \mathfrak{t} .*

After [16] was published we knew the following earlier results. Bott [1] showed that $M \cap \mathfrak{t}$ is an orbit of the Weyl group of \mathfrak{g} and Takeuchi [13] showed that the Weyl group acts transitively on the great antipodal set of M .

Take a maximal abelian subalgebra \mathfrak{t} of \mathfrak{k} . Since the involution $e^{\pi \text{ad}J}$ is of inner type, \mathfrak{t} is also a maximal abelian subalgebra of \mathfrak{g} . Since J commutes any element of \mathfrak{k} , the maximality of \mathfrak{t} implies that J is in \mathfrak{t} . We will use the following lemma, which was suggested by K. Mashimo, in Subsection 4.1.

Lemma 2.2. *Denote by $W(\mathfrak{g})$ and $W(\mathfrak{k})$ the Weyl groups of \mathfrak{g} and \mathfrak{k} with respect to \mathfrak{t} respectively. Then $W(\mathfrak{g})J = W(\mathfrak{g})/W(\mathfrak{k})$. In particular, $\#(W(\mathfrak{g})J) = \#(W(\mathfrak{g}))/\#(W(\mathfrak{k}))$.*

Proof. The isotropy subgroup of $W(\mathfrak{g})$ at J is equal to

$$\{s \in W(\mathfrak{g}) \mid sJ = J\} = W(\mathfrak{g}) \cap \text{Ad}_G(K) = W(\mathfrak{k}).$$

□

By definition, a *real form* is a fixed point set of an involutive anti-holomorphic isometry of M . A real form of M is a connected totally geodesic Lagrangian submanifold of M . Leung [9] and Takeuchi [12] classified real forms L of irreducible Hermitian symmetric spaces M of compact type. The list is as follows.

M	L
$G_k(\mathbb{C}^n)$	$G_k(\mathbb{R}^n)$
$G_{2k}(\mathbb{C}^{2n})$	$G_k(\mathbb{H}^n)$
$G_n(\mathbb{C}^{2n})$	$U(n)$
$Q_n(\mathbb{C})$	$S^{k,n-k}$
$SO(4n)/U(2n)$	$U(2n)/Sp(n)$
$SO(2n)/U(n)$	$SO(n)$
$Sp(2n)/U(2n)$	$Sp(n)$
$Sp(n)/U(n)$	$U(n)/O(n)$
$E_6/S^1 \cdot Spin(10)$	$G_2(\mathbb{H}^4)/\mathbb{Z}_2$
$E_6/S^1 \cdot Spin(10)$	$P_2(\text{Cay}) = F_4/Spin(9)$
$E_7/S^1 \cdot E_6$	$(SU(8)/Sp(4))/\mathbb{Z}_2$
$E_7/S^1 \cdot E_6$	$S^1 \cdot E_6/F_4$

In this list above we denote by $G_k(\mathbb{K}^n)$ the Grassmann manifold consisting of \mathbb{K} -subspaces of \mathbb{K} -dimension k in \mathbb{K}^n for $\mathbb{K} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ and by $Q_n(\mathbb{C})$ the complex hyperquadric of complex dimension n in the complex projective space $\mathbb{C}P^{n+1}$, which is holomorphically isometric to the real oriented Grassmann manifold $\tilde{G}_2(\mathbb{R}^{n+2})$. We regard $\tilde{G}_2(\mathbb{R}^{n+2})$ as a submanifold in $\bigwedge^2 \mathbb{R}^{n+2}$ in a natural way and define a real form $S^{p,q}$ of $\tilde{G}_2(\mathbb{R}^{n+2})$ for p, q with $p + q = n$ by

$$S^{p,q} = S^p(\mathbb{R}e_1 \oplus \cdots \oplus \mathbb{R}e_{p+1}) \wedge S^q(\mathbb{R}e_{p+2} \oplus \cdots \oplus \mathbb{R}e_{n+2}),$$

where e_1, \dots, e_{n+2} is the standard orthonormal basis of \mathbb{R}^{n+2} . The real form $S^{p,q}$ is diffeomorphic to $(S^p \times S^q)/\mathbb{Z}_2$.

3. THE FIXED POINT SET OF A HOLOMORPHIC ISOMETRY

In this section we show a necessary and sufficient condition that the fixed point set of a holomorphic isometry of a Hermitian symmetric space of compact type is discrete and prove that the discrete fixed point set is an antipodal set. We use the notation described in the last section.

For a set X and a map $\phi : X \rightarrow X$ we denote

$$F(\phi, X) = \{x \in X \mid \phi(x) = x\}.$$

We use this notation throughout the paper. For any element g in a connected compact semisimple Lie group G with Lie algebra \mathfrak{g} we have $\dim F(\text{Ad}(g), \mathfrak{g}) \geq \text{rank}(G)$, because there exists a maximal torus of G containing g . If $\dim F(\text{Ad}(g), \mathfrak{g}) = \text{rank}(G)$, we call g a *regular element* of G . We can see that the set of all regular elements of G is open and dense in G .

We denote by $A(M)$ the group of all holomorphic isometries of a Hermitian symmetric space M of compact type and by $A_0(M)$ its identity component. If M is equal to $\text{Ad}(G)J \subset \mathfrak{g}$ for a connected compact semisimple Lie group G with Lie algebra \mathfrak{g} , the identity component $A_0(M)$ coincides with $\{\text{Ad}(g)|_M \mid g \in G\}$. Without loss of generality we can suppose that the action of each simple factor of G on M is not trivial.

Theorem 3.1. *Let $M = \text{Ad}(G)J \subset \mathfrak{g}$ be a Hermitian symmetric space of compact type. The fixed point set $F(\text{Ad}(g), M)$ is discrete if and only if g is a regular element of G . In the case $F(\text{Ad}(g), M)$ is a great antipodal set of M .*

Proof. We take a maximal abelian subalgebra \mathfrak{t} of \mathfrak{g} containing J . Then we have $\mathfrak{t} \subset \mathfrak{k}$ by the definition of \mathfrak{k} . We denote by $\mathfrak{g}^{\mathbb{C}}$ the complexification of \mathfrak{g} . For $\alpha \in \mathfrak{t}$ we define the *root space*

$$\mathfrak{g}_{\alpha} = \{X \in \mathfrak{g}^{\mathbb{C}} \mid [H, X] = \sqrt{-1}\langle \alpha, H \rangle X \ (H \in \mathfrak{t})\}$$

and the *root system* $\Delta = \{\alpha \in \mathfrak{t} - \{0\} \mid \mathfrak{g}_{\alpha} \neq \{0\}\}$. Then we have the *root space decomposition*

$$\mathfrak{g}^{\mathbb{C}} = \mathfrak{t}^{\mathbb{C}} + \sum_{\alpha \in \Delta} \mathfrak{g}_{\alpha}.$$

We define a lexicographic order on \mathfrak{t} and write $\Delta_+ = \{\alpha \in \Delta \mid \alpha > 0\}$. We obtain

$$\mathfrak{g} = \mathfrak{t} + \sum_{\alpha \in \Delta_+} \mathfrak{g} \cap (\mathfrak{g}_{\alpha} + \mathfrak{g}_{-\alpha}).$$

We write $T = \exp \mathfrak{t}$, which is a maximal torus of G , hence there exists $g_1 \in G$ such that $a = g_1 g g_1^{-1} \in T$. Since

$$\mathrm{Ad}(g_1)F(\mathrm{Ad}(g), M) = F(\mathrm{Ad}(a), M),$$

we consider the condition that $F(\mathrm{Ad}(a), M)$ is discrete for $a \in T$. From the definition of \mathfrak{g}_α we obtain the following lemma.

Lemma 3.2. *For $a = \exp H \in T$ with $H \in \mathfrak{t}$ we have*

$$F(\mathrm{Ad}(a), \mathfrak{g}) = \mathfrak{t} + \sum_{\substack{\alpha \in \Delta, \\ \langle \alpha, H \rangle \in 2\pi\mathbb{Z}}} \mathfrak{g} \cap (\mathfrak{g}_\alpha + \mathfrak{g}_{-\alpha}).$$

Proof. By Lemma 3.1 of Chapter VI in Helgason [5] we can see that for $\alpha \in \Delta$ there exists a basis F_α, G_α of $\mathfrak{g} \cap (\mathfrak{g}_\alpha + \mathfrak{g}_{-\alpha})$ which satisfies

$$[H, F_\alpha] = \langle \alpha, H \rangle G_\alpha, \quad [H, G_\alpha] = -\langle \alpha, H \rangle F_\alpha.$$

These imply

$$\begin{aligned} \mathrm{Ad}(\exp H)F_\alpha &= \cos\langle \alpha, H \rangle F_\alpha + \sin\langle \alpha, H \rangle G_\alpha, \\ \mathrm{Ad}(\exp H)G_\alpha &= -\sin\langle \alpha, H \rangle F_\alpha + \cos\langle \alpha, H \rangle G_\alpha. \end{aligned}$$

Therefore we obtain

$$F(\mathrm{Ad}(a), \mathfrak{g}) = \mathfrak{t} + \sum_{\substack{\alpha \in \Delta, \\ \langle \alpha, H \rangle \in 2\pi\mathbb{Z}}} \mathfrak{g} \cap (\mathfrak{g}_\alpha + \mathfrak{g}_{-\alpha}).$$

□

Corollary 3.3. *For $H \in \mathfrak{t}$ the following three conditions are equivalent.*

- (1) *The element $\exp H$ is regular.*
- (2) *$F(\mathrm{Ad}(\exp H), \mathfrak{g}) = \mathfrak{t}$.*
- (3) *$\langle \alpha, H \rangle \notin 2\pi\mathbb{Z}$, for any $\alpha \in \Delta$.*

Using these preliminaries we prove the theorem. We consider the case where g is a regular element of G . In this case $a = g_1 g g_1^{-1} \in T$ is also regular. By Corollary 3.3 we have $F(\mathrm{Ad}(a), \mathfrak{g}) = \mathfrak{t}$ and hence

$$F(\mathrm{Ad}(a), M) = M \cap \mathfrak{t},$$

which is discrete and a great antipodal set of M .

Next we consider the case where g is not a regular element. In this case $a = g_1 g g_1^{-1} \in T$ is not regular. We write $a = \exp H$ for $H \in \mathfrak{t}$. By Corollary 3.3 there exists $\alpha \in \Delta$ which satisfies $\langle \alpha, H \rangle \in 2\pi\mathbb{Z}$.

Let \mathfrak{g}_i ($1 \leq i \leq n$) be simple ideals of \mathfrak{g} and $\mathfrak{g} = \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_n$ is a direct sum decomposition of \mathfrak{g} . There exists i such that $\alpha \in \mathfrak{t}_i = \mathfrak{t} \cap \mathfrak{g}_i$. We denote by W_i the Weyl group of \mathfrak{g}_i with respect to \mathfrak{t}_i and by J_i the \mathfrak{g}_i -component of J . Since the action of the Lie subgroup corresponding to \mathfrak{g}_i is not trivial, we have $J_i \neq 0$. The Weyl group W_i acts transitively

on the set of all long roots of \mathfrak{g}_i and the set of all short roots, thus we have

$$\text{span}_{\mathbb{R}}\{w\alpha \mid w \in W_i\} = \mathfrak{t}_i.$$

Hence there exists $w \in W_i$ which satisfies $\langle \alpha, wJ_i \rangle \neq 0$. We can replace J with wJ . Then we have $\langle \alpha, J \rangle \neq 0$ and the decomposition

$$J = \frac{\langle \alpha, J \rangle}{\langle \alpha, \alpha \rangle} \alpha + \left(J - \frac{\langle \alpha, J \rangle}{\langle \alpha, \alpha \rangle} \alpha \right)$$

We take F_α in the proof of Lemma 3.2 and consider

$$\begin{aligned} \text{Ad}(\exp tF_\alpha)J &= \text{Ad}(\exp tF_\alpha) \frac{\langle \alpha, J \rangle}{\langle \alpha, \alpha \rangle} \alpha + \text{Ad}(\exp tF_\alpha) \left(J - \frac{\langle \alpha, J \rangle}{\langle \alpha, \alpha \rangle} \alpha \right) \\ &= \text{Ad}(\exp tF_\alpha) \frac{\langle \alpha, J \rangle}{\langle \alpha, \alpha \rangle} \alpha + \left(J - \frac{\langle \alpha, J \rangle}{\langle \alpha, \alpha \rangle} \alpha \right). \end{aligned}$$

Since $[F_\alpha, G_\alpha] = \|F_\alpha\|^2 \alpha = \|G_\alpha\|^2 \alpha$, the first term is

$$\text{Ad}(\exp tF_\alpha) \frac{\langle \alpha, J \rangle}{\langle \alpha, \alpha \rangle} \alpha = \cos(\|F_\alpha\| \|\alpha\| t) \frac{\langle \alpha, J \rangle}{\langle \alpha, \alpha \rangle} \alpha - \sin(\|F_\alpha\| \|\alpha\| t) \frac{\langle \alpha, J \rangle}{\|F_\alpha\| \|\alpha\|} G_\alpha$$

and it is a circle in $\text{span}_{\mathbb{R}}\{\alpha, G_\alpha\} \subset \mathfrak{t} + \mathfrak{g} \cap (\mathfrak{g}_\alpha + \mathfrak{g}_{-\alpha})$. The second term is contained in \mathfrak{t} , hence we have $\text{Ad}(\exp tF_\alpha)J \in F(\text{Ad}(a), M)$, which means that $F(\text{Ad}(a), M)$ is not discrete. Therefore $F(\text{Ad}(g), M)$ is not discrete. This completes the proof of the theorem. \square

Remark 3.4. We note that $\langle \alpha, J \rangle = 0, \pm 1$ for any $\alpha \in \Delta$ since $(\text{ad}J)^3 = -\text{ad}J$. This fact will be used in Subsection 4.1.

We consider the fixed point set of an element of $A(M) - A_0(M)$. We recall the results on $A(M)/A_0(M)$ obtained by Takeuchi [11].

Lemma 3.5 ([11]). *Let M be an irreducible Hermitian symmetric space of compact type. Then $A(M)/A_0(M)$ are as follows.*

- (A) *If $M = Q_{2m}(\mathbb{C}) (m \geq 2)$ or $M = G_m(\mathbb{C}^{2m}) (m \geq 2)$, then $A(M)/A_0(M) \cong \mathbb{Z}_2$.*
- (B) *Otherwise, $A(M) = A_0(M)$.*

In the case where M is irreducible, it is sufficient to consider the cases where $M = Q_{2m}(\mathbb{C}), G_m(\mathbb{C}^{2m}) (m \geq 2)$.

In the case where $M = Q_{2m}(\mathbb{C}) (m \geq 2)$, we can suppose that $G = SO(2m+2)$ and regard $M = SO(2m+2)/(SO(2) \times SO(2m))$ as a submanifold in $\bigwedge^2 \mathbb{R}^{2m+2}$ in a natural way. We take

$$\phi = \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & & & -1 \end{bmatrix} \in O(2m+2).$$

Then we have $A(M) = A_0(M) \cup \text{Ad}(\phi)A_0(M)$ as we showed in the proof of Proposition 2.2 in [17]. Hence

$$A(M) - A_0(M) = \text{Ad}(\phi)A_0(M) = \text{Ad}(\{g \in O(2m+2) \mid \det g = -1\}).$$

For any $g \in O(2m+2)$ there exists $g_1 \in O(2m+2)$ which satisfies

$$(3.1) \quad g_1 g g_1^{-1} = \begin{bmatrix} R(\theta_1) & & & & & \\ & \ddots & & & & \\ & & R(\theta_m) & & & \\ & & & 1 & & \\ & & & & & -1 \end{bmatrix},$$

where

$$R(\theta_i) = \begin{bmatrix} \cos \theta_i & -\sin \theta_i \\ \sin \theta_i & \cos \theta_i \end{bmatrix} \quad (1 \leq i \leq m).$$

Let e_i be the i -th column vector of g_1^{-1} . Then e_1, \dots, e_{2m+2} is an orthonormal basis of \mathbb{R}^{2m+2} and

$$(3.2) \quad \begin{aligned} g[e_{2i-1} \ e_{2i}] &= [e_{2i-1} \ e_{2i}]R(\theta_i) \quad (1 \leq i \leq m) \\ g[e_{2m+1} \ e_{2m+2}] &= [e_{2m+1} \ e_{2m+2}] \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \end{aligned}$$

Using these we can obtain the following theorem on the fixed point set of an element of $A(M) - A_0(M)$.

Theorem 3.6. *Let $M = Q_{2m}(\mathbb{C})(m \geq 2)$. Any element of $A(M) - A_0(M)$ is equal to $\text{Ad}(g)$ for $g \in O(2m+2)$ with $\det g = -1$. $F(\text{Ad}(g), M)$ is discrete if and only if there exists $g_1 \in O(2m+2)$ such that (3.1) holds and that $R(\theta_1), \dots, R(\theta_m)$ are different from each other. In the case*

$$F(\text{Ad}(g), M) = W(SO(2m+1))J,$$

where $SO(2m+1)$ is the stabilizer of e_{2m+2} . $F(\text{Ad}(g), M)$ is an antipodal set of M and

$$\#F(\text{Ad}(g), M) = 2m < 2m+2 = \#_2 M.$$

Proof. By (3.2) we have

$$\{\pm e_1 \wedge e_2, \pm e_3 \wedge e_4, \dots, \pm e_{2m-1} \wedge e_{2m}\} \subset F(\text{Ad}(g), M).$$

We suppose that there exist i, j ($i < j$) satisfying $R(\theta_i) = R(\theta_j)$. For $\xi \in \mathbb{R}$ we write

$$C(\xi) = \begin{bmatrix} \cos \xi & 0 \\ 0 & \cos \xi \end{bmatrix}, \quad S(\xi) = \begin{bmatrix} \sin \xi & 0 \\ 0 & \sin \xi \end{bmatrix}.$$

Then we have

$$g[e_{2i-1} \ e_{2i} \ e_{2j-1} \ e_{2j}] \begin{bmatrix} C(\xi) \\ S(\xi) \end{bmatrix} = [e_{2i-1} \ e_{2i} \ e_{2j-1} \ e_{2j}] \begin{bmatrix} C(\xi) \\ S(\xi) \end{bmatrix} R(\theta_i),$$

hence g acts on the 2-dimensional subspace spanned by

$$[e_{2i-1} \ e_{2i} \ e_{2j-1} \ e_{2j}] \begin{bmatrix} C(\xi) \\ S(\xi) \end{bmatrix}$$

as the rotation with angle θ_i . Therefore

$$(\cos \xi e_{2i-1} + \sin \xi e_{2j-1}) \wedge (\cos \xi e_{2i} + \sin \xi e_{2j}) \in F(\text{Ad}(g), M)$$

for any $\xi \in \mathbb{R}$ and $F(\text{Ad}(g), M)$ is not discrete.

Next we suppose that $R(\theta_i)$ and $R(\theta_j)$ are different for any different i, j . Let V_i be the 2-dimensional subspace spanned by e_{2i-1} and e_{2i} . Then we have an orthogonal direct sum decomposition

$$\mathbb{R}^{2m+2} = V_1 \oplus \cdots \oplus V_{m+1}.$$

We take $u \wedge v \in F(\text{Ad}(g), M)$ where u and v are orthonormal. We decompose u and v as follows.

$$\begin{aligned} u &= u_1 + \cdots + u_{m+1} & (u_i \in V_i, \ 1 \leq i \leq m+1) \\ v &= v_1 + \cdots + v_{m+1} & (v_i \in V_i, \ 1 \leq i \leq m+1). \end{aligned}$$

The element g acts on these as follows.

$$\begin{aligned} gu &= R(\theta_1)u_1 + \cdots + R(\theta_m)u_m + gu_{m+1} \\ gv &= R(\theta_1)v_1 + \cdots + R(\theta_m)v_m + gv_{m+1}. \end{aligned}$$

On the other hand $u \wedge v \in F(\text{Ad}(g), M)$, hence g acts on the 2-dimensional subspace spanned by u and v as a rotation. There exists $\xi \in \mathbb{R}$ which satisfies

$$g[u \ v] = [u \ v] \begin{bmatrix} \cos \xi & -\sin \xi \\ \sin \xi & \cos \xi \end{bmatrix}$$

and we obtain

$$\begin{aligned} gu &= \cos \xi u + \sin \xi v \\ gv &= -\sin \xi u + \cos \xi v. \end{aligned}$$

Thus we get

$$\begin{aligned} R(\theta_i)u_i &= \cos \xi u_i + \sin \xi v_i \quad (1 \leq i \leq m), \\ gu_{m+1} &= \cos \xi u_{m+1} + \sin \xi v_{m+1}. \end{aligned}$$

The action of g on V_{m+1} is not a rotation, we have $u_{m+1} = 0$. Since $R(\theta_i) \neq R(\theta_j)$ for different i, j , there exists k such that

$$u_k \neq 0, \quad u_i = 0 \quad (i \neq k)$$

and we have $u \wedge v = \pm e_{2k-1} \wedge e_{2k}$. Therefore

$$F(\text{Ad}(g), M) = \{\pm e_1 \wedge e_2, \pm e_3 \wedge e_4, \dots, \pm e_{2m-1} \wedge e_{2m}\}$$

and $F(\text{Ad}(g), M)$ is discrete. The above description of $F(\text{Ad}(g), M)$ shows that it is an orbit of $W(SO(2m+1))$ through $J = e_1 \wedge e_2$ and an antipodal set of M . \square

In the case where $M = G_m(\mathbb{C}^{2m})(m \geq 2)$, we can suppose that $G = SU(2m)$. We take

$$J_m = \begin{bmatrix} & 1_m \\ -1_m & \end{bmatrix}$$

in G , where 1_m denotes the $m \times m$ identity matrix. We regard M as the submanifold $\text{Ad}(G)J$ in $\mathfrak{g} = \mathfrak{su}(2m)$, where

$$J = \frac{\sqrt{-1}}{2} \begin{bmatrix} 1_m & \\ & -1_m \end{bmatrix} \in \mathfrak{g}.$$

We define an involutive automorphism ϕ of G by $\phi(g) = J_m \bar{g} J_m^{-1}$ for $g \in G$. The fixed point set $F(\phi, G)$ is equal to $Sp(m)$ and ϕ defines a symmetric pair $(SU(2m), Sp(m))$. The differential map $d\phi$ of ϕ is represented by $d\phi(X) = J_m \bar{X} J_m^{-1}$ for $X \in \mathfrak{g}$. So we simply write $\phi(X) = J_m \bar{X} J_m^{-1}$ for $X \in \mathfrak{g}$. The automorphism ϕ of G also induces a holomorphic isometry of M defined by

$$\text{Ad}(g)J \mapsto \text{Ad}(\phi(g))J \quad (g \in G).$$

Since $\phi(J) = J$, we have

$$\text{Ad}(\phi(g))J = \text{Ad}(\phi(g))\phi(J) = \phi(\text{Ad}(g)J).$$

Thus the holomorphic isometry of M induced by ϕ is the restriction of $\phi : \mathfrak{g} \rightarrow \mathfrak{g}$ to $M \subset \mathfrak{g}$. So we also denote it by ϕ .

The holomorphic isometry ϕ is contained in $A(M) - A_0(M)$, which is showed in [17]. We take a maximal torus T of $Sp(m)$.

Lemma 3.7. *For any $h \in A(M) - A_0(M)$ there exist $t \in T$ and $g \in G$ such that $h = \text{Ad}(g)\text{Ad}(t)\text{Ad}(\phi(g^{-1}))\phi$ and*

$$F(h, M) = \text{Ad}(g)F(\text{Ad}(t)\phi, M).$$

Proof. We define two involutive automorphisms θ_1, θ_2 of $G \times G$ by

$$\theta_1(g, h) = (h, g), \quad \theta_2(g, h) = (\phi^{-1}(h), \phi(g)) \quad ((g, h) \in G \times G).$$

For a general automorphism ϕ of G , the automorphism θ_2 is involutive. Since ϕ is involutive, θ_1 and θ_2 are commutative. The fixed point sets

of them are

$$\begin{aligned} K_1 &= F(\theta_1, G \times G) = \{(g, g) \mid g \in G\}, \\ K_2 &= F(\theta_2, G \times G) = \{(g, \phi(g)) \mid g \in G\} \end{aligned}$$

and we obtain two direct sum decompositions of $\mathfrak{g} \times \mathfrak{g}$ as follows

$$\mathfrak{g} \times \mathfrak{g} = \mathfrak{k}_1 \oplus \mathfrak{m}_1 = \mathfrak{k}_2 \oplus \mathfrak{m}_2,$$

where

$$\begin{aligned} \mathfrak{k}_1 &= \{(X, X) \mid X \in \mathfrak{g}\}, & \mathfrak{m}_1 &= \{(X, -X) \mid X \in \mathfrak{g}\}, \\ \mathfrak{k}_2 &= \{(X, \phi(X)) \mid X \in \mathfrak{g}\}, & \mathfrak{m}_2 &= \{(X, -\phi(X)) \mid X \in \mathfrak{g}\}. \end{aligned}$$

These imply

$$\begin{aligned} \mathfrak{k}_1 \cap \mathfrak{k}_2 &= \{(X, X) \mid X \in \mathfrak{sp}(m)\}, \\ \mathfrak{m}_1 \cap \mathfrak{m}_2 &= \{(X, -X) \mid X \in \mathfrak{sp}(m)\}. \end{aligned}$$

We denote by \mathfrak{t} the Lie algebra of T . The subspace

$$\mathfrak{a} = \{(H, -H) \mid H \in \mathfrak{t}\}$$

is a maximal abelian subspace of $\mathfrak{m}_1 \cap \mathfrak{m}_2$. We identify $(g, \phi(g)) \in K_2$ with $g \in G$ and $(g_1, g_2)K_1 \in (G \times G)/K_1$ with $g_1g_2^{-1} \in G$. Then the action of K_2 on $(G \times G)/K_1$ is equivalent with the action of G on G defined by

$$g \cdot x = gx\phi(g^{-1}) \quad (g, x \in G).$$

This action is a Hermann action and θ_1, θ_2 are commutative. In particular the action is a hyperpolar action with section T by [4]. So we have

$$G = \bigcup_{g \in G} gT\phi(g^{-1}).$$

Since $A(M) = A_0(M) \cup A_0(M)\phi$, for any $h \in A(M) - A_0(M)$ there exist $g \in G$ and $t \in T$ such that

$$h = \text{Ad}(g)\text{Ad}(t)\text{Ad}(\phi(g^{-1}))\phi.$$

Using this we get

$$F(h, M) = \{X \in M \mid \text{Ad}(g)\text{Ad}(t)\text{Ad}(\phi(g^{-1}))\phi(X) = X\}.$$

We put $X = \text{Ad}(g)Y$. Since $\phi(\text{Ad}(g)Y) = \text{Ad}(\phi(g))\phi Y$, we obtain

$$F(h, M) = \text{Ad}(g)F(\text{Ad}(t)\phi, M).$$

□

for any $H \in \mathfrak{t}$. We second decompose \mathfrak{m} into a direct sum of weight spaces with respect to \mathfrak{t} . We define

$$X_{ij}^- = \begin{bmatrix} E_{ij} - E_{ji} & \\ & -(E_{ij} - E_{ji}) \end{bmatrix}, Y_{ij}^- = \sqrt{-1} \begin{bmatrix} E_{ij} + E_{ji} & \\ & E_{ij} + E_{ji} \end{bmatrix},$$

$$X_{ij}^+ = \begin{bmatrix} & E_{ij} - E_{ji} \\ E_{ij} - E_{ji} & \end{bmatrix}, Y_{ij}^+ = \sqrt{-1} \begin{bmatrix} & E_{ij} - E_{ji} \\ -(E_{ij} - E_{ji}) & \end{bmatrix}$$

for $1 \leq i < j \leq m$ and

$$V(\mathfrak{m}) = \{X \in \mathfrak{m} \mid [X, \mathfrak{t}] = \{0\}\}.$$

We get

$$\mathfrak{m} = V(\mathfrak{m}) \oplus \sum_{i < j} (\mathbb{R}X_{ij}^- \oplus \mathbb{R}Y_{ij}^-) \oplus \sum_{i < j} (\mathbb{R}X_{ij}^+ \oplus \mathbb{R}Y_{ij}^+)$$

and

$$[H, X_{ij}^\pm] = \langle e_i \pm e_j, H \rangle Y_{ij}^\pm, \quad [H, Y_{ij}^\pm] = -\langle e_i \pm e_j, H \rangle X_{ij}^\pm$$

for any $H \in \mathfrak{t}$. Using the above decompositions we obtain

$$(3.3) \quad F(\text{Ad}(\exp H) \circ \phi, \mathfrak{g})$$

$$= \mathfrak{t} \oplus \sum_{\langle e_i + e_j, H \rangle \in 2\pi\mathbb{Z}} (\mathbb{R}F_{ij}^+ \oplus \mathbb{R}G_{ij}^+) \oplus \sum_{\langle e_i - e_j, H \rangle \in 2\pi\mathbb{Z}} (\mathbb{R}F_{ij}^- \oplus \mathbb{R}G_{ij}^-)$$

$$\oplus \sum_{\langle e_i + e_j, H \rangle \in \pi + 2\pi\mathbb{Z}} (\mathbb{R}X_{ij}^+ \oplus \mathbb{R}Y_{ij}^+) \oplus \sum_{\langle e_i - e_j, H \rangle \in \pi + 2\pi\mathbb{Z}} (\mathbb{R}X_{ij}^- \oplus \mathbb{R}Y_{ij}^-)$$

and the following theorem.

Theorem 3.8. *Let $M = G_m(\mathbb{C}^{2m})(m \geq 2)$. Any element h of $A(M) - A_0(M)$ is equal to $\text{Ad}(g)\text{Ad}(\exp H)\text{Ad}(\phi(g^{-1}))\phi$ for $g \in SU(2m)$, $H \in \mathfrak{t}$. Its fixed point set $F(h, M)$ is discrete if and only if $\langle e_i \pm e_j, H \rangle \notin \pi\mathbb{Z}$ for any $i \neq j$ and $\langle e_i, H \rangle \notin \pi\mathbb{Z}$ for any i . In the case*

$$F(\text{Ad}(\exp H) \circ \phi, M) = W(\text{Sp}(m))J,$$

which is an antipodal set of M and

$$\#F(h, M) = 2^m < \binom{2m}{m} = \#_2 M.$$

Proof. Lemma 3.7 implies the description of h and it is sufficient to consider $F(\text{Ad}(\exp H)\phi, M)$. According to (3.3), $F(\text{Ad}(\exp H)\phi, \mathfrak{g}) = \mathfrak{t}$ if and only if $\langle e_i \pm e_j, H \rangle \notin \pi\mathbb{Z}$ for any $i \neq j$ and $\langle e_i, H \rangle \notin \pi\mathbb{Z}$ for any i . In this case we have

$$F(\text{Ad}(\exp H) \circ \phi, M) = M \cap \mathfrak{t} = W(\text{Sp}(m))J$$

and $\#F(h, M) = 2^m$.

In order to prove the theorem we have to show that $F(\text{Ad}(\exp H)\phi, M)$ is not discrete if $F(\text{Ad}(\exp H)\phi, \mathfrak{g}) \neq \mathfrak{t}$. We first consider the case where there exist i, j such that $\langle e_i + e_j, H \rangle \in 2\pi\mathbb{Z}$. Since $W(\text{Sp}(m))J$ spans \mathfrak{t} , there exists $X \in W(\text{Sp}(m))J$ satisfying $\langle e_i + e_j, X \rangle \neq 0$. Using $[F_{ij}^+, G_{ij}^+] = |F_{ij}^+|^2(e_i + e_j)$ obtained from their definitions, we get

$$\begin{aligned} \text{Ad}(\exp tF_{ij}^+)X &= \left(X - \frac{\langle e_i + e_j, X \rangle}{\|e_i + e_j\|^2}(e_i + e_j) \right) \\ &+ \frac{\langle e_i + e_j, X \rangle}{\|e_i + e_j\|} \left(\cos(\|F_{ij}^+\| \|e_i + e_j\| t) \frac{e_i + e_j}{\|e_i + e_j\|} - \sin(\|F_{ij}^+\| \|e_i + e_j\| t) \frac{G_{ij}^+}{\|G_{ij}^+\|} \right) \end{aligned}$$

for $t \in \mathbb{R}$. Therefore $\text{Ad}(\exp tF_{ij}^+)X \in F(\text{Ad}(\exp H) \circ \phi, M)$ for $t \in \mathbb{R}$ and $F(\text{Ad}(\exp H) \circ \phi, M)$ is not discrete. We second consider the case where there exist $i \neq j$ such that $\langle e_i - e_j, H \rangle \in \pi + 2\pi\mathbb{Z}$. There exists $X \in W(\text{Sp}(m))J$ satisfying $\langle e_i - e_j, X \rangle \neq 0$. Using $[X_{ij}^-, Y_{ij}^-] = \|X_{ij}^-\|^2(e_i - e_j)$ obtained from their definitions, we get

$$\begin{aligned} \text{Ad}(\exp tX_{ij}^-)X &= \left(X - \frac{\langle e_i - e_j, X \rangle}{\|e_i - e_j\|^2}(e_i - e_j) \right) \\ &+ \frac{\langle e_i - e_j, X \rangle}{\|e_i - e_j\|} \left(\cos(\|X_{ij}^-\| \|e_i - e_j\| t) \frac{e_i - e_j}{\|e_i - e_j\|} - \sin(\|X_{ij}^-\| \|e_i - e_j\| t) \frac{Y_{ij}^-}{\|Y_{ij}^-\|} \right) \end{aligned}$$

for $t \in \mathbb{R}$. Therefore $\text{Ad}(\exp tX_{ij}^-)X \in F(\text{Ad}(\exp H) \circ \phi, M)$ for $t \in \mathbb{R}$ and $F(\text{Ad}(\exp H) \circ \phi, M)$ is not discrete. In the other cases we can see that $F(\text{Ad}(\exp H) \circ \phi, M)$ is not discrete in a similar way. This completes the proof of the theorem. \square

Remark 3.9. When a Hermitian symmetric space M of compact type is not irreducible, $F(h, M)$ for $h \in A(M)$ is obtained from the fixed point sets of holomorphic isometries of irreducible Hermitian symmetric spaces of compact type, which we have already known above. Let $M = M_1 \times \cdots \times M_k$ be the decomposition of M to the product of irreducible factors. We take $\phi \in A(M)$. In order to investigate $F(\phi, M)$ it is sufficient to consider the case where $M_1 = \cdots = M_k$ and

$$\phi(x_1, \dots, x_k) = (\phi_k(x_k), \phi_1(x_1), \dots, \phi_{k-1}(x_{k-1})) \quad (x_i \in M_i),$$

where $\phi_i : M_i \rightarrow M_{i+1}$ ($1 \leq i \leq k-1$) and $\phi_k : M_k \rightarrow M_1$ are holomorphically isometric maps. In this case we have

$$\begin{aligned} F(\phi, M) &= \{(x_1, \phi_1(x_1), \phi_2\phi_1(x_1), \dots, \phi_{k-1} \cdots \phi_1(x_1)) \mid x_1 \in F(\phi_k \cdots \phi_1, M_1)\}. \end{aligned}$$

Hence $F(\phi, M)$ is discrete if and only if $F(\phi_k \cdots \phi_1, M_1)$ is discrete. In the case $F(\phi, M)$ is an antipodal set of M and

$$\#F(\phi, M) = \#F(\phi_k \cdots \phi_1, M_1).$$

4. THE INTERSECTION OF TWO REAL FORMS

4.1. Characteristic elements associated with a root system. As we mentioned in Remark 3.4 the complex structure J of a Hermitian symmetric space of compact type satisfies $\langle \alpha, J \rangle = 0, \pm 1$ for any root α . Based on the fact we define a characteristic element of a root system as follows.

Let R be a root system of a finite dimensional vector space \mathfrak{a} with an inner product $\langle \cdot, \cdot \rangle$. Then $J \in \mathfrak{a} - \{0\}$ is a *characteristic element of the first kind* or simply a *characteristic element* associated with R if $\langle \lambda, J \rangle = 0, \pm 1$ for any $\lambda \in R$. We denote by $W(R)$ the Weyl group of R . If J is a characteristic element associated with R , then so are $-J$ and sJ for any $s \in W(R)$. In the sequel we assume that R is irreducible. For a characteristic element J we can take a fundamental system $\Pi = \{\alpha_1, \dots, \alpha_r\}$ of R such that $\langle \alpha_i, J \rangle = 0, 1$ for any α_i . Denote by $\delta = \sum m_i \alpha_i$ the highest root of R . When the type of R is one of E_8, F_4 and G_2 , there does not exist a characteristic element since $m_i \geq 2$ for any i . In order to describe $W(R)J$ in detail we give the definition of a two-point homogeneous space, which appears in the following proposition. For a group A which isometrically acts on a metric space (X, d) if for any two pairs $x, y \in X$ and $x', y' \in X$ satisfying $d(x, y) = d(x', y')$ there exists $a \in A$ such that $ax = x', ay = y'$, then we call X a *two-point homogeneous space* by the action of A . We prove the following proposition using the classification of an irreducible root system.

Proposition 4.1. *The orbit $W(R)J$ of a characteristic element J associated with an irreducible root system R is a two-point homogeneous space by the action of $W(R)$.*

Proof. Define a set $\{d_1, \dots, d_t\}$ ($0 < d_1 < \dots < d_t$) by

$$\{d_1, \dots, d_t\} = \{\|sJ - J\| \mid s \in W(R)\} - \{0\}.$$

The condition for $W(R)J$ to be two-point homogeneous is equivalent to the condition that the isotropy subgroup $\{s \in W(R) \mid sJ = J\}$ acts transitively on each $\{sJ \mid \|sJ - J\| = d_i, s \in W(R)\}$. Denote by $Ch(R)$ the set of all characteristic elements associated with R . We examine the condition above for each characteristic element J associated with each $R = A_r, B_r, C_r, BC_r, D_r, E_6$ and E_7 in Examples 4.2, 4.3, 4.4, 4.5,

4.6, 4.8, 4.10. If R is the root system of \mathfrak{g} in Section 2, according to Theorem 2.1 the orbit $W(R)J$ is a great antipodal set of the Hermitian symmetric space of compact type associated with J , so we also calculate its cardinality $\#(W(R)J)$. We follow the same notations of the set of positive roots in [2]. Denote by $\{e_1, \dots, e_r\}$ the standard orthonormal basis of \mathbb{R}^r .

Example 4.2. In the case where $R = B_r = \{\pm e_i \pm e_j, \pm e_i\}$, set $J = e_1$. Then we have

$$Ch(R) = W(R)J = \{\pm e_1, \dots, \pm e_r\}.$$

Thus $\#(W(R)J) = 2r$, $t = 2$, $d_1 = \sqrt{2}$ and $d_2 = 2$. We can verify that $W(R)J$ is two-point homogeneous.

Example 4.3. In the case where $R = C_r = \{\pm e_i \pm e_j, \pm 2e_i\}$, set

$$J = \frac{1}{2}(e_1 + e_2 + \dots + e_r).$$

Then we have

$$Ch(R) = W(R)J = \left\{ \frac{1}{2} \sum_{i=1}^r \epsilon_i e_i \mid \epsilon_i = \pm 1 \right\}.$$

Thus $\#(W(R)J) = 2^r$, $t = r$ and $d_i = \sqrt{i}$ ($1 \leq i \leq r$). We can verify that $W(R)J$ is two-point homogeneous.

Example 4.4. In the case where $R = BC_r = \{\pm e_i \pm e_j, \pm e_i, \pm 2e_i\}$, Examples 4.2 and 4.3 imply that there does not exist a characteristic element.

Example 4.5. In the case where $R = D_r = \{\pm e_i \pm e_j\}$, we define characteristic elements J_1 , J_2 and J_3 by

$$J_1 = e_1, \quad J_2 = \frac{1}{2} \left(\sum_{j=1}^{r-1} e_j - e_r \right), \quad J_3 = \frac{1}{2} \sum_{j=1}^r e_j.$$

Then

$$\begin{aligned} W(R)J_1 &= \{\pm e_1, \dots, \pm e_r\}, \\ W(R)J_2 &= \left\{ \frac{1}{2} \sum_{j=1}^r \epsilon_j e_j \mid \epsilon_j = \pm 1, \epsilon_1 \cdots \epsilon_r = -1 \right\}, \\ W(R)J_3 &= \left\{ \frac{1}{2} \sum_{j=1}^r \epsilon_j e_j \mid \epsilon_j = \pm 1, \epsilon_1 \cdots \epsilon_r = 1 \right\} \end{aligned}$$

and $Ch(R) = W(R)J_1 \cup W(R)J_2 \cup W(R)J_3$. Thus $\#(W(R)J_1) = 2r$, $\#(W(R)J_2) = \#(W(R)J_3) = 2^{r-1}$. We can verify that $W(R)J_i$ is two-point homogeneous.

Example 4.6. In the case where $R = A_r = \{\pm(e_i - e_j)\}$, we define characteristic elements J_1, \dots, J_r by

$$J_i = (e_1 + \dots + e_i) - \frac{i}{r+1} \sum_{j=1}^{r+1} e_j.$$

Then

$$W(R)J_i = \left\{ \sum_{j \in A} e_j - \frac{i}{r+1} \sum_{j=1}^{r+1} e_j \mid A \in P_i(r+1) \right\},$$

where we put $P_i(r+1) = \{A \subset \{1, 2, \dots, r+1\} \mid \#A = i\}$. Thus $Ch(R) = W(R)J_1 \cup \dots \cup W(R)J_r$ and $\#(W(R)J_i) = \binom{r+1}{i}$. We can verify that $W(R)J_i$ is two-point homogeneous.

Remark 4.7. In the case of A_2 , the orbit $W(R)J_1$ consists of three vertices of an equilateral triangle whose center of mass is the origin. Since $W(R)J_2 = -W(R)J_1$, the above statement holds for $W(R)J_2$. These will be used later.

Example 4.8. In the case where $R = E_6$, we define characteristic elements J_1 and J_2 by

$$J_1 = \frac{2}{3}(e_8 - e_7 - e_6) = \frac{1}{3}(4\alpha_1 + 3\alpha_2 + 5\alpha_3 + 6\alpha_4 + 4\alpha_5 + 2\alpha_6),$$

$$J_2 = \frac{1}{3}(e_8 - e_7 - e_6) + e_5 = \frac{1}{3}(2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 6\alpha_4 + 5\alpha_5 + 4\alpha_6).$$

Then

$$Ch(R) = W(R)J_1 \cup W(R)J_2, \quad W(R)(-J_2) = W(R)J_1.$$

Lemma 2.2 implies that

$$\#(W(R)J_i) = \frac{\#(W(\mathfrak{e}_6))}{\#(W(\mathfrak{so}(10) + \mathbb{R}))} = \frac{2^7 \cdot 3^4 \cdot 5}{2^4 \cdot 5!} = 3^3.$$

We can verify that $t = 2$, $d_1 = 2$, $d_2 = 4$ and $W(R)J_i$ is two-point homogeneous.

Remark 4.9. There exist exactly five subsets Δ_i ($1 \leq i \leq 5$) of $W(R)J_1$ such that each Δ_i contains J_1 and consists of three vertices of an equilateral triangle whose center of mass is equal to the origin. If we set $\mathfrak{a}^{(i)} = \text{span}_{\mathbb{R}}(\Delta_i)$ ($1 \leq i \leq 5$), then

$$\Delta_i = \mathfrak{a}^{(i)} \cap W(R)J_1.$$

These will be used later.

Example 4.10. In the case where $R = E_7$, we define a characteristic element J by

$$\langle \alpha_7, J \rangle = 1, \quad \langle \alpha_i, J \rangle = 0 \quad (i \neq 7).$$

Then $Ch(R) = W(R)J$. Lemma 2.2 implies that

$$\#(W(R)J) = \frac{\#(W(\mathfrak{e}_7))}{\#(W(\mathfrak{e}_6 + \mathbb{R}))} = \frac{2^{10} \cdot 3^4 \cdot 5 \cdot 7}{2^7 \cdot 3^4 \cdot 5} = 2^3 \cdot 7.$$

We can verify that $t = 3$, $d_1 = \sqrt{2}$, $d_2 = 2$, $d_3 = \sqrt{6}$ and $W(R)J$ is two-point homogeneous.

Hence we complete the proof of Proposition 4.1. \square

Theorem 2.1 and Proposition 4.1 imply the following theorem.

Theorem 4.11. *A great antipodal set of an irreducible Hermitian symmetric space of compact type is a two-point homogeneous space.*

4.2. Symmetric triads. In this subsection we review some results on symmetric triads obtained in [6] and [7]. These results will be used in Section 4.3 and Section 5.

Let \mathfrak{a} be a finite dimensional vector space over \mathbb{R} with an inner product $\langle \cdot, \cdot \rangle$. A triple $(\tilde{\Sigma}, \Sigma, W)$ is a *symmetric triad* of \mathfrak{a} , if it satisfies the following six conditions:

- (1) $\tilde{\Sigma}$ is an irreducible root system of \mathfrak{a} , and $\tilde{\Sigma}$ spans \mathfrak{a} .
- (2) Σ is a root system of \mathfrak{a} .
- (3) W is a nonempty subset of \mathfrak{a} , which is invariant under the multiplication by -1 , and $\tilde{\Sigma} = \Sigma \cup W$.
- (4) $\Sigma \cap W$ is a nonempty subset. If we put $l = \max\{\|\alpha\| \mid \alpha \in \Sigma \cap W\}$, then $\Sigma \cap W = \{\alpha \in \tilde{\Sigma} \mid \|\alpha\| \leq l\}$.
- (5) For $\alpha \in W$, $\lambda \in \Sigma - W$, $2 \frac{\langle \alpha, \lambda \rangle}{\|\alpha\|^2}$ is odd if and only if $s_\alpha \lambda \in W - \Sigma$, where we set $s_\alpha \lambda = \lambda - 2 \frac{\langle \alpha, \lambda \rangle}{\|\alpha\|^2} \alpha$.
- (6) For $\alpha \in W$, $\lambda \in W - \Sigma$, $2 \frac{\langle \alpha, \lambda \rangle}{\|\alpha\|^2}$ is odd if and only if $s_\alpha \lambda \in \Sigma - W$.

If $(\tilde{\Sigma}, \Sigma, W)$ is a symmetric triad of \mathfrak{a} , then Σ spans \mathfrak{a} . In fact, using (4) we have

$$\mathfrak{a} \supset \text{span}(\Sigma) \supset \text{span}(\Sigma \cap W) \supset \text{span}\{\text{the shortest roots in } \tilde{\Sigma}\} = \mathfrak{a}.$$

For a symmetric triad $(\tilde{\Sigma}, \Sigma, W)$ of \mathfrak{a} , take a fundamental system $\tilde{\Pi}$ of $\tilde{\Sigma}$. Denote by $\tilde{\Sigma}^+$ the set of positive roots in $\tilde{\Sigma}$ with respect to $\tilde{\Pi}$. If

we put $\Sigma^+ = \Sigma \cap \tilde{\Sigma}^+$ and $W^+ = W \cap \tilde{\Sigma}^+$, then $\Sigma = \Sigma^+ \cup (-\Sigma^+)$ and $W = W^+ \cup (-W^+)$. We define a nonempty subset \mathfrak{a}_r in \mathfrak{a} by

$$\mathfrak{a}_r = \bigcap_{\substack{\lambda \in \Sigma \\ \alpha \in W}} \left\{ H \in \mathfrak{a} \mid \langle \lambda, H \rangle \notin \pi\mathbb{Z}, \langle \alpha, H \rangle \notin \frac{\pi}{2} + \pi\mathbb{Z} \right\}.$$

Then \mathfrak{a}_r is an open dense subset of \mathfrak{a} . A point in \mathfrak{a}_r is called a *regular point*.

Let G be a connected compact simple Lie group and (G, F_1, F_2) a compact symmetric triad: There exist two involutions θ_1 and θ_2 on G such that the closed subgroup F_i of G lies between $F(\theta_i, G)$ and its identity component $F(\theta_i, G)_0$. We denote by \mathfrak{g} , \mathfrak{f}_1 and \mathfrak{f}_2 the Lie algebras of G , F_1 and F_2 respectively. We assume that $\theta_1\theta_2 = \theta_2\theta_1$ and that θ_1 cannot be transformed to θ_2 by an inner automorphism of G . We denote the differential of θ_i by the same symbol θ_i . We have two canonical decompositions of \mathfrak{g} :

$$\mathfrak{g} = \mathfrak{f}_1 \oplus \mathfrak{p}_1 = \mathfrak{f}_2 \oplus \mathfrak{p}_2,$$

where $\mathfrak{p}_i = F(-\theta_i, \mathfrak{g})$. Since $\theta_1\theta_2 = \theta_2\theta_1$, we have

$$\mathfrak{g} = (\mathfrak{f}_1 \cap \mathfrak{f}_2) \oplus (\mathfrak{p}_1 \cap \mathfrak{p}_2) \oplus (\mathfrak{f}_1 \cap \mathfrak{p}_2) \oplus (\mathfrak{f}_2 \cap \mathfrak{p}_1).$$

Take a maximal abelian subspace \mathfrak{a} of $\mathfrak{p}_1 \cap \mathfrak{p}_2$. The isometric action of F_1 on a Riemannian symmetric space G/F_2 of compact type is called a Hermann action. Since the action is a hyperpolar action whose section is the orbit of $A = \exp \mathfrak{a}$ through the origin, we have $G = F_1 A F_2$. For each $\alpha \in \mathfrak{a}$ define a subspace $\mathfrak{g}(\mathfrak{a}, \alpha)$ of $\mathfrak{g}^{\mathbb{C}}$ by

$$\mathfrak{g}(\mathfrak{a}, \alpha) = \{X \in \mathfrak{g}^{\mathbb{C}} \mid [H, X] = \sqrt{-1}\langle \alpha, H \rangle X \ (H \in \mathfrak{a})\}$$

and set $\tilde{\Sigma} = \{\alpha \in \mathfrak{a} - \{0\} \mid \mathfrak{g}(\mathfrak{a}, \alpha) \neq \{0\}\}$. For $\epsilon = \pm 1$ define a subspace $\mathfrak{g}(\mathfrak{a}, \alpha, \epsilon)$ of $\mathfrak{g}(\mathfrak{a}, \alpha)$ by

$$\mathfrak{g}(\mathfrak{a}, \alpha, \epsilon) = \{X \in \mathfrak{g}(\mathfrak{a}, \alpha) \mid \theta_1\theta_2 X = \epsilon X\}.$$

Since $\mathfrak{g}(\mathfrak{a}, \alpha)$ is $\theta_1\theta_2$ -invariant, we have

$$\mathfrak{g}(\mathfrak{a}, \alpha) = \mathfrak{g}(\mathfrak{a}, \alpha, 1) \oplus \mathfrak{g}(\mathfrak{a}, \alpha, -1).$$

Set $\Sigma = \{\alpha \in \tilde{\Sigma} \mid \mathfrak{g}(\mathfrak{a}, \alpha, 1) \neq \{0\}\}$ and $W = \{\alpha \in \tilde{\Sigma} \mid \mathfrak{g}(\mathfrak{a}, \alpha, -1) \neq \{0\}\}$. Then the triple $(\tilde{\Sigma}, \Sigma, W)$ is a symmetric triad of \mathfrak{a} . Define closed subgroups G_{12} and F_{12} by $G_{12} = F(\theta_1\theta_2, G)$ and $F_{12} = \{g \in G_{12} \mid \theta_1(g) = g\}$. Then the Lie algebras of G_{12} and F_{12} are given by

$$\mathfrak{g}_{12} = (\mathfrak{f}_1 \cap \mathfrak{f}_2) \oplus (\mathfrak{p}_1 \cap \mathfrak{p}_2), \quad \mathfrak{f}_{12} = \mathfrak{f}_1 \cap \mathfrak{f}_2,$$

respectively. The restricted root system of the compact symmetric pair (G_{12}, F_{12}) with respect to \mathfrak{a} coincides with Σ . For $\lambda \in \Sigma$, we define subspaces \mathfrak{p}_λ in $\mathfrak{p}_1 \cap \mathfrak{p}_2$ and \mathfrak{f}_λ in $\mathfrak{f}_1 \cap \mathfrak{f}_2$ as follows:

$$\begin{aligned}\mathfrak{p}_\lambda &= \{X \in \mathfrak{p}_1 \cap \mathfrak{p}_2 \mid [H, [H, X]] = -\langle \lambda, H \rangle^2 X \ (H \in \mathfrak{a})\}, \\ \mathfrak{f}_\lambda &= \{X \in \mathfrak{f}_1 \cap \mathfrak{f}_2 \mid [H, [H, X]] = -\langle \lambda, H \rangle^2 X \ (H \in \mathfrak{a})\}.\end{aligned}$$

Take a maximal abelian subalgebra \mathfrak{t} in \mathfrak{g}_{12} containing \mathfrak{a} . Denote by \tilde{R} the root system of \mathfrak{g}_{12} with respect to \mathfrak{t} . Let $\mathfrak{t} \rightarrow \mathfrak{a}$; $H \mapsto \bar{H}$ be the orthogonal projection and set $\tilde{R}_0 = \{\alpha \in \tilde{R} \mid \bar{\alpha} = 0\}$. Define a subalgebra \mathfrak{f}_0 in $\mathfrak{f}_1 \cap \mathfrak{f}_2$ by

$$\mathfrak{f}_0 = \{X \in \mathfrak{f}_1 \cap \mathfrak{f}_2 \mid [\mathfrak{a}, X] = \{0\}\}.$$

Take a compatible ordering of \mathfrak{t} . Then we have the following lemma.

Lemma 4.12. (1) *We have orthogonal direct sum decompositions:*

$$\mathfrak{f}_1 \cap \mathfrak{f}_2 = \mathfrak{f}_0 \oplus \sum_{\lambda \in \Sigma^+} \mathfrak{f}_\lambda, \quad \mathfrak{p}_1 \cap \mathfrak{p}_2 = \mathfrak{a} \oplus \sum_{\lambda \in \Sigma^+} \mathfrak{p}_\lambda.$$

(2) *For each $\alpha \in \tilde{R}^+ - \tilde{R}_0$ there exist $S_\alpha \in \mathfrak{f}_1 \cap \mathfrak{f}_2$ and $T_\alpha \in \mathfrak{p}_1 \cap \mathfrak{p}_2$ such that*

$$\{S_\alpha \mid \alpha \in \tilde{R}^+, \bar{\alpha} = \lambda\}, \quad \{T_\alpha \mid \alpha \in \tilde{R}^+, \bar{\alpha} = \lambda\}$$

are respectively orthonormal bases of \mathfrak{f}_λ and \mathfrak{p}_λ , and for $H \in \mathfrak{a}$

$$[H, S_\alpha] = \langle \alpha, H \rangle T_\alpha, \quad [H, T_\alpha] = -\langle \alpha, H \rangle S_\alpha, \quad [S_\alpha, T_\alpha] = \bar{\alpha},$$

$$\text{Ad}(\exp H)S_\alpha = \cos\langle \alpha, H \rangle S_\alpha + \sin\langle \alpha, H \rangle T_\alpha,$$

$$\text{Ad}(\exp H)T_\alpha = -\sin\langle \alpha, H \rangle S_\alpha + \cos\langle \alpha, H \rangle T_\alpha.$$

Define subspaces of $\mathfrak{f}_1 \cap \mathfrak{p}_2$ and $\mathfrak{p}_1 \cap \mathfrak{f}_2$ by

$$V(\mathfrak{f}_1 \cap \mathfrak{p}_2) = \{X \in \mathfrak{f}_1 \cap \mathfrak{p}_2 \mid [\mathfrak{a}, X] = \{0\}\},$$

$$V(\mathfrak{p}_1 \cap \mathfrak{f}_2) = \{X \in \mathfrak{p}_1 \cap \mathfrak{f}_2 \mid [\mathfrak{a}, X] = \{0\}\},$$

$$V^\perp(\mathfrak{f}_1 \cap \mathfrak{p}_2) = \{X \in \mathfrak{f}_1 \cap \mathfrak{p}_2 \mid X \perp V(\mathfrak{f}_1 \cap \mathfrak{p}_2)\},$$

$$V^\perp(\mathfrak{p}_1 \cap \mathfrak{f}_2) = \{X \in \mathfrak{p}_1 \cap \mathfrak{f}_2 \mid X \perp V(\mathfrak{p}_1 \cap \mathfrak{f}_2)\}.$$

For $\alpha \in W$ define subspaces $V_\alpha^\perp(\mathfrak{f}_1 \cap \mathfrak{p}_2)$ in $V^\perp(\mathfrak{f}_1 \cap \mathfrak{p}_2)$ and $V_\alpha^\perp(\mathfrak{p}_1 \cap \mathfrak{f}_2)$ in $V^\perp(\mathfrak{p}_1 \cap \mathfrak{f}_2)$ by

$$V_\alpha^\perp(\mathfrak{f}_1 \cap \mathfrak{p}_2) = \{X \in V^\perp(\mathfrak{f}_1 \cap \mathfrak{p}_2) \mid [H, [H, X]] = -\langle \alpha, H \rangle^2 X \ (H \in \mathfrak{a})\},$$

$$V_\alpha^\perp(\mathfrak{p}_1 \cap \mathfrak{f}_2) = \{X \in V^\perp(\mathfrak{p}_1 \cap \mathfrak{f}_2) \mid [H, [H, X]] = -\langle \alpha, H \rangle^2 X \ (H \in \mathfrak{a})\}.$$

Then we have the orthogonal direct sum decompositions:

$$V^\perp(\mathfrak{f}_1 \cap \mathfrak{p}_2) = \sum_{\alpha \in W^+} V_\alpha^\perp(\mathfrak{f}_1 \cap \mathfrak{p}_2), \quad V^\perp(\mathfrak{p}_1 \cap \mathfrak{f}_2) = \sum_{\alpha \in W^+} V_\alpha^\perp(\mathfrak{p}_1 \cap \mathfrak{f}_2).$$

For $\lambda \in \Sigma$ and $\alpha \in W$, set

$$m(\lambda) = \dim_{\mathbb{C}} \mathfrak{g}(\mathfrak{a}, \lambda, 1), \quad n(\alpha) = \dim_{\mathbb{C}} \mathfrak{g}(\mathfrak{a}, \alpha, -1).$$

Lemma 4.13. (1) For any $\alpha \in W^+$,

$$[\mathfrak{a}, V_{\alpha}^{\perp}(\mathfrak{f}_1 \cap \mathfrak{p}_2)] = V_{\alpha}^{\perp}(\mathfrak{p}_1 \cap \mathfrak{f}_2),$$

$$[\mathfrak{a}, V_{\alpha}^{\perp}(\mathfrak{p}_1 \cap \mathfrak{f}_2)] = V_{\alpha}^{\perp}(\mathfrak{f}_1 \cap \mathfrak{p}_2).$$

(2) There exist orthonormal bases $\{X_{\alpha,i}\}_{1 \leq i \leq n(\alpha)}$ and $\{Y_{\alpha,i}\}_{1 \leq i \leq n(\alpha)}$ of $V_{\alpha}^{\perp}(\mathfrak{f}_1 \cap \mathfrak{p}_2)$ and $V_{\alpha}^{\perp}(\mathfrak{p}_1 \cap \mathfrak{f}_2)$ respectively such that, for any $H \in \mathfrak{a}$,

$$[H, X_{\alpha,i}] = \langle \alpha, H \rangle Y_{\alpha,i}, \quad [H, Y_{\alpha,i}] = -\langle \alpha, H \rangle X_{\alpha,i},$$

$$[X_{\alpha,i}, Y_{\alpha,i}] = \alpha,$$

$$\text{Ad}(\exp H)X_{\alpha,i} = \cos \langle \alpha, H \rangle X_{\alpha,i} + \sin \langle \alpha, H \rangle Y_{\alpha,i},$$

$$\text{Ad}(\exp H)Y_{\alpha,i} = -\sin \langle \alpha, H \rangle X_{\alpha,i} + \cos \langle \alpha, H \rangle Y_{\alpha,i}.$$

Lemmas 4.12 and 4.13 imply that

$$(4.4) \quad W(\tilde{\Sigma}) \subset \{\text{Ad}(g)|_{\mathfrak{a}} \mid g \in G, \text{Ad}(g)\mathfrak{a} = \mathfrak{a}\},$$

where $W(\tilde{\Sigma})$ denotes the Weyl group of $\tilde{\Sigma}$. See Corollary 4.17 and Lemma 4.4 in [6] for the detail.

Take a maximal abelian subspace \mathfrak{a}_i of \mathfrak{p}_i containing \mathfrak{a} . The maximality of \mathfrak{a} implies that $\mathfrak{a} = \mathfrak{a}_1 \cap \mathfrak{a}_2$. Denote by R_i the restricted root system of (G, F_i) with respect to \mathfrak{a}_i . If $\mathfrak{a} = \mathfrak{a}_1$, then $\tilde{\Sigma} = R_1$. We list some (G, F_1, F_2) 's, their symmetric triad $(\tilde{\Sigma}, \Sigma, W)$'s, the restricted root system (R_i) 's, and the structure of \mathfrak{f}_0 's as a compact Lie algebra, which we will need in Subsection 4.3 and Section 5.

	(G, F_1, F_2)	$(\tilde{\Sigma}, \Sigma, W)$
(1)	$(SU(2n), S(U(n) \times U(n)), SO(2n))$	(I- C_n)
(2)	$(Sp(2m), Sp(m) \times Sp(m), U(2m))$	(III- C_m)
(3)	$(SO(4m), U(2m), SO(2m) \times SO(2m))$	(I- C_m)
(4)	$(E_7, S^1 \cdot E_6, SU(8))$	(I- C_3)
(5)	$(SO(r+s+t), SO(r) \times SO(s+t),$ $SO(r+s) \times SO(t))$ ($s > 0, r < t$)	(I- B_r)
(6)	$(E_6, F_4, Sp(4))$	(III- A_2)
(7)	$(SU(2(m+q)), Sp(m+q), SO(2(m+q)))$	(III- A_{m+q-1})
(8)	$(SU(4m), Sp(2m), S(U(2m) \times U(2m)))$	(III- C_m)

	R_1	R_2	\mathfrak{f}_0
(1)	C_n	A_{2n-1}	$\{0\}$
(2)	C_m	C_{2m}	\mathbb{R}^m
(3)	C_m	D_{2m}	\mathbb{R}^m
(4)	C_3	E_7	$\mathfrak{su}(2)^4$
(5)	B_r	$B_{\min\{r+s,t\}}$ (if $r+s \neq t$) D_t (if $r+s = t$)	$\mathfrak{so}(s) \oplus \mathfrak{so}(t-r)$
(6)	A_2	E_6	$\mathfrak{sp}(1)^4$
(7)	A_{m+q-1}	$A_{2(m+q)-1}$	\mathbb{R}^{m+q}
(8)	A_{2m-1}	C_{2m}	$\mathfrak{sp}(1)^m$

We explain the notations in the table above. In column of $(\tilde{\Sigma}, \Sigma, W)$'s we used the following notations.

	$\tilde{\Sigma}$	Σ	W
(III- $\tilde{\Sigma}$)	$\tilde{\Sigma}$	$\tilde{\Sigma}$	$\tilde{\Sigma}$
(I'- C_n)	C_n	D_n	C_n
(I- C_m)	C_m	C_m	D_m
(I- B_r)	B_r	B_r	$\{\pm e_i\}$

In the table above, $\mathfrak{f}_0 = \mathbb{R}^m$ means that \mathfrak{f}_0 is an abelian Lie algebra of dimension m . We used table 1 in Tamaru [14] and a table of Section 4 in Matsuki [10] to determine the structure of \mathfrak{f}_0 when G is of exceptional type. Note that $\mathfrak{so}(s)$ is abelian if and only if $s \leq 2$.

4.3. The intersection of two real forms. Let $M = \text{Ad}(G)J \subset \mathfrak{g}$ be an irreducible Hermitian symmetric space of compact type. In this subsection we study a necessary and sufficient condition that the intersection of two real forms of M is discrete, and describe the intersection when it is discrete. Any two real forms of M always intersect. Let $L_1 = F(\tau_1, M)$ and $L_2 = F(\tau_2, M)$ be two real forms of M , where τ_i is an involutive anti-holomorphic isometry of M . Define an involution θ_i of G by $\theta_i(g) = \tau_i g \tau_i^{-1}$. If we set $F_i = F(\theta_i, G)$, then (G, F_1, F_2) is a compact symmetric triad. In order to study $L_1 \cap \text{Ad}(a)L_2$ for $a \in G$, we may assume that $\tau_1 \tau_2 = \tau_2 \tau_1$ by the classification of real forms in irreducible Hermitian symmetric spaces of compact type. We use the same notation in Subsection 4.2. Take a maximal abelian subspace \mathfrak{a} of $\mathfrak{p}_1 \cap \mathfrak{p}_2$ which contains J . We may assume that a is in $\exp \mathfrak{a}$ since $G = F_1(\exp \mathfrak{a})F_2$. By Theorem 4.3 in [16], we have

$$\begin{aligned} L_1 &= M \cap \mathfrak{p}_1, & \text{Ad}(a)L_2 &= M \cap \text{Ad}(a)\mathfrak{p}_2, \\ L_1 \cap \text{Ad}(a)L_2 &= M \cap (\mathfrak{p}_1 \cap \text{Ad}(a)\mathfrak{p}_2). \end{aligned}$$

If L_1 is congruent to L_2 , which means that there exists $g \in G$ such that $L_2 = \text{Ad}(g)L_1$, then we may assume that $L_1 = L_2$. We divide into the following two cases:

- (1) $L_1 = L_2$.
- (2) L_1 is not congruent to L_2 .

First we assume that $L_1 = L_2$. Set $\tau = \tau_1 = \tau_2$, $L = L_1 = L_2$ and so on. Denote by R the restricted root system of (G, F) with respect to \mathfrak{a} . Then we have root space decompositions of \mathfrak{f} and \mathfrak{p} :

$$\mathfrak{f} = \mathfrak{f}_0 \oplus \sum_{\lambda \in R_+} \mathfrak{f}_\lambda, \quad \mathfrak{p} = \mathfrak{a} \oplus \sum_{\lambda \in R_+} \mathfrak{p}_\lambda,$$

where R_+ is the set of positive roots in R with respect to a lexicographic ordering. The complex structure J is a characteristic element associated with R . Since

$$\mathfrak{m} = [J, \mathfrak{g}] = \sum_{\substack{\lambda \in R_+ \\ \langle \lambda, J \rangle \neq 0}} (\mathfrak{f}_\lambda \oplus \mathfrak{p}_\lambda),$$

we have

$$(4.5) \quad \dim M = 2 \sum_{\substack{\lambda \in R_+ \\ \langle \lambda, J \rangle \neq 0}} m_R(\lambda),$$

where we denote by $m_R(\lambda)$ the multiplicity of λ . If we set $a = \exp H$ for $H \in \mathfrak{a}$, then

$$\mathfrak{p} \cap \text{Ad}(a)\mathfrak{p} = \mathfrak{a} \oplus \sum_{\substack{\lambda \in R_+ \\ \langle \lambda, H \rangle \in \pi\mathbb{Z}}} \mathfrak{p}_\lambda.$$

Theorem 4.14. *The intersection $L \cap \text{Ad}(a)L$ is discrete if and only if $\langle \lambda, H \rangle \notin \pi\mathbb{Z}$ for any $\lambda \in R$. In this case,*

$$(4.6) \quad L \cap \text{Ad}(a)L = M \cap \mathfrak{a} = W(R)J,$$

where $M \cap \mathfrak{a}$ is a great antipodal set of L .

Proof. If $\langle \lambda, H \rangle \notin \pi\mathbb{Z}$ for any $\lambda \in R$, then we have (4.6), since $\mathfrak{p} \cap \text{Ad}(a)\mathfrak{p} = \mathfrak{a}$. Here the second equality follows from Proposition 2.2 in [5, Ch.VII].

If there exists $\lambda \in R$ such that $\langle \lambda, H \rangle \in \pi\mathbb{Z}$, then there exists $X \in W(R)J$ such that $\langle \lambda, X \rangle \neq 0$. There exist unit vectors $S_\lambda \in \mathfrak{f}_\lambda$ and $T_\lambda \in \mathfrak{p}_\lambda$ such that for any $H' \in \mathfrak{a}$

$$[H', S_\lambda] = \langle \lambda, H' \rangle T_\lambda, \quad [H', T_\lambda] = -\langle \lambda, H' \rangle S_\lambda, \quad [S_\lambda, T_\lambda] = \lambda.$$

Then

$$\begin{aligned} \text{Ad}(\exp tS_\lambda)X &= X + \frac{\langle \lambda, X \rangle}{\|\lambda\|^2}(\cos(t\|\lambda\|) - 1)\lambda - \frac{\langle \lambda, X \rangle}{\|\lambda\|} \sin(t\|\lambda\|)T_\lambda \\ &\in L_1 \cap \text{Ad}(a)L_2. \end{aligned}$$

Hence $L_1 \cap \text{Ad}(a)L_2$ is not discrete. \square

Example 4.15. If $(M, L) = (Sp(r)/U(r), U(r)/O(r))$, then $R = C_r$ and $\#(W(R)J) = 2^r$.

Proof. The assertion immediately follows from the table in Subsection 4.2 and Example 4.3. \square

Example 4.16. If $(M, L) = (G_k(\mathbb{C}^n), G_k(\mathbb{R}^n))$, then $R = A_{n-1}$ and $\#(W(R)J) = \binom{n}{k}$.

Proof. Since $(G, F) = (SU(n), SO(n))$, we have $R = A_{n-1}$ and the multiplicity of any root in R is equal to 1. (4.5) implies that $J = J_k$ or $J = J_{n-k}$ in Example 4.6. Thus $\#(W(R)J) = \binom{n}{k}$ by Example 4.6. \square

Example 4.17. If $(M, L) = (SO(2r)/U(r), SO(r))$, then $R = D_r$ and $\#(W(R)J) = 2^{r-1}$.

Proof. Since $(G, F) = (SO(2r), S(O(r) \times O(r)))$, we have $R = D_r$ and the multiplicity of any root in R is equal to 1. (4.5) implies that $J = J_2$ in Example 4.5. Thus $\#(W(R)J) = 2^{r-1}$ by Example 4.5. \square

Next we assume that L_1 is not congruent to L_2 . Denote by $(\tilde{\Sigma}, \Sigma, W)$ the symmetric triad associated with (G, F_1, F_2) . By Lemmas 4.12 and 4.13 J is a characteristic element associated with $\tilde{\Sigma}$. Lemmas 4.12 and 4.13 also imply that

$$\mathfrak{p}_1 \cap \text{Ad}(a)\mathfrak{p}_2 = \mathfrak{a} \oplus \sum_{\substack{\lambda \in \Sigma^+ \\ \langle \lambda, H \rangle \in \pi\mathbb{Z}}} \mathfrak{p}_\lambda \oplus \sum_{\substack{\alpha \in W^+ \\ \langle \alpha, H \rangle \in \frac{\pi}{2} + \pi\mathbb{Z}}} V_\alpha^\perp(\mathfrak{p}_1 \cap \mathfrak{f}_2).$$

Denote by R_i the restricted root system of (G, F_i) with respect to \mathfrak{a}_i .

Theorem 4.18. *The intersection $L_1 \cap \text{Ad}(a)L_2$ ($a = \exp H$) is discrete if and only if H is a regular point of $(\tilde{\Sigma}, \Sigma, W)$.*

Proof. If H is a regular point of $(\tilde{\Sigma}, \Sigma, W)$, then $\mathfrak{p}_1 \cap \text{Ad}(a)\mathfrak{p}_2 = \mathfrak{a}$. Thus

$$(4.7) \quad L_1 \cap \text{Ad}(a)L_2 = (M \cap \mathfrak{a}_1) \cap (M \cap \mathfrak{a}_2) = W(R_1)J \cap W(R_2)J.$$

We assume that H is not a regular point of $(\tilde{\Sigma}, \Sigma, W)$. Then (i) there exists $\lambda \in \Sigma$ such that $\langle \lambda, H \rangle \in \pi\mathbb{Z}$, or, (ii) there exists $\alpha \in W$ such that $\langle \alpha, H \rangle \in \frac{\pi}{2} + \pi\mathbb{Z}$.

In the case of (i) we can prove that $L_1 \cap \text{Ad}(a)L_2$ is not discrete in a similar manner of the proof of Theorem 4.14. In the case of (ii) there exists $X \in W(\Sigma)J$ such that $\langle \alpha, X \rangle \neq 0$ since $W(\Sigma)\alpha$ spans \mathfrak{a} . Lemma 4.13 implies that

$$\begin{aligned} \text{Ad}(\exp tX_{\alpha,i})X &= X + \frac{\langle \alpha, X \rangle}{\|\alpha\|^2}(\cos(t\|\alpha\|) - 1)\alpha - \frac{\langle \alpha, X \rangle}{\|\alpha\|} \sin(t\|\alpha\|)Y_{\alpha,i} \\ &\in L_1 \cap \text{Ad}(a)L_2. \end{aligned}$$

Hence $L_1 \cap \text{Ad}(a)L_2$ is not discrete. \square

Note that H is a regular point of $(\tilde{\Sigma}, \Sigma, W)$ if and only if F_2 -orbit through $aF_1 \in G/F_1$ is a regular orbit.

In the sequel we assume that $L_1 \cap \text{Ad}(a)L_2$ is discrete. (4.4) and (4.7) imply

$$W(\tilde{\Sigma})J \subset M \cap \mathfrak{a} = L_1 \cap \text{Ad}(a)L_2 \subset W(R_i)J \cap \mathfrak{a}.$$

Based on the fact, we prove the following theorem.

Theorem 4.19. *Assume that $L_1 \cap \text{Ad}(a)L_2$ is discrete. Then*

$$L_1 \cap \text{Ad}(a)L_2 = W(\tilde{\Sigma})J = W(R_1)J \cap \mathfrak{a} = W(R_2)J \cap \mathfrak{a}.$$

Proof. It is sufficient to prove that $W(R_i)J \cap \mathfrak{a} \subset W(\tilde{\Sigma})J$. The type of $\tilde{\Sigma}$ is A , B or C by the classification of (M, L_1, L_2) . If $\tilde{\Sigma}$ is of type B or C , then $W(\tilde{\Sigma}) = \text{Ch}(\tilde{\Sigma})$ by Examples 4.2 and 4.3. Any $X \in W(R_i)J \cap \mathfrak{a}$ satisfies $(\text{ad}X)^3 = -\text{ad}X$. Hence X is in $\text{Ch}(\tilde{\Sigma})$. If $\tilde{\Sigma}$ is of type A , then $(M, L_1, L_2) = (E_6/S^1 \cdot \text{Spin}(10), F_4/\text{Spin}(9), G_2(\mathbb{H}^4)/\mathbb{Z}_2)$ or $(G_{2q}(\mathbb{C}^{2(m+q)}), G_q(\mathbb{H}^{m+q}), G_{2q}(\mathbb{R}^{2(m+q)}))$. In these cases we will prove $W(\tilde{\Sigma})J = W(R_i)J \cap \mathfrak{a}$ below.

Example 4.20. If

$$(M, L_1, L_2) = (E_6/S^1 \cdot \text{Spin}(10), F_4/\text{Spin}(9), G_2(\mathbb{H}^4)/\mathbb{Z}_2),$$

then

$$\mathfrak{a}_1 = \mathfrak{a}, \quad \tilde{\Sigma} = R_1 = A_2, \quad R_2 = E_6$$

and

$$\begin{aligned} W(\tilde{\Sigma})J &= W(R_1)J = W(R_2)J \cap \mathfrak{a}, \\ \#(W(\tilde{\Sigma})J) &= 3, \quad \#(W(R_2)J) = 3^3. \end{aligned}$$

Proof. Since $(G, F_1, F_2) = (E_6, F_4, Sp(4))$ we have $\tilde{\Sigma} = R_1 = A_2$ and $R_2 = E_6$ by (6) of the table in Subsection 4.2. Hence $W(\tilde{\Sigma})J = W(R_1)J \cap \mathfrak{a} = W(R_1)J$. From Example 4.6 we have $\#(W(\tilde{\Sigma})J) = 3$. There exists i ($1 \leq i \leq 5$) such that $\mathfrak{a}_2 = \mathfrak{a}^{(i)}$ by Remarks 4.7 and 4.9.

Hence $\#(W(R_2)J \cap \mathfrak{a}) = 3$ by Remark 4.9. From Example 4.8 we have $\#(W(R_2)J) = 3^3$. \square

Example 4.21. If

$$(M, L_1, L_2) = (G_{2q}(\mathbb{C}^{2(m+q)}), G_q(\mathbb{H}^{m+q}), G_{2q}(\mathbb{R}^{2(m+q)})),$$

then

$$\mathfrak{a}_1 = \mathfrak{a}, \quad \tilde{\Sigma} = R_1 = A_{m+q-1}, \quad R_2 = A_{2(m+q)-1}$$

and

$$\begin{aligned} W(\tilde{\Sigma})J &= W(R_1)J = W(R_2)J \cap \mathfrak{a}, \\ \#(W(\tilde{\Sigma})J) &= \binom{m+q}{q}, \quad \#(W(R_2)J) = \binom{2m+2q}{2m}. \end{aligned}$$

Proof. We have $\mathfrak{a}_1 = \mathfrak{a}$, $\tilde{\Sigma} = R_1 = A_{m+q-1}$, $R_2 = A_{2(m+q)-1}$ by (7) of the table in Subsection 4.2. Hence $W(\tilde{\Sigma})J = W(R_1)J$. Since J is a characteristic element associated with R_1 , there exists J_i ($1 \leq i \leq m+q-1$) in Example 4.6 such that $J = J_i$. Since the multiplicity of any root in R_1 is equal to 4, (4.5) implies that

$$8mq = \dim M = 2 \cdot 4 \cdot i(m+q-i).$$

Hence $i = m$ or $i = q$. In any case we have

$$\#(W(\tilde{\Sigma})J) = \binom{m+q}{q}$$

by Example 4.6. Since J is also a characteristic element associated with R_2 , there exists J_k ($1 \leq k \leq 2m+2q-1$) in Example 4.6 such that $J = J_k$. Since the multiplicity of any root in R_2 is equal to 1, (4.5) implies that

$$8mq = \dim M = 2 \cdot k(2m+2q-k).$$

Hence $k = 2m$ or $k = 2q$. In any case we have

$$\#(W(R_2)J) = \binom{2m+2q}{2m}$$

by Example 4.6. To show $W(\tilde{\Sigma})J = W(R_2)J \cap \mathfrak{a}$, we identify \mathfrak{a}_2 and \mathfrak{a} with the following subspaces.

$$\begin{aligned} \mathfrak{a}_2 &\cong \left\{ \sum_{i=1}^{p+q} (x_i e_i + y_i e_{i+p+q}) \in \mathbb{R}^{2(m+q)} \mid \sum_{i=1}^{p+q} (x_i + y_i) = 0 \right\}, \\ \mathfrak{a} &\cong \left\{ \sum_{i=1}^{p+q} x_i (e_i + e_{i+p+q}) \in \mathbb{R}^{2(m+q)} \mid \sum_{i=1}^{p+q} x_i = 0 \right\}. \end{aligned}$$

Under the identification we have $R_2^+ \cong \{e_i - e_j \mid 1 \leq i < j \leq 2m + 2q\}$. Since $J = J_{2m}$ or $J = J_{2q}$, set $J = J_{2m}$ for instance. In order to describe $W(R_2)J$ we use the following notations. Denote by $P_a(m+q)$ the set consisting of all subsets of cardinality a in $\{1, 2, \dots, m+q\}$. For $A \in P_a(m+q)$ denote by \bar{A} the complement of A in $\{1, \dots, m+q\}$ and set $A' = \{x + m + q \mid x \in A\}$. For any $X \in W(R_2)J$ there exist $A \in P_a(m+q)$ and $B \in P_b(m+q)$ such that $a + b = 2m$ and that

$$\begin{aligned} X &= \sum_{j \in A} e_j + \sum_{k \in B'} e_k - \frac{m}{m+q} \left(\sum_{j \in A} e_j + \sum_{j \in \bar{A}} e_j + \sum_{k \in \bar{B}} e_k + \sum_{k \in B'} e_k \right) \\ &= \frac{q}{m+q} \left(\sum_{j \in A} e_j + \sum_{k \in \bar{B}} e_k \right) - \frac{m}{m+q} \left(\sum_{j \in \bar{A}} e_j + \sum_{k \in B'} e_k \right). \end{aligned}$$

Hence X is in \mathfrak{a} if and only if $a = b = m$, $A = B$. In this case

$$X = \frac{q}{m+q} \sum_{j \in A} (e_j + e_{j+m+q}) - \frac{q}{m+q} \sum_{j \in \bar{A}} (e_j + e_{j+m+q}).$$

Hence we have

$$\#(W(R_2)J \cap \mathfrak{a}) = \binom{m+q}{q}.$$

Thus $W(\tilde{\Sigma})J = W(R_2)J \cap \mathfrak{a}$. When $J = J_{2q}$, we get the same conclusion. \square

This completes the proof of Theorem 4.19. \square

We give another example.

Example 4.22. If

$$(M, L_1, L_2) = (G_{2m}(\mathbb{C}^{4m}), G_m(\mathbb{H}^{2m}), U(2m)),$$

then $\tilde{\Sigma} = C_m$, $R_1 = A_{2m-1}$, $R_2 = C_{2m}$ and

$$\#(L_1 \cap \text{Ad}(a)L_2) = 2^m, \quad \#(W(R_1)J) = \binom{2m}{m}, \quad \#(W(R_2)J) = 2^{2m}.$$

Proof. We have $\tilde{\Sigma} = C_m$, $R_1 = A_{2m+1}$ and $R_2 = C_m$ by (8) of the table in Subsection 4.2. Theorem 4.19 and Example 4.3 imply

$$\#(L_1 \cap \text{Ad}(a)L_2) = 2^m, \quad \#(W(R_2)J) = 2^{2m}.$$

Since J is a characteristic element associated with R_1 , there exists J_i ($1 \leq i \leq 2m - 1$) in Example 4.6 such that $J = J_i$. Since the multiplicity of any root is equal to 4, (4.5) implies that

$$8m^2 = \dim M = 8 \cdot i(m - i).$$

Hence $i = m$. Thus $\#(W(R_1)J) = \binom{2m}{m}$ by Example 4.6. \square

The following results immediately follows from Theorem 4.19.

Corollary 4.23. *Assume that $L_1 \cap \text{Ad}(a)L_2$ is discrete. If $\mathfrak{a} = \mathfrak{a}_1$ then $\tilde{\Sigma} = R_1$ and $L_1 \cap \text{Ad}(a)L_2 = W(R_1)J$.*

The following examples satisfy the assumption $\mathfrak{a} = \mathfrak{a}_1$ of Corollary 4.23.

Example 4.24. If

$$(M, L_1, L_2) = (Sp(2m)/U(2m), Sp(m), U(2m)/O(2m)),$$

then $\tilde{\Sigma} = R_1 = C_m$, $R_2 = C_{2m}$ and

$$\begin{aligned} L_1 \cap \text{Ad}(a)L_2 &= W(R_1)J = W(R_2)J \cap \mathfrak{a}, \\ \#(L_1 \cap \text{Ad}(a)L_2) &= 2^m, \quad \#(W(R_2)J) = 2^{2m}. \end{aligned}$$

Proof. We have $\tilde{\Sigma} = R_1 = C_m$ and $R_2 = C_{2m}$ by (2) of the table in Subsection 4.2. Theorem 4.19 and Example 4.3 imply $\#(L_1 \cap \text{Ad}(a)L_2) = 2^m$ and $\#(W(R_2)J) = 2^{2m}$. \square

Example 4.25. If

$$(M, L_1, L_2) = (E_7/S^1 \cdot E_6, S^1 \cdot E_6/F_4, (SU(8)/Sp(4))/\mathbb{Z}_2),$$

then $\tilde{\Sigma} = R_1 = C_3$, $R_2 = E_7$ and

$$\begin{aligned} L_1 \cap \text{Ad}(a)L_2 &= W(R_1)J = W(R_2)J \cap \mathfrak{a}, \\ \#(L_1 \cap \text{Ad}(a)L_2) &= 2^3, \quad \#(W(R_2)J) = 2^3 \cdot 7. \end{aligned}$$

Proof. We can verify $\tilde{\Sigma}$, R_1 and R_2 by (4) of the table in Subsection 4.2. Theorem 4.19 and Example 4.3 imply $\#(L_1 \cap \text{Ad}(a)L_2) = 2^3$. Example 4.10 implies $\#(W(R_2)J) = 2^3 \cdot 7$. \square

Example 4.26. If

$$(M, L_1, L_2) = (G_n(\mathbb{C}^{2n}), U(n), G_n(\mathbb{R}^{2n})),$$

then $\tilde{\Sigma} = R_1 = C_n$, $R_2 = A_{2n-1}$ and

$$\begin{aligned} L_1 \cap \text{Ad}(a)L_2 &= W(R_1)J = W(R_2)J \cap \mathfrak{a}, \\ \#(L_1 \cap \text{Ad}(a)L_2) &= 2^n, \quad \#(W(R_2)J) = \binom{2n}{n}. \end{aligned}$$

Proof. We can verify $\tilde{\Sigma}$, R_1 and R_2 by (1) of the table in Subsection 4.2. Theorem 4.19 and Example 4.3 imply $\#(L_1 \cap \text{Ad}(a)L_2) = 2^n$. Since J is a characteristic element associated with R_2 , there exists J_i ($1 \leq i \leq 2n-1$) in Example 4.6 such that $J = J_i$. Since the multiplicity of any root in R_2 is equal to 1, (4.5) implies that

$$2n^2 = \dim M = 2(2n - i)i.$$

Hence $i = n$. Thus $\#(W(R_2)J) = \binom{2n}{n}$ by Example 4.6. \square

Example 4.27. If

$$(M, L_1, L_2) = (Q_{r+s+t-2}(\mathbb{C}), S^{r-1, s+t-1}, S^{r+s-1, t-1}) \quad (s > 0, r < t),$$

then

$$\tilde{\Sigma} = R_1 = B_1, \quad R_2 = \begin{cases} B_{\min\{r+s, t\}} & (r + s \neq t) \\ D_t & (r + s = t) \end{cases}$$

and

$$\begin{aligned} L_1 \cap \text{Ad}(a)L_2 &= W(R_1)J = W(R_2)J \cap \mathfrak{a}, \\ \#(L_1 \cap \text{Ad}(a)L_2) &= 2r, \quad \#(W(R_2)J) = 2 \min\{r + s, t\}. \end{aligned}$$

Proof. We can verify $\tilde{\Sigma}$, R_1 and R_2 by (5) of the table in Subsection 4.2. Theorem 4.19 and Example 4.2 imply $\#(L_1 \cap \text{Ad}(a)L_2) = 2^r$. If $r + s \neq t$, then $\#(W(R_2)J) = 2 \min\{r + s, t\}$ by Example 4.2. If $r + s = t$, then $J = J_1$ in Example 4.5, since the multiplicity of any root in R_2 is equal to 1. Thus $\#(W(R_2)J) = 2t$ by Example 4.5. \square

Example 4.28. If

$$(M, L_1, L_2) = (SO(4m)/U(2m), U(2m)/Sp(m), SO(2m)),$$

then $\tilde{\Sigma} = R_1 = C_m$, $R_2 = D_m$ and

$$\begin{aligned} L_1 \cap \text{Ad}(a)L_2 &= W(R_1)J = W(R_2)J \cap \mathfrak{a}, \\ \#(L_1 \cap \text{Ad}(a)L_2) &= 2^m, \quad \#(W(R_2)J) = 2^{2m+1}. \end{aligned}$$

Proof. We can verify $\tilde{\Sigma}$, R_1 and R_2 by (3) of the table in Subsection 4.2. Theorem 4.19 and Example 4.3 imply $\#(L_1 \cap \text{Ad}(a)L_2) = 2^m$. Since the multiplicity of any root in R_2 is equal to 1, (4.5) implies that $J = J_2$ in Example 4.5. Thus $\#(W(R_2)J) = 2^{2m+1}$ by Example 4.5. \square

Examples 4.20, \dots , 4.28 exhaust all two real forms L_1 and L_2 which are not congruent to each other in an irreducible Hermitian symmetric space M of compact type. For each real form L in M we can determine

the cardinality of $L \cap \text{Ad}(a)L$ described in Theorem 4.14 by the use of Examples in this section. The result is as follows.

M	L	$\#(L \cap \text{Ad}(a)L)$	Example
$G_k(\mathbb{C}^n)$	$G_k(\mathbb{R}^n)$	$\binom{n}{k}$	4.16
$G_{2k}(\mathbb{C}^{2n})$	$G_k(\mathbb{H}^n)$	$\binom{n}{k}$	4.21
$G_n(\mathbb{C}^{2n})$	$U(n)$	2^n	4.26
$Q_n(\mathbb{C})$	$S^{k,n-k}$	$2k + 2$	4.27
$SO(4n)/U(2n)$	$U(2n)/Sp(n)$	2^n	4.28
$SO(2n)/U(n)$	$SO(n)$	2^{n-1}	4.17
$Sp(2n)/U(2n)$	$Sp(n)$	2^n	4.24
$Sp(n)/U(n)$	$U(n)/O(n)$	2^n	4.15
$E_6/S^1 \cdot Spin(10)$	$G_2(\mathbb{H}^4)/\mathbb{Z}_2$	3^3	4.20
$E_6/S^1 \cdot Spin(10)$	$F_4/Spin(9)$	3	4.20
$E_7/S^1 \cdot E_6$	$(SU(8)/Sp(4))/\mathbb{Z}_2$	$2^3 \cdot 7$	4.25
$E_7/S^1 \cdot E_6$	$S^1 \cdot E_6/F_4$	2^3	4.25

We explain how to determine $\#(L \cap \text{Ad}(a)L)$ in one line in the above list. We can similarly determine $\#(L \cap \text{Ad}(a)L)$ in the other lines. In the case where $M = E_6/S^1 \cdot Spin(10)$ and $L = G_2(\mathbb{H}^4)/\mathbb{Z}_2$, this real form is described as L_2 in Example 4.20. Theorem 4.14 and Example 4.20 show that $\#(L \cap \text{Ad}(a)L) = \#(W(R_2)J) = 3^3$.

5. THE FIXED POINT SET AND THE INTERSECTION

Let $L_1 = F(\tau_1, M)$ and $L_2 = F(\tau_2, M)$ be two real forms in an irreducible Hermitian symmetric space $M = \text{Ad}(G)J$ of compact type. Since $\text{Ad}(a)L_2 = F(a\tau_2a^{-1}, M)$ for $a \in G$, we have

$$L_1 \cap \text{Ad}(a)L_2 \subset F(\tau_1^{-1}a\tau_2a^{-1}, M).$$

In this section we study the relation between $L_1 \cap \text{Ad}(a)L_2$ and the fixed point set $F(\tau_1^{-1}a\tau_2a^{-1}, M)$ of a holomorphic isometry $\tau_1^{-1}a\tau_2a^{-1}$ when $L_1 \cap \text{Ad}(a)L_2$ is discrete. We may assume that $\tau_1\tau_2 = \tau_2\tau_1$ and a is in $\exp \mathfrak{a}$. After some preparations we will prove the following theorem.

Theorem 5.1. *When $L_1 \cap \text{Ad}(a)L_2$ is discrete, then*

$$F(\tau_1^{-1}a\tau_2a^{-1}, M) = (\mathfrak{a} \oplus \mathfrak{f}_0) \cap M, \quad F(\tau_1^{-1}a\tau_2a^{-1}, M) \cap \mathfrak{a} = L_1 \cap \text{Ad}(a)L_2.$$

Further

(1) *If \mathfrak{f}_0 is abelian, then*

$$F(\tau_1^{-1}a\tau_2a^{-1}, M) = W(\mathfrak{g}_{12})J.$$

(2) *If $\mathfrak{f}_0 = \{0\}$, then*

$$F(\tau_1^{-1}a\tau_2a^{-1}, M) = L_1 \cap \text{Ad}(a)L_2.$$

Lemma 5.2. *Set $a = \exp H$ for some $H \in \mathfrak{a}$.*

(1) *If $L_1 = L_2$, then*

$$F(\tau_1^{-1}a\tau_2a^{-1}, M) = \left(\mathfrak{a} \oplus \mathfrak{f}_0 \oplus \sum_{\substack{\lambda \in R^+ \\ \langle \lambda, H \rangle \in \pi\mathbb{Z}}} (\mathfrak{f}_\lambda \oplus \mathfrak{p}_\lambda) \right) \cap M.$$

(2) *If L_1 is not congruent to L_2 , then*

$$\begin{aligned} & F(\tau_1^{-1}a\tau_2a^{-1}, M) \\ &= \left(\mathfrak{a} \oplus \mathfrak{f}_0 \oplus \sum_{\substack{\lambda \in R^+ \\ \langle \lambda, H \rangle \in \pi\mathbb{Z}}} (\mathfrak{f}_\lambda \oplus \mathfrak{p}_\lambda) \oplus \sum_{\substack{\alpha \in W^+ \\ \langle \alpha, H \rangle \in \frac{\pi}{2} + \pi\mathbb{Z}}} V_\alpha^\perp(\mathfrak{f}_1 \cap \mathfrak{p}_2) \oplus V_\alpha^\perp(\mathfrak{f}_2 \cap \mathfrak{p}_1) \right) \cap M. \end{aligned}$$

Proof. We only give the proof of (2) since we can prove (1) in a similar manner to the proof of (2).

Since $\theta_i = -\tau_i$ on $M \subset \mathfrak{g}$ ([16]), we have

$$F(\tau_1^{-1}a\tau_2a^{-1}, M) = F(\text{Ad}(a^{-2})\theta_1\theta_2, M).$$

Since $\mathfrak{g} = \mathfrak{g}_{12} \oplus (\mathfrak{f}_1 \cap \mathfrak{p}_2) \oplus (\mathfrak{f}_2 \cap \mathfrak{p}_1)$, we have

$$\begin{aligned} & F(\text{Ad}(a^{-2})\theta_1\theta_2, \mathfrak{g}) \\ &= F(\text{Ad}(a^2), \mathfrak{g}_{12}) \oplus F(-\text{Ad}(a^2), (\mathfrak{f}_1 \cap \mathfrak{p}_2) \oplus (\mathfrak{f}_2 \cap \mathfrak{p}_1)). \end{aligned}$$

Lemma 4.12 implies that

$$F(\text{Ad}(a^2), \mathfrak{g}_{12}) = \mathfrak{a} \oplus \mathfrak{f}_0 \oplus \sum_{\substack{\lambda \in R^+ \\ \langle \lambda, H \rangle \in \pi\mathbb{Z}}} (\mathfrak{f}_\lambda \oplus \mathfrak{p}_\lambda).$$

Lemma 4.13 implies that

$$\begin{aligned} & F(-\text{Ad}(a^2), (\mathfrak{f}_1 \cap \mathfrak{p}_2) \oplus (\mathfrak{f}_2 \cap \mathfrak{p}_1)) \\ &= \sum_{\substack{\alpha \in W^+ \\ \langle \alpha, H \rangle \in \frac{\pi}{2} + \pi\mathbb{Z}}} V_\alpha^\perp(\mathfrak{f}_1 \cap \mathfrak{p}_2) \oplus V_\alpha^\perp(\mathfrak{f}_2 \cap \mathfrak{p}_1). \end{aligned}$$

Hence we get the assertion. \square

Proof of Theorem 5.1. The first part follows from Theorems 4.14, 4.18 and Lemma 5.2. In order to prove (1) we assume that \mathfrak{f}_0 is abelian. Then the maximality of \mathfrak{a} implies that $\mathfrak{a} \oplus \mathfrak{f}_0$ is a maximal abelian subalgebra of \mathfrak{g}_{12} . Thus (1) holds. (2) is clear from the first part of this theorem. \square

In the case where $L_1 = L_2$ we know the structure of \mathfrak{f}_0 by Tamaru's table([14]). In the case where L_1 is not congruent to L_2 we know the structure of \mathfrak{f}_0 by the table in Subsection 4.2.

Example 5.3. We consider a real form $L = U(n)$ in $M = G_n(\mathbb{C}^{2n})$. If $L \cap \text{Ad}(a)L$ is discrete, then

$$L \cap \text{Ad}(a)L = W(C_n)J, \quad F(\tau^{-1}a\tau a^{-1}, M) = W(\mathfrak{su}(2n))J.$$

In particular,

$$\#(L \cap \text{Ad}(a)L) = 2^n, \quad \#F(\tau^{-1}a\tau a^{-1}, M) = \binom{2n}{n}.$$

REFERENCES

- [1] R. Bott, The geometry and representation theory of compact Lie groups, Representation theory of Lie groups, 1979, 65–90, London Mathematical Society Lecture Note Series 34.
- [2] N. Bourbaki, Groupes et algebres de Lie, Hermann, Paris, 1975.
- [3] B.-Y. Chen and T. Nagano, A Riemannian geometric invariant and its applications to a problem of Borel and Serre, Trans. Amer. Math. Soc. **308** (1988), 273–297.
- [4] E. Heintze, R. S. Palais, C.-L. Terng and G. Thorbergsson, Hyperpolar actions on symmetric spaces, Geometry, topology, & physics, 214–245, Conf. Proc. Lecture Notes Geom. Topology, IV, Int. Press, Cambridge, MA, 1995.
- [5] S. Helgason, Differential Geometry, Lie groups, and Symmetric spaces, Academic Press, New York, 1978.
- [6] O. Ikawa, The geometry of symmetric triad and orbit spaces of Hermann actions, J. Math. Soc. Japan, **63** (2011) 79–136.
- [7] O. Ikawa, A note on symmetric triad and Hermann action, Proceedings of the workshop on differential geometry and submanifolds and its related topics, Saga, August 4–6, 2012, 220–229.
- [8] O. Ikawa, M. S. Tanaka and H. Tasaki, The fixed point set of a holomorphic isometry and the intersection of two real forms in the complex Grassmann manifold, Springer Proceedings in Mathematics & Statistics 106, Y.J. Suh et al. (eds.), ICM Satellite Conference on "Real and Complex Submanifolds", (2014), 319–327.
- [9] D. P. S. Leung, Reflective submanifolds. IV, Classification of real forms of Hermitian symmetric spaces, J. Differential Geom. **14** (1979), 179–185.
- [10] T. Matsuki, Classification of two involutions on compact semisimple Lie groups and root systems, Journal of Lie theory, **12** (2002), 41–68.
- [11] M. Takeuchi, On the fundamental group and the group of isometries of a symmetric space, J. Fac. Sci. Univ. Tokyo, Sect. 1 **10** (1964), 88–123.
- [12] M. Takeuchi, Stability of certain minimal submanifolds of compact Hermitian symmetric spaces, Tohoku Math. J., (2) **36** (1984), 293–314.
- [13] M. Takeuchi, Two-number of symmetric R -spaces, Nagoya Math. J., **115** (1989), 43–46.

- [14] H. Tamaru, The local orbit types of symmetric spaces under the action of the isotropy subgroups, *Differential geometry and its applications*, **11** (1999), 29–38.
- [15] M. S. Tanaka and H. Tasaki, The intersection of two real forms in Hermitian symmetric spaces of compact type, *J. Math. Soc. Japan* **64** (2012), 1297–1332.
- [16] M. S. Tanaka and H. Tasaki, Antipodal sets of symmetric R -spaces, *Osaka J. Math.* **50** (2013), 161–169.
- [17] M. S. Tanaka and H. Tasaki, The intersection of two real forms in Hermitian symmetric spaces of compact type II, *J. Math. Soc. Japan* **67** (2015), 275–291.
- [18] M. S. Tanaka and H. Tasaki, Correction to: “The intersection of two real forms in Hermitian symmetric spaces of compact type”, to appear in *J. Math. Soc. Japan*.

DEPARTMENT OF MATHEMATICS AND PHYSICAL SCIENCES, FACULTY OF ARTS
AND SCIENCES, KYOTO INSTITUTE OF TECHNOLOGY, SAKYOKU, KYOTO, 606-
8585 JAPAN

E-mail address: ikawa@kit.ac.jp

FACULTY OF SCIENCE AND TECHNOLOGY, TOKYO UNIVERSITY OF SCIENCE,
NODA, CHIBA, 278-8510 JAPAN

E-mail address: tanaka_makiko@ma.noda.tus.ac.jp

DIVISION OF MATHEMATICS, FACULTY OF PURE AND APPLIED SCIENCES, UNI-
VERSITY OF TSUKUBA, TSUKUBA, IBARAKI, 305-8571 JAPAN

E-mail address: tasaki@math.tsukuba.ac.jp