

Mechanism of dominance of the Breit interaction in dielectronic recombination

著者別名	全 曉民
journal or publication title	Journal of physics. B, Atomic, molecular and optical physics
volume	48
number	14
page range	144002
year	2015-07
権利	(C) 2015 IOP Publishing Ltd. This is an author-created, un-copied version of an article accepted for publication in Journal of Physics B: Atomic, Molecular and Optical Physics. The publisher is not responsible for any errors or omissions in this version of the manuscript or any version derived from it. The Version of Record is available online at doi:10.1088/0953-4075/48/14/144002.
URL	http://hdl.handle.net/2241/00126222

doi: 10.1088/0953-4075/48/14/144002

Mechanism of dominance of the Breit interaction in dielectronic recombination

Xiao-Min Tong^{1‡}, Zhimin Hu^{2,3,4}, Yueming Li⁵, Xiaoyin Han⁵, Daiji Kato⁶, Hirofumi Watanabe⁷, and Nobuyuki Nakamura²

¹ Graduate School of Pure and Applied Sciences, and Center for Computational Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8573, Japan

² Institute for Laser Science, The University of Electro-Communications, Chofu, Tokyo 182-8585, Japan

³ Research Center of Laser Fusion, China Academy of Engineering Physics, Mianyang 621900, China

⁴ Physics Institute, Heidelberg University, Im Neuenheimer Feld 226, 69120 Heidelberg, Germany

⁵ Institute of Applied Physics and Computational Mathematics, P. O. Box 8009, Beijing 100088, China

⁶ Department of Fusion Science, The Graduate University for Advanced Studies, Sokendai, Toki, Gifu 509-5292, Japan

⁷ Center of Applied Superconductivity and Sustainable Energy Research, Chubu University, Kasugai, Aichi 487-8501, Japan

Abstract. The recent theoretical and experimental studies show that the Breit interaction plays a dominant role in the dielectronic recombination for some particular transitions. The detailed mechanism of why the Breit interaction is dominant for such a process is still unknown. In this work, we performed a simulation and decomposed each individual term in the transition matrix level and found that the Breit interaction is dominant when the leading term ($1/r_{>}$ with $r_{>}$ the larger of r_1 and r_2) contribution of the two-electron Coulomb interaction is vanished. Based on this mechanism, we explained why the dielectronic capture strength to $1s2s^22p_{1/2} J_d = 1$ state is much stronger than the one to $1s2s2p_{1/2}^2 J_d = 1$ as well as why the Breit interaction plays a dominant role in the anisotropic parameters. Furthermore, the present finding may guide us to search some physical processes in which the Breit interaction is dominant by simply analyzing the coupling coefficients for a given isoelectronic sequence.

Keywords: Breit interaction, Coulomb interaction, dielectronic recombination, anisotropic parameter

Submitted to: *J. Phys. B: At. Mol. Phys.*

‡ email: tong@ims.tsukuba.ac.jp

37 1. Introduction

38 Since the discovery of the Breit interaction [1, 2], its contributions to the atomic
 39 structure have been studied extensively and systematically with various theoretical
 40 methods [3, 4, 5], also as reviewed by Grant [6]. Although the Breit interaction is
 41 getting important for high-Z atomic ions, its contribution to the total energy is still
 42 much smaller. Namely for nobelium (Z=102), the Breit contribution to the total
 43 energy is just of the order of 10^{-3} [7]. For total energy, the dominant contributor
 44 is the electron-nucleus Coulomb interaction, then the electron-electron Coulomb and
 45 Breit interactions. Thus, in most cases, the Breit interaction is treated perturbatively
 46 or simply ignored even for high-Z atoms [8]. The Breit interaction does not only
 47 contribute to the total energy but also to dynamical processes involving two-electron
 48 interactions, like Auger decay or its inverse process dielectronic capture (DC). For
 49 such a dynamical process, the Breit interaction directly competes with the electron-
 50 electron Coulomb interaction and in most cases, the Breit interaction may modify the
 51 process quantitatively not qualitatively. The Breit interaction plays an important role
 52 for high energy electrons colliding with a highly charged ion [9, 10, 11] as well.

53 The effect of the Breit interaction on dielectronic recombination (DR) has been
 54 studied in highly charged ions and it was found experimentally [12] that the Breit
 55 interaction is comparable to the Coulomb interaction in lithium-like ions [12] for the
 56 autoionization state $1s2s^22p_{1/2} J_d = 1$. Although the Breit interaction is important
 57 in such a DR process, the conclusion – the DC strength to $1s2s^22p_{1/2} J_d = 1$ state is
 58 much stronger than the one to $1s2s2p_{1/2}^2 J_d = 1$ state either with or without taking
 59 the Breit interaction into account – is still hold.

60 Later on, Fritzsche *et al.*, [13] found that the Breit interaction may change the
 61 angular distribution of the emitted X-ray in the lithium-like DR process involving the
 62 autoionization state $1s2s^22p_{1/2} J_d = 1$ in a theoretical study. For example, for lithium-
 63 Like Au ions, without the Breit interaction, the emitted X-ray is mainly along the
 64 electron beam direction while with the Breit interaction, the emitted X-ray is mainly
 65 along the perpendicular direction to the electron beam. This theoretical prediction
 66 was confirmed by experiment [14].

67 Now the question is why the Breit interaction plays such a dominant role in
 68 these processes or under what conditions the Breit interaction could be important.
 69 To answer these questions, we performed a systematic study on the DR processes
 70 involving the captures to $1s2s^22p_{1/2} J_d = 1$ and $1s2s2p_{1/2}^2 J_d = 1$ two states. Instead
 71 of just analyzing the DR rates and anisotropic parameters obtained in the simulation,
 72 we also analyzed each term of the transition matrix elements and found that the Breit
 73 interaction is important when the leading term ($1/r_>$ with $r_>$ the larger of r_1 and
 74 r_2 where \mathbf{r}_1 and \mathbf{r}_2 are the space coordinates of electron 1 and electron 2 involved in
 75 the DR transitions) contribution of the two-electron Coulomb interaction is vanished.
 76 We call this leading term the Coulomb monopole. Based on the mechanism, we can
 77 explain the experimental observations where the Breit interaction is important for
 78 both the DR rate and the anisotropic parameter.

79 2. Theory

80 Theory on atomic dielectronic recombination can be found in many literatures
 81 [15, 16, 17, 18] and the DR rates can be calculated by commonly used atomic packages,
 82 like, GRASP (a general-purpose relativistic atomic structure package) [5]. Here we

83 only present the necessary equations which will be used for the discussion in this
84 paper. Atomic units $\hbar = m_e = e = 1$ are used throughout unless otherwise stated.

85 Dielectronic recombination involves two steps: a free electron is captured by an
86 atomic ion with exciting another inner-shell electron to an excited state and forming
87 an autoionization state, followed by the radiative stabilization (RS) by emitting a
88 photon from the autoionization state. The first step is an inverse process of Auger
89 decay. The Auger decay rate can be expressed as

$$R_A = 2\pi \sum_{lj} |\langle \Psi_{J_d} || V || \Psi_{J_i} \psi_{lj} \rangle|^2 \quad (1)$$

90 with Ψ_{J_i} the wave function of the initial state for the N-electron system, Ψ_{J_d} the wave
91 function of the autoionization state of the (N+1)-electron system and ψ_{lj} the continue
92 electron wave function with angular moment l and total angular momentum j . V is
93 the two-electron interaction, which includes the electron-electron Coulomb and Breit
94 interactions. The autoionization state (or intermediate state) Ψ_{J_d} can decay to a final
95 state Ψ_{J_f} radiatively by emitting a photon. J_i, J_d, J_f are the total angular momenta
96 of the initial, intermediate and final states, respectively. The angular distribution of
97 the emitted photon is written as

$$\frac{d\sigma}{d\Omega} = \frac{\sigma_T}{4\pi} (1 + \beta P_2(\cos \theta)), \quad (2)$$

98 with σ_T the total DR cross section, θ the angle between the emitted photon and
99 the electron beam directions, and $P_2(\cos \theta)$ the second order Legendre polynomial.
100 Note here we focus on the dipole transition and the general expression for multipole
101 transitions can be found in Refs. [19, 20]. The anisotropic parameter β is given by

$$\beta = (-1)^{1+J_d+J_f} \left[\frac{3(2J_d+1)}{2} \right]^{1/2} \begin{Bmatrix} 1 & 1 & 2 \\ J_d & J_d & J_f \end{Bmatrix} \frac{P_{J_i J_d}^{(2)}}{P_{J_i J_d}^{(0)}}, \quad (3)$$

102 with

$$P_{J_i J_d}^{(L)} = \frac{(-1)^{J_i+L+J_d-1/2}}{2(2J_i+1)} \sum_{ij'j'l'l'} i^{l-l'} [j, j', l, l', L]^{1/2} \begin{bmatrix} l & l' & L \\ 0 & 0 & 0 \end{bmatrix} \begin{Bmatrix} j' & j & L \\ l & l' & 1/2 \end{Bmatrix} \\ \times \begin{Bmatrix} J_d & J_d & L \\ j & j' & J_i \end{Bmatrix} \langle \Psi_{J_d} || V || \Psi_{J_i} \psi_{lj} \rangle \langle \Psi_{J_i} \psi_{l'j'} || V || \Psi_{J_d} \rangle, \quad (4)$$

103 where the Wigner $3j, 6j$ symbols are used. Here we focus on two transition lines of lithium-like
104 isoelectronic ions. They are

$$\begin{aligned} \text{Line a: } & |1s^2 2s \psi_{lj} \rangle \xrightarrow{\text{DC}} |1s 2s^2 2p_{1/2}(J_d = 1) \rangle \xrightarrow{\text{RS}} |1s^2 2s^2 (J_f = 0) \rangle, \\ \text{Line b: } & |1s^2 2s \psi_{lj} \rangle \xrightarrow{\text{DC}} |1s 2s 2p_{1/2}^2 (J_d = 1) \rangle \xrightarrow{\text{RS}} |1s^2 2s 2p_{1/2} (J_f = 0, 1) \rangle. \end{aligned}$$

105 For Line b, due to the selection rule, the initial continue state can only be an s partial wave
106 and thus the anisotropic parameter β is zero. For Line a, the anisotropic parameter can be
107 explicitly expressed as

$$\beta = -\frac{1}{2} \frac{(2\sqrt{2}V_1 + V_2) \cdot V_2}{V_1^2 + V_2^2}, \quad (5)$$

108 with

$$\begin{aligned} V_1 &= V_{p_{1/2}} = \langle 1s 2s^2 2p_{1/2}(J_d = 1) || V_C + V_{Breit} || 1s^2 2s \epsilon p_{1/2} \rangle = V_1^C + V_1^B(6) \\ V_2 &= V_{p_{3/2}} = \langle 1s 2s^2 2p_{1/2}(J_d = 1) || V_C + V_{Breit} || 1s^2 2s \epsilon p_{3/2} \rangle = V_2^C + V_2^B(7) \end{aligned}$$




Figure 1. (Color online) Reduced matrix elements of dielectronic capture for Lines a and b. V_1^C (thick solid line), V_2^C (thick dashed line) and V^C (thick dotted line) are defined in Eqs. (6)-(8) with the Coulomb interaction only. V_1^B (thin solid line), V_2^B (thin dashed line) and V^B (thin dotted line) are defined in Eqs. (6)-(8) with the generalized Breit interaction only.

109 Here V_C, V_{Breit} are the electron-electron Coulomb and Breit interactions. Note that the
 110 anisotropic parameter depends on the DC process so we will analyze the DC transition
 111 matrix V_1, V_2 in detail. For the convenience of discussion, we also give the reduced transition
 112 matrix element for Line b as

$$V = V_{s_{1/2}} = \langle 1s2s2p_{1/2}^2(J_d = 1) || V_C + V_{Breit} || 1s^2s\epsilon s_{1/2} \rangle = V^C + V^B. \quad (8)$$

113 3. Results and discussion

114 We first calculated the single-electron wave function from the relativistic density functional
 115 theory with the self-interaction correction [21] and then calculated the reduced matrix
 116 elements. The coupling coefficients were calculated by using the ANCO package [22] and
 117 the detailed numerical method can be found in Ref. [23]. The reduced matrix elements were
 118 calculated using the single configuration approximation as well as configuration interactions
 119 and the two results are close to each other within two or three digits. Therefore to simplify
 120 the discussion, we only present the results from the single configuration simulation. We also
 121 present the results from the generalized Breit interaction (GBI) [24] and the Breit interaction
 122 (BI0) in the zero-frequency limit. To study lithium-like atomic ions systematically, we
 123 calculated all the ions with $Z = 40 - 100$. Note that to calculate the reduced matrix element,
 124 there is an arbitrary phase factor and we chose the phase factor in such a way that V_1^C is
 125 positive.

126 3.1. Reduced matrix elements for dielectronic capture

127 Figure 1 shows the reduced matrix elements for the Coulomb interaction only and the GBI
 128 defined in Eqs. (6)-(8). From a scaling law, the Coulomb interaction is scaled by atomic
 129 number Z and the GBI is scaled by (Z^3/c^2) with c the velocity of light. Indeed, V_1^C is
 130 the largest one among the transition matrix elements and it increases monotonically as Z
 131 increases. V_1^B increases rapidly and is comparable but still smaller than V_1^C even for very

Table 1. Coupling coefficients of the reduced matrix elements of dielectronic captures for Lines a and b.

Term	k	X^k	Term	k	X^k
$V_1 : (2s2p_{1/2} 1s\epsilon p_{1/2})$	0	-1/2	$V_2 : (2s2p_{1/2} 1s\epsilon p_{3/2})$	1	-1/3
$V_1 : (2s2p_{1/2} 1s\epsilon p_{1/2})$	1	-1/6	$V_2 : (2s2p_{1/2} \epsilon p_{3/2}1s)$	1	1/3
$V_1 : (2s2p_{1/2} \epsilon p_{1/2}1s)$	1	-1/3			
$V : (2p_{1/2}^2 1s\epsilon s)$	0	1/2	$V : (2p_{1/2}^2 1s\epsilon s)$	1	1/2

high Z ions. This means that for the DC strength, the Coulomb interaction is still a dominant contributor for Line a. V_2^B follows the trend of V_1^B but smaller than V_1^B and V^B is almost zero. Interestingly, V_2^C, V^C are almost constants and smaller than V_1^C . Especially for V_2^C , it has an opposite sign with V_2^B . The absolute value of V_2^B is smaller than V_2^C for low Z atomic ions and reaches the same magnitude at Z=73, then is larger than V_2^C for high-Z atomic ions. Overall, from the reduced transition matrix elements, we see that the Coulomb interaction is a dominant contributor to the DC process and the GBI contribution is comparable but still smaller than the one of the Coulomb interaction. The key step to understand why the Breit interaction plays a dominant role for anisotropic parameters is to understand why V_2^C is smaller.

Both the Coulomb and Breit interactions are two-electron operators and they can be evaluated as a product of coupling coefficients and radial integrals as

$$\langle \Psi_{J_d} || V_C || \Psi_{J_i} \phi_{lj} \rangle = \sum_k X^k \cdot R_C^k, \quad (9)$$

$$\langle \Psi_{J_d} || V_{\text{Breit}} || \Psi_{J_i} \phi_{lj} \rangle = \sum_k X^k \cdot R_B^k, \quad (10)$$

where X^k is the coupling coefficient which is independent from the atomic number Z and R_C^k, R_B^k are the radial integrals for an irreducible tensor operator of rank k . Note that we followed the convention of the ANCO program [22] and the coefficients are listed in Table 1. For the Coulomb interaction, R_C^k is decided by the integration of the operator

$$\begin{aligned} V_C &= \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} = \sum_k \left(\frac{r_{<}}{r_{>}} \right)^k \frac{1}{r_{>}} P_k(\cos \gamma) \\ &= \sum_k \left(\frac{r_{<}}{r_{>}} \right)^k \frac{1}{r_{>}} \frac{4\pi}{2k+1} \sum_m Y_{-m}^k(\hat{\mathbf{r}}_1) Y_m^k(\hat{\mathbf{r}}_2), \end{aligned} \quad (11)$$

where $r_{<}$ ($r_{>}$) stands for the smaller (larger) one of $|\mathbf{r}_1|$, $|\mathbf{r}_2|$, P_k the k -order of Legendre polynomials, γ the angle between $\mathbf{r}_1, \mathbf{r}_2$, and Y^k the k -order of spherical harmonics. R_C^k can be written as

$$R_C^k \propto \langle a || Y^k || c \rangle \langle b || Y^k || d \rangle, \quad (12)$$

where a, b, c, d are the four orbits involved in the two-electron operator as $(ab|cd)$ in Table 1. For example, the Coulomb contribution of V_2 can be evaluated as

$$\begin{aligned} V_2^C &\propto -\frac{1}{3} \langle 2s || Y^{k=1} || 1s \rangle \langle 2p_{1/2} || Y^{k=1} || \epsilon p_{3/2} \rangle \\ &\quad + \frac{1}{3} \langle 2s || Y^{k=1} || \epsilon p_{3/2} \rangle \langle 2p_{1/2} || Y^{k=1} || \epsilon 1s \rangle \end{aligned} \quad (13)$$

$$= \frac{1}{3} \langle 2s || Y^{k=1} || \epsilon p_{3/2} \rangle \langle 2p_{1/2} || Y^{k=1} || \epsilon 1s \rangle. \quad (14)$$

The first term of right hand in Eq. (13) is zero because $\langle 2s || Y^{k=1} || 1s \rangle = 0$.

154 For the generalized Breit interaction [24], R_B^k is decided by the integration of the
155 operator

$$V_{\text{Breit}} = -\boldsymbol{\alpha}_1 \cdot \boldsymbol{\alpha}_2 \frac{\cos(\omega r)}{r} + (\boldsymbol{\alpha}_1 \cdot \nabla_1)(\boldsymbol{\alpha}_2 \cdot \nabla_2) \frac{\cos(\omega r) - 1}{\omega^2 r}, \quad (15)$$

156 where $r = |\mathbf{r}_1 - \mathbf{r}_2|$ and ω is the virtual photon energy divided by c , and $\boldsymbol{\alpha}$ the Dirac matrixes.
157 If we choose $\omega = 0$, we get the Breit interaction (BI0) in the zero-frequency limit. Similar to
158 R_C^k , R_B^k can be evaluated as

$$R_B^k \propto \langle a || T^k || c \rangle \langle b || T^k || d \rangle, \quad (16)$$

159 with T^k the tensor of rank k . The general expression of T^k is complicate and can be found
160 in Ref. [3] for details. The tensor operator of the first term of right hand in Eq. (15) can be
161 written

$$T^k = \sum_{k'} \boldsymbol{\alpha} Y^{k'} \quad (17)$$

162 as a vector spherical harmonics of order of k [25]. For example, the Breit contribution to
163 V_2 involves the tensor product of $\boldsymbol{\alpha}$ (tensor of rank 1) with the space tensor $Y^{k'}$ (spherical
164 harmonics) and it dose not vanish for $Y^{k'=0}$. This should be the largest one for the Breit
165 interaction.

166 Generally speaking, for Coulomb interaction, the high order (k) multipole contributions
167 are getting smaller because $(r_</r_>)^k$ decreases as k increases. For V_1^C , the monopole ($k = 0$)
168 contribution is not vanished and it is the major contributor to the DC transition matrix
169 elements as shown in Table 1. For V_2^C , the monopole is vanished and this results in smaller
170 value of V_2^C . Although the coupling coefficient of V is not zero for $k = 0$, the reduced matrix
171 element of the Coulomb interaction is zero because for a single electron irreducible tensor
172 operator of rank 0, $s \rightarrow p_{1/2}$ ($\langle 2p_{1/2} || Y^0 || 1s \rangle$) matrix element is zero due to the parity
173 conservation. For Breit interaction, the leading contribution should be the transition matrix
174 elements with the tensor of rank $k = 1$ with the spherical harmonic of $k' = 0$. From above
175 analysis and the simulation data shown in Fig. 1, we conclude that the relative weak strength
176 of Line b comparing with Line a is due to the lack of Coulomb monopole contributions.

177 3.2. Ratio of the dielectronic capture strengths

178 Since directly measuring the anisotropic parameters is still difficult in the Tokyo EBIT
179 (electron beam ion trap) [26], the anisotropic parameter was extracted from the measured
180 ratio of DC strengths of Lines a and b, and the emitted X-ray at 90° to the electron beam.
181 Therefore we first analyze the ratio of DC strengths between Lines a and b as shown in Fig. 2.
182 Without the Breit interaction, the ratio increases slowly from 3 at $Z=40$ to 9 at $Z=100$. With
183 the BI0, the ratio increases rapidly from 3.5 at $Z=40$ to 23 at $Z=100$. With the GBI, the
184 ratio is slightly higher than the one of the BI0. The results show that the Breit interaction
185 enhanced the ratio by more than a factor of two for a very high Z atomic ion, while in Fig. 1
186 V_1^C is always the largest one. The ratio enhanced by a factor of two does not mean the Breit
187 interaction is the dominant one because the DC strength is proportional to the square of the
188 reduced matrix elements $|V_1^C + V_1^B|^2$ and if V_1^B is just about 50% of the V_1^C , the DC strength
189 can be doubled. The results are consistent with the experiments [12, 14, 27], which showed
190 that the Breit interaction enhances the ratio by a factor of two. Although the GBI modifies
191 the Ratio slightly over the BI0, the ratio of the GBI is more closer to the measured ones.
192 Discrepancies between the simulation and measurement may call further theoretical studies
193 as well as experiments with better resolution.

194 3.3. Dominance of the Breit interaction for anisotropic parameters

195 The reduced matrix elements in Fig. 1 show that the Coulomb interaction is *always* the
196 dominant one for the total DC strengths. For anisotropic parameters, the Breit interaction




Figure 2. (Color online) Ratios of dielectronic capture strengths of Lines a and b. Solid line: the theoretical results with the generalized Breit interaction, dashed line: the theoretical results with the Breit interaction in the zero-frequency limit, and dotted line: the theoretical results without the Breit interaction. The data of Exp.(a) are from Ref. [27] and the ones of Exp.(b) are from Ref. [12].

197 may change the angular distribution of the emitted X-ray from peaked on the forward
 198 direction along the electron beam ($\beta > 0$) with the Coulomb interaction only to peaked
 199 on the perpendicular direction to the electron beam ($\beta < 0$) with the Breit interaction for
 200 high-Z ions as shown in Fig. 3. To understand the mechanism, we approximate the anisotropic
 201 parameter in Eq. (5) as

$$\beta \approx -\sqrt{2} \frac{V_2}{V_1}, \quad (18)$$

202 because V_1 is positive and always larger than the value of $|V_2|$ as shown in Fig. 1. In the figure,
 203 we see that the Breit interaction for V_2 is comparable with the Coulomb interaction for low-Z
 204 ions and is larger than the Coulomb interaction for high Z ions with $Z > 72$. This is mainly
 205 because of the vanishing of the Coulomb monopole for V_2 . For this particular transition, Line
 206 a, the anisotropic parameter is almost linearly proportional to the reduced matrix element
 207 V_2 , in which the Breit interaction plays a dominant role for high Z ions. Our results of the
 208 Coulomb only (dash-dotted line) and Coulomb plus BI0 (dotted line) are in good agreements
 209 with the results (open circles and filled circles) from Fritzsche *et al.*, [13]. Again, we see that
 210 the BGI modifies the anisotropic parameters quantitatively not qualitatively over the BI0.
 211 Both the BI0 and GBI's results are in reasonable agreement with the available measurements
 212 [14, 27].

213 4. Summary and Conclusions

214 To summarize, we have investigated the mechanism of the dominance of the Breit interaction
 215 in the angular distribution of the emitted X-ray in dielectronic recombination of lithium-
 216 like isoelectronic ions for $Z = 40 - 100$. By analyzing the reduced matrix elements in
 217 the dielectronic capture when off-diagonal elements are involved, we found that the Breit
 218 interaction could be important when the Coulomb monopole interaction is vanished. Based
 219 on the mechanism, we have explained the anomalous Breit interaction enhanced ratio of
 220 dielectronic capture strengths for Lines a and b as well as the dominance of the Breit




Figure 3. (Color online) Anisotropic parameters of Line a. Solid line (C+GBI): the theoretical results with the generalized Breit interaction, dashed line (C+BI0): the theoretical results with the Breit interaction in the zero-frequency limit, dash-dotted line (Coulomb): the theoretical results with the Coulomb interaction only; filled and open circles: from Ref. [13]. The experimental data of Pr(Z=59) and Ho(Z=67) are from ref. [27] and the datum of Au(Z=79) is from ref. [14].

221 interaction in the angular distribution of the emitted X-ray in dielectronic recombination.
 222 By comparing the results of the generalized Breit interaction and the Breit interaction and
 223 the zero-frequency limit, we conclude that the modification of GBI over the BI0 is smaller
 224 and can be neglected in most cases. The present work may provide an effective way to
 225 search a physical process in which the Breit interaction is important simply by examining the
 226 coupling coefficients which are the same for a given isoelectronic sequence under the single
 227 configuration approximation.

228 5. Acknowledgments

229 XMT was supported by a Grand-in-Aid for Scientific Research (C24540421) from the Japan
 230 Society for the Promotion of Science. ZH acknowledges the support of the GermanResearch
 231 Foundation (DFG) within the Emmy Noether program under Contract No. TA 740/1-1. DK
 232 was supported by JSPS-NRF-NSFC A3 Foresight Program in the field of Plasma Physics
 233 (NSFC: No.11261140328, NRF: 2012K2A2A6000443) .

234 References

- 235 [1] Breit G 1930 *Phys. Rev.* **36** 383–397
 236 [2] Breit G 1932 *Phys. Rev.* **39** 616–624
 237 [3] Grant I P and Pyper N C 1976 *J. Phys. B: At. Mol. Phys.* **9** 761–774
 238 [4] Sapirstein J 1998 *Rev. Mod. Phys.* **70** 55–76
 239 [5] Jonsson P, He X, Fischer C F and and I P G 2007 *Comput. Phys. Commun.* **177** 597–622
 240 [6] Grant I P 2010 *J. Phys. B: At. Mol. Opt. Phys.* **43** 074033
 241 [7] Mann J B and Johnson W R 1971 *Phys. Rev. A* **4** 41–51
 242 [8] Karasiev V V, Ludena E V and Shukruto O A 2004 *Phys. Rev. A* **69** 052509
 243 [9] Marrs R E, Elliott S R and Scofield J H 1997 *Phys. Rev. A* **56** 1338–1345
 244 [10] Sampson D H, Zhang H L and Fontes C J 2009 *Phys. Rep.* **477** 111 – 214

- 245 [11] Gumberidze A, Thorn D B, Fontes C J, Najjari B, Zhang H L, Surzhykov A, Voitkiv A, Fritzsche
246 S, Banaś D, Beyer H, Chen W, DuBois R D, Geyer S, Grisenti R E, Hagmann S, Hegewald
247 M, Hess S, Kozhuharov C, Martin R, Orban I, Petridis N, Reuschl R, Simon A, Spillmann U,
248 Trassinelli M, Trotsenko S, Weber G, Winters D F A, Winters N, Yu D and Stöhlker T 2013
249 *Phys. Rev. Lett.* **110** 213201
- 250 [12] Nakamura N, Kavanagh A P, Watanabe H, Sakaue H A, Li Y, Kato D, Currell F J and Ohtani
251 S 2008 *Phys. Rev. Lett.* **100** 073203
- 252 [13] Fritzsche S, Surzhykov A and Stöhlker T 2009 *Phys. Rev. Lett.* **103** 113001
- 253 [14] Hu Z, Han X, Li Y, Kato D, Tong X and Nakamura N 2012 *Phys. Rev. Lett.* **108**(7) 073002
254 URL <http://link.aps.org/doi/10.1103/PhysRevLett.108.073002>
- 255 [15] Inal M K and Dubau J 1987 *J. Phys. B: At. Mol. Phys.* **20** 4221
- 256 [16] Chen M H, Reed K J, Guo D S and Savin D W 1998 *Phys. Rev. A* **58** 4539–4547
- 257 [17] Qu Y Z, Wang J G, Yuan J K and Li J M 1998 *Phys. Rev. A* **57** 1033–1037
- 258 [18] Fritzsche S, Kabachnik N M and Surzhykov A 2008 *Phys. Rev. A* **78** 032703
- 259 [19] Berezhko E G and Kabachnik N M 1977 *J. Phys. B: At. Mol. Phys.* **10** 2467
- 260 [20] Tseng H K, Pratt R H, Yu S and Ron A 1978 *Phys. Rev. A* **17** 1061–1079
- 261 [21] Tong X M and Chu S I 1998 *Phys. Rev. A* **57**(1) 452–461 URL <http://link.aps.org/doi/10.1103/PhysRevA.57.452>
- 262 [22] Gaigalas G and Fritzsche S 2002 *Comput. Phys. Commun.* **148** 349–351
- 263 [23] Tong X M, Nakamura N, Ohtani S, Watanabe T and Toshima N 2009 *Phys. Rev. A* **80**(4) 042502
264 URL <http://link.aps.org/doi/10.1103/PhysRevA.80.042502>
- 265 [24] Grant I P and McKenzie B J 1980 *J. Phys. B: At. Mol. Phys.* **13** 2671–2681
- 266 [25] Tong X M, Li J M, Kissel L and Pratt R H 1990 *Phys. Rev. A* **42**(3) 1442–1449 URL
267 <http://link.aps.org/doi/10.1103/PhysRevA.42.1442>
- 268 [26] Nakamura N, Asada J, Currell F J, Fukami T, Hirayama T, Motohashi K, Nagata T, Nojikawa
269 E, Ohtani S, Okazaki K, Sakurai M, Shiraiishi H, Tsurubuchi S and Watanabe H 1997 *Phys.*
270 *Scr. T.* **73** 362–364
- 271 [27] Hu Z, Li Y, Han X, Kato D, Tong X, Watanabe H and Nakamura N 2014 *Phys. Rev. A* **90**
272 062702
273