

# Ni-Catalyzed Synthesis of Fluoroarenes via [2+2+2] Cycloaddition Involving $\alpha$ -Fluorine Elimination

著者別名	藤田 健志, 市川 淳士
journal or publication title	Chemistry : a European journal
volume	21
number	38
page range	13225-13228
year	2015-09
権利	(C) 2015 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim This is the accepted version of the following article: Chemistry 21(38):13225-8, 2015, which has been published in final form at DOI:10.1002/chem.201502744
URL	<a href="http://hdl.handle.net/2241/00129781">http://hdl.handle.net/2241/00129781</a>

doi: 10.1002/chem.201502744

# Ni-Catalyzed Synthesis of Fluoroarenes via [2+2+2] Cycloaddition Involving $\alpha$ -Fluorine Elimination

Takeshi Fujita, Yota Watabe, Tomohiro Ichitsuka, and Junji Ichikawa\*


**Abstract:** A method for direct synthesis of tetrasubstituted fluoroarenes via nickel-catalyzed [2+2+2] cycloaddition is presented. The reaction combines one molecule of 1,1-difluoroethylene with two molecules of alkynes and involves sequential cleavage of the C–F and C–H bonds in difluoroethylene. The catalytic cycle is established by reduction of the intermediary Ni(II) fluoride with a triethylborane-based borate.

Application of fluoroarenes in the fields of pharmaceutical and materials sciences has resulted in considerable efforts toward development of methods for their synthesis.<sup>[1]</sup> Traditionally, fluoroarenes are synthesized via decomposition of aryl diazonium tetrafluoroborates (the Balz–Schiemann reaction)<sup>[1b,2]</sup> or nucleophilic substitution of electron-deficient aryl halides with fluoride salts (the Halex reaction).<sup>[1b,3]</sup> Recent developments in fluorinating reagents have given rise to other alternatives for the synthesis of fluoroarenes, including (i) reactions of aryl nucleophiles with electrophilic fluorine sources<sup>[1b,4]</sup> and (ii) reactions of aryl electrophiles with nucleophilic fluorine sources.<sup>[5]</sup> Furthermore, the use of transition metal catalysts has enabled directing group-assisted aromatic C–H bond fluorination,<sup>[1b,c,e,6]</sup> fluorination of arylmetals,<sup>[1c,7]</sup> and fluorination of nonactivated aryl halides (Scheme 1a).<sup>[1b–e,8]</sup> Thus, conventional methods for synthesizing fluoroarenes require regioselective prefunctionalization on benzene rings.


Recently, we reported the nickel-mediated [3+2] cycloaddition of 2-trifluoromethyl-1-alkenes and alkynes.<sup>[9]</sup> During this cycloaddition, two C–F bonds are cleaved via  $\beta$ -fluorine elimination, a characteristic feature of fluorine-containing organometallic compounds and two C–C bonds are constructed instead.<sup>[9,10]</sup> Despite the loss of two fluorine substituents from the trifluoromethyl group, the surviving fluorine substituent is regioselectively installed in the final cyclopentadiene products. On the other hand, when 1,1-difluoroethylene was reacted as a fluorinated component with alkynes in the presence of Ni(0) complex, fluoroarenes were directly obtained. Here, we demonstrate that the combination of 1,1-difluoroethylene and alkynes via nickel-catalyzed [2+2+2] cycloaddition is an alternative method for the synthesis of fluoroarenes (Scheme 1b).<sup>[11,12]</sup> Although transition-metal-catalyzed [2+2+2] cycloaddition is a facile method for the construction of benzene ring from three alkyne molecules,<sup>[11]</sup> similar syntheses involving

alkene substrates require an additional oxidation<sup>[12a,c]</sup> or elimination<sup>[13]</sup> step to ensure aromatization. By contrast, our protocol allows for direct construction of benzene ring through nickel-promoted  $\alpha$ -fluorine elimination<sup>[14,15]</sup> as the key elementary step.

## (a) Conventional methods



## (b) This work




**Scheme 1.** Transition-metal-catalyzed syntheses of fluoroarenes.

First, we sought suitable conditions for the [2+2+2] cycloaddition reaction of 1,1-difluoroethylene (**1**, 2.3 mmol) and diphenylacetylene (**2a**, 0.50 mmol) in the presence of a catalytic amount of a Ni(0) complex (Table 1). The choice of ligands used with Ni(cod)<sub>2</sub> (5 mol% based on the amount of **2a**) was critical for the efficiency of the reaction. Use of IMes·HCl with KH (5 mol% each) or P(*t*Bu)<sub>3</sub> (10 mol%) afforded 1-fluoro-2,3,4,5-tetraphenylbenzene (**3a**), albeit in low yields (Entries 2 and 4). Among the ligands examined, PCy<sub>3</sub> (5 mol%) was found to be the best (Entry 6). No improvement in reaction yields was observed using bases, such as Hünig's base and DBU, which suggested that fluoroarene **3a** might be formed directly and not via a HF elimination after the formation of **4a** (Entries 9 and 10). Addition of Et<sub>3</sub>SiH, *i*PrOLi, or *i*PrOBpin (Entries 11–13) was only marginally effective in increasing the yield of the reaction. However, Et<sub>3</sub>B indicated potential for regeneration of the nickel catalyst (Entry 14). To activate Et<sub>3</sub>B, *i*PrOLi was added to improve the turnover number of the Ni(0) complex significantly (Entry 15). Finally, the use of equimolar quantities of **1** and **2a** afforded **3a** in 82% isolated yield (Entry 16).

**Table 1.** Optimization of Reaction Conditions for the Ni-Catalyzed [2+2+2] Cycloaddition<sup>[a]</sup>

[\*] Dr. T. Fujita, Y. Watabe, Dr. T. Ichitsuka, Prof. Dr. J. Ichikawa  
Division of Chemistry  
Faculty of Pure and Applied Sciences  
University of Tsukuba, Tsukuba, Ibaraki 305-8571 (Japan)  
Fax: (+81) 29-853-4237  
E-mail: junji@chem.tsukuba.ac.jp

Supporting information for this article is given via a link at the end of the document.



Entry	Ligand (mol%)	Additive	Time [h]	<b>3a</b> [%] <sup>[c]</sup>
1	–	–	14	N.D. <sup>[b]</sup>
2	IMes·HCl (5) <sup>[d]</sup>	–	14	4
3	dppp (5)	–	22	N.D. <sup>[b]</sup>
4	P( <i>t</i> Bu) <sub>3</sub> (10)	–	14	3
5	PCy <sub>3</sub> (10)	–	22	9
6	PCy <sub>3</sub> (5)	–	22	8
7	PCy <sub>3</sub> (5)	LiOAc	22	9
8	PCy <sub>3</sub> (5)	Et <sub>3</sub> N	22	7
9	PCy <sub>3</sub> (5)	<i>i</i> Pr <sub>2</sub> NEt	7	10
10	PCy <sub>3</sub> (5)	DBU	7	9
11	PCy <sub>3</sub> (5)	Et <sub>3</sub> SiH	9	20
12	PCy <sub>3</sub> (5)	<i>i</i> PrOLi	15	37
13	PCy <sub>3</sub> (5)	<i>i</i> PrOBpin	31	26
14	PCy <sub>3</sub> (5)	Et <sub>3</sub> B	22	44
15	PCy <sub>3</sub> (5)	Et <sub>3</sub> B + <i>i</i> PrOLi	12	83 (80)
16 <sup>[e]</sup>	PCy <sub>3</sub> (5)	Et <sub>3</sub> B + <i>i</i> PrOLi	12	84 (82)

[a] Molar percentages of Ni(cod)<sub>2</sub>, ligands, and additives are based on the amount of **2a**. Reaction conditions: Ni(cod)<sub>2</sub> (0.025 mmol), **1** (2.3 mmol), **2a** (0.50 mmol), and toluene (2.0 mL). [b] N.D. = Not detected. [c] Yield was determined by <sup>19</sup>F NMR measurement using PhCF<sub>3</sub> as an internal standard. Isolated yield is shown in parentheses. [d] KH (5 mol%) was added. [e] **2a** (2.3 mmol).

With the optimal conditions in hand, the scope of the reaction was investigated using various alkynes (Table 2). Diarylacetylenes **2b–e** bearing electron-donating groups (*m*-Me, *p*-Me, *p*-Bu, and *p*-OMe) underwent cycloaddition effectively to afford the corresponding tetraarylated fluorobenzenes **3b–e** in 72%, 81%, 79%, and 80% yields (Entries 2–5), respectively. The reaction of diarylacetylene **2f** bearing electron-withdrawing CF<sub>3</sub> groups also proceeded to give **3f** in 76% yield (Entry 6). Chlorine-substituted diarylacetylene **2g** underwent catalytic cycloaddition without loss of Cl groups (Entry 7). Aliphatic alkyne **2h** participated in the reaction to provide tetraalkylated fluorobenzene **3h** in 79% yield (Entry 8). Ester, benzyl ether, acetal, and silyl ether moieties on dialkylacetylenes **2i–l** were tolerant in this reaction, which effectively afforded the corresponding fluorobenzenes **3i–l** (Entries 9–12). The cycloaddition of unsymmetrical alkynes **2m** and **2n** proceeded with substantial regioselectivities (84:16 and 85:15) to afford *o*-

terphenyl derivatives **3m** and **3n** as major products (Entries 13 and 14), respectively.<sup>[16,17]</sup>


**Table 2.** Ni-Catalyzed Synthesis of Fluoroarenes **3** from 1,1-Difluoroethylene (**1**) and Alkynes **2**<sup>[a]</sup>

Entry	<b>2</b>	R <sup>1</sup> , R <sup>2</sup>	Time [h]	<b>3</b>	Yield [%]
1	<b>2a</b>	Ph, Ph	12	<b>3a</b>	82
2	<b>2b</b>	C <sub>6</sub> H <sub>4</sub> ( <i>m</i> -Me), C <sub>6</sub> H <sub>4</sub> ( <i>m</i> -Me)	16	<b>3b</b>	72
3	<b>2c</b>	C <sub>6</sub> H <sub>4</sub> ( <i>p</i> -Me), C <sub>6</sub> H <sub>4</sub> ( <i>p</i> -Me)	12	<b>3c</b>	81
4	<b>2d</b>	C <sub>6</sub> H <sub>4</sub> ( <i>p</i> -Bu), C <sub>6</sub> H <sub>4</sub> ( <i>p</i> -Bu)	12	<b>3d</b>	79
5	<b>2e</b>	C <sub>6</sub> H <sub>4</sub> ( <i>p</i> -OMe), C <sub>6</sub> H <sub>4</sub> ( <i>p</i> -OMe)	12	<b>3e</b>	80
6	<b>2f</b>	C <sub>6</sub> H <sub>4</sub> ( <i>p</i> -CF <sub>3</sub> ), C <sub>6</sub> H <sub>4</sub> ( <i>p</i> -CF <sub>3</sub> )	15	<b>3f</b>	76
7	<b>2g</b>	C <sub>6</sub> H <sub>4</sub> ( <i>m</i> -Cl), C <sub>6</sub> H <sub>4</sub> ( <i>m</i> -Cl)	12	<b>3g</b>	45
8	<b>2h</b>	Pr, Pr	15	<b>3h</b>	79
9	<b>2i</b>	(CH <sub>2</sub> ) <sub>2</sub> CO <sub>2</sub> <i>t</i> Bu, (CH <sub>2</sub> ) <sub>2</sub> CO <sub>2</sub> <i>t</i> Bu	16	<b>3i</b>	39
10	<b>2j</b>	(CH <sub>2</sub> ) <sub>3</sub> OCH <sub>2</sub> Ph, (CH <sub>2</sub> ) <sub>3</sub> OCH <sub>2</sub> Ph	14	<b>3j</b>	58
11	<b>2k</b>	(CH <sub>2</sub> ) <sub>3</sub> OTHP, (CH <sub>2</sub> ) <sub>3</sub> OTHP	14	<b>3k</b>	67
12	<b>2l</b>	(CH <sub>2</sub> ) <sub>3</sub> OSiMe <sub>3</sub> , (CH <sub>2</sub> ) <sub>3</sub> OSiMe <sub>3</sub>	14	<b>3l</b>	62
13	<b>2m</b>	Me, Ph	18	<b>3m</b>	59 <sup>[b]</sup>
14	<b>2n</b>	Pr, C <sub>6</sub> H <sub>4</sub> ( <i>p</i> -OMe)	14	<b>3n</b>	60 <sup>[c]</sup>

[a] Molar percentages of Ni(cod)<sub>2</sub>, PCy<sub>3</sub>, Et<sub>3</sub>B, and *i*PrOLi are based on the amount of **2**. Reaction conditions: Ni(cod)<sub>2</sub> (0.12 mmol), PCy<sub>3</sub> (0.12 mmol), **1** (2.3 mmol), **2** (2.3 mmol), Et<sub>3</sub>B (2.3 mmol), *i*PrOLi (2.3 mmol), and toluene (4.7 mL). [b] Isolated yield. [c] The regioisomer ratio (84:16) was determined by <sup>19</sup>F NMR measurement. [d] The regioisomer ratio (85:15) was determined by <sup>19</sup>F NMR measurement.

To gain some insights into the reaction mechanism, the initial rate of the formation of product **3a** ((Δ[**3a**]/Δ*t*)<sub>*t*=0</sub>) was measured. We first monitored the dependency of (Δ[**3a**]/Δ*t*)<sub>*t*=0</sub> by changing the partial pressure of **1** (*p*(**1**); 0.3–1.0 atm). A linear correlation between *p*(**1**) and (Δ[**3a**]/Δ*t*)<sub>*t*=0</sub> (Figure 1a) was obtained. Furthermore, a straight line (slope = 1.12) provided a good fit for the log-log plot of (Δ[**3a**]/Δ*t*)<sub>*t*=0</sub> against *p*(**1**), indicating that the reaction has a nearly first-order dependence on the concentration of **1** in the solution (Figure 1b). Next, the dependency of the initial rate ((Δ[**3a**]/Δ*t*)<sub>*t*=0</sub>) on the initial concentration of alkyne **2a** ([**2a**]<sub>0</sub>; 0.2–0.9 M) under a constant pressure (1.0 atm) of **1** was examined. A linear correlation between the two was observed (Figure 1c). The linear fitting of the log-log plot with a slope of 1.02 shows a first-order dependence of the reaction rate on [**2a**]<sub>0</sub> (Figure 1d). Furthermore, the dependency of (Δ[**3a**]/Δ*t*)<sub>*t*=0</sub> on the initial


concentration of the Ni(0) complex ( $[\text{Ni}]_0$ ; 0.013–0.11 M) was estimated by reactions with a constant concentration (0.25 M) of **2a** under a constant pressure (1.0 atm) of **1** in the absence of  $\text{Et}_3\text{B}$  and  $i\text{PrOLi}$ . Not only a linear correlation between  $[\text{Ni}]_0$  and  $(\Delta[\mathbf{3a}]/\Delta t)_{t=0}$  (Figure 1e) but also the linear fitting of the log-log plot (slope: 1.09) clearly exhibits a first-order dependence of the reaction rate on  $[\text{Ni}]_0$  (Figure 1f). These results suggest that the initial rate-limiting oxidative cyclization proceeds with the involvement of one component each of **1**, **2a**, and Ni(0).<sup>[18]</sup>




**Figure 1.** (a) Initial reaction rate versus  $p(\mathbf{1})$  and (b) the corresponding log-log plot. Reaction conditions:  $\text{Ni}(\text{cod})_2$  (0.025 mmol),  $\text{PCy}_3$  (0.025 mmol), **1** (partial pressure: 0.30–1.0 atm), **2a** (0.50 mmol),  $\text{Et}_3\text{B}$  (0.50 mmol),  $i\text{PrOLi}$  (0.50 mmol), and toluene (1.0 mL) at 40 °C for 15 min. (c) Initial reaction rate versus  $[\mathbf{2a}]_0$  and (d) the corresponding log-log plot. Reaction conditions:  $\text{Ni}(\text{cod})_2$  (0.025 mmol),  $\text{PCy}_3$  (0.025 mmol), **1** (excess in balloon, 1 atm), **2a** (0.20–0.90 mmol),  $\text{Et}_3\text{B}$  (0.50 mmol),  $i\text{PrOLi}$  (0.50 mmol), and toluene (1.0 mL) at 40 °C for 15 min. (e) Initial reaction rate versus  $[\text{Ni}]_0$  and (f) the corresponding log-log plot. Reaction conditions:  $\text{Ni}(\text{cod})_2$  (0.013–0.11 mmol),  $\text{PCy}_3$  (0.013–0.11 mmol), **1** (excess in balloon, 1 atm), **2a** (0.25 mmol), and toluene (1.0 mL) at 40 °C for 5 min.

To elucidate the mechanism further, the stoichiometric reaction of 1,1-difluoro-1,6-enyne **5** and alkyne **2a** with Ni(0) was performed. Treatment of equimolar amounts of enyne **5** and alkyne **2a** with stoichiometric amounts of  $\text{Ni}(\text{cod})_2$  and  $\text{PCy}_3$  afford the ring-monofluorinated indane **6** in 60% yield [Eq. (1)]. Since one difluoroalkene moiety and two alkyne components were involved in the formation of **6**, we presumed that this reaction proceeded through a cycloaddition process similar to that in the case of the reaction involving **1** and **2**. The formation of **6** is likely to involve the initial intramolecular oxidative cyclization of difluoroalkene and alkyne moieties in **5** with Ni(0).<sup>[19,20]</sup> This cyclization mode would be consistent with the

forementioned result of the reaction between **1** and **2a**. Correspondingly, the oxidative cyclization of **1**, **2**, and Ni(0) probably affords the intermediate nickelacyclopentenes, wherein the difluoromethylene moiety regioselectively occupies the position  $\alpha$  to the nickel atom.




Taken together, these observations suggest a plausible mechanism for the Ni(0)-catalyzed cycloaddition (Scheme 2). This reaction starts with oxidative cyclization, rate-limiting chemo- and regioselective formation of nickelacyclopentenes **A**, resulting from the combination of one molecule each of **1** and **2**. The nickelacyclopentenes **A** thus formed facilitates the insertion of another molecule of **2** to generate nickelheptadienes **B**. Subsequent  $\alpha$ -fluorine elimination from **B** gives cyclohexadienylnickel(II) fluorides **C**.<sup>[14,15]</sup> Finally,  $\beta$ -hydrogen elimination affords fluoroarenes **3** and nickel(II) hydrofluoride **D**, which can then be reduced to Ni(0) complex via transmetalation with the borate derived from  $\text{Et}_3\text{B}$  and  $i\text{PrOLi}$ .<sup>[21]</sup>



**Scheme 2.** Plausible reaction mechanism.

The obtained tetraarylated fluoroarene products can serve as building blocks in further transformations. For example, treatment of fluoroarene **3d** with excess  $\text{FeCl}_3$  led to ring fusions via three oxidative C–H/C–H couplings<sup>[22]</sup> to afford tribenzoperylene **7** with the fluorine substituent in 74% yield [Eq. (2)].<sup>[23]</sup> The resulting pinpoint-fluorinated planar  $\pi$ -conjugated system can be a promising candidate as organic electronic material.<sup>[24]</sup>

In summary, we developed a nickel-catalyzed method for direct synthesis of fluoroarenes via  $\alpha$ -fluorine elimination. This method for fluoroarene synthesis complements conventional methods, which install fluorine on preformed benzene rings. With proper choice of alkyne substrates, our method enables modular synthesis of diversely substituted fluoroarenes<sup>[25]</sup> from 1,1-difluoroethylene, an industrial material.



## Experimental Section

Typical procedure for the synthesis of fluoroarenes **3** via nickel-catalyzed [2+2+2] cycloaddition: In an argon-purged 50-mL test tube equipped with a PTFE cap (EYELA, PPS25-TC) were placed *n*PrOH (179  $\mu$ L, 2.3 mmol) and toluene (2.4 mL). To the mixture was slowly added *n*BuLi (1.58 M in hexane, 1.47 mL, 2.32 mmol) at 0 °C. After stirring for 10 min at 0 °C, BEt<sub>3</sub> (1.0 M in hexane, 2.32 mL, 2.3 mmol) was added to the reaction mixture at the same temperature. The reaction mixture was warmed to room temperature, and was stirred for another 30 min. To the reaction mixture were added diphenylacetylene (**2a**, 414 mg, 2.32 mmol), Ni(cod)<sub>2</sub> (32 mg, 0.12 mmol), PCy<sub>3</sub> (33 mg, 0.12 mmol), and toluene (2.3 mL). The reaction vessel was evacuated, filled with 1,1-difluoroethylene (**1**, 1.0 atm, 56 mL, 2.3 mmol) through a balloon, and then sealed by closing the stopcock of the PTFE cap. After stirring for 12 h at 40 °C, the reaction mixture was filtered through a pad of silica gel (EtOAc). The filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/EtOAc = 20:1) to give fluoroarene **3a** as a white solid (380 mg, 82%).

## Acknowledgements

This work was supported by the Grant-in-Aid for Scientific Research (B) (No. 25288016) and the Grant-in-Aid for Young Scientists (B) (No. 26870079) from JSPS. This work was also supported by The Asahi Glass Foundation. We acknowledge Tosoh F-Tech, Inc. for generous gifts of 1,1-difluoroethylene and 3-bromo-3,3-difluoropropene.

**Keywords:** C–F bond activation • fluoroarenes • Ni • fluoroalkenes • alkynes

[1] For reviews, see: a) V. V. Grushin, *Acc. Chem. Res.* **2010**, *43*, 160–171; b) T. Furuya, J. E. M. N. Klein, T. Ritter, *Synthesis* **2010**, 1804–1821; c) T. Furuya, A. S. Kamlet, T. Ritter, *Nature* **2011**, *473*, 470–477; d) C. Hollingworth, V. Gouverneur, *Chem. Commun.* **2012**, *48*, 2929–2942; e) Y. Li, Y. Wu, G.-S. Li, X.-S. Wang, *Adv. Synth. Catal.* **2014**, *356*, 1412–1418; f) A. F. Brooks, J. J. Topczewski, N. Ichiishi, M. S. Sanford, P. J. H. Scott, *Chem. Sci.* **2014**, *5*, 4545–4553; g) M. G. Campbell, T. Ritter, *Chem. Rev.* **2015**, *115*, 612–633.

[2] a) A. Roe, *Org. React.* **1949**, *5*, 193–228; b) K. K. Laali, V. J. Gettewert, *J. Fluorine Chem.* **2001**, *107*, 31–34, and references cited therein.

[3] a) D. J. Adams, J. H. Clark, *Chem. Soc. Rev.* **1999**, *28*, 225–231; b) M.-A. Lacour, M. Zablocka, C. Duhayon, J.-P. Majoral, M. Taillefer, *Adv. Synth. Catal.* **2008**, *350*, 2677–2682, and references cited therein.

[4] For recent reports, see: a) C. Cazorla, E. Métay, B. Andrioletti, M. Lemaire, *Tetrahedron Lett.* **2009**, *50*, 3936–3938; b) S. Yamada, A. Gavryushin, P. Knochel, *Angew. Chem.* **2010**, *122*, 2261–2264; *Angew. Chem. Int. Ed.* **2010**, *49*, 2215–2218; c) P. Anbarasan, H. Neumann, M. Beller, *Angew. Chem.* **2010**, *122*, 2265–2268; *Angew. Chem. Int. Ed.* **2010**, *49*, 2219–2222.

[5] For recent reports, see: a) V. V. Grushin, W. J. Marshall, *Organometallics* **2008**, *27*, 4825–4828; b) B. Wang, L. Qin, K. D. Neumann, S. Uppaluri, R. L. Cerny, S. G. DiMango, *Org. Lett.* **2010**, *12*, 3352–3355; c) M. Saito, K. Miyamoto, M. Ochiai, *Chem. Commun.* **2011**, *47*, 3410–3412; d) P. Tang, W. Wang, T. Ritter, *J. Am. Chem. Soc.* **2011**, *133*, 11482–11484; e) H. Yoshida, R. Yoshida, K. Takaki, *Angew. Chem.* **2013**, *125*, 8791–8794; *Angew. Chem. Int. Ed.* **2013**, *52*, 8629–8632.

[6] a) K. L. Hull, W. Q. Anani, M. S. Sanford, *J. Am. Chem. Soc.* **2006**, *128*, 7134–7135; b) T. Truong, K. Klimovica, O. Daugulis, *J. Am. Chem. Soc.* **2013**, *135*, 9342–9345.

[7] P. Tang, T. Furuya, T. Ritter, *J. Am. Chem. Soc.* **2010**, *132*, 12150–12154.

[8] D. A. Watson, M. Su, G. Teverovskiy, Y. Zhang, J. Garcia-Fortanet, T. Kinzel, S. L. Buchwald, *Science* **2009**, *325*, 1661–1664.

[9] T. Ichitsuka, T. Fujita, T. Arita, J. Ichikawa, *Angew. Chem.* **2014**, *126*, 7694–7698; *Angew. Chem. Int. Ed.* **2014**, *53*, 7564–7568.

[10] For transition-metal-mediated reactions via  $\beta$ -fluorine elimination, see: a) W. Heitz, A. Knebelkamp, *Makromol. Chem. Rapid Commun.* **1991**, *12*, 69–75; b) M. Fujiwara, J. Ichikawa, T. Okauchi, T. Minami, *Tetrahedron Lett.* **1999**, *40*, 7261–7265; c) K. Sakoda, J. Mihara, J. Ichikawa, *Chem. Commun.* **2005**, 4684–4686; d) J. Ichikawa, K. Sakoda, J. Mihara, N. Ito, *J. Fluorine Chem.* **2006**, *127*, 489–504; e) J. Ichikawa, R. Nadano, N. Ito, *Chem. Commun.* **2006**, 4425–4427; f) T. Miura, Y. Ito, M. Murakami, *Chem. Lett.* **2008**, *37*, 1006–1007; g) M. Hu, Z. He, B. Gao, L. Li, C. Ni, J. Hu, *J. Am. Chem. Soc.* **2013**, *135*, 17302–17305; h) D. J. Harrison, G. M. Lee, M. C. Leclerc, I. Korobkov, R. T. Baker, *J. Am. Chem. Soc.* **2013**, *135*, 18296–18299; i) Z. Zhang, Q. Zhou, W. Yu, T. Li, G. Wu, T. Zhang, J. Wang, *Org. Lett.* **2015**, *17*, 2474–2477.

[11] For reviews on transition-metal-catalyzed [2+2+2] cycloaddition, see: a) M. Lautens, W. Klute, W. Tam, *Chem. Rev.* **1996**, *96*, 49–92; b) P. A. Inglesby, P. A. Evans, *Chem. Soc. Rev.* **2010**, *39*, 2791–2805; c) G. Domínguez, J. Pérez-Castells, *Chem. Soc. Rev.* **2011**, *40*, 3430–3444.

[12] For Ni-catalyzed [2+2+2] reactions between one molecule of alkenes and two molecules of alkynes, see: a) S.-i. Ikeda, N. Mori, Y. Sato, *J. Am. Chem. Soc.* **1997**, *119*, 4779–4780; b) S.-i. Ikeda, H. Watanabe, Y. Sato, *J. Org. Chem.* **1998**, *63*, 7026–7029; c) T. Sambaiyah, L.-P. Li, D.-J. Huang, C.-H. Lin, D. K. Rayabarapu, C.-H. Cheng, *J. Org. Chem.* **1999**, *64*, 3663–3670; d) Z. Qiu, Z. Xie, *Angew. Chem.* **2009**, *121*, 5839–5842; *Angew. Chem. Int. Ed.* **2009**, *48*, 5729–5732; e) H. Horie, T. Kurahashi, S. Matsubara, *Chem. Commun.* **2010**, *46*, 7229–7231.

[13] a) H. Hara, M. Hirano, K. Tanaka, *Org. Lett.* **2008**, *10*, 2537–2540; b) H. Tsuji, K.-i. Yamagata, T. Fujimoto, E. Nakamura, *J. Am. Chem. Soc.* **2008**, *130*, 7792–7793; c) Y. Kuninobu, M. Nishi, S. S. Yudha, K. Takai, *Org. Lett.* **2008**, *10*, 3009–3011; d) H. Hara, M. Hirano, K. Tanaka, *Org. Lett.* **2009**, *11*, 1337–1340; e) K. Zhang, J. Louie, *J. Org. Chem.* **2011**, *76*, 4686–4691.

[14] For a review on transition-metal-mediated  $\alpha$ -fluorine elimination, see: R. P. Hughes, *Eur. J. Inorg. Chem.* **2009**, 4591–4606.

[15] M. Takachi, Y. Kita, M. Tobisu, Y. Fukumoto, N. Chatani, *Angew. Chem.* **2010**, *122*, 8899–8902; *Angew. Chem. Int. Ed.* **2010**, *49*, 8717–8720.

[16] Structures of major and minor regioisomers were characterized by 2D NMR measurements. Minor products were thus found to be *m*-terphenyl derivatives. For details, see Supporting Information. For regioselectivity on nickel-catalyzed coupling reactions of alkynes via


- oxidative cyclization, see: P. Liu, P. McCarren, P. H.-Y. Cheong, T. F. Jamison, K. N. Houk, *J. Am. Chem. Soc.* **2010**, *132*, 2050–2057.
- [17] Reactions with terminal alkynes such as 1-hexyne, phenylacetylene, and trimethylsilylacetylene afforded the corresponding fluoroarenes, albeit in 5%, 3%, and 5% yields (determined by  $^{19}\text{F}$  NMR measurement using  $\text{PhCF}_3$  as an internal standard), respectively.
- [18] A stepwise oxidative cyclization model satisfactorily illustrates the experimental results. This stepwise model consists of (i) rapid pre-equilibrium between the reactants ( $\text{Ni}(0)$  and **1**) and the intermediary nickelacyclopropane (see ref. 9) and (ii) subsequent slow insertion of **2** into the nickelacyclopropane. For details, see Supporting Information.
- [19] The proposed oxidative cyclization is supported by Hoberg's and Chatani's reports. See: a) H. Hoberg, D. Guhl, *J. Organomet. Chem.* **1989**, *378*, 279–292; b) Ref. 15. See also, M. Takachi, N. Chatani, *Org. Lett.* **2010**, *12*, 5132–5134.
- [20] In the case of the reaction of 1,1-difluoro-1,6-enyne **5** with diphenylacetylene (**2a**),  $\beta$ -hydrogen elimination yielding the corresponding fluoroarene **6** was sluggish, probably due to the rigid bicyclic system of the intermediate. Thus, in the presence of the reductant,  $\text{Et}_3\text{B-}i\text{PrOLi}$ , transmetalation from the intermediary cyclohexadienylnickel(II) fluoride corresponding to **C** in Scheme 2 preferably occurred rather than  $\beta$ -hydrogen elimination, leading to the cyclohexadiene. To avoid confusion, we herein demonstrate that the stoichiometric reaction, conducted in the absence of the reductant, selectively afforded the corresponding fluoroarene **6** (eq 1).
- [21] Since generation of ethylene and dihydrogen during the reaction was confirmed by each gas detector, a  $\beta$ -hydrogen elimination–reductive elimination sequence definitely occurred as a route from  $\text{Ni}(\text{II})$  to  $\text{Ni}(0)$ .
- [22] For a review on oxidative C–H/C–H coupling, see: a) A. A. O. Sarhan, C. Bolm, *Chem. Soc. Rev.* **2009**, *38*, 2730–2744; b) M. Grzybowski, K. Skonieczny, H. Butenschön, D. T. Gryko, *Angew. Chem.* **2013**, *125*, 10084–10115; *Angew. Chem. Int. Ed.* **2013**, *52*, 9900–9930.
- [23] M. Danz, R. Tonner, G. Hilt, *Chem. Commun.* **2012**, *48*, 377–379.
- [24] K. Fuchibe, T. Morikawa, K. Shigeno, T. Fujita, J. Ichikawa, *Org. Lett.* **2015**, *17*, 1126–1129.
- [25] For a modular synthesis of substituted fluoroarenes, see: Y. Wang, D. J. Burton, *Tetrahedron Lett.* **2006**, *47*, 9279–9281.

Entry for the Table of Contents (Please choose one layout)

Layout 1:

## COMMUNICATION

A method for direct synthesis of tetrasubstituted fluoroarenes via nickel-catalyzed [2+2+2] cycloaddition is presented. The reaction combines one molecule of 1,1-difluoroethylene with two molecules of alkynes and involves sequential cleavage of the C–F and C–H bonds in difluoroethylene. The catalytic cycle is established by reduction of the intermediary Ni(II) fluoride with a triethylborane-based borate.



T. Fujita, Y. Watabe, T. Ichitsuka, J. Ichikawa\*

Page No. – Page No.

**Ni-Catalyzed Synthesis of Fluoroarenes via [2+2+2] Cycloaddition Involving  $\alpha$ -Fluorine Elimination**