

## IL-17F induces IL-6 via TAK1-NF B pathway in airway smooth muscle cells

著者別名	川口 未央, 森島 祐子, 石井 幸雄, 佐藤 浩昭, 坂本 透, 檜澤 伸之
journal or publication title	Immunity, Inflammation and Disease
volume	5
number	2
page range	124-131
year	2017-06
権利	(C) 2017 The Authors. Immunity, Inflammation and Disease Published by John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.
URL	<a href="http://hdl.handle.net/2241/00146404">http://hdl.handle.net/2241/00146404</a>

doi: 10.1002/iid3.149

## ORIGINAL RESEARCH

**IL-17F induces IL-6 via TAK1-NF $\kappa$ B pathway in airway smooth muscle cells**

Masayuki Nakajima<sup>1</sup>, Mio Kawaguchi<sup>1</sup>, Kyoko Ota<sup>1</sup>, Junichi Fujita<sup>1</sup>, Satoshi Matsukura<sup>2</sup>, Shau-Ku Huang<sup>3,4</sup>, Yuko Morishima<sup>1</sup>, Yukio Ishii<sup>1</sup>, Hiroaki Satoh<sup>1</sup>, Tohru Sakamoto<sup>1</sup>, & Nobuyuki Hizawa<sup>1</sup>

<sup>1</sup>Department of Pulmonary Medicine, Faculty of Medicine, University of Tsukuba, Ibaraki, Japan

<sup>2</sup>Respiratory Disease Center, Showa University Northern Yokohama Hospital, Kanagawa, Japan

<sup>3</sup>Asthma and Allergy Center, Johns Hopkins University, Baltimore, Maryland, USA

<sup>4</sup>National Health Research Institutes, Taipei, Taiwan

**Keywords**

IL-17F, NF- $\kappa$ B, TAK1

**Correspondence**

Mio Kawaguchi, Department of Pulmonary Medicine, Faculty of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 3058575, Japan.  
Tel: +81-298533144; Fax: +81-298533144;  
E-mail: mkawaguchi@md.tsukuba.ac.jp

**Funding information**

This work was supported by MEXT KAKENHI Grants-in-Aid for Scientific Research (C) 15K09209. S. K. Huang was supported, in part, by National Institute of Health (AI-052468) and National Health Research Institutes Taiwan.

Received: 16 September 2016; Revised: 17 November 2016; Accepted: 9 December 2016  
Final version published online 3 March 2017.

*Immunity, Inflammation and Disease*  
2017; 5(2): 124–131

doi: 10.1002/iid3.149

**Ethical statement:** No humans or animals were involved in this study—no ethical approval was required for this manuscript.

**Introduction**

Asthma is characterized by the association of airway inflammation, airway obstruction, increased airway hyper-reactivity, and remodeling. Airway smooth muscle is

**Abstract**

**Introduction:** Interleukin (IL)-17F plays a critical role in the pathophysiology of asthma. However, the precise role of IL-17F in airway smooth muscle cells (ASMCs) and its regulatory mechanisms remain to be defined. Therefore, we sought to investigate the expression of IL-6 by IL-17F and the involvement of transforming growth factor  $\beta$ -activated kinase 1 (TAK1) and nuclear factor (NF)- $\kappa$ B by in ASMCs.

**Methods:** ASMCs were cultured in the presence or absence of IL-17F. The expression of IL-6 gene and protein was analyzed using real-time PCR and ELISA, and the activation of TAK1 and NF- $\kappa$ B was detected by Western blotting. The effect of TAK1 inhibitor 5Z-7-oxozeaenol and NF- $\kappa$ B inhibitor BAY 11-7082 on the expression of IL-6 was investigated. Finally, the short interfering RNAs (siRNAs) targeting TAK1 and a subunit of NF- $\kappa$ B, p65 were transfected into ASMCs.

**Results:** The expression of IL-6 gene and protein was significantly induced by IL-17F. IL-17F activated TAK1 and NF- $\kappa$ B in ASMCs. Transfection of siRNAs targeting TAK1 abolished IL-17F-induced phosphorylation of p65. Both 5Z-7-oxozeaenol and BAY 11-7082 significantly inhibited IL-17F-induced IL-6 production in a dose-dependent manner. Similarly, transfection of the cells with siRNAs targeting TAK1 and p65 inhibited the expression of IL-6.

**Conclusions:** Collectively, these results provided evidence supporting the potential importance of the Th17-ASMCs crosstalk via the IL-17F-IL-6 axis in airway inflammation and as a candidate pharmacological target for airway inflammatory diseases such as asthma.

believed to contribute to the pathophysiology of asthma including direct causation of airflow obstruction via contraction. However, recent emerging evidences have demonstrated that airway smooth muscle plays a pivotal role in regulation of allergic airway inflammation via their

ability to produce many inflammatory molecules including IL-6 [1, 2]. IL-6 is known to be a potential contributor of asthma pathogenesis. Intranasal administration of a blocking anti-IL-6 receptor antibody in a mouse model of asthma has been shown to be able to decrease eosinophilia, the expression of Th2 cytokines, and airway hyperresponsiveness [3]. Hence, IL-6 might be a suitable target for a new approach to asthma therapy. But, how ASMCS-derived pro-inflammatory mediators such as IL-6 are regulated remain to be fully defined. Thus, further understanding of the mechanisms through which ASMCS-derived IL-6 is regulated would be important to uncover the pathogenic mechanisms of asthma.

Discovery of a novel molecule involving allergic airway inflammation and identification of its signaling mechanisms might help to clarify the pathogenesis of asthma. We and other groups cloned human IL-17F gene [4–6]. The IL-17F gene is evidently upregulated in the airway from patients with asthma [4]. Moreover, we have demonstrated that a single nucleotide polymorphism in IL-17F gene that results in a loss-of-function mutation is inversely related to asthma risk and is a natural IL-17F antagonist [7, 8]. These findings indicate that IL-17F is one of the important cytokines involved in the etiology of asthma. In addition, the level of IL-17F expression is correlated with the disease severity of asthma [9]. Although IL-17F is able to act several cell types to induce various cytokines, the role of IL-17F in ASMCS remains unclear [10–12]. ASMCS express a receptor for IL-17F, a heterodimer of IL-17RA and IL-17RC [13]. However, its intercellular signaling pathway is not well defined. Recent studies have shown that transforming growth factor  $\beta$ -activated kinase-1 (TAK1), a member of the mitogen-activated protein kinase kinase kinase family, is a pivotal signaling molecule leading to the activation of the transcription factors nuclear factor- $\kappa$ B (NF- $\kappa$ B) [14, 15]. TAK1 regulates the pathogenesis of innate and adaptive immunity including airway inflammation. In bronchial epithelial cells, TAK1 mediates the signaling of environmental stimuli such as respiratory viral infection which is a major cause of acute exacerbations [16]. However, the inducible factors of TAK1 in ASMCS remain unclear. In this study, we demonstrated, for the first time, that IL-17F is able to activate the TAK1-NF- $\kappa$ B signaling pathway to induce IL-6 expression in ASMCS.

## Methods

### Cell culture

ASMCS were purchased from Lonza (Walkersville, MD, USA) and cultured in SmBM medium with SmGM-2 SingleQuots (Lonza, Tokyo, Japan) containing insulin, fibroblast growth factor, gentamicin, 5% fetal bovine serum, and epidermal

growth factor at 37°C with 5% CO<sub>2</sub> in humidified air. Confluent cells at passages 2–4 were used.

### Analysis of IL-6 gene expression

Total RNA was extracted using RNeasy Mini Kit (Qiagen, Chatsworth, CA, USA) from  $1 \times 10^6$  ASMCS at 4, 12, 24, and 48 h after stimulation with 10 and 100 ng/mL of IL-17F (R&D Systems, Tokyo, Japan). cDNAs were synthesized from 1  $\mu$ g of total RNA using the ReverTra Ace qPCR RT Kit (TOYOBO, Tokyo, Japan), followed by real-time PCR. The sequences of primers for IL-6 are as follows: forward, 5'-AAAGAGGCACTGGCAGAAAA-3', reverse, 5'-CACCAGGCAAGTCTCCTCAT-3'; G3PDH: forward, 5'-ACCACAGTC-CATGCCATCAC-3', reverse, 5'-TCCACCACCCTGTTGCTGTA-3'. Real-time PCR was done using a THUNDERBIRD SYBR qPCR Mix (TOYOBO), gene-specific primers, and an ABI 7500 real-time PCR system. The data were shown as fold induction relative to the control group. The values are expressed as mean  $\pm$  SEM ( $n = 6$  experiments).

### Analysis of IL-6 protein production

Cell supernatants in ASMCS were harvested from cultures in the absence or presence of 10 and 100 ng/mL of IL-17F at 4, 12, 24, and 48 h. Alternatively, the cells were also stimulated by 100 ng/mL of IL-17A (R&D Systems) for 24 h. IL-6 protein levels in the supernatants of IL-17F-stimulated cells were determined with a commercially available ELISA kit (R&D Systems) according to the manufacturer's instruction, and expressed as the amount recovered per  $10^6$  cells. The values are expressed as mean  $\pm$  SEM ( $n = 6$  experiments).

### Detection of TAK1 and NF- $\kappa$ B by Western blotting

For analysis of activation of TAK1 and NF- $\kappa$ B, ASMCS were treated with IL-17F (100 ng/mL) and in some cases with or without transfection with the siRNAs targeting TAK1, p65 and control using HiPerFect Transfection Reagent (Qiagen). The total cellular extracts ( $1 \times 10^6$  cell equivalents/lane) were subjected to 7.5–15% Tris-glycine gel electrophoresis (DRC, Tokyo, Japan), followed by transfer onto polyvinylidene difluoride membranes (Bio-Rad, Tokyo, Japan). The Abs used were anti-TAK1 Ab, anti-phospho-TAK1 Ab, anti-phospho-p65 Ab (Cell Signaling Technology, Danvers, MA, USA), anti-p65 (Rel A) Ab (Santa Cruz Biotechnology, Santa Cruz, CA, USA).

### Effect of TAK1 inhibition on the expression of IL-6

For analysis of the involvement of TAK1, ASMCS were treated in the presence or absence of a TAK1 inhibitor, 5Z-7-

Oxozaenol (Sigma–Aldrich, Tokyo, Japan) at varying doses, and a vehicle control, 0.1% DMSO, for 3 h before treatment with IL-17F (100 ng/mL). The supernatants were harvested at 48 h after stimulation for analyses with ELISA. IL-6 protein levels in the supernatants were determined as described above. These values are expressed as mean  $\pm$  SEM ( $n = 6$  experiments). Simultaneously, pre-designed siRNAs for TAK1 (Santa Cruz Biotechnology) and control siRNAs (Ambion, Tokyo, Japan) were used. The siRNA transfection into ASMCs by HiPerFect Transfection Reagent (Qiagen) was performed according to the manufacturer's instruction. The supernatants were then harvested at 24 h after stimulation with 100 ng/mL of IL-17F and subjected to analysis by ELISA, respectively. IL-6 protein levels in the supernatants are expressed as mean  $\pm$  SEM ( $n = 6$  experiments).

### Effect of NF- $\kappa$ B inhibition on the expression of IL-6

For the analysis of involvement of NF- $\kappa$ B, ASMCs were treated in the presence or absence of a NF- $\kappa$ B inhibitor, BAY 11-7082 (Calbiochem, Tokyo, Japan) at varying doses, and a vehicle control, 0.1% DMSO, for 1 h before treatment with IL-17F (100 ng/mL). The supernatants were harvested at 24 h after stimulation for analyses with ELISA. IL-6 protein levels in the supernatants were determined as described above. These values are expressed as mean  $\pm$  SEM ( $n = 6$  experiments). Pre-designed siRNAs for p65 (Santa Cruz Biotechnology) were also used for transfection into ASMCs as described above. The supernatants were then harvested at 24 h after stimulation with 100 ng/mL of IL-17F and subjected to analysis by ELISA, respectively. IL-6 protein levels in the supernatants are expressed as mean  $\pm$  SEM ( $n = 6$  experiments).

### Data analysis

The statistical significance of differences was determined by analysis of variance (ANOVA). The values are expressed as mean  $\pm$  SEM from independent experiments. Any difference with  $P$ -values less than 0.05 was considered significant. When ANOVA indicated a significant difference, the Scheffe  $F$ -test was used to determine the difference between groups.

## Results

### Expression of IL-6 gene and protein

A 100 ng/mL of IL-17F significantly induced IL-6 gene expression when compared with control at 4, 12, and 24 h time points (Fig. 1A). IL-17F significantly induced its expression in a dose-dependent manner at 4 and 12 h after stimulation. The levels of IL-6 protein production were

analyzed by ELISA (Fig. 1B). IL-6 protein were detected in unstimulated cells, but its levels in supernatants were significantly increased, peaking at 24 h time point, in ASMCs stimulated with IL-17F when compared with control at every time point. Similarly, IL-17F significantly induced IL-6 production in a dose-dependent manner at 12, 24, and 48 h after stimulation. Other IL-17 cytokine family, IL-17A showed the similar potency to induce IL-6 expression as IL-17F at 24 h time point (Fig. 1C).

### Activation of TAK1 by IL-17F

TAK1 was equally detected at all-time points (Fig. 2). In contrast, a transient phosphorylation of TAK1 was seen on stimulation of the cells with IL-17F, reaching the maximum at 10–20 min after stimulation and returned to baseline levels by 120 min in ASMCs.

### Effect of TAK1 inhibition on IL-17F-induced IL-6 expression


Pretreatment of the cells with 0.1, 0.5, and 1.0  $\mu$ M of TAK1 inhibitor, 5Z-7-Oxozaenol significantly blocked IL-17F-induced IL-6 production in a dose-dependent manner, while 3-h pretreatment of the cells with vehicle alone (DMSO) did not affect its production in ASMCs (Fig. 3A). To further confirm whether TAK1 plays a role in IL-17F-induced IL-6 expression, total TAK1 expression was diminished in the cells by transfecting with siRNAs targeting TAK1 (Fig. 3B). As shown in Figure 4C, IL-6 production induced by IL-17F was significantly inhibited in cells transfected with siRNAs targeting TAK1 when compared with cells transfected with a control siRNAs.

### Activation of NF- $\kappa$ B (p65) by IL-17F

A subunit of NF- $\kappa$ B, p65, was detected at all-time points (Fig. 4A). Phosphorylation of p65 was evidently detected at 10–60 min, and returned to baseline levels by 120 min in IL-17F-stimulated ASMCs (Fig. 4A). To establish the interrelationship between TAK1 and NF- $\kappa$ B, the cells were transfected with siRNAs targeting TAK1 before the stimulation with IL-17F. Transfection with the siRNAs clearly diminished the activation of p65 induced by IL-17F, while transfection of control siRNAs did not affect the activation of p65 (Fig. 4B).

### Effect of NF- $\kappa$ B inhibition on IL-17F-induced IL-6 expression


Pretreatment of ASMCs for 1 h with 0.5, 1.0 and 5.0  $\mu$ M of a NF- $\kappa$ B inhibitor, BAY 11-7082 significantly decreased the levels of IL-17F-induced IL-6 protein production in a dose-dependent



**Figure 1.** The expression of IL-6 gene and protein by IL-17F in ASMCs. (A) Gene expression of IL-6 by IL-17F. Real-time PCR was performed as described in Materials and Methods. ASMCs were stimulated with IL-17F (100 ng/mL) for 4, 12, and 24 h (n = 6). \* $P < 0.05$  versus medium control. (B) IL-6 protein expression by IL-17F. ELISA was performed as described in Materials and Methods. The cells were stimulated with IL-17F (100 ng/mL) for 4–48 h (n = 6). \* $P < 0.05$  versus medium control. # $P < 0.05$  versus 10 ng/mL of IL-17F-stimulated cells. (C) IL-6 protein levels induced by IL-17A and IL-17F in supernatants (n = 6). The cells were stimulated with 100 ng/mL of IL-17A or IL-17F for 24 h (n = 6). \*\* $P < 0.05$  versus medium control.

manner, while 1 h pretreatment of the cells with vehicle alone (DMSO) did not affect IL-6 production (Fig. 5A). Finally, to further confirm whether NF- $\kappa$ B plays a role in IL-17F-induced IL-6 expression, it was found that when NF- $\kappa$ B p65 was knocked

down in the cells with siRNAs (Fig. 5B), IL-6 production induced by IL-17F was significantly inhibited when compared with cells transfected with control siRNAs (Fig. 5C).




**Figure 2.** Activation of TAK1 by IL-17F in ASMCs. The cells were incubated with IL-17F (100 ng/mL) for different time points as indicated. Western blotting analysis was performed with Abs against total (t)-TAK1 and phosphorylated (p)-TAK1 as indicated. These results shown are representative of three separate experiments.

## Discussion

In this study, we demonstrated that IL-17F significantly induced the expression of IL-6 in ASMCs via the activation of TAK1-NF- $\kappa$ B signaling pathway. These findings indicate the likely importance of the IL-17F-IL-6 axis in airway inflammation as a consequence of the Th17-ASMC crosstalk.


Current study demonstrated that IL-17F signal mediates TAK1-NF- $\kappa$ B pathway. This suggests that this signaling pathway is pivotal for IL-17F-induced IL-6 expression in




**Figure 3.** Effect of the inhibition for TAK1 on IL-6 expression. ASMCs were pretreated with 5Z-7-Oxozeaenol (0.1, 0.5, and 1.0 μM) for 3 h before the 24 h-stimulation of IL-17F (100 ng/mL), and then IL-6 protein levels in supernatants were measured by ELISA. The values are expressed as means ± SEM (n = 6). \*P < 0.05 versus IL-17F-stimulated cells in the absence of the inhibitor. \*\*P < 0.05 versus the presence of individual inhibitor. (B) The validation of blocking by siRNAs targeting TAK1 was performed by Western blotting. These results shown are representative of three separate experiments. (C) After transfection of the siRNAs, ASMCs were stimulated with IL-17F (100 ng/mL) for 24 h. The levels of IL-6 protein production in the supernatants were measured by ELISA. The values are expressed as means ± SEM (n = 6). \*P < 0.05 versus IL-17F-stimulated cells transfected with a control siRNA.

ASMCs. TAK1 is for the first time, identified as a novel signaling molecule involved in the function of IL-17F. Consistent with the previous studies, TAK1 is located upstream of NF-κB, since a TAK1 inhibitor 5Z-7-

Oxozeaenol diminished the phosphorylation of NF-κB [15]. Moreover, the activation of NF-κB is crucial for IL-6 expression by IL-17F, since NF-κB inhibitor and the specific siRNAs abrogated its expression. Recent reports demonstrated that TAK1 is clearly involved in the pathogenesis of airway inflammation. TAK1 activation in bronchial epithelial cells was induced by several inflammatory stimuli such as RS virus, *Pseudomonas aeruginosa* and diesel exhaust particles that are able to exacerbate airway inflammation [16–18]. Interestingly, TAK1 is involved in steroid responsiveness in asthma [19]. Blocking of TAK1 recovers cellular response to steroids in the presence of pathogenic bacteria via the regulation of MAPK phosphatase-activation. However, the functional role of TAK1 in ASMCs still remains. It is known that TAK1 contributes to cigarette smoke-induced IL-8 production and the causing of airway remodeling through the induction of growth factor-induced proliferation of ASMCs [20, 21]. Here, we reported that IL-17F induced phosphorylation of TAK1, indicating that TAK1 is a potentially pharmacological target for the IL-17F-mediated airway inflammatory diseases. However, the inhibitors and siRNAs targeting TAK1 or NFκB-p65 did not completely inhibit IL-17F-induced IL-6 expression in this study. This suggests that the involvement of other signaling molecules. Further study is needed to identify novel signaling pathways of IL-17F and its functional impact on IL-6 expression.



**Figure 4.** Activation of NF-κB by IL-17F. (A) Kinetic activation of NF-κB by IL-17F in ASMCs. The cells were incubated with IL-17F (100 ng/mL) for different time points as indicated. Western blotting was performed with Abs against total (t)-p65 and phosphorylated (p)-p65. (B) Effect of siRNAs targeting TAK1 on IL-17F-induced phosphorylation of p65. The cells were transfected with siRNAs targeting TAK1 or control siRNAs, and then ASMCs were stimulated with IL-17F for 30 min. Western blotting analysis was performed with Abs against t-p65 and p-p65. The results shown are representative of three separate experiments.



**Figure 5.** Effect of NF- $\kappa$ B inhibition on IL-17F-induced IL-6 expression. (A) ASM cells were pretreated with NF- $\kappa$ B inhibitor, BAY 11-7082 for 1 h before the 24 h-stimulation of IL-17F (100 ng/mL), and then IL-6 protein levels in supernatants were measured by ELISA. The values are expressed as means  $\pm$  SEM (n = 6). \* $P$  < 0.05 versus IL-17F-stimulated cells in the absence of the inhibitor. \*\* $P$  < 0.05 versus the presence of individual inhibitor. (B) The validation of blocking by siRNAs targeting p65 was performed by Western blotting. These results shown are representative of three separate experiments. (C) After transfection of the siRNAs, ASM cells were stimulated with IL-17F (100 ng/mL for 24 h). The levels of IL-6 protein production in the supernatants were measured by ELISA. The values are expressed as means  $\pm$  SEM (n = 6). \* $P$  < 0.05 versus IL-17F-stimulated cells transfected with control siRNAs.

IL-17F shows several distinctive features of asthma such as airway remodeling, goblet cell hyperplasia, and increasing airway hyperreactivity. Overexpression of IL-17F following Ag challenge in the airway of mice resulted in the induction of goblet cell hyperplasia and, gene expression of MUC5AC and significant increase in airway hyperreactivity [22]. These findings suggest that IL-17F can provide an additive effect on antigen-induced allergic inflammatory responses. In addition, IL-17F is associated with airway neutrophilia. Although neutrophilic inflammation is one of the features of severe asthma, its precise mechanism has not been clarified. Overexpression of IL-17F in the mouse airways results in an increased number of neutrophils in bronchoalveolar lavage fluid (BALF) [22, 23]. Interestingly, IL-17F is more involved in airway neutrophilic inflammation to *Aspergillus oryzae* when compared with IL-17A [24]. Moreover, IL-17F is able to induce CXC chemokines, such as IL-8, ENA-78, and GRO $\alpha$ , that are potent chemoattractants for neutrophils [4, 25]. Hence, Neutrophil recruitment into the airway may be regulated via, at least partially, IL-17F-induced CXC chemokines. IL-17F-producing cells are known to come from many cell types, such as bronchial epithelial cells, basophils, mast cells,  $\gamma\delta$ T cells, CD8<sup>+</sup>T cells, and Th17 cells [4, 26–28]. Especially, Th17 cells are deeply involved in the pathogenesis of a diverse group of immune-mediated diseases, including asthma [29]. Th17 cells are

the major cell source of IL-17F, and our current study demonstrated that IL-17F could induce IL-6 expression. IL-6 is necessary for Th17 development through the induction of a transcriptional factor, retinoic acid receptor-related orphan nuclear receptor (ROR $\gamma$ t) [30]. Although the cell source of IL-6 in airway inflammation has not been fully understood, the current study suggests that IL-6 is derived from, at least partially, ASM cells in response to IL-17F. Taken together, it is possible that as a mode of the Th17-ASM cell crosstalk, IL-17F-induced IL-6 further promotes Th17 cell differentiation, which, in turn, establishes a positive feedback loop resulting in the amplification of the Th17 response. Further in vivo study is needed to clarify the importance of the IL-17F/IL-6 axis in asthma.

The role of ASM cells in the pathogenesis of airway inflammation has become increasingly clear. ASM cells are able to induce a wide range of cytokines and chemokines that orchestrate airway inflammation [31]. However, the functional role of IL-17F in ASM cells has not been clarified. A few studies have demonstrated that IL-17F induces the migration and proliferation of ASM cells [13, 32]. These findings suggest the involvement of IL-17F in airway remodeling. Interestingly, unlike IL-17A, IL-17F did not affect ASM cell contraction [33]. However, the reason of this difference has not been dissolved yet. Although both IL-17A and IL-17F bind to same receptor the heterodimeric complex of IL-17RA and IL-17RC, the binding affinity of

IL-17A and IL-17F is quite different [34]. IL-17RA effectively binds to IL-17A. In contrast, it binds to IL-17F with extremely lower affinity. The relative binding affinity of IL-17F to IL-17RC is more potent than to IL-17RA. These findings may affect their different biological activity for ASM cell contraction. Further study is needed in the future.

On the other hand, IL-6 is a key factor in the pathophysiologic events of asthma [35]. Increased levels of IL-6 are observed in serum, BALF, and induced sputum from asthmatic patients. The levels of IL-6 in sputum and serum are inversely correlated with lung function in asthmatic patients [36, 37]. Besides inducing Th17 cell development, IL-6 is a critical regulator of promoting IL-4 production during Th2 differentiation, inhibiting Th1 differentiation, and is a co-factor in IL-4-dependent IgE synthesis [38, 39]. Moreover, IL-6 is involved in airway mucus hypersecretion that is one of the features of asthma [40]. Hence, ASM cells are important target cells for IL-17F, and contribute to the pathogenesis of allergic airway inflammation, at least partially, via the induction of IL-6 expression.

In conclusion, we demonstrated that IL-17F activates TAK1-NF- $\kappa$ B signaling pathway to induce IL-6 expression. IL-17F/IL-6 axis might be involved in the pathophysiology of allergic airway inflammation, and targeting IL-17F and its signaling pathways could be a novel strategy for asthma.

## Conflict of Interest

None declared.

## References

- Perry, M. M., J. E. Baker, D. S. Gibeon, I. M. Adcock, and K. F. Chung. 2014. Airway smooth muscle hyperproliferation is regulated by microRNA-221 in severe asthma. *Am. J. Respir. Cell. Mol. Biol.* 50:7–17.
- Elias, J. A., Y. Wu, T. Zheng, and R. Panettieri. 1997. Cytokine- and virus-stimulated airway smooth muscle cells produce IL-11 and other IL-6-type cytokines. *Am. J. Physiol.* 273:L648–L655.
- Doganci, A., T. Eigenbrod, N. Krug, G. T. De Sanctis, M. Hausding, V. J. Erpenbeck, el-B. Haddad, H. A. Lehr, E. Schmitt, T. Bopp, et al. 2005. The IL-6R alpha chain controls lung CD4+ CD25+ Treg development and function during allergic airway inflammation in vivo. *J. Clin. Invest.* 115:313–325.
- Kawaguchi, M., L. F. Onuchic, X. D. Li, D. M. Essayan, J. Schroeder, H. Q. Xiao, M. C. Liu, G. Krishnaswamy, G. Germino, and S. K. Huang. 2001. Identification of a novel cytokine, ML-1, and its expression in subjects with asthma. *J. Immunol.* 167:4430–4435.
- Hymowitz, S. G., E. H. Filvaroff, J. P. Yin, J. Lee, L. Cai, P. Risser, M. Maruoka, W. Mao, J. Foster, R. F. Kelley, et al. 2001. IL-17s adopt a cystine knot fold: structure and activity of a novel cytokine, IL-17F, and implications for receptor binding. *EMBO J.* 20:5332–5341.
- Starnes, T., M. J. Robertson, G. Sledge, S. Kelich, H. Nakshatri, H. E. Broxmeyer, and R. Hromas. 2001. IL-17F, a novel cytokine selectively expressed in activated T cells and monocytes, regulates angiogenesis and endothelial cell cytokine production. *J. Immunol.* 167:4137–4140.
- Kawaguchi, M., D. Takahashi, N. Hizawa, S. Suzuki, S. Matsukura, F. Kokubu, Y. Maeda, Y. Fukui, S. Konno, S. K. Huang, et al. 2006. IL-17F sequence variant (His161Arg) is associated with protection against asthma and antagonizes wild-type IL-17F activity. *J. Allergy Clin. Immunol.* 117:795–801.
- Hizawa, N., M. Kawaguchi, S. K. Huang, and M. Nishimura. 2006. Role of interleukin-17F in chronic inflammatory and allergic lung disease. *Clin. Exp. Allergy* 36:1109–1114.
- Al-Ramli, W., D. Préfontaine, F. Chouiali, J. G. Martin, R. Olivenstein, C. Lemièrre, and Q. Hamid. 2009. T(H)17-associated cytokines (IL-17A and IL-17F) in severe asthma. *J. Allergy Clin. Immunol.* 123:1185–1187.
- Kawaguchi, M., F. Kokubu, J. Fujita, S. K. Huang, and N. Hizawa. 2009. Role of interleukin-17F in asthma. *Inflamm. Allergy Drug Targets* 8:383–389.
- Kawaguchi, M., J. Fujita, F. Kokubu, G. Ohara, S. K. Huang, S. Matsukura, Y. Ishii, M. Adachi, H. Satoh, and N. Hizawa. 2010. Induction of insulin-like growth factor-I by interleukin-17F in bronchial epithelial cells. *Clin. Exp. Allergy* 40:1036–1043.
- Kawaguchi, M., M. Adachi, N. Oda, F. Kokubu, and S. K. Huang. 2004. IL-17 cytokine family. *J. Allergy Clin. Immunol.* 114:1265–1273.
- Chang, Y., L. Al-Alwan, P. A. Risse, L. Roussel, S. Rousseau, A. J. Halayko, J. G. Martin, Q. Hamid, and D. H. Eidelman. 2011. TH17 cytokines induce human airway smooth muscle cell migration. *J. Allergy Clin. Immunol.* 127:1046–1053.
- Yamaguchi, K., K. Shirakabe, H. Shibuya, K. Irie, I. Oishi, N. Ueno, T. Taniguchi, E. Nishida, and K. Matsumoto. 1995. Identification of a member of the MAPKKK family as a potential mediator of TGF-beta signal transduction. *Science* 270:2008–2011.
- Sakurai, H., H. Miyoshi, W. Toriumi, and T. Sugita. 1999. Functional interactions of transforming growth factor beta-activated kinase 1 with IkappaB kinases to stimulate NF-kappaB activation. *J. Biol. Chem.* 274:10641–10648.
- Dey, N., T. Liu, R. P. Garofalo, and A. Casola. 2011. TAK1 regulates NF- $\kappa$ B and AP-1 activation in airway epithelial cells following RSV infection. *Virology* 418:93–101.
- Martel, G., J. Bérubé, and S. Rousseau. 2013. The protein kinase TPL2 is essential for ERK1/ERK2 activation and cytokine gene expression in airway epithelial cells exposed to pathogen-associated molecular patterns (PAMPs). *PLoS ONE* 8:e59116.



18. Yun, Y. P., J. D. Joo, J. Y. Lee, H. Y. Nam, Y. H. Kim, K. H. Lee, C. S. Lim, H. J. Kim, Y. G. Lim, and Y. Lim. 2005. Induction of nuclear factor-kappaB activation through TAK1 and NIK by diesel exhaust particles in L2 cell lines. *Toxicol. Lett.* 155:337–342.
19. Goleva, E., L. P. Jackson, J. K. Harris, C. E. Robertson, E. R. Sutherland, C. F. Hall, J. T. Good, Jr, E. W. Gelfand, R. J. Martin, and D. Y. Leung. 2013. The effects of airway microbiome on corticosteroid responsiveness in asthma. *Am. J. Respir. Crit. Care Med.* 188:1193–1201.
20. Pera, T., C. Atmaj, M. van der Vegt, A. J. Halayko, J. Zaagsma, and H. Meurs. 2012. Role for TAK1 in cigarette smoke-induced proinflammatory signaling and IL-8 release by human airway smooth muscle cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 303:L272–L278.
21. Pera, T., R. Sami, J. Zaagsma, and H. Meurs. 2011. TAK1 plays a major role in growth factor-induced phenotypic modulation of airway smooth muscle. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 301:L822–L828.
22. Oda, N., P. B. Canelos, D. M. Essayan, B. A. Plunkett, A. C. Myers, and S. K. Huang. 2005. Interleukin-17F induces pulmonary neutrophilia and amplifies antigen-induced allergic response. *Am. J. Respir. Crit. Care Med.* 171: 12–18.
23. Hurst, S. D., T. Muchamuel, D. M. Gorman, J. M. Gilbert, T. Clifford, S. Kwan, S. Menon, B. Seymour, C. Jackson, T. T. Kung, et al. 2002. New IL-17 family members promote Th1 or Th2 responses in the lung: in vivo function of the novel cytokine IL-25. *J. Immunol.* 169:443–453.
24. Yang, X. O., S. H. Chang, H. Park, R. Nurieva, B. Shah, L. Acero, Y. H. Wang, K. S. Schluns, R. R. Broaddus, Z. Zhu, et al. 2008. Regulation of inflammatory responses by IL-17F. *J. Exp. Med.* 205:1063–1075.
25. Kawaguchi, M., F. Kokubu, S. Matsukura, K. Ieki, M. Odaka, S. Watanabe, S. Suzuki, M. Adachi, and S. K. Huang. 2003. Induction of C-X-C chemokines, growth-related oncogene alpha expression, and epithelial cell-derived neutrophil-activating protein-78 by ML-1 (interleukin-17F) involves activation of Raf1-mitogen-activated protein kinase kinase-extracellular signal-regulated kinase 1/2 pathway. *J. Pharmacol. Exp. Ther.* 307:1213–1220.
26. O'Brien, R. L., C. L. Roark, and W. K. Born. 2009. IL-17-producing gammadelta T cells. *Eur. J. Immunol.* 39:662–666.
27. Ciric, B., M. El-behi, R. Cabrera, G. X. Zhang, and A. Rostami. 2009. IL-23 drives pathogenic IL-17-producing CD8+ T cells. *J. Immunol.* 182:5296–5305.
28. Bettelli, E., Y. Carrier, W. Gao, T. Korn, T. B. Strom, M. Oukka, H. L. Weiner, and V. K. Kuchroo. 2006. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* 441:235–238.
29. Newcomb, D. C., and R. S. Peebles, Jr. 2013. Th17-mediated inflammation in asthma. *Curr. Opin. Immunol.* 25:755–760.
30. Ghoreschi, K., A. Laurence, X. P. Yang, C. M. Tato, M. J. McGeachy, J. E. Konkel, H. L. Ramos, L. Wei, T. S. Davidson, N. Bouladoux, et al. 2010. Generation of pathogenic T(H)17 cells in the absence of TGF- $\beta$  signalling. *Nature* 467:967–971.
31. Ozier, A., B. Allard, I. Bara, P. O. Girodet, T. Trian, R. Marthan, and P. Berger. 2011. The pivotal role of airway smooth muscle in asthma pathophysiology. *J. Allergy (Cairo)* 2011:742710.
32. Chang, Y., L. Al-Alwan, P. A. Risse, A. J. Halayko, J. G. Martin, C. J. Baglole, D. H. Eidelman, and Q. Hamid. 2012. Th17-associated cytokines promote human airway smooth muscle cell proliferation. *FASEB J.* 26:5152–5160.
33. Kudo, M., A. C. Melton, C. Chen, M. B. Engler, K. E. Huang, X. Ren, Y. Wang, X. Bernstein, J. T. Li, K. Atabai, et al. 2012. IL-17A produced by  $\alpha\beta$  T cells drives airway hyper-responsiveness in mice and enhances mouse and human airway smooth muscle contraction. *Nat. Med.* 18:547–554.
34. Kuestner, R. E., D. W. Taft, A. Haran, C. S. Brandt, T. Brender, K. Lum, B. Harder, S. Okada, C. D. Ostrander, J. L. Kreindler, et al. 2007. Identification of the IL-17 receptor related molecule IL-17RC as the receptor for IL-17F. *J. Immunol.* 179:5462–5473.
35. Rincon, M., and C. G. Irvin. 2012. Role of IL-6 in asthma and other inflammatory pulmonary diseases. *Int. J. Biol. Sci.* 8:1281–1290.
36. Dixon, A. E., D. M. Raymond, B. T. Suratt, L. M. Bourassa, and C. G. Irvin. 2008. Lower airway disease in asthmatics with and without rhinitis. *Lung* 186:361–368.
37. Dixon, A. E., D. M. Shade, R. I. Cohen, G. S. Skloot, J. T. Holbrook, L. J. Smith, J. J. Lima, H. Allayee, C. G. Irvin, R. A. Wise, American Lung Association-Asthma Clinical Research Centers. 2006. Effect of obesity on clinical presentation and response to treatment in asthma. *J. Asthma* 43:553–558.
38. Dienz, O., and M. Rincon. 2009. The effects of IL-6 on CD4 T cell responses. *Clin. Immunol.* 130:27–33.
39. Vercelli, D., H. H. Jabara, K. Arai, T. Yokota, and R. S. Geha. 1989. Endogenous interleukin 6 plays an obligatory role in interleukin 4-dependent human IgE synthesis. *Eur. J. Immunol.* 19:1419–1424.
40. Neveu, W. A., J. B. Allard, O. Dienz, M. J. Wargo, G. Ciliberto, L. A. Whittaker, and M. Rincon. 2009. IL-6 is required for airway mucus production induced by inhaled fungal allergens. *J. Immunol.* 183:1732–1738.