

Synthesis of Pinpoint-Fluorinated Polycyclic Aromatic Hydrocarbons: Benzene Ring Extension Cycle Involving Microwave-Assisted SNAr Reaction

著者別名	市川 淳士, 湊辺 耕平
journal or publication title	Chemistry : an Asian journal
volume	12
number	18
page range	2359-2363
year	2017-09
権利	(C) 2017 Wiley-VCH Verlag GmbH &Co. KGaA, Weinheim
URL	http://hdl.handle.net/2241/00148493

doi: 10.1002/asia.201700870

Synthesis of Pinpoint-Fluorinated Polycyclic Aromatic Hydrocarbons: Benzene Ring Extension Cycle Involving Microwave-Assisted S_NAr Reaction


Kohei Fuchibe, Hisanori Imaoka, and Junji Ichikawa*

Abstract: Fluoroarenes bearing no electron-withdrawing groups (non-activated fluoroarenes) readily underwent nucleophilic aromatic substitution with α -cyanocarbanions under microwave irradiation. The sequence (i) formylalkylation involving the cyanoalkylation of fluoroarenes, (ii) difluorovinylideneation, and (iii) Friedel–Crafts-type cyclization, afforded one benzene ring-extended fluoroarenes in one cycle. Furthermore, the performance of multiple cycles successfully provided higher order pinpoint-fluorinated polycyclic aromatic hydrocarbons (F-PAHs).

Pinpoint-fluorinated PAHs (regioselectively fluorinated polycyclic aromatic hydrocarbons) are promising organic semiconducting materials¹ because of the unique properties of the fluorine substituent.² The high electronegativity of fluorine leads to the enhanced resistance of PAHs to aerial oxidation by lowering the energy level of their HOMO. In addition, the repulsive interaction between lone pairs in the fluorine 2p orbitals and the adjacent π -electrons in the carbon 2p orbitals perturbs the electron density of the extended π -systems,³ which would render these compounds highly soluble in polar organic solvents, leading to printable organic electronics.⁴ It is also worth mentioning the low steric impact of fluorine, whose introduction into PAH molecules should not change their molecular shape, would not affect much their π - π stacking in the solid structure.^{5,6} Therefore, the effects of installing a single fluorine substituent in PAH skeletons would lead to advantageous semiconducting materials, such as THF-soluble fluorinated picones, which exhibit p-type semiconducting behavior.^{5,7}


To facilitate the efficient synthesis of pinpoint-fluorinated PAHs, we have recently developed metal-catalyzed cyclizations of fluoroalkenes on the basis of the following benzene ring construction strategy (Scheme 1): (a) 1,1-difluoroalkenes, readily prepared via difluorovinylideneation of aldehydes,⁸ undergo a Friedel–Crafts-type cyclization in the presence of a catalytic amount of indium(III) bromide.⁹ Subsequent DDQ (2,3-dichloro-5,6-dicyano-*p*-benzoquinone) dehydrogenation affords pinpoint-fluorinated aromatic compounds (fluoroarenes). (b) These compounds were also accessible from 1,1-difluoroalkenes via a cationic palladium(II)-catalyzed cyclization (Scheme 1).^{5,10,11}

As the preparation of the substrates for the indium(III)-catalyzed synthesis of pinpoint-fluorinated PAHs, i.e.,



Scheme 1. F-PAH Syntheses.

difluoroalkenes, involved fluoroarenes as starting materials, we envisioned a cyclic benzene ring extension strategy that would comprise the following steps (Scheme 2): (i) starting fluoroarenes would be subjected to formylalkylation via S_NAr reaction to provide arylaldehydes; (ii) the aldehydes would be then subjected to difluorovinylideneation,⁸ finally, (iii) the Friedel–Crafts-type cyclization of the obtained 1,1-difluoroalkenes⁹ would furnish the benzene ring-extended fluoroarenes.



Scheme 2. Benzene Ring Extension Cycle.

However, S_NAr reactions of fluoroarenes have been conducted on those activated by electron-withdrawing groups, including halogenes and trifluoromethyl groups,¹² and the S_NAr reaction of non-activated fluoroarenes have been limited to those on benzene π systems.¹³ In the course of this study, we found that microwave irradiation¹⁴ significantly facilitated the desired S_NAr reaction of non-activated fluoroarenes with naphthalene or higher order PAH π systems (Table 1).¹⁵ When fluoronaphthalene **1a** and propionitrile (4.0 equiv) were heated in refluxing THF using a conventional oil bath in the presence of

[a] Dr. K. Fuchibe, H. Imaoka, and Prof. Dr. J. Ichikawa
Division of Chemistry, Faculty of Pure and Applied Sciences,
University of Tsukuba
Tsukuba, Ibaraki 305-8571 (Japan)
E-mail: junji@chem.tsukuba.ac.jp
Homepage: <http://www.chem.tsukuba.ac.jp/junji/index.html>

potassium bis(trimethylsilyl)amide (KHMDS, 4.0 equiv, entry 1), the desired alkylated naphthalene **2a** was obtained albeit in low yields, i.e., 6–17% (Table 1, entries 1 and 2). In contrast, microwave irradiation at 80 °C dramatically increased the yield of **2a** to 70% (entry 3). Further optimization afforded yields up to 82% after isolation (entries 4 and 5).^{16,17}




Table 1. Optimization for S_NAr Reaction.^[a]

Entry	Heating	Concentration of 1a [M]	Conditions	Yield of 2a [%]
1	Oil Bath	0.25	65 °C, 24 h	6
2	Oil Bath	0.25	80 °C, ^[b] 16 h	17
3	Microwave	0.25	80 °C, ^[b] 1 h	70
4	Microwave	0.25	80 °C, ^[b] 1.5 h	77
5	Microwave	0.50	80 °C, ^[b] 1.5 h	88 (82)

[a] ¹H NMR yield based on an internal standard CH₂Br₂. Isolated yield in parentheses. [b] The reaction was conducted in a sealed vessel.

Acetonitrile (Table 2, entry 1) reacted with **1a** in a manner similar to that of propionitrile, affording the corresponding primary nitrile **2b** in 53% yield under microwave irradiation. In contrast, **2b** was not obtained when performing the S_NAr reaction of acetonitrile under oil bath heating (THF, reflux, 18 h). Decanonitrile, whose C8 alkyl chain is expected to enhance the solubility of PAHs, also afforded the corresponding naphthalene **2c** in 90% yield (entry 2). Thus, as well as




Figure 1. List of Fluoroarenes with Extended π Systems.




Table 2. Microwave-assisted S_NAr Reaction^[a]


Entry	1 ^[b]	R ¹ , R ²	Nitrile 2	Yield [%] ^[c]
1	1a	H, H		2b 53 (ND)
2	1a	<i>n</i> -C ₈ H ₁₇ , H		2c 90 (10)
3	1a	CH ₃ , CH ₃		2d 92 (70)
4	1b ^[d]	CH ₃ , H		2e 54 (ND)
5	1c ^[d]	CH ₃ , H		2f 76
6	1d	CH ₃ , H		2g 93
7	1e ^[d]	CH ₃ , H		2h 70
8	1f ^[d]	CH ₃ , H		2i 89
9	1g ^[e]	CH ₃ , H		2j 92

[a] The reaction was conducted in a sealed vessel. [b] Unless otherwise noted, concentration of **1** was 0.50 M. [c] Isolated yield. The yield obtained under oil bath heating (THF, 65 °C) is shown in parentheses. [d] **1** 0.25 M. [e] **1g** 0.008 M. ND = Not detected by ¹H NMR spectroscopy.

secondary nitriles,^{13a} acetonitrile and primary nitriles were demonstrated to react with fluoronaphthalene under microwave irradiation. Isobutyronitrile, which is an effective substrate for the nucleophilic aromatic substitution, gave **2d** in 92% yield (entry 3).

The microwave-assisted S_NAr reaction was applicable to a wide variety of fluoroarenes (Figure 1). Thus, 3-fluoropyridine (**1b**) afforded the corresponding product **2e** in 54% yield (Table 2, entry 4). Substitution of fluorophenanthrenes **1c** and **1d** also worked well to afford the corresponding **2f** and **2g** in 76% and 93% yields, respectively (entries 5 and 6). Fluorochrysene ([4]phenacene) **1e** and fluoro[4]helicene **1f** were successfully transformed into **2h** and **2i** in 70% and 89% yields, respectively (entries 7 and 8), proving that the substitution proceeded smoothly even at the bay region of the phenanthrene substructure of **1e**. Finally, fluoropicene ([5]phenacene) **1g** underwent the substitution under identical conditions to give **2j** in 92% yield (entry 9). Thus, the scope of microwave-assisted S_NAr reaction was expanded to include non-activated fluoroarenes with extended π systems.

With the microwave-assisted alkylation method in hand, we sought to explore its application to the planned benzene ring extension (Scheme 3). Nitrile **2a**, obtained as detailed above from **1a** (Table 1, entry 5: step 1, 82% yield), was subjected to half reduction with diisobutylaluminum hydride (DIBAL) to afford aldehyde **3a** in 81% yield (step 2). Difluorovinylideneation of **3a** afforded 1,1-difluoroallene **4a** in 73% yield in two steps (steps 3 and 4). Then **4a** underwent Friedel–Crafts-type cyclization under indium(III) catalysis to provide benzene ring-extended fluorophenanthrene **1h** in 94% yield (step 5), thus completing the first cycle. In a similar manner, the benzene ring extension starting from **2c** afforded the corresponding **1i** in good yield.



Scheme 3. Benzene Ring Extension of 1-Fluoronaphthalene (1st Cycle).

By repeating the same protocol with fluoroarene **1h**, higher order pinpoint-fluorinated PAHs were synthesized (Table 3). Thus, **1h** was subjected to the second ring extension to afford pinpoint-fluorinated chrysene **1j** (entry 1). The third ring extension of **1j** was performed to afford pinpoint-fluorinated picene **1k** (entry 2).





Table 3. Benzene Ring Extension of Fluoroarenes (2nd and 3rd Cycles) ^{[a][b]}

Entry	Cycle	Starting Compd	Yield [%]			
			Step 1)	Step 2)	Steps 3,4) ^[c]	Step 5)
1	2nd	1h	71 ^[d]	85	64	87, 1j
2	3rd	1j	64 ^[e]	79	36	79, 1k

[a] Isolated. [b] For steps 1) to 5), see Scheme 3. [c] Two-step yield. [d] **1h** 0.5 M. [e] **1j** 0.25 M.

The benzene ring extension was successfully applied to not only terminal fluoroarenes but also internal ones (Scheme 4). Pinpoint-fluorinated triphenylene **1l** (top) was successfully synthesized from fluorophenanthrene **1c** via nitrile **2f**, whereas pinpoint-fluorinated triphenylene **1m** (bottom) was similarly obtained from fluoro[4]helicene **1f** via **2i**.¹⁸



Scheme 4. Benzene Ring Extension of Internal Fluoroarenes.

It is worth noting that the benzene ring extension using primary nitriles facilitated the synthesis of PAHs bearing a substituent in the bay region (Schemes 3, 4 and Table 3). Although effects of bay substitution are of importance not only from the viewpoint of organic devices but also from the viewpoint of cancer research,^[19] synthetic methods applicable to the molecules have been rare.^[20] The benzene ring extension thus contributes to a wide range of research areas through providing the bay-substituted PAHs.

In summary, we have achieved a microwave-assisted S_NAr reaction of non-activated fluoroarenes with extended π systems. This S_NAr reaction along with difluorovinylidenation and cationic cyclization facilitated a novel benzene ring extension cycle, by which terminal and internal fluoroarenes were converted to the corresponding one benzene ring-extended fluoroarenes in good yields. Thus, the cycle increases the variety of pinpoint-fluorinated PAHs.

Experimental Section

Microwave-assisted S_NAr reaction of non-activated fluoroarenes (step 1): synthesis of nitrile **2a** is described as a typical procedure. To a THF solution (2.4 mL) of KHMDS (1.03 g, 4.80 mmol) was added 1-fluoronaphthalene (**1a**, 155 μ L, 1.20 mmol) at room temperature. After propionitrile (340 μ L, 4.75 mmol) was added, microwave (65W) was irradiated at 80 °C for 90 min. The reaction mixture was poured into aqueous hydrochloric acid (2 M, 5 mL) at room temperature. Organic materials were extracted with ethyl acetate three times. The combined extracts were washed with brine and dried over anhydrous sodium sulfate.

To the residue (396 mg) were added dibromomethane (109 mg) and α,α,α -trifluorotoluene (24 mg) as internal standards. Analysis by NMR spectroscopy indicated that nitrile **2a** was generated in 88% yield (by ¹H NMR based on CH₂Br₂) and fluoronaphthalene **1a** was consumed completely (by ¹⁹F NMR based on CF₃Ph). The residue was purified by column chromatography on silica gel (hexane/ethyl acetate, 10:1) to give nitrile **2a** (178 mg, 82% yield) as a colorless liquid.

Half-reduction of nitriles (step 2): preparation of aldehyde **3a** is described as a typical procedure. To a toluene solution (150 mL) of nitrile **2a** (28.2 g, 156 mmol) was added a toluene solution of diisobutylaluminum hydride (DIBAL, 1.00 M, 184 mL, 181 mmol) at -78 °C. After being stirred for 3 h at the temperature, aqueous hydrochloric acid (6 M, 300 mL) was added and the mixture was allowed to warm to 0 °C. The mixture was filtered through a pad of celite using ethyl acetate as an eluent. Organic materials were extracted with ethyl acetate three times. The combined extracts were washed with brine and dried over anhydrous sodium sulfate. After removal of the solvent under reduced pressure, the residue was purified by column chromatography on silica gel (hexane/ethyl acetate, 4:1) to give aldehyde **3a** (19.9 g, 81% yield) as a yellow liquid.

Difluorovinylidenation of aldehydes (steps 3 and 4): preparation of 1,1-difluoroallene **4a** is described as a typical procedure. To a THF solution (350 mL) of lithium diisopropylamide (LDA, 479 mmol), prepared from diisopropylamine (48.3 mL, 479 mmol) and butyllithium (1.60 M in hexane, 300 mL, 480 mmol) at -78 °C for 30 min, was added 1,1,1-trifluoro-2-iodoethane (23.4 mL, 240 mmol) at -95 °C. After being stirred for 30 min at the temperature, a THF solution (300 mL) of aldehyde **3a** (368 mg, 200 mmol) was added. The mixture was stirred at -90 °C for 1 h and allowed to warm to -50 °C. Acetic anhydride (28.3 mL, 299 mmol) was added. After the mixture was allowed to warm to 0 °C and stirred for 5 h, the reaction was quenched with saturated aqueous ammonium

chloride (150 mL). Organic materials were extracted with ethyl acetate three times. The combined extracts were washed with brine and dried over anhydrous sodium sulfate. After removal of the solvent under reduced pressure, a crude 3,3-difluoro-2-iodoallyl acetate was obtained. This material was used without further purification for the next step.

To a DMF suspension (120 mL) of zinc powder (2.61 g, 399 mmol) was added a DMF solution (180 mL) of the crude acetate at -20 °C. After being stirred for 2 h at the temperature, the mixture was allowed to warm to room temperature. The mixture was filtered through a pad of celite using ether as an eluent, and saturated aqueous ammonium chloride (100 mL) was added. Organic materials were extracted with ether three times. The combined extracts were washed with brine and dried over anhydrous sodium sulfate. After removal of the solvent under reduced pressure, the residue was purified by column chromatography on silica gel (hexane) to give 1,1-difluoroallene **4a** (33.6 g, 73% yield, 2 steps) as a colorless liquid.

In(III)-catalyzed cyclization of difluoroallenes (step 5): synthesis of fluoroarene **1h** is described as a typical procedure. To a 1,2-dichloroethane suspension (4 mL) of indium(III) bromide (4 mg, 0.01 mmol) was added a 1,2-dichloroethane solution (10 mL) of 1,1-difluoroallene **4a** (120 mg, 0.523 mmol) at room temperature. After being stirred for 1 h, the mixture was then poured into pH 7 phosphate buffer (7 mL). Organic materials were extracted with dichloromethane three times. The combined extracts were washed with brine and dried over anhydrous sodium sulfate. After removal of the solvent under reduced pressure, the residue was purified by column chromatography on silica gel (hexane) to give fluoroarene **1h** (103 mg, 94% yield) as colorless crystals.

Acknowledgements

This research was supported by JSPS KAKENHI Grant Number JP16H04105 (J.I.), MEXT KAKENHI Grant Number JP16H01002 (J.I.), and JSPS KAKENHI Grant Number JP15K05414 (K.F.). Tosoh F-Tech, Inc. is acknowledged for the generous gift of 1,1,1-trifluoro-2-iodoethane.


Keywords: aromatic substitution • fluorine • indium • microwave chemistry • ring extension

- [1] For fluorine-free PAHs as organic semiconducting materials, including phenacenes, see: a) H. Okamoto, N. Kawasaki, Y. Kaji, Y. Kubozono, A. Fujiwara, M. Yamaji, *J. Am. Chem. Soc.* **2008**, *130*, 10470–10471; b) J. E. Anthony, *Angew. Chem. Int. Ed.* **2008**, *47*, 452–483; c) Y. Yamashita, *Sci. Technol. Adv. Mater.* **2009**, *10*, 024313; d) Y. Kubozono, X. He, S. Hamao, K. Teranishi, H. Goto, R. Eguchi, T. Kambe, S. Gohda, Y. Nishihara, *Eur. J. Inorg. Chem.* **2014**, *2014*, 3806–3819.
- [2] a) B. E. Smart, in *Organofluorine Chemistry, Principles and Commercial Applications* (Eds.: R. E. Banks, B. E. Smart, J. C. Tatlow), Plenum Press, New York, **1994**, pp. 57–88; b) T. Hiyama, in *Organofluorine Compounds: Chemistry and Applications*, Springer, Berlin, **2000**, pp. 1–23; c) K. Uneyama, in *Organofluorine Chemistry*, Blackwell, Oxford, **2006**, pp. 1–100.
- [3] S. D. Sharma, S. Doraiswamy, *Indian J. Pure Appl. Phys.* **1983**, *21*, 445–452.
- [4] a) *Printed Organic and Molecular Electronics* (Eds.: D. Gamota, P. Brazis, K. Kalyanasundaram, J. Zhang), Springer, Berlin, **2005**; b) M. Singh, H. M. Haverinen, P. Dhagat, G. E. Jabbour, *Adv. Mater.* **2010**, *22*, 673–685.
- [5] a) K. Fuchibe, T. Morikawa, K. Shigeno, T. Fujita, J. Ichikawa, *Org. Lett.* **2015**, *17*, 1126–1129; b) K. Fuchibe, T. Morikawa, R. Ueda, T. Okauchi, J. Ichikawa, *J. Fluorine Chem.* **2015**, *179*, 106–115.

- [6] F. B. Mallory, C. W. Mallory, C. K. Regan, R. J. Aspden, A. B. Ricks, J. M. Racowski, A. I. Nash, A. V. Gibbons, P. J. Carroll, J. M. Bohlen, *J. Org. Chem.* **2013**, *78*, 2040–2045.
- [7] Polyfluorinated PAHs, typically bearing tetrafluorobenzo moieties, have been prepared for developing organic n-type semiconducting materials during this decade. See: a) F. Babudri, G. M. Farinola, F. Naso, R. Ragni, *Chem. Commun.* **2007**, 1003–1022; b) M. L. Tang, Z. Bao, *Chem. Mater.* **2011**, *23*, 446–455.
- [8] a) M. Yokota, K. Fuchibe, M. Ueda, Y. Mayumi, J. Ichikawa, *Org. Lett.* **2009**, *11*, 3994–3997; b) K. Oh, K. Fuchibe, J. Ichikawa, *Synthesis* **2011**, 881–886; c) K. Oh, K. Fuchibe, M. Yokota, J. Ichikawa, *Synthesis* **2012**, *44*, 857–861; d) K. Fuchibe, M. Abe, K. Oh, J. Ichikawa, *Org. Synth.* **2016**, *93*, 352–366.
- [9] a) K. Fuchibe, Y. Mayumi, N. Zhao, S. Watanabe, M. Yokota, J. Ichikawa, *Angew. Chem. Int. Ed.* **2013**, *52*, 7825–7828; b) K. Fuchibe, Y. Mayumi, M. Yokota, H. Aihara, J. Ichikawa, *Bull. Chem. Soc. Jpn.* **2014**, *87*, 942–949.
- [10] Syntheses of pinpoint-fluorinated PAHs via ring construction other than ours (refs 5 and 9) have been mainly performed by oxidative photocyclization. See: a) S. Mirsadeghi, G. K. B. Prasad, N. Whittaker, D. R. Thakker, *J. Org. Chem.* **1989**, *54*, 3091–3096; b) H. Li, K.-H. He, J. Liu, B.-Q. Wang, K.-Q. Zhao, P. Hu, Z.-J. Shi, *Chem. Commun.* **2012**, *48*, 7028–7030; c) Z. Li, R. J. Twieg, *Chem. Eur. J.* **2015**, *21*, 15534–15539; d) S. Banerjee, S. Sinha, P. Pradhan, A. Caruso, D. Liebowitz, D. Parrish, M. Rossi, B. Zajc, *J. Org. Chem.* **2016**, *81*, 3983–3993. See also: e) K. Kamikawa, I. Takemoto, S. Takemoto, H. Matsuzaka, *J. Org. Chem.* **2007**, *72*, 7406–7408; f) H. Tsuji, Y. Ueda, L. Iliès, E. Nakamura, *J. Am. Chem. Soc.* **2010**, *132*, 11854–11855.
- [11] For reviews on synthesis of fluorinated arenes via fluorine introduction, see: a) M. G. Campbell, T. Ritter, *Chem. Rev.* **2015**, *115*, 612–633; b) P. A. Champagne, J. Desroches, J.-D. Hamel, M. Vandamme, J.-F. Paquin, *Chem. Rev.* **2015**, *115*, 9073–9174.
- [12] For reviews on S_NAr reactions, see: a) H. Amii, K. Uneyama, *Chem. Rev.* **2009**, *109*, 2119–2183; b) C. Liu, B. Zhang, *Chem. Rec.* **2016**, *16*, 667–687.
- [13] [Carbon nucleophiles] a) S. Caron, E. Vazquez, J. M. Wojcik, *J. Am. Chem. Soc.* **2000**, *122*, 712–713; b) X. Ji, T. Huang, W. Wu, F. Liang, S. Cao, *Org. Lett.* **2015**, *17*, 5096–5099. [Phosphine nucleophiles] c) M. Tepper, O. Stelzer, T. Häusler, W. S. Sheldrick, *Tetrahedron Lett.* **1997**, *38*, 2257–2258; d) M. W. Haenel, S. Oevers, J. Bruckmann, J. Kuhnigk, C. Krüger, *Synlett* **1998**, 301–303. [Nitrogen nucleophiles] e) F. Diness, D. P. Fairlie, *Angew. Chem. Int. Ed.* **2012**, *51*, 8012–8016.
- [14] For reviews on microwave-assisted organic synthesis, see: a) C. O. Kappe, *Angew. Chem. Int. Ed.* **2004**, *43*, 6250–6284; b) S. Caddick, R. Fitzmaurice, *Tetrahedron* **2009**, *65*, 3325–3355.
- [15] For specific examples of the effect of microwave irradiation on S_NAr reaction, see: a) S.-W. Yang, G. Ho, D. Tulshian, W. J. Greenlee, J. Anthes, X. Fernandez, R. L. McLeod, J. A. Hey, X. Xu, *J. Med. Chem.* **2009**, *52*, 5323–5329. See also: b) R. Y. Iliashenko, N. Y. Gorobets, A. O. Doroshenko, *Tetrahedron Lett.* **2011**, *52*, 5086–5089; c) R. Singh, B. K. Allam, D. S. Raghuvanshi, K. N. Singh, *Tetrahedron* **2013**, *69*, 1038–1042.
- [16] See Supporting Information for details.
- [17] The substitution of **1a** with ethyl propionate did not proceed under the identical conditions to Table 1, Entry 5. The substitution of 1-bromo- or 1-chloronaphthalene with propionitrile hardly proceeded to afford **2a** in 3% yield each.
- [18] For reviews on triphenylene syntheses, see: a) D. Pérez, E. Guitián, *Chem. Soc. Rev.* **2004**, *33*, 274–283; b) D. Pérez, D. Peña, E. Guitián, *Eur. J. Org. Chem.* **2013**, *2013*, 5981–6013.
- [19] a) R. E. Lehr, S. Kumar, W. Levin, A. W. Wood, R. L. Chang, A. H. Conney, H. Yagi, J. M. Sayer, D. M. Jerina, in *Polycyclic Hydrocarbons and Carcinogenesis* (Ed.: R. D. Harvey), American Chemical Society, **1985**, Vol. 283, pp. 63–84; b) S. S. Hecht, S. Amin, A. A. Melikian, E. J. Lavoie, D. Hoffmann, in *Polycyclic Hydrocarbons and Carcinogenesis* (Ed.: R. D. Harvey), American Chemical Society, **1985**, Vol. 283, pp. 85–105.
- [20] a) R. J. Olsen, S. R. Pruet, *J. Org. Chem.* **1985**, *50*, 5457–5460; b) J. Carreras, G. Gopakumar, L. Gu, A. Gimeno, P. Linowski, J. Petušková, W. Thiel, M. Alcarazo, *J. Am. Chem. Soc.* **2013**, *135*, 18815–18823. See also: c) T. Fujita, I. Takahashi, M. Hayashi, J. Wang, K. Fuchibe, J. Ichikawa, *Eur. J. Org. Chem.* **2017**, 262–265.

Entry for the Table of Contents

COMMUNICATION



Fluoroarenes bearing no electron-withdrawing groups (non-activated fluoroarenes) readily underwent nucleophilic aromatic substitution with α -cyanocarbanions under microwave irradiation. The sequence (i) formylalkylation involving the cyanoalkylation of fluoroarenes, (ii) difluorovinylideneation, and (iii) Friedel-Crafts-type cyclization, afforded one benzene ring-extended fluoroarenes in one cycle. Furthermore, the performance of multiple cycles successfully provided higher order pinpoint-fluorinated polycyclic aromatic hydrocarbons (F-PAHs).

Kohei Fuchibe, Hisanori Imaoka, and
Junji Ichikawa*

Page No. – Page No.

**Synthesis of Pinpoint-Fluorinated
Polycyclic Aromatic Hydrocarbons:
Benzene Ring Extension Cycle
Involving Microwave-Assisted S_NAr
Reaction**