

Effect of *Chlorella*-derived multicomponent supplementation on maximal oxygen uptake and serum vitamin B₂ concentration in young men

Asako Zempo-Miyaki,¹ Seiji Maeda² and Takeshi Otsuki^{1,*}

¹Faculty of Sport and Health Sciences, Ryutsu Keizai University, 120 Ryugasaki, Ibaraki 301-8555, Japan

²Faculty of Health and Sport Sciences, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki 305-8577, Japan

(Received 18 April, 2017; Accepted 12 June, 2017; Published online 18 August, 2017)

Chlorella is a unicellular green alga that contains high levels of proteins, vitamins and minerals. The present study investigated the effects of a 4-week *Chlorella*-derived multicomponent supplementation on maximal oxygen uptake and circulating vitamin B₂ levels in healthy men. Thirty-four participants were randomly divided into two groups: placebo or *Chlorella*. Prior to the intervention, we observed that the intake of several minerals and soluble vitamins did not satisfy the nutrient requirements of either group by assessing the frequency of daily food intake. There was a significant negative relationship between the pre-intervention maximal oxygen uptake and serum vitamin B₂ concentrations in all subjects ($r = -0.372$). Maximal oxygen uptake significantly increased after *Chlorella* supplementation (before vs after, 42.1 ± 1.5 vs 44.9 ± 1.6 ml/kg/min), while serum vitamin B₂ concentrations did not (14.6 ± 0.9 vs 14.0 ± 0.9 µg/L). In conclusion, *Chlorella*-derived multicomponent supplementation increases maximal oxygen uptake in individuals with an insufficient micronutrient status, although there was no association between the increase in aerobic capacity and serum levels of vitamin B₂.

Key Words: aerobic capacity, *Chlorella*, daily food intake frequency, maximal oxygen uptake, vitamin B₂

A well-balanced diet is one of the most important factors for being fit and leading a healthy life. In the last few decades, dietary social problems, including skipping breakfast, eating meals irregularly and consuming fast food, have become increasingly common and have led to malnutrition in young individuals.⁽¹⁻³⁾ In particular, university students living away from home are at a high risk of poor nutrition.⁽⁴⁾ The National Health and Nutrition Examination Survey by the Ministry of Health, Labour and Welfare of Japan reported that many young Japanese men have poor dietary habits, including skipping breakfast or having a low vegetable intake, as compared with women and/or other generations. In particular, the average vegetable intake among Japanese men aged 20–29 years in 2014 was 237 g/day, despite the recommended intake being 350 g/day; only 19% of young Japanese men achieved the recommended level. Fujii *et al.*⁽⁵⁾ reported that Japanese college students who regularly skip breakfast also tend to have a poor vegetable intake. Such dietary problems are not unique to Japan, but also occur throughout the world.^(1,2,6) As a solution to such dietary problems in young individuals, dietary supplements may be effective at overcoming nutrient deficiencies by providing a better nutrient balance.

As a result of an evidence-based analysis, the Academy of Nutrition and Dietetics, Dietitians of Canada and American College of Sports Medicine have stated that well-balanced nutrition is necessary for maximal exercise performance. Exercise

training increases micronutrient requirements, and a daily multivitamin and multimineral supplement would benefit athletes who have a poor micronutrient status.⁽⁷⁾ In our previous cross-over study, a *Chlorella*-derived multicomponent supplementation increased the aerobic exercise capacity of healthy young individuals.⁽⁸⁾ However, our previous study did not explore the nutritional status of the participants or the mechanisms underlying the increase in aerobic exercise capacity.

Chlorella is a unicellular green alga that contains high levels of proteins, chlorophylls, vitamins, minerals and dietary fiber. A lack of vitamin B₁, B₂, B₆ and C for 10 weeks decreased the peak oxygen uptake in young individuals.⁽⁹⁾ In addition, vitamin B₂ intake is an important factor regarding endurance exercise capacity.⁽¹⁰⁾ Therefore, we hypothesized that *Chlorella*-derived supplementation would increase maximal oxygen uptake ($\dot{V}O_{2max}$) by improving the levels of vitamin B₂ and other micronutrients in individuals with an insufficient micronutrient status. To test this hypothesis, we assessed daily food intake frequency and examined the effects of a 4-week *Chlorella*-derived supplementation on the $\dot{V}O_{2max}$ and circulating vitamin B₂ levels in healthy young men using a double-blinded study design.

Materials and Methods

Subjects and experimental design. A total of 34 young men volunteered to participate in this study. After pre-supplementation tests, the subjects were randomly divided into the placebo ($n = 17$) and *Chlorella* ($n = 17$) groups in a double-blinded manner. During the trial, the participants self-administered 30 tablets of either *Chlorella* or the placebo per day (15 tablets twice daily, after breakfast and dinner) for 4 weeks. This dosage was in accordance with our previous studies^(8,11,12) and the general recommended dosage for Japanese consumers. The participants were asked not to modify their lifestyle during the trial period. Post-supplementation measurements and blood sampling were performed one day after the final tablet intake.

This study was approved by the Ethics Committee of Ryutsu Keizai University. The protocol of this study conformed to the principles of the Declaration of Helsinki. All participants provided written informed consent prior to study initiation.

A power calculation was performed for Pearson's correlation coefficients in all participants and for repeated measures two-way analysis of variance (ANOVA) using G*Power 3.⁽¹³⁾ The sample size of this study was enough to detect the correlation and effects of supplementation at 90% power and with an α of 5% when effect

*To whom correspondence should be addressed.
E-mail: takeshi.otsuki17@gmail.com

size was assumed as medium.⁽¹⁴⁾

The *Chlorella* (SunChlorella A; Sun Chlorella Corp., Kyoto, Japan) and placebo tablets used in this study were the same as those used in our previous studies.^(8,11,12) Briefly, the *Chlorella* tablet comprised dried *Chlorella pyrenoidosa* powder as the main ingredient. The mass of each *Chlorella* tablet was 200 mg. The nutritional values per 100 g of the tablet were as follows: energy, 399 kcal; water, 5.3 g; protein, 60.8 g; lipid, 9.2 g; saccharide, 6.3 g; dietary fiber, 11.9 g; and ash, 6.5 g (including 5.61 mg of vitamin B₂). The main ingredient of the placebo tablet was lactose. The color and shape of the placebo tablet were similar to that of the *Chlorella* tablet, but the protein (2.0 g), dietary fiber (1.1 g) and ash (2.2 g) contents in the placebo tablet were markedly lower than those in the *Chlorella* tablet.

Maximal exercise testing. Subjects refrained from alcohol consumption starting on the day before testing, intense physical activity and caffeine consumption starting at 9 p.m. on the day before testing, and drinking and eating excluding water for 1 h before the testing.

Maximal exercise testing involved incremental cycling to exhaustion (4 min at 70 W, with a 30 W increase every min), with monitoring of breath-by-breath oxygen uptake and carbon dioxide production (AE300S; Minato Medical Science, Osaka, Japan), heart rate (HR) calculation using 3-lead electrocardiography (LRR-03; GMS, Tokyo, Japan) and ratings of perceived exertion (RPE, Borg's 6–20 scale). We considered a participant as having achieved maximal exertion when at least two of the following four criteria were met: 1) a plateau in oxygen uptake with increasing exercise intensity (<100 ml/min); 2) achievement of age-predicted maximal HR (± 10 bpm); 3) a respiratory exchange ratio of at least 1.15; and 4) an RPE of at least 18 units.⁽⁸⁾ In our laboratory, the day-to-day coefficient of variation for $\dot{V}O_{2\max}$ was $4.7 \pm 4.0\%$.

Serum vitamin B₂ analysis. Serum vitamin B₂ concentration was measured using a microbiological test. The subjects refrained from alcohol consumption from the day before testing. Blood samples were collected from the antecubital vein by venipuncture after an overnight fast, placed in chilled tubes and centrifuged at 3,000 rpm for 15 min at 4°C. The plasma was stored at -80°C prior to analysis. Serum concentrations of vitamin B₂ were determined using a commercial kit including a microtiter plate coated with *Lactobacillus rhamnosus*, according to the manufacturer's protocol (ID-Vit[®] Vitamin B₂; Immundiagnostik AG, Bensheim, Germany).

Dietary intake. Dietary habits were assessed using a validated, brief, self-administered dietary history questionnaire (BDHQ).⁽¹⁵⁾ The BDHQ is a four-page structured questionnaire that enquires about the consumption frequency of 58 food and beverage items, with specified serving sizes described in terms of the natural portion or the standard weight and volume measurement of servings commonly consumed in the general Japanese population. Validation of the BDHQ was performed using 16-day weighed dietary records as the gold standard.⁽¹⁵⁾

With regard to protein, calcium, magnesium, iron, zinc, copper, retinol, vitamin B₁, vitamin B₂, niacin, vitamin B₆, vitamin B₁₂ and vitamin C, the intake amounts are expressed as the ratio to the recommended dietary amount (RDA) of the Dietary Reference Intake for Japanese 2015. Sodium intake is expressed as the ratio to the estimated average requirement (EAR), as the RDA of sodium was not established in the Dietary Reference Intake. For nutrients without a RDA or an EAR (n-6 fatty acid, n-3 fatty acid, potassium, phosphorus, manganese, vitamin D, vitamin K, α -tocopherol and pantothenic acid), the sufficiency rates were estimated based on the adequate intake (AI). The RDA and EAR indicate the amount that would meet the requirement for 97–98 and 50% of the population, respectively. The AI was developed when EAR and RDA could not be set because of insufficient scientific evidence and indicates the amount of nutrients required to maintain a healthy status.

Body mass and fat mass. The subjects refrained from alcohol consumption from the day before testing. Body mass was measured and body fat mass was assessed by bioelectrical impedance analysis after an overnight fast (InBody 430; InBody Japan, Tokyo, Japan).

Statistical analysis. Results are presented as the mean \pm SE. Inter-group comparisons for patient variables before supplementation were performed using unpaired *t* tests. Relationships between two variables were assessed using Pearson's correlation coefficient. To compare the effects of *Chlorella* tablet intake with placebo supplementation, repeated-measures two-way ANOVA was used. If a significant *F* value was found, a post-hoc Fisher's exact test was conducted to identify the effects of placebo and *Chlorella* tablet intake. *P* values <0.05 were considered statistically significant.

Results

Table 1 shows the anthropometric characteristics of the participants. There were no significant differences in the pre-measurement data between the *Chlorella* and placebo groups. Moreover, the height, body mass and body fat mass of both groups did not change significantly after the 4-week intervention.

Table 2 shows dietary habits of the two groups before the intervention and the nutrient intakes including *Chlorella* supplementation. There were no differences in the daily nutrient intake between the groups before the tablet intake period. Several nutrients did not satisfy the nutrient demand (i.e., RDAs, EARs or AIs) in both groups, as follows: potassium, calcium, magnesium, zinc, manganese, vitamin A, vitamin B₁, vitamin B₂, vitamin B₆ and vitamin C. On the other hand, sodium was above the nutrient demand in both groups. The energy intake was lower than the demand in both groups, but the intake of protein and fat was higher than the demand. The sufficiency rate of iron, vitamin B₂, D and K, and niacin increased by $\geq 20\%$ with *Chlorella* supplementation.

There was a significant negative relationship between the $\dot{V}O_{2\max}$ and serum vitamin B₂ concentration in all participants before the intervention (Fig. 1). The $\dot{V}O_{2\max}$ prior to and following the intervention is shown in Fig. 2. There was a significant interaction in the changes between the *Chlorella* and control groups. The $\dot{V}O_{2\max}$ significantly increased following *Chlorella*-derived supplementation. Changes in vitamin B₂ concentration before and after the intervention are shown in Fig. 3; there was no significant interaction between the *Chlorella* and control groups. Additionally, the relationship between the change in the $\dot{V}O_{2\max}$ and that of the serum vitamin B₂ levels did not reach statistical significance in the *Chlorella* group ($r = -0.311$, $p = 0.22$).

Discussion

We investigated the effects of a 4-week *Chlorella*-derived multicomponent supplementation on the $\dot{V}O_{2\max}$ and circulating vitamin B₂ levels in healthy young men. Prior to the intervention,

Table 1. Anthropometric characteristics

		Pre	Post
Age (years)	Placebo	20.1 \pm 0.3	—
	<i>Chlorella</i>	19.6 \pm 0.3	—
Height (cm)	Placebo	172.8 \pm 1.8	—
	<i>Chlorella</i>	172.4 \pm 1.5	—
Body mass (kg)	Placebo	70.0 \pm 3.5	69.6 \pm 3.5
	<i>Chlorella</i>	69.2 \pm 2.7	68.8 \pm 2.8
Body fat mass (kg)	Placebo	16.0 \pm 1.3	15.3 \pm 0.9
	<i>Chlorella</i>	14.6 \pm 0.9	14.1 \pm 0.9

Data are shown as the mean \pm SE.

Table 2. Nutrient intake and sufficiency rate

Nutrient	Placebo group		<i>Chlorella</i> group			
			Baseline		With supplement	
	Mean ± SE	SR (%)	Mean ± SE	SR (%)	Mean ± SE	SR (%)
Energy intake (kcal/d)	2,221 ± 281	82.4	2,055 ± 107	75.9	2,078 ± 107*	76.7
Protein (g/d)	77 ± 13	128.7	64 ± 4	106.6	67 ± 4*	112.3
Fat (g/d)	58 ± 9	—	53 ± 3	—	54 ± 3*	—
n-6 (g/d)	11.3 ± 1.5	102.8	10.6 ± 0.8	96.5	10.7 ± 0.8*	97.3
n-3 (g/d)	2.6 ± 0.3	131.8	2.3 ± 0.3	116.3	2.4 ± 0.3*	121.0
Carbohydrate (g/d)	325 ± 43	—	317 ± 21	—	318 ± 21*	—
Na (mg/d)	5,530 ± 785	921.6	4,444 ± 268	740.6	4,446 ± 268*	740.9
K (mg/d)	2,175 ± 318	87.0	1,976 ± 233	79.0	2,030 ± 233*	81.2
Ca (mg/d)	459 ± 78	57.3	400 ± 40	50.0	418 ± 40*	52.3
Mg (mg/d)	242 ± 36	71.2	202 ± 17	59.4	223 ± 17*	65.5
P (mg/d)	1,048 ± 164	104.8	888 ± 52	88.8	982 ± 52*	98.2
Fe (mg/d)	8.0 ± 1.2	114.7	6.6 ± 0.6	94.5	14.4 ± 0.7*	205.9
Zn (mg/d)	9.5 ± 1.4	95.2	8.4 ± 0.5	84.4	8.5 ± 0.5*	85.0
Cu (mg/d)	1.29 ± 0.17	143.7	1.17 ± 0.08	130.2	1.18 ± 0.08*	131.6
Mn (mg/d)	3.4 ± 0.3	85.6	3.2 ± 0.2	79.8	3.2 ± 0.2	79.8
Retinol equivalent (µg/d)	735 ± 123	86.4	586 ± 110	68.9	611 ± 110*	71.9
Vitamin D (µg/d)	10.3 ± 1.6	186.7	8.0 ± 1.4	145.6	26.3 ± 1.5*	478.7
Vitamin K (µg/d)	312 ± 55	208.3	287 ± 56	191.4	359 ± 56*	239.3
α-Tocopherol (mg/d)	6.8 ± 1.0	105.2	6.3 ± 0.6	97.3	6.5 ± 0.6*	99.6
Vitamin B ₁ (mg/d)	0.8 ± 0.1	57.1	0.7 ± 0.1	51.7	0.8 ± 0.1*	59.1
Vitamin B ₂ (mg/d)	1.2 ± 0.2	75.4	1.1 ± 0.1	66.7	1.4 ± 0.1*	86.2
Niacin (mg/d)	16.5 ± 2.7	109.9	13.2 ± 1.2	87.8	16.2 ± 1.3*	107.8
Vitamin B ₆ (mg/d)	1.2 ± 0.2	84.5	1.0 ± 0.1	72.2	1.1 ± 0.1*	80.6
Vitamin B ₁₂ (µg/d)	8.6 ± 1.46	357.0	6.1 ± 0.9	255.2	6.2 ± 0.9*	256.1
Folic acid (µg/d)	305.9 ± 41.3	127.5	264.0 ± 40.3	110.0	264.2 ± 40.3*	110.1
Pantothenic acid (mg/d)	6.9 ± 1.1	137.7	6.1 ± 0.4	122.5	6.4 ± 0.4*	127.9
Vitamin C (mg/d)	89.1 ± 14.7	89.1	88.3 ± 21.1	88.3	88.5 ± 21.1*	88.5

SR, sufficiency rate. * $p < 0.05$ vs baseline. There were no differences in the daily nutrient intake between the groups before the tablet intake period. The SR of iron, vitamin B₂, D and K, and niacin increased by $\geq 20\%$ with *Chlorella* supplementation (shown as bold font).




Fig. 1. Relationship between maximal oxygen uptake and serum vitamin B₂ concentrations in all participants before the intervention. There was a negative relationship between the maximal oxygen uptake and serum vitamin B₂ concentrations in all participants before the intervention.




Fig. 2. Maximal oxygen uptake pre and post intervention of the placebo and *Chlorella* groups. Values are mean ± SE. The maximal oxygen uptake was increased following *Chlorella*-derived multicomponent supplementation.

the daily intake of calcium, magnesium, vitamin B₁, vitamin B₂, vitamin B₆ and vitamin C did not satisfy nutrient demands in both groups. However, vitamin B₂ intake was markedly increased in

the *Chlorella* group following supplementation. There was a significant negative relationship between the pre-intervention VO₂max value and the serum levels of vitamin B₂ in all subjects.




Fig. 3. Serum vitamin B₂ concentrations pre and post intervention of the placebo and *Chlorella* groups. Values are mean \pm SE.

In addition, in the *Chlorella* group, the $\dot{V}O_{2\max}$ was significantly increased following the supplementation, and serum vitamin B₂ levels showed a decreasing trend. Therefore, we concluded that *Chlorella*-derived supplementation increases the $\dot{V}O_{2\max}$ in individuals with an insufficient micronutrient status, although there was no association between the increased $\dot{V}O_{2\max}$ and blood levels of vitamin B₂.

In a previous crossover study, we demonstrated that *Chlorella*-derived supplementation increased the $\dot{V}O_{2\max}$ in young men ($n = 7$) and women ($n = 3$).⁽⁸⁾ In this study, we investigated the effects of a 4-week *Chlorella*-derived supplementation on aerobic exercise capacity in 34 young men, thus increasing the number of participants and eliminating any confounding effects associated with the female menstrual cycle. In addition, the current study assessed the daily food intake frequency in order to clarify the participants' dietary habits. This study supports the results of the previous study,⁽⁸⁾ as a 4-week *Chlorella*-derived supplementation significantly increased the $\dot{V}O_{2\max}$ in young men with inadequate dietary habits.

In the present cross-sectional analysis of pre-supplementation values, we showed for the first time that there was a significant negative correlation between the $\dot{V}O_{2\max}$ and serum vitamin B₂ concentrations. These results suggest that individuals with a higher aerobic capacity metabolize more vitamin B₂ than do those with a lower aerobic capacity. Previously, Belko *et al.*^(16,17) reported that various lifestyle modifications, including dietary modification and/or regular exercise, would lead to increased vitamin B₂ demands. Moreover, Manore⁽¹⁸⁾ demonstrated that considerably more vitamin B₂ was required when dieting was combined with exercise. It is possible that the lower circulating levels of vitamin B₂ in individuals with a higher aerobic capacity were due to the higher constitutive usage of vitamin B₂.

Before the intervention, the sufficiency rate of vitamin B₂ was 67% in the *Chlorella* group. However, after the *Chlorella*-derived supplementation, this rate was increased to 86%. No other nutrients demonstrated such a dramatic increase in the sufficiency rate. Therefore, this study subsequently investigated serum vitamin B₂ concentrations. Vitamin B₂ is necessary for the synthesis of two important coenzymes, flavin mononucleotide and flavin adenine dinucleotide. These coenzymes are especially important for the metabolism of glucose, fatty acids, glycerol and amino acids. In a previous study, reduced intake of vitamins B₁, B₂, B₆ and C was

associated with a decreased aerobic exercise capacity in young men.⁽⁹⁾ In addition, Peluchetti *et al.*⁽¹⁰⁾ demonstrated that vitamin B₂ supplementation dramatically increased aerobic exercise capacity (before: 10.5–14.4 ml/kg/min; after: 30.5–39.3 ml/kg/min) in patients with multiple acyl-coenzyme A dehydrogenase deficiency. These results suggest that vitamin B₂ and other soluble vitamins are important factors associated with the $\dot{V}O_{2\max}$. However, the present study did not show a significant change in circulating vitamin B₂ levels following *Chlorella* supplementation. Dietary modifications and/or regular exercise increase the vitamin B₂ demand,^(16–18) and the circulating level of vitamin B₂ was lower in participants with a higher aerobic capacity in the present study. Therefore, the additional vitamin B₂ supplied by the *Chlorella*-derived tablet and the increased vitamin B₂ demand due to an increase in aerobic capacity may have counteracted each other, resulting in no change in the vitamin B₂ concentration.

The National Health and Nutrition Examination Survey by the Ministry of Health, Labour and Welfare of Japan has reported that vegetable intake in Japan has declined in recent years across all generations, and the decline has been particularly pronounced in men. Indeed, some vitamins and minerals (i.e., potassium, calcium, magnesium, zinc, manganese, retinol, vitamin B₁, vitamin B₂, vitamin B₆ and vitamin C) were deficient in the participants of the present study. The Academy of Nutrition and Dietetics, Dietitians of Canada and American College of Sports Medicine⁽⁷⁾ have suggested that calcium, iron, zinc, magnesium, vitamin B, vitamin C and vitamin E are important nutrients for athletes. The *Chlorella*-derived supplement used in the present study contained many vitamins and minerals (e.g., vitamin A, B vitamins, vitamin C, vitamin E, calcium, folate, iron, pantothenic acid, potassium, magnesium and zinc). In particular, it should be noted that the sufficiency rate of vitamin B₂ was essentially improved following self-administration of *Chlorella*-derived tablets.

It is unclear whether vitamin B₂ directly affects aerobic capacity. Future investigations should focus on whether the bioavailability of vitamin B₂ is associated with the observed increase in the $\dot{V}O_{2\max}$ following *Chlorella*-derived tablet intake. Since B vitamins are water-soluble, changes in blood and urine vitamin B₂ concentrations should be examined.

In conclusion, we found a significant negative relationship between the $\dot{V}O_{2\max}$ and serum vitamin B₂ concentrations before the intervention. In addition, we observed that a 4-week *Chlorella*-derived multicomponent supplementation increased the $\dot{V}O_{2\max}$ in male university students with insufficient dietary habits, although there was no association between the increase in the $\dot{V}O_{2\max}$ and circulating levels of vitamin B₂.

Abbreviations

AI	adequate intake
ANOVA	analysis of variance
BDHQ	brief, self-administered dietary history questionnaire
EAR	estimated average requirement
HR	heart rate
RDA	recommended dietary amount
RPE	ratings of perceived exertion
$\dot{V}O_{2\max}$	maximal oxygen uptake

Conflict of Interest

Sun Chlorella Corp. provided funding for the study and supplied the test supplements used in the study. TO has received speaker's honorarium and payment for manuscripts not involving this study from Sun Chlorella Corp. AZ-M and SM have no competing interests.

References

- 1 Lien L. Is breakfast consumption related to mental distress and academic performance in adolescents? *Public Health Nutr* 2007; **10**: 422–428.
- 2 Paeratakul S, Ferdinand DP, Champagne CM, Ryan DH, Bray GA. Fast-food consumption among US adults and children: dietary and nutrient intake profile. *J Am Diet Assoc* 2003; **103**: 1332–1338.
- 3 Tanaka M, Mizuno K, Fukuda S, Shigihara Y, Watanabe Y. Relationships between dietary habits and the prevalence of fatigue in medical students. *Nutrition* 2008; **24**: 985–989.
- 4 Shimbo S, Zhang ZW, Matsuda-Inoguchi N, *et al*. Effects of life away from home and physical exercise on nutrient intake and blood/serum parameters among girl students in Japan. *Tohoku J Exp Med* 2004; **203**: 275–286.
- 5 Fujii H, Nakano T, Muto T, Aikawa K. Skipping breakfast is associated with poor vegetable intake among college students in Japan. *Dokkyo J Med Sci* 2010; **37**: 47–54.
- 6 Al Ani MF, Al Subhi LK, Bose S. Consumption of fruits and vegetables among adolescents: a multi-national comparison of eleven countries in the Eastern Mediterranean Region. *Br J Nutr* 2016; **115**: 1092–1099.
- 7 American Dietetic Association; Dietitians of Canada; American College of Sports Medicine, Rodriguez NR, Di Marco NM, Langley S. American College of Sports Medicine position stand. Nutrition and athletic performance. *Med Sci Sports Exerc* 2009; **41**: 709–731.
- 8 Umemoto S, and Otsuki. *Chlorella*-derived multicomponent supplementation increases aerobic endurance capacity in young individuals. *J Clin Biochem Nutr* 2014; **55**: 143–146.
- 9 van der Beek EJ, van Dokkum W, Wedel M, Schrijver J, van den Berg H. Thiamin, riboflavin and vitamin B6: impact of restricted intake on physical performance in man. *J Am Coll Nutr* 1994; **13**: 629–640.
- 10 Peluchetti D, Antozzi C, Roi S, DiDonato S, Cornelio F. Riboflavin responsive multiple acyl-CoA dehydrogenase deficiency: functional evaluation of recovery after high dose vitamin supplementation. *J Neurol Sci* 1991; **105**: 93–98.
- 11 Otsuki T, Shimizu K, Maeda S. Changes in arterial stiffness and nitric oxide production with *Chlorella*-derived multicomponent supplementation in middle-aged and older individuals. *J Clin Biochem Nutr* 2015; **57**: 228–232.
- 12 Otsuki T, Shimizu K, Zempo-Miyaki A, Maeda S. Changes in salivary flow rate following *Chlorella*-derived multicomponent supplementation. *J Clin Biochem Nutr* 2016; **59**: 45–48.
- 13 Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007; **39**: 175–191.
- 14 Cohen J. A power primer. *Psychol Bull* 1992; **112**: 155–159.
- 15 Kobayashi S, Murakami K, Sasaki S, *et al*. Comparison of relative validity of food group intakes estimated by comprehensive and brief-type self-administered diet history questionnaires against 16 d dietary records in Japanese adults. *Public Health Nutr* 2011; **14**: 1200–1211.
- 16 Belko AZ, Obarzanek E, Roach R, *et al*. Effects of aerobic exercise and weight loss on riboflavin requirements of moderately obese, marginally deficient young women. *Am J Clin Nutr* 1984; **40**: 553–561.
- 17 Belko AZ, Meredith MP, Kalkwarf HJ, *et al*. Effects of exercise on riboflavin requirements: biological validation in weight reducing women. *Am J Clin Nutr* 1985; **41**: 270–277.
- 18 Manore MM. Effect of physical activity on thiamine, riboflavin, and vitamin B-6 requirements. *Am J Clin Nutr* 2000; **72** (2 Suppl): 598S–606S.