

Purkinje Arrhythmia Origin Made Easy

著者別名	野上 昭彦
journal or publication title	Circulation. Arrhythmia and electrophysiology
volume	10
number	11
page range	e005889
year	2017-11
権利	(C) 2018 American Heart Association, Inc.
URL	http://hdl.handle.net/2241/00150841

doi: 10.1161/CIRCEP.117.005889

1 **Running title:** Purkinje arrhythmia origin "made easy"

2

3 **Authors' names, academic degrees, and affiliations**

4 Ahmed Karim Talib, MD, PhD; Akihiko Nogami, MD, PhD

5 Department of Cardiology, Faculty of Medicine, University of Tsukuba, Japan

6

7 **Name and address for correspondence**

8 Akihiko Nogami, MD, PhD

9 Department of Cardiology, Faculty of Medicine, University of Tsukuba.

10 1-1-1 Tennodai, Tsukuba 305-8575, Japan

11 E-mail: anogami@md.tsukuba.ac.jp

12 TEL +81-29-853-3142 FAX +81-29-853-3143

13

14 **Total words:** 1,408, **Figures:** 1

15

16

1 Tachycardia originating from the fascicles or His-Purkinje system includes a wide
2 spectrum of arrhythmias such as ventricular fibrillation (VF) and ventricular tachycardia (VT)
3 which could be monomorphic or polymorphic with various substrates, locations and
4 mechanisms.¹ The most common type of reentrant fascicular tachycardia is left posterior
5 fascicular VT (LPF-VT) which was recognized as an electrocardiographic entity by Zipes et al.²
6 who defined its diagnostic triad as following: (1) induction with atrial pacing, (2) right bundle
7 branch block (RBBB) and a left-axis configuration, and (3) manifestation in patients without
8 structural heart disease. In 1981, Belhassen et al.³ reported verapamil sensitivity, as a fourth
9 identifying feature of this tachycardia.

10 To ablate LPF-VT, there are two different successful sites defined by two different
11 strategies (**Figure 1**). The first strategy was reported by Nakagawa et al.⁴ who targeted the
12 earliest presystolic Purkinje potentials that can be recorded at the apical-inferior septum of
13 the left ventricle during VT. The second strategy was reported by Tsuchiya et al.⁵ who
14 emphasized the significance of a late diastolic potential at the basal septal region close to
15 the main trunk of the left bundle branch (LBB). To delineate the reentry circuit, we
16 performed left ventricular septal mapping using an octapolar electrode catheter in 20
17 patients with LPF-VT.⁶ In 15 of 20 patients, 2 distinct sequences of potentials other than LV
18 muscular potential were recorded during VT (**Figure 1**). We named these potentials P1, a
19 mid-diastolic potential recorded earlier from proximal rather than distal electrodes, and P2,
20 the fused presystolic Purkinje potential recorded earlier from distal electrodes. Because the
21 diastolic potential (P1) has been proven to be a critical potential in the VT circuit, this
22 potential can be targeted to cure the tachycardia. Any P1 in the VT circuit can be targeted
23 for catheter ablation. We usually target the apical third of the septum, to avoid the creation
24 of left bundle branch block (LBBB) or atrioventricular block.

1 In the current issue of *Circulation: Arrhythmia and Electrophysiology*, Ma and
2 coworkers⁷ interestingly describe the relationship between 12-lead ECG morphology and
3 the location of the exit of LPF-VT in addition to the H-V interval during tachycardia. They
4 retrospectively analyzed the electrocardiographic and electrophysiological characteristics of
5 41 patients who underwent successful catheter ablation of LPF-VT. Patients were divided
6 into 3 subgroups: proximal, mid, and distal left posterior fascicular groups according to the
7 12-lead ECG morphology, QRS duration and H-V interval during VT. Quick prediction of
8 tachycardia exit and successful ablation sites in LPF-VT is practically useful to guide the
9 direction of further mapping. However, the study results would be practical and made easy
10 when Purkinje fibers are constructed in homogenous 2-D orientation with equal conduction
11 velocity in all parts of the heart. In animal models, conduction velocity of the His-Purkinje
12 system is heterogenous at various parts of the heart and local changes in bundle branch
13 architecture is the most likely cause of significant conduction velocity reduction in the
14 midseptum in addition to presence of regional differences in expression pattern and
15 distribution of gap junction proteins.⁸ Furthermore, there is intense branching of bundle
16 branch fibers in the midseptum resulting in load mismatch or increased path length. In
17 canine hearts, conduction velocity is faster in proximal bundle branch areas versus that in
18 the distal network areas and these variations in delay and conduction velocity may serve as
19 the physiologic substrate for fascicular VT.⁹

20 This complexity of both functional and anatomical aspects of His-Purkinje system
21 results in complex tachycardia circuits and difficulties in predicting its exact origin. The
22 investigators used the H-V interval during LPF-VT as a guide to identify LPF-VT origin
23 (proximal, mid and distal LPF origin).⁷ Theoretically, the upper turn around point of LPF-VT is
24 undetermined while long H-V interval during LPF-VT may indicate the lower turn around

1 point is located basally, conversely; short the H-V interval during LPF-VT indicates that the
2 lower turn-around point is more apical far from His bundle.

3 It is important to consider that, the relative duration of the H-V interval recorded
4 during VT as compared to sinus rhythm would depend on 2 factors: (a) the balance between
5 antegrade and retrograde conduction times from the upper turnaround point of the reentry
6 circuit; and (b) the site of His bundle recording relative to the upper turnaround point.¹⁰

7 Conduction delay in the antegrade limb of the circuit would tend to prolong H-V interval
8 during VT, while retrograde conduction delay to the His bundle recording site as well as the
9 use of relatively proximal His bundle recording site (far from the turnaround point) would
10 tend to shorten it. These facts represent a major study limitation. Another important
11 consideration is due to the fact that there may be arborization of Purkinje fibers beyond the
12 region of reentry, there may be several early myocardial sites with relatively late sites in
13 between. In conclusion, the study conducted by Ma et al. provides excellent pre-operative
14 information to guide mapping and ablation of LPF-VT, however; potential limitations of H-V
15 interval value during tachycardia and surface ECG morphology should be taken into
16 consideration when predicting the origin of LPF-VT.

17

18

1 References

- 2 1. Nogami A. Purkinje-Related Arrhythmias Part I: Monomorphic Ventricular Tachycardias.
3 *Pacing Clin Electrophysiol.* 2011; 34:624–650.
- 4 2. Zipes DP, Foster PR, Troup PJ, Pedersen DH. Atrial induction of ventricular tachycardia:
5 reentry versus triggered automaticity. *Am J Cardiol* 1979; 44: 1-8.
- 6 3. Belhassen B, Rotmensch HH, Laniado S: Response of recurrent sustained ventricular
7 tachycardia to verapamil. *Br Heart J* 1981; 46: 679–682.
- 8 4. Nakagawa H, Beckman KJ, McClelland JH, Wang X, Arruda M, Santoro I, Hazlitt HA,
9 Abdalla I, Singh A, Gossinger H. Radiofrequency catheter ablation of idiopathic left
10 ventricular tachycardia guided by a Purkinje potential. *Circulation.* 1993; 88:2607–2617.
- 11 5. Tsuchiya T, Okumura K, Honda T, et al: Significance of late diastolic potential preceding
12 Purkinje potential in verapamil-sensitive idiopathic left ventricular tachycardia.
13 *Circulation* 1999; 99: 2408–2413.
- 14 6. Nogami A, Naito S, Tada H, Taniguchi K, Okamoto Y, Nishimura S, Yamauchi Y, Aonuma K,
15 Goya M, Iesaka Y, Hiroe M. Demonstration of diastolic and presystolic Purkinje potentials
16 as critical potentials in a macroreentry circuit of verapamil-sensitive idiopathic left
17 ventricular tachycardia. *J Am Coll Cardiol.* 2000; 36:811–823.
- 18 7. Ma W, Lu F, Shehata M, Wang X, Zhang F, Chen B, Wu D, He L, Sun S, Cheng A, Ding J,
19 Cong H, Xu J. Catheter Ablation of Idiopathic Left Posterior Fascicular Ventricular
20 Tachycardia: Predicting the Site of Origin via Mapping and Electrocardiography. *Circ*
21 *Arrhythm Electrophysiol.* 2017; 10: e005240.
- 22 8. van Veen TA, van Rijen HV, van Kempen MJ, Miquerol L, Opthof T, Gros D, Vos
23 MA, Jongsma HJ, de Bakker JM. Discontinuous conduction in mouse bundle branches is

- 1 caused by bundle-branch architecture. *Circulation* 2005; 112:2235-44.
- 2 9. Miquerol L, Meysen S, Mangoni M, Bois P, van Rijen HV, Abran P, Jongsma H, Nargeot
- 3 J, Gros D. Architectural and functional asymmetry of the His-Purkinje system of the
- 4 murine heart. *Cardiovasc Res* 2004; 63:77-86.
- 5 10. Tchou P, Jazayeri M, Denker S, Dongas J, Caceres J, Akhtar M. Transcatheter electrical
- 6 ablation of right bundle branch: A method of treating macroreentrant ventricular
- 7 tachycardia attributed to bundle branch reentry. *Circulation* 1988; 78:246-257.

8

9

10

1 **Figure legends**

2 **Figure 1.**

3 Schematic representation of our hypothetical circuit of reentrant LPF-VT. During VT,
4 diastolic potential (P1) is propagated antegradely. In the distal third of the septum, the
5 antegrade wavefront penetrates the posterior fascicle of the left bundle branch system at
6 the so-called *lower turn around point*. This creates diverging wavefronts traveling to the
7 myocardial exit site as well as activating the posterior fascicle in a retrograde fashion,
8 represented by presystolic potential (P2), until the wavefront reaches the basal septum and
9 exits the fascicle close to the main left bundle then the His bundle. Then propagation
10 wavefront re-enters the zone of slow conduction via so called *upper turn around point*. The
11 dotted lines indicate the speculated upper turnaround and its connection to the His-Purkinje
12 system. Successful ablation can be achieved by targeting the earliest P2 or any P1 in the
13 circuit, preferably in the apical third of the septum to avoid the creation of left bundle
14 branch block (LBBB) or atrioventricular block.

15

16

