

Exploration of Potential Prognostic Biomarkers in Aflibercept Plus FOLFIRI in Japanese Patients With Metastatic Colorectal Cancer

著者 (英)	Tetsuya Hamaguchi, Tadamichi Denda, Toshihiro Kudo, Naotoshi Sugimoto, Takashi Ura, Kentaro Yamazaki, Hirofumi Fujii, Takeshi Kajiwara, Takako Eguchi Nakajima, Shin Takahashi, Satoshi Otsu, Yoshito Komatsu, Fumio Nagashima, Toshikazu MORIWAKI, Taito Esaki, Takeo Sato, Michio Itabashi, Eiji Oki, Toru Sasaki, Marielle Chiron, Takayuki Yoshino
journal or publication title	Cancer Science
volume	110
number	11
page range	3565-3572
year	2019-12
権利	(C) 2019 The Authors. Cancer Science published by John Wiley & Sons Australia, Ltd on behalf of Japanese Cancer Association. This is an open access article under the terms of the Creative Commons Attribution NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes
URL	http://hdl.handle.net/2241/00160172

doi: 10.1111/cas.14198



Exploration of potential prognostic biomarkers in aflibercept plus FOLFIRI in Japanese patients with metastatic colorectal cancer

Tetsuya Hamaguchi¹ | Tadamichi Denda² | Toshihiro Kudo³ | Naotoshi Sugimoto⁴ | Takashi Ura⁵ | Kentaro Yamazaki⁶ | Hirofumi Fujii⁷ | Takeshi Kajiwara⁸ | Takako Eguchi Nakajima⁹ | Shin Takahashi¹⁰ | Satoshi Otsu¹¹ | Yoshito Komatsu¹² | Fumio Nagashima¹³ | Toshikazu Moriwaki¹⁴ | Taito Esaki¹⁵ | Takeo Sato¹⁶ | Michio Itabashi¹⁷ | Eiji Oki¹⁸ | Toru Sasaki¹⁹ | Marielle Chiron²⁰ | Takayuki Yoshino²¹

¹Department of Gastrointestinal Medical Oncology, National Cancer Center Hospital, Tokyo, Japan

²Department of Gastroenterology, Chiba Cancer Center, Chiba, Japan

³Osaka University Graduate School of Medicine, Suita, Japan

⁴Department of Medical Oncology, Osaka Medical Center for Cancer and Cardiovascular Disease, Osaka, Japan

⁵Department of Clinical Oncology, Aichi Cancer Center Hospital, Nagoya, Japan

⁶Department of Gastrointestinal Oncology, Shizuoka Cancer Center, Shizuoka, Japan

⁷Department of Clinical Oncology, Jichi Medical University Hospital, Shimotsuke, Japan

⁸Department of Gastrointestinal Medical Oncology, Shikoku Cancer Center, Matsuyama, Japan

⁹Department of Medical Oncology, St. Marianna University School of Medicine Hospital, Kawasaki, Japan

¹⁰Department of Medical Oncology, Tohoku University Hospital, Sendai, Japan

¹¹Department of Medical Oncology, Oita University Hospital, Yufu, Japan

¹²Department of Cancer Center, Hokkaido University Hospital, Sapporo, Japan

¹³Department of Medical Oncology, Kyorin University Hospital, Mitaka, Japan

¹⁴Department of Gastroenterology, University of Tsukuba Hospital, Tsukuba, Japan

¹⁵Department of Gastrointestinal and Medical Oncology, National Kyushu Cancer Center, Fukuoka, Japan

¹⁶Department of Gastrointestinal Surgery, Kitasato University East Hospital, Sagami, Japan

¹⁷Department of Surgery 2, Tokyo Women's Medical University Hospital, Tokyo, Japan

¹⁸Department of Gastrointestinal Surgery, Kyushu University Hospital, Fukuoka, Japan

¹⁹Research and development, Sanofi K. K., Tokyo, Japan

²⁰Biomarkers & Imaging Genomics, Sanofi, Alfortville, France

²¹Department of Gastrointestinal Oncology, National Cancer Center Hospital East, Kashiwa, Japan

Abbreviations: 5-FU, 5-fluorouracil; ANG-2, angiotensin-2; CV, coefficient of variation; EN-RAGE, extracellular newly identified receptor for advanced glycation end-products binding protein; FDR, false discovery rate; FOLFIRI, 5-FU/levofolinate/irinotecan; HE4, human epididymis protein 4; IGF1, insulin-like growth factor-1; IL, interleukin; ITAC, interferon-inducible T-cell alpha chemoattractant; mCRC, metastatic colorectal cancer; mFOLFOX6, modified folinic acid, fluorouracil, and oxaliplatin; OS, overall survival; PFS, progression-free survival; PlGF, placental growth factor; SP-D, surfactant protein D; TIMP-1, tissue inhibitor of metalloproteinases 1; TN-C, tenascin-C; TNFR2, tumor necrosis factor receptor 2; VEGF, vascular endothelial growth factor; VEGFR, vascular endothelial growth factor receptor.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors. *Cancer Science* published by John Wiley & Sons Australia, Ltd on behalf of Japanese Cancer Association.

Correspondence

Takayuki Yoshino, Department of Gastroenterology and Gastrointestinal Oncology, National Cancer Center Hospital East, Kashiwa, Japan.
Email: tyoshino@east.ncc.go.jp

Present address

Toshihiro Kudo and Naotoshi Sugimoto, Department of Medical Oncology, Osaka International Cancer Institute, Osaka, Japan

Takashi Ura, Department of Clinical Oncology, National Hospital Organization Kyoto Medical Center, Kyoto, Japan

Michio Itabashi, Department of Surgery, Institute of Gastroenterology, Tokyo Women's Medical University, Tokyo, Japan

Funding information

Sanofi

Abstract

Aflibercept plus 5-fluorouracil/levofolinate/irinotecan (FOLFIRI) is a second-line treatment for metastatic colorectal cancer. This ancillary exploratory analysis of data in Japanese people was aimed at exploring the relationship between a set of potential prognostic biomarkers and efficacy endpoints following aflibercept plus FOLFIRI therapy. Sixty-two patients with metastatic colorectal cancer received aflibercept (4 mg/kg) plus FOLFIRI every 2 weeks. Seventy-eight potential protein biomarkers were chosen for analysis based on their roles in angiogenesis, tumor progression, and tumor-stroma interaction. Plasma levels of biomarkers at baseline and at pre-dose 3 (day 1 of treatment cycle 3) were measured in all patients by ELISA. Relationships between these levels and efficacy endpoints were assessed. Ten potential biomarkers had a $\pm 30\%$ change from baseline to pre-dose 3 (adjusted $P < .001$), with the greatest changes occurring in placental growth factor (median: +4716%) and vascular endothelial growth factor receptor 1 (+2171%). Baseline levels of eight potential biomarkers correlated with overall survival in a univariate Cox regression analysis: extracellular matrix metalloproteinase inducer, insulin-like growth factor-binding protein 1, interleukin-8, kallikrein 5, pulmonary surfactant-associated protein D, tissue inhibitor of metalloproteinases 1, tenascin-C, and tumor necrosis factor receptor 2. None correlated with progression-free survival or maximum tumor shrinkage. Pre-dose 3 levels did not correlate with any efficacy endpoints. Preliminary data show that these eight biomarkers could be associated with overall survival. ClinicalTrials.gov identifier: NCT01882868.

KEYWORDS

aflibercept, biomarker, colorectal cancer, FOLFIRI, metastasis

1 | INTRODUCTION

Colorectal cancer is one of the most common malignancies in the Japanese population.¹ The 5-year survival rate for mCRC is approximately 12%.²

Aflibercept is an anti-cancer drug that has been used in combination with FOLFIRI as a second-line treatment for mCRC. Aflibercept acts by binding to and sequestering VEGF-A and -B and PlGF, thereby preventing downstream events such as angiogenesis and metastasis. To date, the largest phase III trial on aflibercept plus FOLFIRI (EFC10262, or VELOUR)³ was an international randomized double-blind study conducted outside of Japan and consisting of 1226 patients with mCRC who had previously been treated with oxaliplatin with or without bevacizumab. The researchers found that aflibercept plus FOLFIRI significantly improved both overall survival (OS; 13.50 vs 12.06 months, $P = .0032$) and progression-free survival (PFS; 6.90 vs 4.67 months, $P = .0001$) compared to placebo plus FOLFIRI. Another multinational randomized double-blind phase III study (EFC11338) consisting of 332 patients in Asia yielded similar results: aflibercept plus FOLFIRI improved both OS (14.59 vs 11.93 months) and PFS (6.93 vs 5.59 months) compared to placebo plus FOLFIRI.⁴

Early studies have identified biomarkers that may predict response to treatment with aflibercept plus FOLFIRI. For example, analysis of plasma samples from the phase III VELOUR study led to identification of several biomarkers potentially predictive or prognostic of OS: VEGF-A, VEGF receptors 2 and 3, IL-8, macrophage migration inhibitory factor (MIF), and SP-D.⁵ Lambrechts et al⁶ found that high plasma levels of IL-8 at baseline together with increased levels of IL-8 during treatment were significantly associated with reduced PFS. However, there are currently no established biomarkers that predict treatment response to aflibercept plus FOLFIRI.

The objective of the present study was to explore the relationship between a set of potential prognostic biomarkers and efficacy endpoints following aflibercept plus FOLFIRI therapy.

2 | MATERIALS AND METHODS

2.1 | Patients

This study included Japanese patients with mCRC whose cancer had failed to respond to a prior oxaliplatin treatment regimen. Further inclusion and exclusion criteria have been described in detail elsewhere.⁷ All patients gave informed consent.

2.2 | Study design

The study design has been described in detail elsewhere.⁷ A brief description follows:

This study was a prospective, multicenter, open-label, single-arm study. The first patient was enrolled on July 30, 2013, and the last data were collected on August 28, 2015.

The study consisted of three time periods: a baseline period, a treatment period, and a post-treatment follow-up period. During the baseline period, tumors were imaged and baseline data were collected. During the treatment period, the following treatment regimen was given once every 2 weeks until disease progression, unacceptable toxicity, or patient withdrawal: aflibercept (4 mg/kg) by i.v. infusion; then levofolinate (200 mg/m²) plus irinotecan (180 mg/m²), simultaneously by i.v. infusion; then a bolus of 5-FU (400 mg/m²); and then 5-FU (2400 mg/m²) by continuous i.v. infusion. Tumors were imaged every 6 ± 1 weeks and at the end-of-treatment visit 30 ± 3 days after the last study treatment. In the post-treatment follow-up period, tumors were imaged every 6 weeks, and survival status was determined every 2 months.

Primary endpoint (objective response rate) and secondary endpoints (PFS, OS, and pharmacokinetics) have been described previously.⁷ The focus of the current article is an exploratory endpoint: plasma levels of potential prognostic biomarkers.

2.3 | Assessments

A total of 109 potential protein biomarkers, including cytokines and angiogenic factors (Tables S1 and S2), were chosen for analysis based on their roles in tumor progression. Plasma was collected from all 62 patients at baseline and before infusion of aflibercept in treatment cycles 1 and 3. Plasma levels of the potential biomarkers were assessed by Myriad RBM using their multiplexed immunoassays, which are based on Luminex bead technology: Assays were carried out on the bead surface and then read in a compact analyzer using multiple lasers or light-emitting diodes and high-speed digital-signal processors.

Relationships between biomarker levels (at baseline and in treatment) and efficacy variables (PFS, OS, maximum shrinkage in tumor size, and best overall response) were assessed. Biomarker levels were also analyzed in patient groups stratified by whether or not the patients had previously received bevacizumab.

2.4 | Statistical analysis

All analyses of plasma protein biomarker variables were defined in an ad hoc method and carried out on all patients. *P* values for biomarker data were adjusted by FDR methods. Welch's test was used to determine *P* values for biomarker levels in patients stratified by whether they had previously received bevacizumab.

Median values of biomarker data were used to examine correlations with PFS and OS. This methodology has previously been shown to be effective for use in exploratory investigations of potential relationships between biomarkers and clinical outcomes.⁸

2.5 | Ethical considerations

As mentioned elsewhere,⁷ the present study was conducted in accordance with the principles of the Declaration of Helsinki, and in compliance with all international and Japanese laws, regulations, and guidelines. All aspects of the study were approved by an independent ethics committee and institutional review board.

3 | RESULTS

Sixty-two patients from 19 clinical sites in Japan were enrolled in this study. Of the 109 biomarkers measured, 31 were excluded from the analysis due to lack of informative data, as detailed in Figure 1.

Upon visual inspection of histograms of the individual biomarkers, 47 of the 78 biomarkers were found to have a non-Gaussian distribution. In order to obtain distributions closer to Gaussian distributions, a logarithmic transformation was applied to measurements for these 47 biomarkers.

Progression-free survival and OS were stratified by above versus below median plasma concentration for each biomarker. Baseline levels of eight biomarkers were found to correlate with OS with an adjusted *P* < .05 (Figures 2 and 3). The most significant differences in OS were found for TIMP-1, IL-8, and EN-RAGE (all *P* < .001), with lower concentrations corresponding to longer OS for all three biomarkers. Spearman's rank correlation coefficient at baseline was 0.80 between TIMP-1 and IL-8; 0.67 between TIMP-1 and EN-RAGE; and 0.77 between IL-8 and EN-RAGE.

Median plasma concentrations used for stratification of OS were 22.50 ng/mL (CV, 295.213) for EN-RAGE, 26.33 ng/mL (87.557) for IGFBP-1, 6.45 pg/mL (131.301) for IL-8, 3.30 ng/mL (33.276) for kallikrein 5, 5.43 ng/mL (67.590) for SP-D, 430.25 ng/mL (57.457) for TN-C, 76.00 ng/mL (43.984) for TIMP-1, and 5.78 ng/mL (34.486) for TNFR2.




FIGURE 1 Flow chart of biomarkers measured, excluded from analysis, and analyzed. LLOQ, lower limit of quantification




FIGURE 2 Progression-free survival (PFS) (A) and overall survival (OS) (B) stratified by above/below median plasma concentrations at baseline of biomarkers with adjusted $P < .05$ for PFS or OS. These biomarkers were extracellular newly identified receptor for advanced glycation end-products binding protein (EN-RAGE), insulin-like growth factor-binding protein 1 (IGFBP-1), interleukin-8 (IL-8), kallikrein 5, pulmonary surfactant-associated protein D (SP-D), tenascin-C (TN-C), tissue inhibitor of metalloproteinases 1 (TIMP-1), and tumor necrosis factor receptor 2 (TNFR2). *** $P < .001$; * $P < .05$; NS, not statistically significant

Baseline levels of none of the biomarkers were found to correlate with PFS, maximum shrinkage in tumor size, or best overall response, using an FDR correction of P value as the indicator. On-treatment levels of none of the biomarkers were found to correlate with any of the efficacy variables.

Ten of the 78 potential biomarkers had a $\pm 30\%$ change in plasma concentration ($P < .001$) from baseline to pre-dose 3 (Table 1). The most significant change was observed in PIGF, which had a 4716% change. In patients stratified by having received prior bevacizumab, significant differences were observed in baseline measurements

of four biomarkers (Figure 4). The largest difference was observed in log-transformed VEGF: median was 6.47 pg/mL in patients who had received prior bevacizumab versus 4.22 pg/mL in patients who had not ($P = 1.4E-14$). Baseline levels of PIGF and decorin were also significantly higher in patients who had received prior bevacizumab versus those who had not, whereas the baseline level of ANG-2 was significantly lower.

For 30 of the 78 biomarkers, two lots of reagent were used. Of these 30 biomarkers, nine had a batch effect with $P < .001$ and/or a value below the lower limit of quantification in one lot: VEGF-C, HE4, carcinoembryonic antigen-related cell adhesion molecule 1, cadherin-13, endoglin, VEGFR-1, hepatocyte growth factor receptor, human epidermal growth factor receptor 2, and ITAC. To eliminate the confounding effect, a correction was applied for all of these biomarkers except for VEGFR-1; for VEGFR-1, a quality problem was suspected in lot 1 prior to treatment cycle 1, which may have led to the apparent lot effect. For HE4 and ITAC, one lot was discarded, and the two biomarkers were excluded due to lack of available data.









4 | DISCUSSION

The objective of the present study was to explore the relationship between a set of potential prognostic biomarkers and efficacy endpoints following second-line treatment of mCRC with aflibercept plus FOLFIRI. We found eight biomarkers that are potentially prognostic for response to second-line aflibercept plus FOLFIRI. It is important to note that because this was a single-arm study, it could not be determined whether these eight biomarkers are actually prognostic, only that they are potentially prognostic for treatment response.

Among the eight potentially prognostic biomarkers found were TIMP-1 and IL-8: a lower baseline concentration of each of these biomarkers correlated with a longer OS. Other researchers have also implicated IL-8 as a potentially prognostic biomarker. Lambrechts et al⁶ found that high plasma levels of IL-8 at baseline together with increased levels of IL-8 during first-line treatment with 4 mg/mL aflibercept plus mFOLFOX6 (85 mg/m² oxaliplatin and 350 mg/m² leucovorin given simultaneously by i.v. infusion over 2 hours, followed by 400 mg/m² 5-FU as a bolus and then 2400 mg/m² 5-FU by i.v. infusion over 46 hours), given every 2 weeks,⁹ were significantly associated with reduced PFS compared to mFOLFOX6 alone. A retrospective analysis of data from the VELOUR study also implicated IL-8 as a potential prognostic biomarker, as well as SP-D,⁵ similar to the current study.

Interleukin-8 has been found to be potentially prognostic for response not only to aflibercept but also to bevacizumab, which,

FIGURE 3 Kaplan-Meier curves of overall survival (OS) stratified by median plasma concentration at baseline for eight biomarkers: (A) advanced glycation end-products binding protein (EN-RAGE), (B) insulin-like growth factor-binding protein 1 (IGFBP-1), (C) interleukin-8 (IL-8), (D) kallikrein 5, (E) pulmonary surfactant-associated protein D (SP-D), (F) tenascin-C (TN-C), (G) tissue inhibitor of metalloproteinases 1 (TIMP-1), and (H) tumor necrosis factor receptor 2 (TNFR2)



HR, hazard ratio

TABLE 1 Potential prognostic biomarkers with a $\pm 30\%$ change in plasma concentration ($P < .001$) from baseline to pre-dose 3

Biomarker	Median plasma level (CV%) [min : max]			Adjusted P^a
	Baseline	Pre-dose 3	% change [min : max]	
PIGF (pg/mL)	26.50 (61.941) [8.0:72.5]	1320.00 (31.579) [72.0:1990.0]	4716.11 [65.5:23 900.0]	1.26E-16
VEGFR-1 (pg/mL)	59.00 (282.331) [59.0:3899.5]	1905.00 (45.372) [693.0:3880.0]	2171.19 [-73.8:5628.8]	1.78E-16
VEGF (pg/mL)	610.75 (64.106) [42.0:1470.0]	881.00 (50.922) [497.0:3090.0]	92.06 [-9.3:3900.0]	2.23E-16
MMP-3 (ng/mL)	9.35 (48.341) [4.4:27.0]	23.00 (68.269) [5.8:92.0]	118.33 [-72.6:613.1]	1.15E-15
VEGFR-3 (ng/mL)	22.00 (57.507) [5.1:71.5]	15.00 (57.449) [3.3:45.0]	-33.64 [-75.4:27.3]	2.15E-11
FRTN (ng/mL)	112.25 (112.920) [9.5:835.5]	186.00 (85.930) [23.0:950.0]	57 [-32.4:817.6]	1.15E-09
PARC (ng/mL)	76.25 (58.546) [26.5:266.5]	94.50 (54.861) [35.0:333.0]	30.19 [-33.3:136.5]	7.56E-08
IGFBP-1 (ng/mL)	26.33 (87.557) [0.3:126.0]	55.00 (72.197) [0.3:175.0]	50.5 [-96.9:2763.5]	1.99E-06
cFib (μ g/mL)	36.50 (70.420) [11.0:189.0]	25.00 (90.614) [7.5:148.0]	-36.09 [-77.6:185.4]	5.79E-05
EN-RAGE (ng/mL)	22.50 (295.213) [1.1:2765.0]	13.50 (265.312) [1.1:1170.0]	-46.94 [-95.0:300.0]	1.31E-04

Abbreviations: cFib, cellular fibronectin; CV, coefficient of variation; EN-RAGE, extracellular newly identified receptor for advanced glycation end-products binding protein; FRTN, ferritin; IGFBP-1, insulin-like growth factor-binding protein 1; PARC, pulmonary and activation regulated chemokine; PIGF, placental growth factor; VEGF, vascular endothelial growth factor; and VEGFR-1, VEGF receptor 1.

^aAdjusted using the false discovery rate method.




FIGURE 4 Baseline biomarker levels in patients stratified by having received prior bevacizumab. N = Patient did not receive prior bevacizumab; Y = Patient did receive prior bevacizumab. P values were determined using Welch's test. ANG-2, angiopoietin-2; PIGF, placental growth factor; VEGF, vascular endothelial growth factor

like aflibercept, is an angiogenesis inhibitor. In one study, relative change in IL-8 was associated with response to bevacizumab in patients with metastatic breast cancer.¹⁰ In other studies, IL-8 levels or changes in IL-8 level were not found to be prognostic for objective response rate, PFS, or OS in patients with colorectal cancer,^{11,12} although in one of those studies, tissue samples rather than serum were studied.¹²

In the current study, 10 biomarkers, including four involved in the VEGF pathway, had a $\pm 30\%$ change in plasma concentration from

baseline to pre-dose 3. The highest increase observed at pre-dose 3 was for VEGF and PIGF which was likely due to stabilization of these two proteins by aflibercept. Aflibercept plasma trough concentrations showed the existence of a certain concentration of bound aflibercept at pre-dose 3. Mean plasma trough concentrations of free and adjusted bound aflibercept at pre-dose 3 were 7.1 μ g/mL (CV, 53.00%) and 4.4 μ g/mL (CV, 15.93%), respectively. Despite these large changes, most of these biomarkers did not correlate with PFS or OS. The only ones that did were IGFBP-1 and EN-RAGE, which

had percent changes of 50.5 (adjusted $P = 1.99E-06$) and -46.94 (adjusted $P = 1.31E-04$), respectively.

Our results are consistent with those for bevacizumab in which bevacizumab was also found to increase circulating PIGF in colorectal cancer¹³ and in ovarian cancer.¹⁴ Its effects on VEGF have been variable.^{13,14}

Baseline levels of four biomarkers were significantly different in patients who had received prior bevacizumab versus those who had not: VEGF, PIGF, and decorin were significantly higher in patients who had received prior bevacizumab, whereas ANG-2 was significantly lower. Similarly, in an exploratory analysis of the VELOUR study, VEGF-A and PIGF were found to correlate with prior bevacizumab therapy¹⁵ in patients who received aflibercept and FOLFIRI and in those who received placebo and FOLFIRI.¹⁶ None of the four biomarkers were found to be potentially prognostic for treatment response.

Thirty-one of the original 109 biomarkers chosen based on their roles in tumor progression were removed from analysis because of insufficient data. It is possible that some of these 31 biomarkers are actually prognostic for response to treatment with aflibercept plus FOLFIRI. This remains to be studied.

One limitation of the present study is that there were a relatively small number of study participants. Another limitation is that it consisted of only a single arm, so comparisons in biomarker level between a study treatment arm and a control arm could not be made. In addition, although the methodology used herein to explore potential correlations between biomarkers and survival times, based on median values, has been validated previously,⁸ future studies may choose to incorporate alternative analysis methods using absolute rather than relative cut-off values. It must also be noted that this study analysis did not identify predictive biomarkers but merely identified potential biomarkers whose levels correlated with overall survival. To determine whether the eight candidate biomarkers found in this study actually predict prognosis, it would be important to do a double-arm prospective study with a larger number of patients. Furthermore, the possibility that the selected biomarkers are false positives cannot be ruled out. Nevertheless, these data would be valuable for biomarker selection in a meta-analysis of data from multiple studies.

In conclusion, eight biomarkers potentially prognostic for OS were identified: TIMP-1, IL-8, EN-RAGE, SP-D, TN-C, IGFBP-1, kallikrein 5, and TNFR2. Further studies are warranted.

ACKNOWLEDGMENTS

The authors would like to thank all patient participants. This study was sponsored by Sanofi writing assistance by Michelle L. Jones, PhD, ELS, and funded by Sanofi. Yoshinori Sunaga, Sanofi K. K., Tokyo, Japan analyzed the data and contributed to interpretation of the study.

DISCLOSURE

T.H. received honoraria from Taiho, Chugai, Takeda, Yakult, and Merck Serono; a consulting fee from NanoCarrier; and research

funding from MSD, Ono, Sanofi, Daiichi-Sankyo, Sumitomo Dainippon Pharma, Taiho, Teijin, and NanoCarrier. T.K. received research funding from Yakult. N.S. received honoraria from Chugai and Eli Lilly, and research funding from Chugai, Eli Lilly, Dainippon Sumitomo, Taiho, MSD, Ono, Daiichi-Sankyo, and Sanofi. T.U. received honoraria from Merck Serono, Taiho, Chugai, and Takeda, and research funding from Sanofi. K.Y. received honoraria from Chugai, Takeda, Yakult, Daiichi-Sankyo, Merck Serono, Bristol, Bayer, Eli Lilly, and Taiho, and research funding from BMS and Sanofi. H.F. received research funding from Sanofi. S.T. received honoraria from Asahikasei and research funding from Merck Serono, Ono, and Sanofi. Y.K. received honoraria from BMS and Sanofi, and research funding from Eli Lilly, BMS, and Sanofi. T.E. received honoraria from Eli Lilly, and research funding from Boehringer, Daiichi-Sankyo, Dainippon Sumitomo, Eli Lilly, Merck Serono, MSD, Novartis, Ono, and Taiho, and scholarship from Ono. S.T. received honoraria from Asahikasei, and research funding from Merck Serono, Ono and Sanofi. E.O. received honoraria from Bayer, Chugai, Eli Lilly, Merck Serono, Taiho, Takeda, and Yakult. T.Y. received honoraria from Chugai, Eli Lilly, Merck Serono, Sanofi, and Taiho, and research funding from Boehringer Ingelheim, Chugai, Dainippon Sumitomo, GlaxoSmithKline, MSD, Novartis, and Sanofi. The affiliated medical institutions of physician authors received study funding from Sanofi. T.S. and M.C. are employees of Sanofi. Funding for this research was provided by Sanofi.

ORCID

Tadamichi Denda  <https://orcid.org/0000-0003-2570-9692>

Toshikazu Moriwaki  <https://orcid.org/0000-0001-7702-122X>

Taito Esaki  <https://orcid.org/0000-0002-9169-8453>

REFERENCES

1. Toukei GT. Cancer Registration Statistics. National Cancer Center Japan. http://ganjoho.jp/reg_stat/statistics/stat/summary.html. Accessed January 18, 2019.
2. American Cancer Society, Inc. Survival rates for colorectal cancer, by stage. <https://www.cancer.org/cancer/colon-rectal-cancer/detection-diagnosis-staging/survival-rates.html>. Accessed November 17, 2018.
3. Van Cutsem E, Tabernero J, Lakomy R, et al. Addition of aflibercept to fluorouracil, leucovorin, and irinotecan improves survival in a phase III randomized trial in patients with metastatic colorectal cancer previously treated with an oxaliplatin-based regimen. *J Clin Oncol*. 2012;30(28):3499-3506.
4. Sanofi. A study of aflibercept versus placebo with FOLFIRI in patients with metastatic colorectal cancer previously treated with an oxaliplatin chemotherapy (AFLAME); 2015. <https://clinicaltrials.gov/ct2/show/results/NCT01661270?sect=X80156>. Accessed November 18, 2018.
5. Sims TN, Gao B, Phillips R, Lowy I. Potential predictive and prognostic biomarkers identified in baseline plasma samples from the VELOUR trial. *J Clin Oncol*. 2015;33(Suppl 3):638.
6. Lambrechts D, Thienpont B, Thuillier V, et al. Evaluation of efficacy and safety markers in a phase II study of metastatic colorectal cancer treated with aflibercept in the first-line setting. *Br J Cancer*. 2015;113(7):1027-1034.

7. Denda T, Sakai D, Hamaguchi T, et al. Phase II trial of aflibercept plus FOLFIRI as a second-line treatment for Japanese patients with metastatic colorectal cancer. *Cancer Sci.* 2019;110(3):1032-1043.
8. Keyvanjah K, DePrimo SE, Harmon CS, et al. Soluble KIT correlates with clinical outcome in patients with metastatic breast cancer treated with sunitinib. *J Transl Med.* 2012;10:165.
9. Folprecht G, Pericay C, Saunders MP, et al. Oxaliplatin and 5-FU/folinic acid (modified FOLFOX6) with or without aflibercept in first-line treatment of patients with metastatic colorectal cancer: the AFFIRM study. *Ann Oncol.* 2016;27(7):1273-1279.
10. Lam SW, Nota NM, Jager A, et al. Angiogenesis- and hypoxia-associated proteins as early indicators of the outcome in patients with metastatic breast cancer given first-line bevacizumab-based therapy. *Clin Cancer Res.* 2016;22(7):1611-1620.
11. Kargi A, Yalcin AD, Erin N, Savas B, Polat HH, Gorczynski RM. Erratum to "IL8 and serum soluble TRAIL levels following anti-VEGF monoclonal antibody treatment in patients with metastatic colon cancer" [Clin Lab. 2012;58:501-505]. *Clin Lab.* 2012;10:1103-1107.
12. Bruhn MA, Townsend AR, Khoon Lee C, et al. Proangiogenic tumor proteins as potential predictive or prognostic biomarkers for bevacizumab therapy in metastatic colorectal cancer. *Int J Cancer.* 2014;135(3):731-741.
13. Hayashi H, Arai T, Matsumoto K, et al. Biomarkers of reactive resistance and early disease progression during chemotherapy plus bevacizumab treatment for colorectal carcinoma. *Oncotarget.* 2014;5(9):2588-2595.
14. Horowitz NS, Penson RT, Duda DG, et al. Safety, efficacy, and biomarker exploration in a phase II study of bevacizumab, oxaliplatin, and gemcitabine in recurrent Müllerian carcinoma. *Clin Ovarian Cancer Other Gynecol Malig.* 2011;4(1):26-33.
15. Taberero J, Paccard C, Chiron M, Dochy E, Van Cutsem E. Placental growth factor and the angiogenic environment based on analysis of baseline plasma biomarkers from the VELOUR trial. *J Clin Oncol.* 2017;35(Suppl 4):592.
16. Van Cutsem E, Dochy E, Paccard C, Chiron M, Taberero J. Impact of prior bevacizumab treatment on VEGFA and PIGF levels and patient outcomes: A retrospective analysis of baseline plasma samples from the VELOUR trial. *Ann Oncol.* 2017;28(Suppl 3):141.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Hamaguchi T, Denda T, Kudo T, et al. Exploration of potential prognostic biomarkers in aflibercept plus FOLFIRI in Japanese patients with metastatic colorectal cancer. *Cancer Sci.* 2019;110:3565-3572. <https://doi.org/10.1111/cas.14198>