

References

- Aono H. 1985. Geologic structure of the Ashio and Yamizo mountains with special reference to its tectonic evolution. *Science Report of the Institute of Geoscience, University of Tsukuba, Section B (Geological Sciences)* **6**: 21-57.
- Beven KJ, Kirkby MJ. 1979. A physically based, variable contributing area model of basin hydrology. *Hydrological Sciences Bulletin* **24**: 43-69.
- Chow VT. 1964. Statistical and probability analysis of hydrologic data, Part I. frequency analysis. In Chow VT ed., *Handbook of Applied Hydrology*: MacGraw-Hill Book: New York; Section 8, 1-42.
- Crouch RJ, Novruzi T. 1989. Threshold conditions for rill initiation on a vertisol, Gunnedah, N.S.W., Australia. *Catena* **16**: 101-110.
- Dietrich WE, Dunne T. 1978. Sediment budget for a small catchment in mountainous terrain. *Zeitschrift für Geomorphologie, Supplementband* **29**: 191-206.
- Dietrich WE, Dunne T. 1993. The channel head. In Beven K, Kirkby MJ eds., *Channel Network Hydrology*: Wiley: Chichester; 175-219.
- Dietrich WE, Reneau SL, Wilson CJ. 1987. Overview: "Zero-order basins" and problems of drainage density, sediment transport and hillslope morphology. In Beschta RL, Blinn T, Grant GE, Ice GG, Swanson FJ eds., *Erosion and Sedimentation in the Pacific Rim, IAHS Publication* **165**: 27-37.
- Dietrich WE, Wilson CJ, Reneau SL. 1986. Hollows, colluvium, and landslides in soil-mantled landscapes. In Abrahams AD ed., *Hillslope Processes*: Allen & Unwin: London; 361-388.

- Dietrich WE, Wilson CJ, Montgomery DR, Mckean J. 1993. Analysis of erosion thresholds, channel networks and landscape morphology using a digital terrain model. *Journal of Geology* **101**: 259-278.
- Dietrich WE, Wilson CJ, Montgomery DR, Mckean J, Bauer R. 1992. Erosion thresholds and land surface morphology. *Geology* **20**: 675-679.
- Dunne T. 1978. Field studies of hillslope flow processes. In Kirkby MJ ed., *Hillslope Hydrology*: Wiley: Chichester; 227-293.
- Ebizuka M, Kondoh A. 1990. Effect of regolith on storm runoff processes. *Journal of Japanese Association of Hydrological Sciences* **20**: 53-64 (In Japanese with English Abstract).
- Freer J, McDonnell J, Beven KJ, Brammer D, Burns D, Hooper RP, Kendall C. 1997. Topographic controls on subsurface storm flow at the hillslope scale for two hydrologically distinct small catchments. *Hydrological Processes* **11**: 1347-1352.
- Freer J, McDonnell JJ, Beven KJ, Peters NE, Burns DA, Hooper RP, Aulenbach B, Kendall C. 2002. The role of bedrock topography on subsurface storm flow. *Water Resources Research* **38**: 1269.
- Genereux DP, Hemond HF, Mulholland PJ. 1993. Spatial and temporal variability in streamflow generation on the West Fork of Walker Branch Watershed. *Journal of Hydrology* **142**: 137-166.
- Gomi T, Sidle RC. 2003. Bed load transport in managed steep-gradient headwater streams of southeastern Alaska. *Water Resources Research* **39**: 1336.
- Hayashi S. 1985. Some relationships between the landslide-area ratio and hydrologic data. *Journal of Japanese Forestry Society* **67**: 209-217 (In Japanese with English Abstract).

- Hirose T. 1996. *Hydrological characteristics and sediment yield in four small catchments with different bedrock types in a humid temperate region*. PhD thesis, University of Tsukuba: 140p.
- Hirose T, Onda Y, Matsukura Y. 1994. Runoff and solute characteristics in four small catchments with different bedrocks in the Abukuma mountains, Japan. *Transactions, Japanese Geomorphological Union* **15A**: 31-48.
- Horton RE. 1945. Erosional development of streams and their drainage basins; Hydrophysical approach to quantitative morphology. *Bulletin of the Geological Society of America* **56**: 275-370.
- Hutchinson DG, Moore RD. 2000. Throughflow variability on a forested hillslope underlain by compacted glacial till. *Hydrological Processes* **14**: 1751-1766.
- Iida T. 1984. A hydrological method of estimation of the topographic effect on the saturated throughflow. *Transactions, Japanese Geomorphological Union* **5**: 1-12 (In Japanese with English Abstract).
- Iida T, Okunishi K. 1983. Development of hillslopes due to landslides. *Zeitschrift für Geomorphologie, Supplementband* **46**: 67-77.
- Istanbulluoglu E, Tarboton DG, Pack RT, Luce C. 2002. A probabilistic approach for channel initiation. *Water Resources Research* **38**: 1325.
- Iwai S, Ishiguro M. 1970. *Applied Hydrological Statistics*. Morikita Shuppan: Tokyo; 370p (In Japanese).
- JSCE (Japan Society of Civil Engineers). 1971. *Hydraulic Formulae*. Maruzen: Tokyo; 619p (In Japanese).
- Kamata Y. 1997. Reconstruction of chert-clastic sequence of the Ashio Terrane in the Kuzu area, central Japan. *Journal of Geological Society of Japan* **103**: 343-356 (In Japanese with English Abstract).

- Kimoto A, Uchida T, Mizuyama T, Changhua L. 2002. Influences of human activities on sediment discharge from devastated weathered granite hills of southern China: effects of 4-year elimination of human activities. *Catena* **48**: 217-233.
- Kirkby MJ. 1980. The stream head as a significant geomorphic threshold. In Coates DR, Vitek JD eds., *Thresholds in Geomorphology*: Allen & Unwin: London; 53-73.
- Kirkby MJ. 1986. A two-dimensional simulation model for slope and stream evolution. In Abrahams AD ed., *Hillslope Processes*: Allen & Unwin: London; 203-222.
- Kirkby MJ. 1987. Modelling some influences of soil erosion, landslides and valley gradient on drainage density and hollow development. In Ahnert F ed., *Geomorphological Models, Catena Supplement* **10**: 1-14.
- Kirkby MJ, Bull LJ, Poesen J, Nachtergaele J, Vandekerckhove L. 2003. Observed and modelled distributions of channel and gully heads-with examples from SE Spain and Belgium. *Catena* **50**: 415-434.
- Kirkby MJ, Chorley RJ. 1967. Throughflow, overland flow and erosion. *International Association of Scientific Hydrology Bulletin* **12**: 5-21.
- Komatsu Y, Onda Y. 1996. Spatial variation in specific discharge of base flow in small catchments, Oe-yama Region, Western Japan. *Journal of Japan Society of Hydrology and Water Resources* **9**: 489-497.
- Machida H, Arai F. 1992. *Atlas of tephra in and around Japan*. University of Tokyo Press: Tokyo; 276p (In Japanese).

- Merz W, Bryan RB. 1993. Critical conditions for rill initiation on sandy loam Brunisols: laboratory and field experiments in southern Ontario, Canada. *Geoderma* **57**: 357-385.
- Montgomery DR, Dietrich WE. 1988. Where do channels begin? *Nature* **336**: 232-234.
- Montgomery DR, Dietrich WE. 1989. Source areas, drainage density, and channel initiation. *Water Resources Research* **25**: 1907-1918.
- Montgomery DR, Dietrich WE. 1994a. A physically based model for the topographic control on shallow landsliding. *Water Resources Research* **30**: 1153-1171.
- Montgomery DR, Dietrich WE. 1994b. Landscape dissection and drainage area-slope thresholds. In Kirkby MJ ed., *Process Models and Theoretical Geomorphology*: Wiley: Chichester; 221-246.
- Montgomery DR, Sullivan K, Greenberg HM. 1998. Regional test of a model for shallow landsliding. *Hydrological Processes* **12**: 943-955.
- Mosley MP. 1979. Streamflow generation in a forested watershed, New Zealand. *Water Resources Research* **15**: 795-806.
- Nachtergaele J, Poesen J, Steegen A, Takken I, Beuselinck L, Vanderkerckhove L, Govers G. 2001. The value of a physically based model versus an empirical approach in the prediction of ephemeral gully erosion for loess-derived soils. *Geomorphology* **40**: 237-252.
- Ohsaka O, Tamura T, Kubota J, Tsukamoto Y. 1992. Process study of the soil stratification on weathered granite slopes. *Journal of the Japan Society of Erosion Control Engineering* **45**(3): 3-12 (In Japanese with English Abstract).

- Okunishi K. 1991. Hydrogeomorphological interactions -A review of approach and strategy-. *Transactions, Japanese Geomorphological Union* **12**: 99-116 (In Japanese with English Abstract).
- Okunishi K. 1994. Concept and methodology of hydrogeomorphology. *Transactions, Japanese Geomorphological Union* **15A**: 5-18.
- O'Loughlin EM. 1981. Saturation regions in catchments and their relations to soil and topographic properties. *Journal of Hydrology* **53**: 229-246.
- O'Loughlin EM. 1986. Prediction of surface saturation zones in natural catchments by topographic analysis. *Water Resources Research* **22**: 794-804.
- Onda Y. 1989. Influence of water storage capacity in regolith zone on runoff characteristics and slope failure on granitic hills in Aichi, Japan. *Transactions, Japanese Geomorphological Union* **10**: 13-26 (In Japanese with English Abstract).
- Onda Y. 1992. Influence of water storage capacity in the regolith zone on hydrological characteristics, slope processes, and slope form. *Zeitschrift für Geomorphologie, Neue Folge* **36**: 165-178.
- Onda Y. 1994a. Contrasting hydrological characteristics, slope processes and topography underlain by Paleozoic sedimentary rocks and granite. *Transactions, Japanese Geomorphological Union* **15A**: 49-65.
- Onda Y. 1994b. Seepage erosion and its implication to the formation of amphitheater valley heads: a case study at Obara, Japan. *Earth Surface Processes and Landforms* **19**: 627-640.
- Prosser IP, Abernethy B. 1996. Predicting the topographic limits to a gully network using a digital terrain model and process thresholds. *Water Resources Research* **32**: 2289-2298.

- Prosser IP, Dietrich WE. 1995. Field experiments on erosion by overland flow and their implication for a digital terrain model of channel initiation. *Water Resources Research* **31**: 2867-2876.
- Prosser IP, Dietrich WE, Stevenson J. 1995. Flow resistance and sediment transport by concentrated overland flow in a grassland valley. *Geomorphology* **13**: 71-86.
- Reneau SL, Dietrich WE, Dorn RI, Berger CR, Rubin M. 1986. Geomorphic and paleoclimatic implications of latest Pleistocene radiocarbon dates from colluvium-mantled hollows, California. *Geology* **14**: 655-658.
- Rodriguez-Iturbe I. 1993. The geomorphological unit hydrograph. In Beven K, Kirkby MJ eds., *Channel Network Hydrology*: Wiley: Chichester; 43-68.
- Schumm SA. 1956. Evolution of drainage systems and slopes in badlands at Perth Amboy, New Jersey. *Bulletin of the Geological Society of America* **67**: 597-646.
- Shimokawa E, Jitousono T, Takano S. 1989. Periodicity of shallow landslide on Shirasu (Ito Pyroclastic flow deposits) steep slopes and prediction of potential landslide sites. *Transactions, Japanese Geomorphological Union* **10**: 267-284 (In Japanese with English Abstract).
- Side RC. 1988. Bed load transport regime of a small forest stream. *Water Resources Research* **24**: 207-218.
- Slattery MC, Bryan RB. 1992. Hydraulic conditions for rill incision under simulated rainfall: a laboratory experiment. *Earth Surface Processes and Landforms* **17**: 127-146.
- Smith TR, Bretherton FP. 1972. Stability and the conservation of mass in drainage basin evolution. *Water Resources Research* **8**: 1506-1529.

- Sonoda M. 1993. Seasonal change of sediment discharge in a forested small basin in Japan. *Transactions, Japanese Geomorphological Union* **14**: 53-70 (In Japanese with English Abstract).
- Sudo S, Makimoto H, Hata M, Unozawa A, Takizawa F, Sakamoto T. 1991. *Geological Map of Japan 1:200,000, Utsunomiya*. Geological Survey of Japan (In Japanese).
- Suzuki H. 1966. Distribution of freeze-thaw days in Japan. *Geographical Review of Japan* **39**: 267-270 (In Japanese with English Abstract).
- Tanaka T. 1982. The role of subsurface water exfiltration in soil erosion processes. In Walling DE ed., *Recent Developments in the Explanation and Prediction of Erosion and Sediment Yield, IAHS Publication* **137**: 73-80.
- Tanaka Y, Matsukura Y, Kim T. 2002. Differences in runoff characteristics between granite and gneiss mountainous basins, Korea. *Transactions, Japanese Geomorphological Union* **23**: 349-355.
- Terajima T, Sakamoto T, Nakai Y, Kitamura K. 1997. Suspended sediment discharge in subsurface flow from the head hollow of a small forested watershed, Northern Japan. *Earth Surface Processes and Landforms* **22**: 987-1000.
- Terajima T, Sakamoto T, Shirai T. 2001. Bed load yield caused by subsurface water discharge in a forested 0-order basin in Hokkaido, Northern Japan. *Transactions, Japanese Geomorphological Union* **22**: 1-22 (In Japanese with English Abstract).

- Tsukamoto Y, Hiramatsu S, Sinohara S. 1973. Study on the growth of stream channel (III) -Relationship between 0 (zero) order channels and landslides-. *Journal of the Japan Society of Erosion Control Engineering* **26**(2): 14-20 (In Japanese).
- Tsukamoto Y, Ohta T, Noguchi H. 1982. Hydrological and geomorphological studies of debris slides on forested hillslopes in Japan. In Walling DE ed., *Recent Developments in the Explanation and Prediction of Erosion and Sediment Yield, IAHS Publication* **137**: 89-98.
- Tucker GE, Bras RL. 1998. Hillslope processes, drainage density, and landscape morphology. *Water Resources Research* **34**: 2751-2764.
- Uchida T, Ohte N, Kimoto A, Mizuyama T, Changhua L. 2000. Sediment yield on a devastated hill in southern China: effects of microbiotic crust on surface erosion process. *Geomorphology* **32**: 129-145.
- Vandaele K, Poesen J, Govers G, van Wesemael B. 1996. Geomorphic threshold conditions for ephemeral gully incision. *Geomorphology* **16**: 161-173.
- Vandekerckhove L, Poesen J, Oostwoud Wijdenes D, Nachtegaele J, Kosmas C, Roxo MJ, de Figueredo T. 2000. Thresholds for gully initiation and sedimentation in Mediterranean Europe. *Earth Surface Processes and Landforms* **25**: 1201-1220.
- Willgoose G, Bras RL, Rodriguez-Iturbe I. 1991. A coupled channel network growth and hillslope evolution model: 1. theory. *Water Resources Research* **27**: 1671-1684.
- Wu W, Sidle RC. 1995. A distributed slope stability model for steep forested basins. *Water Resources Research* **31**: 2097-2110.

- Yamakoshi T, Suwa H. 1998. Responses of surface runoff and sediment discharge to revegetation on the slope of pyroclastic-flow deposits, Mount Unzen, Japan. *Journal of the Japan Society of Erosion Control Engineering* **51**(3): 3-10 (In Japanese with English Abstract).
- Yoshinaga S, Saijo K. 1989. Slope development during Holocene reconstructed from alluvial cone and talus cone aggradation process. *Transactions, Japanese Geomorphological Union* **10**: 285-301 (In Japanese with English Abstract).