

## Acknowledgements

I wish to appreciate to Professor Dr. Yukinori Matsukura (University of Tsukuba) for valuable discussion, guidance, and encouragement. I am very grateful to Associate Professor Dr. Yuichi Onda (University of Tsukuba) for valuable suggestions for determining the field site as well as the offer of instruments. I thank all staffs of the geomorphology research group of University of Tsukuba, Professor Dr. Eiji Matsumoto, Professor Dr. Masamu Aniya, Professor Dr. Norikazu Matsuoka, and Associate Professor Dr. Hiroshi Ikeda, and my seniors Dr. Chiaki T. Oguchi (Japan International Research Center for Agricultural Sciences), Dr. Kuniyasu Mokudai (University of Tsukuba), and Dr. Yosuke Komatsu (National Research Institute for Earth Science and Disaster Prevention) for helpful suggestions.

I appreciate to my colleagues and friends Mr. Yuki Matsushi, Ms. Sachi Wakasa (University of Tsukuba), Mr. Tetsushi Itokazu (Nagoya University), and Mr. Kazuto Unokuchi (MontBell Japan, Co.) for their helps in field works. I also thank to Mr. Hisashi Aoki and all other graduate and undergraduate students in the geomorphology research group of University of Tsukuba, for their supports and encouragement. Thanks are also due to Awano Town office and Awano Town Forest Association for the offer of field sites and lodging facility. The present study has been financially supported by Foundation of River and Watershed Management in year 2000, and JSPS (Japan Society for the Promotion of Science) research fellowships for young scientists in year 2003.