

Rates and mechanism of weathering-rind development on andesite gravel in fluvial terrace deposits

著者	Oguchi Chiaki T
著者別名	小口, 千明
内容記述	Thesis (Ph. D. in Science)--University of Tsukuba, (A), no. 1971, 1998.10.31
発行年	1998
その他のタイトル	河成段丘堆積中の安山岩礫に生じた風化皮膜の発達速度と発達機構
URL	http://hdl.handle.net/2241/6868

References

- Aires-Barros, L., Graça, R. C., and Velez, A. (1975): Dry and wet laboratory tests and thermal fatigue of rocks. *Engineering Geology*, **9**, 249-265.
- Akagi, K. and Koike, K. (1978): Reexamination of formative process of Nasuno-ga-hara alluvial fan. *Proceedings of Geographical Review of Japan*, No.16, 256-257.
- Akutsu, J. (1962): The Quaternary geology of the Nasuno-ga-hara plain, Tochigi prefecture. *Science Report of Utsunomiya University*, No. 12, 73-92.
- Alexander, E. B. (1985): Rates of soil formation from bedrock or consolidated sediments. *Physical Geography*, **6**, 23-42.
- Amit, R., Gerson, R. and Yaalon, D. H. (1993): Stages and rate of the gravel shattering process by salts in desert Reg soil. *Geoderma*, **57**, 295-324.
- Anderson, L. W. and Anderson, D. S. (1981): Weathering rinds on quartzarenite clasts as a relative-age indicator and the glacial chronology of Mount Timpanogos, Wasatch range, Utah. *Arctic and Alpine Research*, **13**, 25-31.
- Aoki, T. (1994): Chronological Study of glacial advances based on the weathering-rind thickness of morainic gravel in the northern part of the Central Japan Alps. *Geographical Review of Japan*. **64A**, 601-618 (in Japanese with English abstract).
- Berner, R. A., Sjoeverg, M. A. Velvel, and Krom, M. D. (1980); Dissolution of pyroxenes and amphiboles during weathering, *Science*, **207**, 1205-1206.
- Birkeland, P. W. (1973): Use of relative age-dating methods in a stratigraphic study of rock glacier deposits, Mt. Sopris, Colorado. *Arctic Alpine Res.*, **5**, 401-416.
- Burke, R. M. and Birkeland, P. W. (1979): Reevaluation of multiparameter relative dating techniques and their application to the glacial sequence along the eastern escarpment of the Sierra Nevada, California. *Quaternary*

- Res.*, **11**, 21-51.
- Caine, N. (1983): *The Mountains of Northern Tasmania*. A. A. Balkema, Rotterdam, 200p.
- Cernohouz, J. and Šolc, (1966): Use of sandstone wanes and weathered basaltic crust in absolute chronology. *Nature*, **212**, 806-807.
- Chesworth, W., Dejou, J. and Larroque, P. (1981): The weathering of basalt and relative mobility of the major elements at Belbex France. *Geochemi. Chosmohimi. Acta.* **45**, 1235-1243.
- Chigira, M. (1990) A mechanism of chemical weathering of mudstone in a mountain area. *Engineering Geology*, **29**, 119-138.
- Chigira, M. (1991) Chemical weathering mechanisms and their effects on engineering properties of soft sandstone and conglomerate cemented by zeolite in a mountainous area. *Engineering Geology*, **30**, 195-219.
- Chinn, T. J. H. (1981): Use of rock weathering-rind thickness for Holocene absolute age-dating in New Zealand. *Arctic and Alpine Research*, **13**, 33-45.
- Colman, S. M. (1981a): Chemical weathering of basalts and andesites: Evidence from weathering rinds. *U. S. Geological Survey Professional Paper*. **1246**.
- Colman, S. M. (1981b): Rock-weathering rates as functions of time. *Quaternary Research*, **15**. 250-264.
- Colman, S. M. (1982): Clay mineralogy of weathering rinds and possible implications concerning the sources of clay minerals in soils. *Geology*, **10**. 370-375.
- Colman, S. M. and Pierce, K. L. (1981): Weathering rinds on andesitic and basaltic stones as a Quaternary age indicator, western United States. *U. S. Geological Survey Professional Paper*, **1210**.
- Cooke, R. U. and Smalley, I. J. (1968): Salt weathering in deserts. *Nature*, **220**, 1227-1227.
- Crabtree, R. W. and Trudgill, S. T. (1985): Chemical denudation on a

- magnesian limestone hillslope, field evidence and implications for modeling. *Earth Surface Processes and Landforms*, **10**, 331-341.
- Craig, D. C. and Loughnan, F. C. (1969): Chemical and mineralogical transformation accompanying the weathering of basic volcanic rocks from New South Wales. *Australian Jour. Soil Res.*, **2**, 218-234.
- Crook, R. Jr. and Gillespie, A. R. (1986): Weathering rates in granitic boulders measured by P-wave speeds. In Colman, S. M. and Dethier, D. P. eds., *Rates of chemical weathering of rocks and minerals*. Academic Press, Orlando, 603p.
- Deere, D. U. and Patton, F. D. (1971): Slope stability in residual soils. *Proc. 4th Pan-American Conf. On Soil Mech. And Found. Engng.*, Puerto Rico, 87-170.
- Drever, J. I. (1997): *The Geochemistry of Natural Waters: Surface and Groundwater Environments-3rd ed.* Prentice-Hall, Inc., New Jersey. 436p.
- Eggleton, R. A., Foudoulis, C. and Varkevisser, D. (1987): Weathering of basalt: Changes in chemistry and mineralogy. *Clay and Clay Miner.*, **35**, 161-169.
- Garrels, R. M. and Mckenzie, F. T. (1967): Origin of the chemical compositions of some springs and lakes. In Stumm, W. ed. *Equilibrium Concepts in Natural Water Systems*. Advance in chemical series, **67**, American Chemical Society. 222-242.
- Gary, M., McAfee, Jr. R., and Wolf, D. eds. (1972): *Glossary of Geology*. American Geological Institute, 805p, Washington D. C.
- Goudie, A., Atkinson, B. W., Gregory, K. J., Simmons, I. G., Stoddart, D. R., and Sugden, D. eds. (1985): *The Encyclopedic Dictionary of Physical Geography*. Blackwell, 611p, Oxford.
- Grandstaff, D. E. (1986): The dissolution rate of forsteritic olivine from Hawaiian beach sand. In Colman, S. M. and Dethier, D. P. eds., *Rates of Chemical Weathering of Rocks and Minerals*. Academic Press, Orlando, 603p

- Friedman, I. and Long, W. (1976): Hydration rate of obsidian. *Science*, **191**, 347-352.
- Friedman, I. and Smith, R. L. (1960): A new dating method using obsidian: part I, the development of the method. *Amer. Artiq.*, **25**, 476-493.
- Haantjens, H. A. and Bleeker, P. (1970): Tropical weathering in the Territory of Papua and New Guinea. *Aust. Jour. Soil Res.*, **8**, 157-177.
- Harrassowitz, H. (1926): Laterit. Material und Versuch erdgeschichtlicher Auswertung. *Forschr. Geolog. Und Paleont*, **4**, 253-566.
- Hay, R. L. (1960): Rates of clay formation and mineral alteration in a 4000-year-old volcanic ash soil on St. Vincent, B. W. I. *Amer. Jour. Sciences*, **258**, 354-368.
- High, C. and Hanna, F. K. (1970): A method for the direct measurement of erosion on rock surfaces: *British Geomorphological Research Group, Technical Bulletin*, **5**, 25p.
- Hirose, T. Hatta, T. and Matsukura, Y. (1995): Dissolution rates of several plutonic rocks and limestone in closed- and batch-system experiments. *Trans. Japanese. Geomorph. Union*, **16**, 43-51 (in Japanese with English abstract). (1995):
- Hirose, T., Onda, Y. and Matsukura, Y. (1994): Runoff and solute characteristics in four small catchments with different bedrocks in the Abukuma Mountains, Japan. *Trans. Japanese. Geomorph. Union*, **15A**, 31-48.
- Hunt R. W. G. (1980): Colour term, symbols, and their usage. In Grum, F. and Bartleson, C. J. eds. *Optical Radiation Measurement*. 11-31, Academic Press, New York.
- Hunt G. R. and Ashley, R. P. (1979): Spectra of altered rocks in visible and near infrared. *Economic Geology*. **74**, 1613-1629.
- Hutchison, C. S. (1974): *Laboratory Handbook of Petrographic Techniques*. John Wiley & Sons, New York, 527p.
- Ichikuni, M. (1972): *Inorganic Geochemistry*, Baihuukan, Tokyo, 148p (in

- Japanese).
- Katsui, Y. and Kondo, Y. (1965): Dating of stone implements by using hydration layer of obsidian. *Japanese Journal of Geology and Geography*, **36**, 45-60.
- Kimiya, K. (1975a): Tensile strength as a physical scale of weathering in granitic rocks. *Journal of Geology of Japan*, **81**, 319-364 (in Japanese with English abstract).
- Kimiya, K. (1975b): Rates of weathering of gravel of granitic rocks in Mikawa and Tomikusa areas. *Journal of Geology of Japan*, **81**, 683-696 (in Japanese with English abstract).
- Knuepfer, P. L. K. (1988): Estimating ages of late Quaternary stream terraces from analysis of weathering rinds and soils. *Geological Society of America Bulletin*, **100**, 1224-1236.
- Koike, K. (1961): Geomorphological developments of the drainage basin of the Naka River. *Geographical Review of Japan*, **34A**, 498-513 (in Japanese with English abstract).
- Koike, K., Iwasaki, T., Danbara, T. and Momose, M. (1985): Fission-Track Ages and their geological meanings of Quaternary volcanic ashes (Shimotsuke-loam) covering the Kitsuregawa Hills, Tochigi Prefecture, Central Japan. *Komazawa Journal of Geography*, No. 21, 39-67 (in Japanese).
- Koizumi, T. and Aoyagi, S. (1993): Debris supply periods estimated from weathering-rind thickness in rubble on the west-facing slope of Mt. Yakushidake, the Northern Japanese Alps. *Geographical Review of Japan*, **66A**, 269-286 (in Japanese with English abstract).
- Koizumi, T. and Seki, H. (1992): Use of rock weathering-rind thickness to redate fossil periglacial slope in Mt. Kisokomagatake, the Central Japanese Alps. *Quaternary Journal of Geography*. **44**, 245-251 (in Japanese with English abstract).
- Kuchitsu, N. (1990): Weathering of stone implements excavated from

- Sunadadai Site, Kanagawa Prefecture — Taphonomy for archeological properties —. *Quaternary Research (Japan)*, **30**, 43-47 (in Japanese).
- Lasaga, A. C. (1984): Chemical kinetics of water-rock interaction. *Journal of Geophysical Research*, **89**, 4009-4025.
- Leneuf, N. and Aubert, G. (1960): Essai d'évaluation de la vitesse de ferrallitisation. *Proc. 7th Int. Conf. Soil Sci.*, 225-228.
- Lumb, P. (1962): The properties of decomposed granite. *Geotechnique*, **12**, 226-243.
- McGreevy, J. P. and Smith, B. J. (1982): Salt weathering in hot deserts: Observations on the design of simulation experiments. *Geografiska Annaler*, **64A**, 161-170
- McGreevy, J. P. and Smith, B. J. (1984): The possible role of clay minerals in salt weathering, *Catena*, **11**, 169-175.
- Machida, H. and Arai, F. (1979): Daisen Kurayoshi pumice: Stratigraphy, chronology, distribution and implication to late Pleistocene events in central Japan. *Journal of Geography*, **88**, 313-330 (in Japanese with English abstract).
- Machida, H. and Arai, F. (1992): *Atlas of Tephra in and around Japan*. Tokyo Univ. Press, Tokyo, 276p (in Japanese).
- Maruyama, A. (1981): Weathering rinds. In Machida *et al.*, eds. *The Encyclopedia of Geomorphology*. Ninomiya-shoten, Tokyo, 767p (in Japanese).
- Matsukura, Y. (1994): A review of the studies on rock control in weathering processes. *Trans. Japanese. Geomorph. Union*, **15**, 203-222 (in Japanese with English abstract).
- Matsukura, Y. and Yatsu, E. (1982): Wet-dry slaking of Tertiary shale and tuff. *Trans. Japanese. Geomorph. Union*, **3**, 25-39 (in Japanese with English abstract).
- Matsukura, Y., Kimata, M. and Yokoyama, S. (1994a): Formation of weathering rinds on andesite blocks under the influence of volcanic gasses

- around the active crater of Aso Volcano, Japan. In Robinson, D. A. and Williams, R. B. G. eds. *Rock weathering and landform evolution*, 89-98, John Wiley & Sons, Chichester.
- Matsukura, Y. Oguchi, C. T. and Hatta, T. (1994b): Preliminary study on Vickers microhardness of weathering rinds. *Annual Report of the Institute of Geoscience, the University of Tsukuba*. No. 20, 15-17.
- Menard, H. W. (1974): *Geology, resources and society*. W. H. Freeman, San Francisco, 621p.
- Matsukura, Y. Maekado, A. Hatta, T. and Yatsu, E. (1983): Vertical changes in mineralogical, physical chemical and mechanical properties due to deep weathering of Inada granitic rocks, *Trans. Japanese. Geomorph. Union*, **4**, 65-80 (in Japanese with English abstract).
- Menard, H. W. (1974): *Geology, Resources and Society*: W. H. Freeman, San Francisco, 621p.
- Nahon, D. and Lappartient, J. R. (1977): Time factor and geochemistry in iron crust genesis. *Catena*, **4**, 249-254.
- Nakada, S. (1987): X-ray fluorescence determination of trace elements in silicate rocks. *Sci. Rep. Geol. Kyusyu Univ.*, **15**, 37-44 (in Japanese with English abstract).
- Nakada, S. Yanagi, T. Maeda, S. Fang, D and Yamaguchi, M. (1985): X-ray fluorescence analysis of major elements in silicate rocks. *Sci. Rep. Geol. Kyusyu Univ.*, **14**, 103-115 (in Japanese with English abstract).
- Nakhla, F. M. (1956): *Economic Geology*. **51**, 811-827.
- Nakashima, S., Miyagi, I., Nakata, E., Sasaki, H., Nittono, S., Hirano, T., Sato, T. and Hayashi, H. (1992): Colour measurement of some natural and synthetic minerals. *Report of Research Institute of Natural Resources, Mining College, Akita University*. **57**, 57-76.
- Nesbitt, H. W. and Young, G. M. (1982): Early proterozoic climates and plate motions inferred from major element chemistry of lutites. *Nature*, **299**, 715-717.

- Nishiyama, K., Yokota, S. and Iwamatsu, A. (submitted): Reddish weathering of gravels within the Quaternary fan deposits and mutual relations among colour index and their physical properties. *Engineering Geol. (Japan)*. (in Japanese with English abstract).
- Oguchi, C. T. and Matsukura, Y. (1996): The changes in Micro structure of porous rhyolite due to weathering and its influence on the reduction of rock strength. *Trans. Japanese. Geomorph. Union*, **17**, 1-15 (in Japanese with English abstract).
- Oguchi, C. T., Hatta, T. and Matsukura, Y. (1994): Changes in rock properties of porous rhyolite through 40,000 years in Kozu-shima Island, Japan. *Geographical Review of Japan*, **67A**, 775-793 (in Japanese with English abstract).
- Oguchi, C. T., Isobe, H., Komuro, K. and Matsukura, Y. (1995): Colour measurements using a visible microspectrometer of weathering rinds on andesite blocks. *Annual Report of the Institute of Geoscience, the University of Tsukuba*. No. 21, 9-13.
- Porter, S. C. (1975): Weathering rinds as a relative-age criterion: Application to subdivision of glacial deposits in the Cascade Range. *Geology*, **3**, 101-104.
- Reiche, P. (1943): Graphic representation of chemical weathering. *Jour. Sed. Petrology*, **13**, 58-68.
- Ruxton, B. F. (1968): Measures of the degree of chemical weathering of rocks. *Jour. Geol.*, **76**, 518-527.
- Saito, T., Abe, M. and Kunori, S. (1974): Study on weathering of volcanic rocks (I). *Butsuri-Tanko (Geophysical exploration)*, **27**, 3-15.
- Sasaki, M., Ajisaka, T. and Okamoto, A. (1958): Hydrogeology of the Nasu plain, Tochigi prefecture. *Journal of Geography*. **67**, 59-73 (in Japanese with English abstract).
- Selby, M. J. (1993): *Hillslope Materials and Processes*, 2nd ed. Oxford Univ. Press, New York, 451p.

- Shiraiwa, T. and Watanabe, T. (1991): Late Quaternary glacial fluctuations in the Langtang Valley, Nepal Himalaya, reconstructed by relative dating methods. *Arctic and Alpine Research*, **23**, 404-416.
- Singer, (1984): Clay formation in saprolites of igneous rocks under semiarid to arid conditions, Negev, Souther Israel. *Soil Sci.*, **137**, 332-240.
- Strahler, A. H. and Strahler, A. N. (1978): *Modern Physical Geography*. John Wiley & Sons, New York, 638p.
- Sugisaki, R., Shimomura, T. and Kazuo, A. (1977): An automatic X-ray fluorescence method for the analysis of silicate rocks. *Jour. Geol. Soc. Japan*, **83**, 725-733 (in Japanese with English abstract).
- Sugisaki, R. Kinoshita, Shimomura, T. and Ando, K. (1981): An automatic X-ray fluorescence method for the trace element analysis in silicate rocks. *Jour. Geol. Soc. Japan*, **87**, 675-688 (in Japanese with English abstract).
- Suzuki, M.(1976): *Kako wo Saguru Kagaku (Research of the Past Time)*. Kodansha, 234p (in Japanese).
- Suzuki, Takasuke and Hachinohe, S. (1995): Weathering rates of bedrock forming marine terraces in Boso peninsula, Japan. *Trans. Japanese. Geomorph. Union*, **16**, 93-113.
- Suzuki, Takasuke and Matsukura, Y. (1992): Pore-size distribution of loess from the Loess Plateau, China. *Trans. Japan. Geomorph. Union*. **13**, 169-183.
- Suzuki, Takasuke, Hirano, M. Takahashi, K. and Yatsu, E. (1977): The interaction between the weathering processes of granites and the evolution of landform in the Rokko Mountains, Japan. — Part 1. Vertical changes in physical, mechanical, mineral and chemical properties of the weathered Rokko granite. *Bull. Fac. Sci. Eng., Chuo Univ.*, **20**, 343-389 (in Japanese with English abstract).
- Suzuki, Takehiko (1992): Stratigraphy of tephra layers from the latter half of middle Pleistocene to late Pleistocene in the Chubu-Kanto area, Central Japan. *Geographical Reports of Tokyo Metropolitan University*. No. 27,

29-53.

- Suzuki, Takehiko and Hayakawa, Y. (1990): The age and stratigraphic horizon of the Omachi Apm Tephra Beds, Middle Pleistocene Time-markers in Central Japan. *The Quaternary Research (Japan)*, **29**, 105-120 (in Japanese with English abstract).
- Takaya, Y., Hatta, T. and Matsukura, Y. (1996): The influence of physical properties of sedimentary and igneous rocks on dissolution in closed-system experiments. *Trans. Japan. Geomorph. Union*, **17**, 193-202 (in Japanese with English abstract).
- Tamari, Y., Tsuji, H. and Kusakabe, Y. (1988): The relationships between rock types and water. *Journal of Geochemical Research (Japan)*, **22**, 139-147.
- Tharp, T. M. (1987): Conditions for crack propagation by frost wedging. *Geological Society of America Bulletin*, **99**, 94-102..
- Trendall, A. F. (1962): The formation of apparent peneplains by a process of combined laterization and surface work. *Zeitschrift für Geomorphologie N. F., Suppl. Bd*, **6**, 183-197.
- Trudgill, S. T. (1975): Measurement of erosional weight-loss of rock tablets. *British Geomorphological Research Group, Technical Bulletin*, **17**, 13-19.
- Trudgill, S. T. (1976): The subaerial and subsoil erosion of limestone on Aldabra Atoll, Indian Ocean. *Zeitschrift für Geomorphologie N. F., Suppl. Bd.*, **26**, 201-210.
- Trudgill, S. T., High, C. J. and Hanna, F. K. (1981): Improvements to the micro-erosion meter. *British Geomorphological Research Group, Technical Bulletin*, **29**, 3-17.
- Wakatsuki, T. and Rasyidin, A. (1992): Rates of weathering and soil formation. *Geoderma*, **52**, 251-263.
- Waragai, T. (1993): Weathering process of andesite lava in Mt. Hachimen, Yabakei, Oita prefecture. *Bull. Fac. Science and literature, Nihon Univ.*, **28**, 15-24 (in Japanese with English abstract).

- Washburn, E. W. (1921): Note on the method of determining the distribution of pore size in a porous material. *Proc. National Academy Science*, **7**, 115-116.
- Watanabe, K. and Sagehashi, N. (1960): Geologic history of the Nasu Plain, Tochigi Prefecture (Hydrology of Nasuno-ga-hara No.3). *Journal of Geology of Japan*, **66**, 147-156 (in Japanese with English abstract).
- Waylen, M. J. (1979): Chemical weathering in a drainage basin underlain by old red sandstone. *Earth Surface Processes*, **4**, 167-178.
- Whitehouse, I. E., McSaveney, M. J., Knuepfer, P. L. K. and Chinn, T. J. H. (1986): Growth of weathering rinds on Torlesse sandstone, southern Alps, New Zealand. In Colman, S. M. and Dethier, D. P. eds., *Rates of Chemical Weathering of Rocks and Minerals*. Academic Press, Orlando, 603p.
- Yatsu, E. (1988): *The Nature of Weathering*. Sozosha, Tokyo, 624p.
- Yoshioka, R. (1975): Estimation of amounts of weathered products through chemical composition of waters in the Kamenose landslide area. *Bull. Disaster Prev. Res. Inst. Kyoto Univ.*, **25**, 1-15.
- Young, B. B. and Millman, A. P. (1964): *Trans. Inst. Min. Metal.*, **73**, 437-476.