

高強度の走行中における無気的エネルギーの動員が 酸素摂取量に及ぼす影響

著者	佐伯 徹郎
著者別名	Saeki Tetsuro
内容記述	筑波大学博士（体育科学）学位論文・平成11年7月23日授与（甲第2212号）
発行年	1999
URL	http://hdl.handle.net/2241/6539

文献 (References)

Aitken, J. C., Bennet, W. M. and Thompson, J. (1989) The effects of high intensity training upon respiratory gas exchanges during fixed term maximal incremental exercise in man. *Eur. J. Appl. Physiol.*, 58 : 717-721.

Aitken, J. C. and Thompson, J. (1989) The effects of dietary manipulation upon respiratory exchange ratio as a maximum oxygen uptake during fixed term maximal incremental exercise in man. *Eur. J. Appl. Physiol.*, 58 : 722-727.

阿久津邦男・杉靖三郎 (1964) Running選手の最大酸素摂取量と最大酸素負債量. *体育学研究*, 8 : 361.

青木積之介 (1976) 長距離走. 金原勇編著, *陸上競技のコーチング* (1). 大修館書店 : 東京, pp.443-513.

青木純一郎 (1989) 運動と呼吸循環. 石河利寛・杉浦正輝編著, *運動生理学*. 建帛社 : 東京, pp.145-182.

有吉正博 (1987) 長距離・マラソン・駅伝. 成美堂 : 東京, pp.124-138.

Astrand, P. O. and Saltin, B. (1961) Oxygen uptake during the first minutes of heavy muscular exercise. *J. Appl. Physiol.*, 16 : 971-976.

Bailey, S. P. and Pate, P. P. (1991) Feasibility of improving running economy. *Sports Med.*, 12 : 228-236.

Bahr, R. (1992) Excess post exercise oxygen consumption - magnitude, mechanisms and practical implications -. *Acta. Physiol.*

Scand., 144, Suppl., 605 : 3-70.

Billat, V., Renoux, J. C., Pinoteau, J., Petit, B. and Koralsztein, J. P. (1994) Reproducibility of running time to exhaustion at $\dot{V}O_{2max}$ in subelite runners. Med. Sci. Sports Exerc., 26 : 254-257.

Billat, V. and Koralsztein, J. P. (1996) Significance of the velocity at $\dot{V}O_{2max}$ and time to exhaustion at this velocity. Sports Med., 22 : 90-108.

Bishop, P. and Martino, M (1993) Blood lactate measurement in recovery as an adjunct to training. -Practical considerations- Sports Med., 16 : 5-13.

Brandon, L. J. and Boileau, R. A. (1992) Influence of metabolic mechanical and physique variables on middle distance running. J. Sports Med., 32 : 1-9.

Brandon, L. J. (1995) Physiological factors associated with middle distance running performance. Sports Med., 19 : 268-277.

Bulbulian, R., Wilcox, A.R. and Darabos, B.L. (1986) Anaerobic contribution to distance running performance of trained cross-country athletes. Med. Sci. Sports Exerc., 18 : 107-113.

Coetzer, P., Noakes, T. D., Sanders, B., Lambert, M. I., Bosch, A. N., Wiggins, T. and Dennis, S. C. (1993) Superior fatigue resistance of elite black south african distance runners. J. Appl. Physiol., 75 : 1822-1827.

Conley, D. L. and Krahenbuhl, G. S. (1980) Running economy and distance running performance of highly trained athletes. Med. Sci. Sports Exerc., 12 : 357-360.

Conley, D. L., Krahenbuhl, G. S., Burkett, L. M., and Millar, A. L. (1984) Following Steve Scott : Physiological changes accompanying training. *Phys. Sportsmed.*, 12 : 103-106.

Costill, D. L., Hoffman, W. M., Kehoe, F., Miller, S. J. and Meyers, W. C. (1968) Maximum anaerobic power among college football players. *J. Sports Med.*, 8 : 103-106.

Costill, D. L. and Fox, E. L. (1969) Energetics of marathon running. *Med. Sci. Sports*, 1 : 81-86.

Costill, D. L. and Winrow, E. (1970) A Comparison of two middle aged ultramarathon runners. *Res. Quart.*, 41 : 135-139.

Costill, D. L., Thomason, H. and Roberts, E. (1973) Fractional utilization of the aerobic capacity during distance running. *Med. Sci. Sports*, 5 : 248-252.

Coyle, E. F. (1995) Physiological factors in endurance performance. *Exerc. Sports Sci. Rev.*, 23 : 25-63.

Crielaard, J. M. and Pirnay, F. (1981) Anaerobic and aerobic power of top athletes. *Eur. J. Appl. Physiol.*, 47 : 295-300.

Cunningham, D. A. and Faulkner, J. A. (1969) The effect of training on aerobic and anaerobic metabolism during a short exhaustive run. *Med. Sci. Sports*, 1 : 65-69.

Cunningham, D. A., Van Waterschost, B. M., Paterson, D. H. Leforce, M. and Sangal, S. P. (1977) Reliability and reproducibility of maximal oxygen uptake measurement in children. *Med. Sci. Sports*, 9 : 104-108.

Daniels, J. (1974) Physiological characteristics of champion male athletes. Res. Quart., 45 : 342-348.

Davies, C. T. M. and Thompson, M. W. (1979) Aerobic performance of female marathon and male ultramarathon athletes. Eur. J. Appl. Physiol., 41 : 233-245.

Dawson, B., Fitzsimous, M., Green, S., Goodman, C., Carey, M. and Cole, K. (1998) Changes in performance, muscle metabolites, enzymes and fibre types after short sprint training. Eur. J. Appl. Physiol., 78 : 163-169.

Edington, D. W. and Edgerton, V. R. : 大平充宣訳 (1988) 運動生理学の基礎. ベースボール・マガジン社 : 東京, pp.173-202.

Eriksson, B. O. (1972) Physical training, oxygen supply and muscle metabolism in 11-13 years old boys. Act. Physiol. Scand. Suppl., 384 : 1-48.

Farrell, P. A., Wilmore, J. H., Coyle, E. F., Billing, J. E. and Costill, D. L. (1979) Plasma lactate accumulation and distance running performance. Med. Sci. Sports, 11 : 338-344.

Fox, E. L. (1979) Sports physiology. W.B. Saunders company : Philadelphia. pp.18-33.

深代千之 (1991) 競技スポーツに科学をどのように生かすか? ⑨長距離選手に対するタイプ別アドバイス. コーチング・クリニック10月号, ベースボール・マガジン社 : 東京, pp.62-65.

Gaesser, G. A. and Brooks, G. A. (1984) Metabolic bases of excess post-exercise oxygen consumption : a review. Med. Sci. Sports Exerc., 16 : 29-43.

Gaesser, G. A. and Poole, D. G. (1996) The slow component of oxygen uptake kinetics in humans. *Exerc. Sport Sci. Rev.*, 24 : 35-70.

1975L

Girandola, R. N. and Henry, F. M. (1974) Individual differences in O₂ deficit and O₂ debt. *Res. Quart.*, 45 : 239-246.

Gore, C. J. and Withers, R. T. (1990) The effect of exercise intensity and duration on the oxygen deficit and excess post-exercise oxygen consumption. *Eur. J. Appl. Physiol.*, 60 : 169-174.

Grant, S., Craig, I., Wilson, J. and Aitchison, T. (1997) The relationship between 3 km running performance and selected physiological variables. *J. Sports Sci.*, 15 : 403-410.

Green, S. and Dawson, B. (1993) Measurement of anaerobic capacities in humans. Definitions, limitations and unsolved problems. *Sports Med.*, 15 : 312-327.

Hagerman, F. C. and Staron, R. S. (1983) Seasonal variations among physiological variables in elite oarsmen. *Can. J. Appl. Sport Sci.*, 8 : 143-148.

八田秀雄 (1997) 乳酸. ブックハウスHD : 東京.

1976

Hewson, D. J. and Hopkins, W. G. (1996) Specificity of training and its relation to the performance of distance runners. *Int. J. Sports Med.*, 17 : 199-204.

Hill, D. W. and Rowell, A. L. (1996) Running velocity at $\dot{V}O_{2max}$. *Med. Sci. Sports Exerc.*, 28 : 114-119.

Holmgren, A. and Astrand, P.-O. (1966) DI and the dimensions and

functional capacities of O₂ transport system in humans. *J. Appl. Physiol.*, 21 : 1463-1470.

Honig, C. R., Connett, R. J. and Gayeski, T. E. J. (1992) O₂ transport and its interaction with metabolism ; a systems view of aerobic capacity. *Med. Sci. Sports Exerc.*, 24 : 47-53.

Houmard, J. A., Costill, D. L., Michell, J. B., Park, S. H. and Chenier, T. C. (1991) The role of anaerobic ability in middle distance running performance. *Eur. J. Appl. Physiol.*, 62 : 40-43.

石河利寛 (1962) スポーツとからだ. 岩波新書 : 東京, p108.

岩垣丞恒 (1989) 運動と代謝・栄養. 石河利寛・杉浦正輝編著, 運動生理学. 建帛社 : 東京, pp.130-132.

Johnson, R. L. Jr., Spicer, W. S., Bishop, J. M. and Foster, R. E. (1960) Pulmonary capillary blood volume, flow and diffusing capacity during exercise. *J. Appl. Physiol.*, 15 : 893-902.

Joyner, M. J. (1991) Modeling : optimal marathon performance on the basis of physiological factors. *J. Appl. Physiol.*, 70 : 683-687

Katch, V. L. and Henry, F. M. (1972) Prediction of running performance from maximal oxygen debt and intake. *Med. Sci. Sports*, 4 : 187-191.

Katch, V. L. and Weltman, A. (1979) Interrelationship between anaerobic power output, anaerobic capacity and aerobic power. *Ergonomics*, 22 : 325-332.

Katch, V. L., Sady, S. S. and Freedson, P. (1982) Biological variability in maximum aerobic power. *Med. Sci. Sports Exerc.*, 14

: 21-25.

勝田茂・宮田浩文・麻場一徳・原田健・永井純（1986）中長距離選手におけるランニング効率とパフォーマンスとの関係について．筑波大学体育科学系紀要，8：45-52.

Keul, J. (1973) The relationship between circulation and metabolism during exercise. *Med. Sci. Sports*, 5 : 209-219.

Kindermann, W. and Keul, J. (1977) Lactate acidosis with different forms activities. *Can. J. Appl. Sport Sci.*, 2 : 177-182.

金原勇・高松薫・阿江通良・伊藤静夫（1973）トレーニング目標とする全身持久性のとらえ方に関する実験的研究．東京教育大学体育学部スポーツ研究所報，11：15-37.

Komi, P. V., Rusko, H., Vos, J. and Vihko, V. (1977) Anaerobic performance capacity in athletes. *Acta. Physiol. Scand.*, 100 : 107-114.

黒田善雄・伊藤静夫・塚越克己・雨宮輝也・鈴木洋児（1973）日本人一流競技選手の最大酸素摂取量並びに最大酸素負債量－第2報－．昭和48年度日本体育協会スポーツ科学研究報告，IX：1-27.

黒川隆志・富樫泰一・野村武男・池上晴夫（1985）最大酸素摂取量，最大酸素負債量および酸素需要量と水泳記録との関係．*体育学研究*，29 : 295-305.

Lacour, J. R., Padilla-Magunacelaya, S., Barthelemy, J. C. and Dormois, D. (1990a) The energetics of middle-distance running. *Eur. J. Appl. Physiol.*, 60 : 38-43.

Lacour, J. R., Bouvat, E. and Barthelemy, J. C. (1990b) Post-

competition blood lactate concentrations as indicators of anaerobic energy expenditure during 400-m and 800-m races. *Eur. J. Appl. Physiol.*, 61 : 172-176.

LaFontaine, T. P., Londeree, B. R. and Spath, W. K. (1981) The maximal steady state versus selected running events. *Med. Sci. Sports Exerc.*, 13 : 190-192.

Lindsay, F. H., Hawley, J. A., Myburgh, K. H., Schomer, H. H., Noakes, T. D. and Dennis, S. C. (1996) Improved athletic performance in highly trained cyclists after interval training. *Med. Sci. Sports Exerc.*, 28 : 1427-1434.

Mackova, E. V., Bass, A., Sprauarova, S., Teisinger, J., Vondra, K. and Bojanovsky, I. (1982) Enzyme activity patterns of energy metabolism in skiers of different performance levels (M. Quadriceps Femoris) . *Eur. J. Appl. Physiol.*, 48 : 315-322.

Maffulli, N., Capasso, G. and Lancia, A (1991) Anaerobic threshold and performance in middle and long distance running. *J. Sports Med.*, 31 : 332-338.

Margaria, R., Aghemo, P. and Rovelli, E. (1966) Measurement of muscular power (anaerobic) in man. *J. Appl. Physiol.*, 21 : 1662-1664.

Maughan, R. J. (1992) Aerobic function. *Sports Science Review*, 1 : 28-42.

Mckenzie, D. C., Parhouse, W. A. and Hearst, W. E. (1982) Anaerobic performance characteristics of elite canadian 800 meter runners. *Can. J. Appl. Sport Sci.*, 7 : 158-160.

Metzger, J. and Moss, R. (1990) Effects on tension and stiffness due to reduced pH in mammalian fast- and slow-twitch skinned skeletal muscle fibres. *J. Physiol.*, 428 : 737-750.

満園良一・丸山敦夫・四元清治 (1985) 中長距離ランナーの酸素摂取能および酸素負債量と走強度との関係. 久留米大学論叢, 34 : 145-152.

宮原清彰・豊岡示朗・吉川潔・足立哲司 (1995) 100mから1万mまでのレース後の血中乳酸濃度. 第7回ランニング学会大会号, p23.

Monod, H. and Scherrer, J. (1965) Work capacity of synergic muscular group. *Ergonomics*, 8 : 329-338.

Morgan, D. W., Baldini, F. D., Martin, P. E. and Kohrt, W. M. (1989) Ten kilometer performance and predicted velocity at $\dot{V}O_{2max}$ among well-trained male runners. *Med. Sci. Sports Exerc.*, 21 : 78-83.

Moritani, T., Nagata, A., deVries, H. A. and Muro, M. (1981) Critical power as a measure of physical work capacity and anaerobic threshold. *Ergonomics*, 24 : 339-350.

Mygind, E., Anderson, L.B. and Rasmussen, B. (1994) Blood lactate and respiratory variables in elite cross-country skiing at racing speeds. *Scand. J. Med. Sci. Sports*, 4 : 243-251.

永井 純 (1989) 中・長距離・障害. ベースボールマガジン社 : 東京, pp.9-10.

中村好男・山本義春 (1993) ATその変遷と新しい理解. ブックハウスHD : 東京.

Noakes, T. D. (1988) Implications of exercise testing for prediction

of athletic performance : a contemporary perspective. *Med. Sci. Sports Exerc.*, 20 : 319-330.

Noakes, T. D., Myburgh, K. H. and Schall, R. (1990) Peak treadmill running velocity during the $\dot{V}O_{2max}$ test predicts running performance. *J. Sports Sci.*, 8 : 35-45.

Noakes, T. D. (1991) *Lore of running*. Human Kinetics Publishers, Inc., Champaign, IL.

沼尻幸吉 (1959) 労働の強さと適正作業量. 労働科学研究所 : 東京, p.250.

尾縣貢・福島洋樹・大山圭悟・安井年文・鍋倉賢治・宮下憲・関岡康雄・永井純 (1998) 下肢の筋持久性と400m走中の疾走速度逓減との関係. *体育学研究*, 42 : 370-379.

Osnes, J. and Hermansen, L. (1972) Acid-base balance after maximal exercise of short duration. *J. Appl. Physiol.*, 32 : 59-63.

Pate, R. R., Sparling, P.B., Wilson, G. E., Cureton, K. J. and Miller, B. J. (1987) Cardiorespiratory and metabolic responses to submaximal and maximal exercise in elite women distance runners. *Int. J. Sport Med.*, 8 : 91-95.

Pate, R. R. and Branch, J. D. (1992) Training for endurance sport. *Med. Sci. Sports Exerc*, 24 : 340-343.

Pollock, M. L., Bohannon, R. L., Cooper, K. H., Ayres, J. J., Ward, A., White, S. R. and Linnerud, A. C. (1976) A comparative analysis of four protocols for maximal treadmill stress testing. *Am. Heart J.*, 92 : 39-46.

Powers, S. K., Dodd, S., Denson, R., Byrd, R. and Mcknight, T. (1983) Ventilatory threshold, running economy and distance running performance of trained athletes. *Res. Quart.*, 54 : 179-182.

Ramsbottom, R., Williams, C., Kerwin, D. G. and Nute, M. L. G. (1992) Physiological and metabolic responses of men and women to a 5-km treadmill time trial. *J. Sports Sci.*, 10 : 119-129.

Robinson, D. M., Robinson, S. M., Hume, P. A. and Hopkins, W. G. (1991) Training intensity of elite male distance runners. *Med. Sci. Sports Exerc.*, 23 : 1078-1082.

Roca, J., Hogan, M. C., Story, D. and Bebout, D. E. (1989) Evidence for tissue diffusion limitation of $\dot{V}O_{2max}$ in normal humans. *J. Appl. Physiol.*, 67 : 291-299.

Rusko, H., Havu, M. and Karvinen, E. (1978) Aerobic performance capacity in athletes. *Eur. J. Appl. Physiol.*, 38 : 151-159.

Sahlín, K. (1986) Muscle fatigue and lactic acid accumulation. *Acta Physiol. Scand.*, 128, Suppl., 556 : 83-91.

Saltin, B., Nazar, K., Costill, D. L., Stein, E., Jansson, E., Essen, B. and Gollnick, P. D. (1976) The nature of the training response ; peripheral and central adaptations to one-legged exercise. *Acta. Physiol. Scand.*, 96 : 289-305.

Saltin, B. and Strange, S. (1992) Maximal oxygen uptake : “old” and “new” arguments for a cardiovascular limitation. *Med. Sci. Sports Exerc.*, 24 : 30-37.

沢木啓祐・高岡郁夫(1993) マラソン. ベースボール・マガジン社 : 東

京, pp.34-36.

Schnabel, A. and Kindermann, W. (1983) Assessment of anaerobic capacity in runners. *Eur. J. Appl. Physiol.*, 52 : 42-46.

Scrimgeour, A. G., Noakes, T. D., Adams, B. and Myburgh, K. (1986) The influence of weekly training distance on fractional utilization of maximum aerobic capacity in marathon and ultramarathon runners. *Eur. J. Appl. Physiol.*, 55 : 202-209.

Scott, C. B., Roby, F. B., Lohman, T. G. and Bunt, J. C. (1991) The maximally accumulated oxygen deficit as indicator of anaerobic capacity. *Med. Sci. Sports Exerc.*, 23 : 618-624.

Scott, B. K. and Houmard, J. A. (1994) Peak running velocity is highly related to distance running performance. *Int. J. Sports Med.*, 15 : 504-507.

Serresse, O., Simoneau, J. A., Bouchard, C. and Boulay, M. R. (1991) Aerobic and anaerobic energy contribution during maximal work output in 90s determined with various ergocycle workloads. *Int. J. Sports Med.*, 12 : 543-547.

Sjodin, B. and Svedenhag, J. (1985) Applied physiology of marathon running. *Sports Med.*, 2 : 83-99.

Skinner, J. S. and Morgan, D. W. (1985) Aspects of anaerobic performance. In: *Limits of human performance*, ed by Clarke, D. H. and Eckert, H. M., Human Kinetics Publishers, Inc. Champaign, IL, pp.31-44.

Stamford, B. A. (1976) Step increment versus constant load tests for determination of maximal oxygen consumption. *Eur. J. Appl.*

Physiol., 35 : 89-93.

Stringer, W., Wasserman, K., Casaburi, R., Porszasz, J., Maehara, K. and French, W. (1994) Lactic acidosis as a facilitator of oxyhemoglobin dissociation during exercise. *J. Appl. Physiol.*, 76 : 1462-1467.

Sutton, J. R. (1992) Limitations to maximal oxygen uptake. *Sports Med.*, 13 : 127-133.

Svedenhag, J. and Sjodin, B. (1984) Maximal and submaximal oxygen uptake and blood lactate levels in elite male middle- and long-distance runners. *Int. J. Sports Med.*, 5 : 255-261.

Svedenhag, J. and Sjodin, B. (1985) Physiological characteristics of elite male runners in and off-season. *Can. J. Appl. Sport Sci.*, 10 : 127-133.

Tanaka, K., Matsuura, Y., Matsuzaka, A., Hirakoba, K., Kumagai, S., Sun-O, S. and Asano, K. (1984) A longitudinal assessment of anaerobic threshold and distance running performance. *Med. Sci. Sports Exerc.*, 16 : 278-282.

Tanaka, H. and Swensen, T. (1998) Impact of resistance training on endurance performance. *Sports Med.*, 25 : 191-200.

Taunton, J. E., Maron, H. and Wilkinson, J. G. (1981) Anaerobic performance in middle and long distance runners. *Can. J. Appl. Sport Sci.*, 6 : 109-113.

Taylor, H. L., Buskirk, H. and Henshel, A. (1955) Maximal oxygen intake as an objective measure of cardio-respiratory performance. *J. Appl. Physiol.*, 8 : 73-80.

Verstappen, F. T. J., Janssen, G. M. E. and Does, R. J. M. M. (1989) Effects of endurance training and competition on exercise tests in relatively untrained people. *Int. J. Sports Med.*, 10 : S126-S131.

Wasserman, K. and McIlroy, M. B. (1964) Detecting the threshold of anaerobic metabolism in cardiac patients during exercise. *Am. J. Cardiol.*, 14 : 844-852.

Wasserman, K., Beaver, W. and Whipp, B. (1986) Mechanisms and patterns blood lactate increase during exercise. *Med. Sci. Sports Exerc.*, 18 : 344-352.

Weltman, A., Moffatt, R. J. and Stamford, B. A. (1978) Supramaximal training in females : Effects on anaerobic power output, anaerobic capacity, and aerobic power. *J. Sports Med.*, 18 : 237-244.

Wilcox, A. R. and Bulbulian, R. (1982) The effect of a cross-country season on running economy (abstract). *Med. Sci. Sports Exerc.*, 14 : 142-143.

Wilmore, J. and Costill, D. L. (1994) Energy for movement. In : *Physiology of sport and exercise*, ed by Wilmore, J. and Costill, D. L., Human Kinetics Publishers, Inc. Champaign, IL, pp.104-107.

山地啓司・宮下充正 (1976) 安静及び作業時の肺拡散能力, 肺胞膜拡散容量及び肺毛細血管量. *体力科学*, 25 : 159-167.

山地啓司 (1990) 最大酸素摂取量から陸上中長距離走、マラソンレースの競技記録を占うことが可能か. *ランニング学研究*, 1 : 7-14.

山地啓司 (1992) 最大酸素摂取量の科学. 杏林書院 : 東京, pp.14-15, 208-225.

山地啓司（1998）最高有酸素的ランニング速度（ $\dot{V}O_2\max$ ）の意義と評価. 日本運動生理学雑誌, 5 : 89-99.

山崎省一・青木純一郎（1977）長距離走者の競技記録と無酸素的能力. 体力科学, 26 : 87-95.

山本正嘉（1989）筋のエネルギー出力. 田畑泉・山本正嘉共著, 身体運動のエナジェティクス. 高文堂：東京, pp.113-205.

矢野徳郎（1985）長・中・短距離走者の無氣的作業閾値, 酸素債および酸素摂取量の反応時間. 体力科学, 34 : 176-182.

吉田敬義（1989）運動開始時の酸素摂取量の働態とシステム同定. 体育の科学, 39 : 871-876.

Yoshida, T., Udo, M., Iwai, K., Chida, M., Ichioka, M., Nakadomo, F. and Yamaguchi, T. (1990) Significance of the contribution of aerobic and anaerobic components to several distance running performances in female athletes. *Eur. J. Appl. Physiol.*, 60 : 249-253.