

**Association of Sleep-Disordered Breathing and Ventricular Arrhythmias in Patients Without Heart Failure**

Yuki Koshino<sup>a,b</sup>, MD, Makoto Satoh, MD, PhD<sup>b</sup>, Yasuko Katayose, MD<sup>b</sup>, Kyo Yasuda, MD, PhD<sup>b</sup>, Takeshi Tanigawa, MD, PhD<sup>c</sup>, Noriyuki Takeyasu, MD, PhD<sup>a</sup>, Shigeyuki Watanabe, MD, PhD<sup>a</sup>, Iwao Yamaguchi, MD, PhD<sup>a</sup>, Kazutaka Aonuma MD, PhD<sup>a</sup>

<sup>a</sup>Department of Cardiology, Institute of Clinical Medicine, <sup>b</sup>Department of Sleep Medicine, Institute of Clinical Medicine, <sup>c</sup>Department of Public Health, Institute of Social and Environmental Medicine, Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki 305-8575, Japan

\*Address for correspondence: Yuki Koshino, Graduate School of Comprehensive Human Science, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki 305-8575, Japan. Tel and Fax: +81-29- 853-3143, E-mail: y-koshino@umin.ac.jp

This study was supported in part by a Research Grant for Cardiovascular Disease (17C-5) from the Ministry of Health, Labor and Welfare, Japan.

**Running head:** Sleep-disordered breathing in ventricular arrhythmia patients

**Abstract:** The prevalence and characteristics of sleep-disordered breathing (SDB) in patients with ventricular arrhythmias, such as premature ventricular complexes (PVCs) and ventricular tachycardia (VT), are unknown. Therefore, we evaluated the prevalence of SDB in patients with severe ventricular arrhythmias and normal left ventricular (LV) function. Thirty-five patients (63% men; mean age:  $57.4 \pm 13.8$  years) underwent a sleep study. All patients had VT or frequent PVCs ( $\geq 300$  PVCs/h) and had been referred to the Cardiology Department for medication, catheter ablation therapy, or implantation of a cardioverter/defibrillator. We excluded heart failure patients with an LV ejection fraction  $< 50\%$ ; in the remaining patients, mean LVEF was  $63.9 \pm 8.0\%$ . Twenty-one patients (60%) had SDB with an apnea-hypopnea index (AHI)  $\geq 5$ , and the average AHI was  $22.7 \pm 17.9$ /h. Twelve patients (34%) had moderate to severe SDB with an average AHI of  $33.6 \pm 16.6$ /h. Central dominant sleep apnea was evident in 3 patients with SDB. Average age and body mass index (BMI) were significantly higher in SDB patients than in non-SDB patients (age:  $62.0 \pm 12.8$  vs  $50.6 \pm 12.7$  years; BMI:  $26.3 \pm 4.0$  vs  $21.2 \pm 2.0$ ). In conclusion, this study found a high prevalence of SDB in patients with ventricular arrhythmias and normal LV function.

**Key Words:** Sleep-disordered breathing, ventricular arrhythmia, normal left ventricular function

## **Introduction**

Some previously reported studies that analyzed the occurrence of sleep-disordered breathing (SDB) with ventricular arrhythmias involved only patients with heart failure.<sup>1-3</sup> In patients with heart failure, ventricular arrhythmias occur frequently not only because of SDB but also cardiac dysfunction itself, resulting in sympathetic nerve activity.<sup>4</sup> Other studies have reported a high occurrence of ventricular arrhythmias in patients with SDB.<sup>5,6</sup> We tested the hypothesis that patients with ventricular arrhythmias are more likely to have SDB even without heart failure. We studied 35 patients with ventricular arrhythmias who did not have major comorbid disorders that could contribute to sleep disruption or desaturation. This study examined the prevalence and severity of SDB in patients having ventricular arrhythmias without heart failure.

## **Methods**

**Study population:** We prospectively studied consecutive patients referred to the Cardiology Department of the Tsukuba University Hospital, Tsukuba, Japan, between December 2005 and September 2007 who had ventricular tachycardia (VT) or frequent premature ventricular complexes (PVCs). Patients were referred for the following reasons: (1) the need for medication to control VT or frequent PVCs, (2)

electrophysiologic study or catheter ablation therapy for VT or frequent PVCs, or (3) implantation of a cardioverter/defibrillator for VT. Patients were excluded from enrollment if they had any of the following conditions: an echocardiographic left ventricular (LV) ejection fraction (EF) <50%, primary severe valvular heart disease, obstructive lung disease, pharyngeal disease, renal disorders, clinical signs of central or peripheral nervous system impairment, or a history of stroke. Thirty-five patients (63% men; mean age:  $57 \pm 14$  years; EF:  $64 \pm 8\%$ ) met the entry criteria. The initial assessment included routine blood tests, 12-lead electrocardiography, and echocardiography. Hypertension was defined as the presence of 1 or more of the following conditions: resting systolic blood pressure of at least 135 mmHg, resting diastolic blood pressure of at least 80 mmHg, or treatment with antihypertensive medication. Hyperlipidemia was defined as total cholesterol of  $\geq 220$  mg/dl or low-density lipoprotein  $\geq 150$  mg/dl, use of lipid-lowering therapy, or a documented diagnosis of hyperlipidemia. The diagnosis of diabetes mellitus and other prevalent chronic diseases were recorded according to the clinical history and use of specific medication, as revealed by review of the patient chart.

Written informed consent was obtained from all subjects, and the study protocol was approved by the Ethical Committee of the University of Tsukuba.

**Arrhythmia analysis:** Isolated and grouped ventricular ectopic activities were analyzed.

VT was defined as three or more PVCs in a row at a rate >100 beats/min. Frequent PVCs were defined as PVCs >300/h on 24-hour Holter electrocardiogram. Day-night patterns of ventricular arrhythmias were evaluated by 24-hour Holter electrocardiograms or bedside monitoring records in the hospital. Day was defined as 06:00–21:00 and night as 21:00–06:00 hours.

**Polysomnography:** Sleep evaluations of all 35 subjects were conducted by a sleep specialist at the Tsukuba University Hospital sleep disorder center. All subjects had undergone standard sleep studies, which were performed by monitoring of electroencephalogram, electrooculogram, electromyogram, electrocardiogram, thoracoabdominal excursions, pulse oximetry, and naso-oral airflow with an Alice 4 (Respironics; Pittsburgh, PA).

Apnea was defined as cessation of inspiration for at least 10 seconds. All such events were counted irrespective of the degree of oxygen desaturation or presence of an arousal. Obstructive apnea was defined as the absence of airflow in the presence of rib cage and/or abdominal excursions. Central apnea was defined as the absence of rib cage and abdominal excursions with absence of airflow. Hypopnea was defined as a reduction in airflow by at least 30% with a decrease in oxygen saturation (SaO<sub>2</sub>) by 4%

or more for at least 10 seconds in the presence of thoracoabdominal ventilatory efforts. The apnea-hypopnea index (AHI) was calculated as the sum of apneic and hypopneic events per hour of sleep.<sup>7</sup> The diagnosis of central sleep apnea required an AHI  $\geq 5/h$ , with more than 50% of the events determined to be central rather than obstructive. We used an AHI of 5/h as the threshold. Mild SDB was defined as  $5/h \leq \text{AHI} < 15/h$ , moderate SDB was defined as  $15 \leq \text{AHI} < 30/h$ , and severe SDB was defined as AHI  $\geq 30/h$ .

**Epworth Sleepiness Scale:** The Epworth Sleepiness Scale (ESS) is a frequently used, 8-item, self-administered subjective measure of sleepiness. The subject rates, on a scale of 0–3, the likelihood that he or she will doze off or fall asleep during 8 different situations commonly encountered in daily life. Scores are tallied across the 8 items to compute an ESS score; an ESS score  $\geq 11$  is considered indicative of subjective sleepiness. This scale has previously shown a high level of internal consistency among its 8 items, high test-retest reliability, and the ability to distinguish patients with excessive daytime sleepiness from normal subjects.

**Statistical analysis:** Results are expressed as mean  $\pm$  SD. Comparisons of continuous variables between groups were made with unpaired *t*-tests where appropriate and otherwise with the Mann-Whitney test. Categorical variables were compared by

Fisher's exact test or the  $\chi^2$  test, depending on which was more appropriate. A value of  $p < 0.05$  was considered significant.

## **Results**

The frequency distribution of the AHI in 5/h- to 15/h-unit intervals for the 35 patients with ventricular arrhythmia is depicted in Figure 1. Twenty-one of 35 patients (60%) had SDB, with mild SDB noted in 26% of patients (mean AHI:  $8.2 \pm 2.5/h$ ), moderate SDB in 14% (mean AHI:  $21.8 \pm 3.7/h$ ), and severe SDB in 20% (mean AHI:  $45.5 \pm 16.0/h$ ). The clinical characteristics of both groups are summarized in Table 1. The difference in sexes between patients with and without SDB was not statistically significant. Patients with SDB were older than those without SDB, and their BMI and waist circumference were significantly higher than those of patients without SDB. No statistical differences were noted in rates of obesity (BMI  $\geq 30$ ), hypertension, diabetes mellitus, hyperlipidemia, and cardiac disease between the 2 groups. Mean ESS score was  $< 11$  in both groups and was not statistically different between the 2 groups. There was no difference in the type of ventricular arrhythmias and the day-night pattern of the arrhythmias between the groups, as shown in Table 2. The results of the sleep study are presented in Table 3. Comparison of patients with and without SDB showed that SDB

patients had significantly increased arousal index and percent SaO<sub>2</sub> <90% and significantly decreased mean and lowest SaO<sub>2</sub>. The clinical characteristics of the 12 patients with moderate to severe SDB are shown in Table 4. Three of these patients had predominantly central sleep apnea: mean central apnea AHI was 17.0 ± 6.6/h. Three patients had ventricular arrhythmia occurring during night rather than day.

## **Discussion**

The novel findings of the present study are that approximately 60% of patients with ventricular arrhythmias had SDB. This is the first report to show a strong association between SDB and ventricular arrhythmias in patients without heart failure. Among such patients, mild SDB was diagnosed in 26% and moderate to severe SDB in 34%. To our knowledge, no data are available regarding the prevalence of SDB in patients with ventricular arrhythmias occurring without heart failure, although several prior studies showed a relation between SDB and ventricular arrhythmia complicated by heart failure.<sup>1-3</sup> Because impaired heart function increases sympathetic activity, resulting in ventricular arrhythmia,<sup>4</sup> heart failure is thought to exacerbate ventricular arrhythmia. However, the contribution of SDB to sympathetic activity in patients with ventricular arrhythmia and normal cardiac function has not been extensively discussed. In this study,



all participants had clinically severe ventricular arrhythmias without heart failure.

Interestingly, there was no difference either in ESS score or number of patients with ESS score  $\geq 11$ . This result showed that patients had no symptoms of daytime sleepiness even with SDB. Among subjects participating in the Sleep Heart Health Study, although there was a strong association of AHI with self-reported sleepiness, the majority of subjects with an AHI  $\geq 5$  did not report excessive sleepiness,<sup>13</sup> indicating that self-reporting measures may underestimate the severity of sleepiness in the setting of hypersomnolence.

Another novel finding of this study involved BMI in the patients with SDB; our patients had a mean BMI  $26.3 \pm 4.0$  kg/m<sup>2</sup>, with 19% having a BMI  $>30$ . Compared with previous reports<sup>8-10</sup> from Western nations, data from patients with SDB in the present study revealed a lower BMI and low rate of obesity. One report has estimated the prevalence of obstructive sleep apnea (OSA) in an Asian population.<sup>11</sup> Kim et al<sup>12</sup> reported the prevalence of SDB in Korea was 27% in men and 16% in women, and mean BMI was  $26.5 \pm 2.9$  in men and  $26.9 \pm 4.1$  in women. These are provocative results because obesity, a strong risk factor for SDB, is prevalent in white populations, but is relatively uncommon in Asian countries.

In addition, 3 of the SDB patients experienced ventricular arrhythmia more often

at night than during the day, suggesting that conditions associated with SDB may lead to ventricular arrhythmias. Gami et al<sup>18</sup> reported that people with SDB have a peak in sudden death from cardiac causes during the sleeping hours, which contrasts strikingly with the nadir of sudden death from cardiac causes during this period in people without SDB. In the present study, one of the patients without SDB was also found to have ventricular arrhythmia more often at night than during the day. This patient showed an arousal index of 28.1/h and an AHI of 18.1/h when we estimated hypopnea as the occurrence of at least 50% reduction in airflow lasting at least 10 seconds with an arousal. Interestingly, PVCs in this patient stopped synchronizing with arousal. No previous report has shown a relation between nocturnal occurrence PVCs and arousal. This finding suggests that not only arterial desaturation but also arousal may be related to the occurrence of ventricular arrhythmias.

Previous reports showed a relation between repetitive intermittent hypoxia and ventricular arrhythmias.<sup>19,20</sup> Alexander<sup>21</sup> reported that at high altitude, normal elderly persons experienced increased heart rate and greater frequency of PVCs and VT when arterial SaO<sub>2</sub> reached 70%. Shepard et al<sup>22</sup> reported that in SDB patients with SaO<sub>2</sub> <60%, a significant increase in PVC frequency was detected with decreasing SaO<sub>2</sub>. In patients with SDB, they suggested that repetitive obstructions to normal breathing

during sleep induce hypoxemia and hypercapnia, which (acting through the chemoreflexes) elicit increased sympathetic activity that induces ventricular arrhythmias.

Altered cardiovascular variability affects predominantly patients with moderate to severe sleep apnea.<sup>23-27</sup> In SDB patients, ventilation and blood pressure increase substantially during hypoxic breathing. Peripheral chemoreceptors, which primarily respond to blood oxygen, are detected with high sensitivity in SDB. In SDB patients, the chemoreflex appears to be a potent mechanism for sympathetic activation, overriding the combined restraining influences of increased blood pressure and increased ventilation. Enhanced chemoreflex sensitivity in SDB may explain the exaggerated sympathetic response during hypoxemic episodes resulting in autonomic activity-dependent arrhythmias.<sup>28-30</sup>

Our study had several limitations. First, due to technical limitations, oxygen saturation monitoring results of the patients with frequent PVCs were excluded because these data were thought to be underestimated. However, hypopnea could be analyzed by the occurrence of at least 50% reduction in airflow lasting at least 10 seconds with an arousal. Second, the diagnosis of sleep apnea was provided by a study based on a single night in a sleep laboratory. However, this is the standard procedure followed for the

diagnosis of sleep apnea in the clinical setting. Third, there is no control group. However, compared to the previous epidemiologic studies,<sup>11,12</sup> prevalence of SDB was revealed to be significantly high in this study.

### **Acknowledgments**

This study was supported in part by a Research Grant for Cardiovascular Disease (17C-5) from the Ministry of Health, Labor and Welfare, Japan.

## References

1. Fichter J, Bauer D, Arampatzis S, Fries R, Heisel A, Sybrecht GW. Sleep-related breathing disorders are associated with ventricular arrhythmias in patients with an implantable cardioverter-defibrillator. *Chest* 2002;122:558-561.
2. Javaheri S. Effects of continuous positive airway pressure on sleep apnea and ventricular irritability in patients with heart failure. *Circulation* 2000;101:392-397.
3. Ryan CM, Usui K, Floras JS, Bradley TD. Effect of continuous positive airway pressure on ventricular ectopy in heart failure patients with obstructive sleep apnoea. *Thorax* 2005;60:781-785.
4. Pogwizd SM, Schlotthauer K, Li L, Yuan W, Bers DM. Arrhythmogenesis and contractile dysfunction in heart failure: Roles of sodium-calcium exchange, inward rectifier potassium current, and residual beta-adrenergic responsiveness. *Circ Res* 2001;88:1159-1167.
5. Guilleminault C, Connolly SJ, Winkle RA. Cardiac arrhythmia and conduction disturbances during sleep in 400 patients with sleep apnea syndrome. *Am J Cardiol* 1983;52:490-494.
6. Mehra R, Benjamin EJ, Shahar E, Gottlieb DJ, Nawabit R, Kirchner HL, Sahadevan J, Redline S. Association of nocturnal arrhythmias with sleep-disordered breathing: The

Sleep Heart Health Study. *Am J Respir Crit Care Med* 2006;173:910-916.

7. Meoli AL, Casey KR, Clark RW, Coleman JA, Jr., Fayle RW, Troell RJ, Iber C. Hypopnea in sleep-disordered breathing in adults. *Sleep* 2001;24:469-470.

8. Marin JM, Carrizo SJ, Vicente E, Agusti AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet* 2005;365:1046-1053.

9. Kanagala R, Murali NS, Friedman PA, Ammash NM, Gersh BJ, Ballman KV, Shamsuzzaman AS, Somers VK. Obstructive sleep apnea and the recurrence of atrial fibrillation. *Circulation* 2003;107:2589-2594.

10. Gami AS, Pressman G, Caples SM, Kanagala R, Gard JJ, Davison DE, Malouf JF, Ammash NM, Friedman PA, Somers VK. Association of atrial fibrillation and obstructive sleep apnea. *Circulation* 2004;110:364-367.

11. Ip MS, Lam B, Tang LC, Lauder IJ, Ip TY, Lam WK. A community study of sleep-disordered breathing in middle-aged Chinese women in Hong Kong: prevalence and gender differences. *Chest* 2004;125:127-134.

12. Kim J, In K, Kim J, You S, Kang K, Shim J, Lee S, Lee J, Lee S, Park C, Shin C. Prevalence of sleep-disordered breathing in middle-aged Korean men and women. *Am J*

*Respir Crit Care Med* 2004;170:1108-1113.

**13.** Gottlieb DJ, Whitney CW, Bonekat WH, Iber C, James GD, Lebowitz M, Nieto FJ, Rosenberg CE. Relation of sleepiness to respiratory disturbance index: the Sleep Heart Health Study. *Am J Respir Crit Care Med* 1999;159:502-707.

**14.** Iwai S, Cantillon DJ, Kim RJ, Markowitz SM, Mittal S, Stein KM, Shah BK, Yarlagadda RK, Cheung JW, Tan VR, Lerman BB. Right and left ventricular outflow tract tachycardias: evidence for a common electrophysiologic mechanism. *J Cardiovasc Electrophysiol* 2006;17:1052-1058.

**15.** Lerman BB, Stein KM, Markowitz SM, Mittal S, Slotwiner DJ. Ventricular arrhythmias in normal hearts. *Cardiol Clin* 2000;18:265-291, vii.

**16.** Chiang BN, Perlman LV, Ostrander LD Jr, Epstein FH. Relationship of premature systoles to coronary heart disease and sudden death in the Tecumseh epidemiologic study. *Ann Intern Med* 1969;70:1159-1166.

**17.** Frolkis JP, Pothier CE, Blackstone EH, Lauer MS. Frequent ventricular ectopy after exercise as a predictor of death. *N Engl J Med* 2003;348:781-790.

**18.** Gami AS, Howard DE, Olson EJ, Somers VK. Day-night pattern of sudden death in obstructive sleep apnea. *N Engl J Med* 2005;352:1206-1214.

**19.** Jyothula SS, Ramachandran S. Reversible ventricular arrhythmia in REM sleep

associated with hypoxic sleep-disordered breathing. *Sleep Med* 2006;7:81-82.

**20.** O'Connor PJ, Merrill GF. Ventricular arrhythmias caused by repeat exposure to hypoxia are dependent on duration of reoxygenation. *FASEB J* 1995;9:387-391.

**21.** Alexander JK. Cardiac arrhythmia at high altitude: the progressive effect of aging. *Tex Heart Inst J* 1999;26:258-263.

**22.** Shepard JW, Jr., Garrison MW, Grither DA, Dolan GF. Relationship of ventricular ectopy to oxyhemoglobin desaturation in patients with obstructive sleep apnea. *Chest* 1985;88:335-340.

**23.** Aytemir K, Deniz A, Yavuz B, Ugur Demir A, Sahiner L, Ciftci O, Tokgozoglu L, Can I, Sahin A, Oto A. Increased myocardial vulnerability and autonomic nervous system imbalance in obstructive sleep apnea syndrome. *Respir Med* 2006.

**24.** Harbison J, O'Reilly P, McNicholas WT. Cardiac rhythm disturbances in the obstructive sleep apnea syndrome: effects of nasal continuous positive airway pressure therapy. *Chest* 2000;118:591-595.

**25.** Usui K, Bradley TD, Spaak J, Ryan CM, Kubo T, Kaneko Y, Floras JS. Inhibition of awake sympathetic nerve activity of heart failure patients with obstructive sleep apnea by nocturnal continuous positive airway pressure. *J Am Coll Cardiol* 2005;45:2008-2011.



- 26.** Aydin M, Altin R, Ozeren A, Kart L, Bilge M, Unalacak M. Cardiac autonomic activity in obstructive sleep apnea: time-dependent and spectral analysis of heart rate variability using 24-hour Holter electrocardiograms. *Tex Heart Inst J* 2004;31:132-136.
- 27.** Roche F, Xuong AN, Court-Fortune I, Costes F, Pichot V, Duverney D, Vergnon JM, Gaspoz JM, Barthelemy JC. Relationship among the severity of sleep apnea syndrome, cardiac arrhythmias, and autonomic imbalance. *Pacing Clin Electrophysiol* 2003;26:669-677.
- 28.** Phillips BG, Somers VK. Neural and humoral mechanisms mediating cardiovascular responses to obstructive sleep apnea. *Respir Physiol* 2000;119:181-187.
- 29.** Narkiewicz K, Montano N, Cogliati C, van de Borne PJ, Dyken ME, Somers VK. Altered cardiovascular variability in obstructive sleep apnea. *Circulation* 1998;98:1071-1077.
- 30.** Narkiewicz K, van de Borne PJ, Pesek CA, Dyken ME, Montano N, Somers VK. Selective potentiation of peripheral chemoreflex sensitivity in obstructive sleep apnea. *Circulation* 1999;99:1183-1189.

### **Figure Legends**

Figure 1. Frequency distribution of AHI in 5- to 15-unit intervals in 35 patients with ventricular arrhythmias. Fourteen patients (40%) showed no evidence of SDB, whereas 21 patients (60%) had SDB with  $AHI \geq 5$ .