

STABLE EQUIVALENCE BETWEEN UNIVERSAL COVERS OF TRIVIAL EXTENSION SELF-INJECTIVE ALGEBRAS

By

Takayoshi WAKAMATSU

Introduction.

Let A be an indecomposable basic artin algebra and T_A a basic tilting module with $B = \text{End}(T_A)$. Let us denote by R and S the trivial extension self-injective algebras $A \ltimes DA$ and $B \ltimes DB$, respectively. In the papers [24] and [22], H. Tachikawa and the author have proved that there is a stably equivalent functor $\mathcal{S}: \underline{\text{mod}}\text{-}R \rightarrow \underline{\text{mod}}\text{-}S$ and the restriction of \mathcal{S} to the tilting torsion class $\mathcal{T} = \{X \in \text{mod}\text{-}A \mid \text{Ext}_A^1(T, X) = 0\}$ coincides with that of the tilting functor $\text{Hom}_A(T, ?)$.

D. Hughes and J. Waschbüsch [18] introduced the following doubly infinite matrix algebra:


$$\hat{A} = \begin{pmatrix} \ddots & \ddots & & & & & & & 0 \\ & A_{n-1} & M_{n-1} & & & & & & \\ & & & A_n & M_n & & & & \\ & & & & & A_{n+1} & M_{n+1} & & \\ & & & & & \ddots & \ddots & \ddots & \\ & & & & & & \ddots & \ddots & \\ & & & & & & & & \ddots \\ 0 & & & & & & & & \ddots \end{pmatrix}$$

in which matrices are assumed to have only finitely many entries different from zero, $A_n = A$ and $M_n = DA$ for all integers n , all the remaining entries are zero, and multiplication is induced from the canonical maps $A \otimes_A DA \simeq DA$, $DA \otimes_A A \simeq DA$ and zero maps $DA \otimes_A DA \rightarrow 0$.

The identity maps $A_n \rightarrow A_{n+1}$, $M_n \rightarrow M_{n+1}$ induce an algebra isomorphism ν_A of \hat{A} . The orbit space \hat{A}/ν_A is easily seen to be the trivial extension algebra R . Similarly, we can consider the orbit space $\hat{A}/(\nu_A)^n$ as a self-injective algebra and it is denoted by R_n for each $n=1, 2, \dots, \infty$. Notice that $R_1 = R$ and $R_\infty = \hat{A}$.

The aim of this article is to prove the existence of a stably equivalent functor $\mathcal{S}_n: \underline{\text{mod}}\text{-}R_n \rightarrow \underline{\text{mod}}\text{-}S_n$ for each n . Here S_n is an orbit space $B/(\nu_B)^n$. The desired functor \mathcal{S}_n will be defined by slightly modifying the definition of the functor $\mathcal{S} = \mathcal{S}_1$ in [24] and [22].

In order to relate the categories $\text{mod-}\hat{A}$ and $\text{mod-}R$, Hughes-Waschbüsch used the exact functor $\Phi: \text{mod-}\hat{A} \rightarrow \text{mod-}R$ which preserves the indecomposability and the composition length of a module and also almost split sequences and irreducible maps. Similarly to the functor Φ , we can define the functors $\Phi_n: \text{mod-}\hat{A} \rightarrow \text{mod-}R_n$ and $\Phi_{m,n}: \text{mod-}R_{m,n} \rightarrow \text{mod-}R_n$. We shall show that the functors $\mathcal{S}_1 = \mathcal{S}, \mathcal{S}_2, \mathcal{S}_3, \dots, \mathcal{S}_\infty$ make the following diagrams commutative:



It should be noted that the functor Φ is not dense in general, though in the case where R is representation-finite or A is hereditary $\Phi = \Phi_1$ is dense and $\mathcal{S} = \mathcal{S}_1$ is induced from \mathcal{S}_∞ .

Recently, D. Happel [15] has proved that $\text{mod-}\hat{A}$ and $\text{mod-}\hat{B}$ are equivalent if $\text{gl. dim. } A < \infty$. But, since Φ is not dense in general even if $\text{gl. dim. } A < \infty$, our results does not follow from his one. At the end of this paper such an example will be given.

Throughout this paper, we fix a commutative artin ring K and all algebras are assumed to be artin K -algebras except R_∞ and S_∞ , and modules are finitely generated over K and morphisms operate on the opposite side of the scalars. The ordinary duality functor is always denoted by D .

1. Preliminaries

In this section, we shall recall some of basic results on tilting theory and trivial extension algebras for the later use.

Let T_A be a tilting module in the sense of Happel-Ringel [16]. Put $B = \text{End}(T_A)$, then ${}_B T$ is again a tilting module with $\text{End}({}_B T) = A$. Let us put $\mathcal{T} = \{X \in \text{mod-}A \mid \text{Ext}_A^1(T, X) = 0\}$, $\mathcal{F} = \{X \in \text{mod-}A \mid \text{Hom}_A(T, X) = 0\}$, $\mathcal{X} = \{Y \in \text{mod-}B \mid Y \otimes_B T = 0\}$ and $\mathcal{Y} = \{Y \in \text{mod-}B \mid \text{Tor}_1^B(Y, T) = 0\}$. Further, let $F = \text{Hom}_A(T, ?)$, $F' = \text{Ext}_A^1(T, ?)$ (resp. $G = (? \otimes_B T)$, $G' = \text{Tor}_1^B(? , T)$) be functors from $\text{mod-}A$ (resp. $\text{mod-}B$) to $\text{mod-}B$ (resp. $\text{mod-}A$). Then there are short exact sequences of functors

$$\begin{aligned}
 0 &\longrightarrow GF \xrightarrow{\varepsilon} 1_{\text{mod-}A} \longrightarrow G'F' \longrightarrow 0, \\
 0 &\longrightarrow F'G' \longrightarrow 1_{\text{mod-}B} \xrightarrow{\eta} FG \longrightarrow 0,
 \end{aligned}$$

where ε and η denote the counit and the unit of the adjunction (F, G) , respectively. Hence the restrictions of the functors F and G (resp. F' and G') give a category equivalence $\mathcal{T} \cong \mathcal{Q}$ (resp. $\mathcal{F} \cong \mathcal{X}$).

We call a short exact sequence $0 \rightarrow X_A \rightarrow V_A \rightarrow L_A \rightarrow 0$ a torsion resolution of X_A if $V \in \mathcal{T}$ and $L \in \text{add}(T_A)$. There is the minimal torsion resolution $0 \rightarrow X \xrightarrow{\alpha_X} V(X) \xrightarrow{\beta_X} T(X) \rightarrow 0$ for any A -module X and every torsion resolution of X is of the form

$$0 \longrightarrow X \longrightarrow V(X) \oplus T_0 \xrightarrow{\begin{pmatrix} \alpha_X & 0 \\ 0 & 1_{T_0} \end{pmatrix}} T(X) \oplus T_0 \longrightarrow 0.$$

Similarly, a short exact sequence $0 \rightarrow W_B \rightarrow U_B \rightarrow Y_B \rightarrow 0$ is said to be a torsion-free resolution of Y_B if $U \in \mathcal{Q}$ and $W \in \text{add}(DT_B)$. It is easy to see that the sequence $0 \rightarrow W_B \rightarrow U_B \rightarrow Y_B \rightarrow 0$ in the category $\text{mod-}B$ is a torsion-free resolution iff the corresponding sequence $0 \rightarrow {}_B DY \rightarrow {}_B DU \rightarrow {}_B DW \rightarrow 0$ in the category $B\text{-mod}$ is a torsion resolution. Therefore, there is the minimal torsion-free resolution $0 \rightarrow W(Y) \xrightarrow{\delta_Y} U(Y) \xrightarrow{\gamma_Y} Y \rightarrow 0$ and every torsion-free resolution is of the form

$$0 \longrightarrow W(Y) \oplus W_0 \xrightarrow{\begin{pmatrix} \delta_Y & 0 \\ 0 & 1_{W_0} \end{pmatrix}} U(Y) \oplus W_0 \xrightarrow{(\gamma_Y, 0)} Y \longrightarrow 0.$$

Any module X_R over the trivial extension self-injective algebra $R = A \ltimes DA$ is defined by giving its underlying A -module X_A and the A -morphism $\phi: X \otimes_A DA \rightarrow X$ such that $\phi \cdot (\phi \otimes DA) = 0$ and any R -morphism $f: X_R = (X_A, \phi) \rightarrow X'_R = (X'_A, \phi')$ can be considered as an A -morphism $f: X_A \rightarrow X'_A$ making the following diagram commutative:

$$\begin{array}{ccc}
 X \otimes_A DA & \xrightarrow{\phi} & X \\
 f \otimes DA \downarrow & & \downarrow f \\
 X' \otimes_A DA & \xrightarrow{\phi'} & X'.
 \end{array}$$

See [12] for details. If the underlying A -module X_A is decomposed as a direct sum $X_0 \oplus X_1$ and the morphism ϕ is of the form $\begin{pmatrix} 0 & 0 \\ \phi_0 & 0 \end{pmatrix}: (X_0 \oplus X_1) \otimes DA \rightarrow (X_0 \oplus X_1)$, we shall denote $X_R = (X_A, \phi)$ by $\begin{matrix} X_0 \\ \overline{X_1} \end{matrix}$. It should be noted that any indecomposable projective (= injective) R -module has to be of the form $fR = \begin{matrix} fA \\ \overline{fDA} \end{matrix}$ with a primi-

tive idempotent $f \in A \subset R$.

Similarly, by the definition, any object X in the category $\text{mod-}R_\infty (R_\infty = \hat{A})$ is defined by giving a family of A -modules $\{X_i\}_{i \in \mathbb{Z}}$ ($X_i \neq 0$ for only finite number of integers $i \in \mathbb{Z}$) and a family of A -morphisms $\{\phi_i: X_i \otimes_A DA \rightarrow X_{i+1}\}_{i \in \mathbb{Z}}$ satisfying $\phi_{i+1} \cdot (\phi_i \otimes_A DA) = 0$ for all $i \in \mathbb{Z}$. Any morphism in the category $\text{mod-}R_\infty$ from $X = \{X_i, \phi_i\}$ to $X' = \{X'_i, \phi'_i\}$ is a family of A -morphisms $\{f_i: X_i \rightarrow X'_i\}$ such that the following diagrams are commutative for all $i \in \mathbb{Z}$:

$$\begin{array}{ccc} X_i \otimes_A DA & \xrightarrow{\phi_i} & X_{i+1} \\ f_i \otimes_A DA \downarrow & & \downarrow f_{i+1} \\ X'_i \otimes_A DA & \xrightarrow{\phi'_i} & X'_{i+1}. \end{array}$$

Similarly to the above also, for any positive integer n , an R_n -module X is defined by giving a family of A -modules X_0, X_1, \dots, X_{n-1} and a family of A -morphisms $\phi_0: X_0 \otimes_A DA \rightarrow X_1, \dots, \phi_{n-2}: X_{n-2} \otimes_A DA \rightarrow X_{n-1}$ and $\phi_{n-1}: X_{n-1} \otimes_A DA \rightarrow X_0$ satisfying $\phi_{i+1} \cdot (\phi_i \otimes_A DA) = 0$ for each $i = 0, 1, \dots, n-1$. An R_n -morphism from $X = \{X_i, \phi_i\}$ to $X' = \{X'_i, \phi'_i\}$ is a family of A -morphisms $f = \{f_i: X_i \rightarrow X'_i\}$ such that $\phi'_i \cdot f_i \otimes_A DA = f_{i+1} \cdot \phi_i$ for each i . Where we put $X_{i+s \cdot n} = X_i$ and $\phi_{i+s \cdot n} = \phi_i$ ($1 \leq i \leq n, s \in \mathbb{N}$), for convenience.

Then the functors $\Phi_n: \text{mod-}R_\infty \rightarrow \text{mod-}R_n$ and $\Phi_{m,n}: \text{mod-}R_{m \cdot n} \rightarrow \text{mod-}R_n$ are defined as follows:

$$\Phi_n(\{X_j, \phi_j\}) = \{Y_i, \phi_i\}_{i=1}^n, \quad Y_i = \bigoplus_{j \equiv i \pmod{n}} X_j$$

and

$$\begin{aligned} \phi_i | X_j \otimes_A DA &= \phi_j \quad \text{for all } j \equiv i \pmod{n}. \\ \Phi_n(\{f_j: X_j \rightarrow X'_j\}) &= \{g_i: \bigoplus_{j \equiv i} X_j \rightarrow \bigoplus_{j \equiv i} X'_j\}, \quad g_i = \bigoplus_{j \equiv i} f_j. \end{aligned}$$

It is easy to verify that the functors $\Phi_n, \Phi_{m,n}$ are exact and preserve the projectivity (= injectivity), indecomposability and composition length of a module and almost split sequences and irreducible maps. Further they make the following commutative diagrams:

$$\begin{array}{ccc} & \text{mod-}R_\infty & \\ \Phi_{m \cdot n} \swarrow & & \searrow \Phi_n \\ \text{mod-}R_{m \cdot n} & \xrightarrow{\Phi_{m,n}} & \text{mod-}R_n \end{array} \quad \text{and} \quad \begin{array}{ccc} & \underline{\text{mod-}R_\infty} & \\ \underline{\Phi}_{m \cdot n} \swarrow & & \searrow \underline{\Phi}_n \\ \underline{\text{mod-}R}_{m \cdot n} & \xrightarrow{\underline{\Phi}_{m,n}} & \underline{\text{mod-}R}_n. \end{array}$$

Here $\underline{\text{mod-}}^*$ denotes the projectively (= injectively) stable category of mod-^* in the sense of M. Auslander, for each self-injective algebra * .

2. The functors $\mathcal{S}_n : \underline{\text{mod}}\text{-}R_n \rightarrow \underline{\text{mod}}\text{-}S_n$ and $\mathcal{Q}_n : \underline{\text{mod}}\text{-}S_n \rightarrow \underline{\text{mod}}\text{-}R_n$

In this section, we shall define the functor $\mathcal{S}_n : \underline{\text{mod}}\text{-}R_n \rightarrow \underline{\text{mod}}\text{-}S_n$ first and then, by making use of this functor \mathcal{S}_n , the functor \mathcal{Q}_n will be defined as the composite $\underline{\text{mod}}\text{-}S_n \xrightarrow{D} S_n\text{-}\underline{\text{mod}} \xrightarrow{\mathcal{S}_n} R_n\text{-}\underline{\text{mod}} \xrightarrow{D} \underline{\text{mod}}\text{-}R_n$. Notice that, since R_n and S_n are self-injective, the duality functor $D : \underline{\text{mod}}\text{-}R_n \xleftrightarrow{\sim} R_n\text{-}\underline{\text{mod}}$, $\underline{\text{mod}}\text{-}S_n \xleftrightarrow{\sim} S_n\text{-}\underline{\text{mod}}$ induces the duality functor $\underline{\text{mod}}\text{-}R_n \xleftrightarrow{\sim} R_n\text{-}\underline{\text{mod}}$, $\underline{\text{mod}}\text{-}S_n \xleftrightarrow{\sim} S_n\text{-}\underline{\text{mod}}$ (we denote this functor also by D). The functor $\mathcal{S}_n : S_n\text{-}\underline{\text{mod}} \rightarrow R_n\text{-}\underline{\text{mod}}$ can be defined similarly to $\mathcal{S}_n : \underline{\text{mod}}\text{-}R_n \rightarrow \underline{\text{mod}}\text{-}S_n$.

For an R_n -module $X = \{X_i, \phi_i\}$, we shall define S_n -modules $\mathcal{A}(X)$ and $\mathcal{B}(X)$ and S_n -monomorphism $u(X) : \mathcal{A}(X) \rightarrow \mathcal{B}(X)$ and the module $\mathcal{S}_n(X)$ is defined as its cokernel $\text{Cok } u(X)$. In order to define those S_n -modules and S_n -morphism, the following lemma is necessary

LEMMA 2.1. ${}_A D A_A \cong {}_A D T \otimes_B T_A$ and ${}_B D B_B \cong {}_B T \otimes_A D T_B$.

PROOF. Since ${}_B T_A$ is a balanced bimodule, we have ${}_A D A_A \cong {}_A D \text{Hom}({}_B T, {}_B T)_A \cong {}_A D T \otimes_B T_A$ and ${}_B D B_B \cong {}_B D \text{Hom}(T_A, T_A)_B \cong {}_B T \otimes_A D T_B$.


In the following, we can identify DA (resp. DB) with $DT \otimes T$ (resp. $T \otimes DT$). Further, from the lemma, it follows that ${}_A D A \otimes_A D T_B \cong {}_A D T \otimes_B T_A$ and ${}_B T \otimes_A D A_A \cong {}_B D B \otimes_B T_A$ and we shall identify these bimodules respectively.

Now let us put $\mathcal{A}(X) = \{X_i \otimes DT, -\phi_i \otimes DT : X_i \otimes DT \otimes DB = X_i \otimes DA \otimes DT \rightarrow X_{i+1} \otimes DT\}$ and $\mathcal{B}(X) = \left\{ F(V(X_{i+1})) \oplus F(V(X_i)) \otimes DB, \begin{pmatrix} 0 & 0 \\ 1_{F(V(X_{i+1})) \otimes DB} & 0 \end{pmatrix} : F(V(X_{i+1})) \otimes DB \oplus F(V(X_i)) \otimes DB \rightarrow F(V(X_{i+2})) \oplus F(V(X_{i+1})) \otimes DB \right\}$. Then it is not hard to see that $\mathcal{A}(X)$ and $\mathcal{B}(X)$ become S_n -modules. We shall define the map $u(X)$ by the following:

$$u(X)_i = \begin{pmatrix} F(\alpha_{X_{i+1}} \cdot -\phi_i) \cdot \eta_{X_i \otimes DT} \\ (\epsilon_{V(X)_i} \otimes DT)^{-1} \cdot \alpha_{X_i} \otimes DT \end{pmatrix};$$

$$\mathcal{A}(X)_i = X_i \otimes DT \longrightarrow F(V(X_{i+1})) \oplus F(V(X_i)) \otimes DB = \mathcal{B}(X)_i.$$

To see that the above map $u(X)$ is an S_n -morphism, it is enough to show the commutativity of the following diagram:



LEMMA 2.2. The above diagram is commutative.

PROOF. From the naturality of the ε and η , we have the following equalities:

$$\begin{aligned}
 & \varepsilon_{V(X_{i+1})} \otimes DT \cdot F(\alpha_{X_{i+1}} \cdot -\phi_i) \otimes DB \cdot \eta_{X_i \otimes DT} \otimes DB \\
 &= (\alpha_{X_{i+1}} \cdot -\phi_i) \otimes DT \cdot \varepsilon_{X_i \otimes DA} \otimes DT \cdot \eta_{X_i \otimes DT} \otimes DB \\
 &= (\alpha_{X_{i+1}} \cdot -\phi_i) \otimes DT \cdot (\varepsilon_{X_i \otimes DT \otimes T} \cdot \eta_{X_i \otimes DT} \otimes T) \otimes DT \\
 &= (\alpha_{X_{i+1}} \cdot -\phi_i) \otimes DT \cdot 1_{X_i \otimes DT \otimes DB} \\
 &= \alpha_{X_{i+1}} \otimes DT \cdot (-\phi_i \otimes DT), \\
 & F(\alpha_{X_{i+2}} \cdot -\phi_{i+1}) \cdot \eta_{X_{i+1} \otimes DT} \cdot -\phi_i \otimes DT \\
 &= F(\alpha_{X_{i+2}} \cdot -\phi_{i+1}) \cdot F(-\phi_i \otimes DA) \cdot \eta_{X_i \otimes DA \otimes DT} \\
 &= F(\alpha_{X_{i+2}}) \cdot F(\phi_{i+1} \cdot \phi_i \otimes DA) \cdot \eta_{X_i \otimes DA \otimes DT} = 0.
 \end{aligned}$$

The desired commutativity follows from the above equalities.

Since $\alpha_{X_i} \otimes DT$ is an injection and $\varepsilon_{V(X_i)} \otimes DT$ is a bijection, $u(X)_i$ is also an injection for each i . Therefore, $u(X)$ is an S_n -monomorphism. Thus we can define the S_n -module $\mathcal{S}_n(X)$ as the cokernel $\text{Cok } u(X)$ of this S_n -monomorphism $u(X)$.

From the definition of \mathcal{S}_n , the following lemma is easily checked.

LEMMA 2.3. For any projective R_n -module P , the S -module $\mathcal{S}_n(P)$ is also

projective.

The remaining part of this section is devoted to the proof of the following proposition.

PROPOSITION 2.4. *The correspondence S_n can be seen as a stable functor from $\underline{\text{mod}}\text{-}R_n$ to $\underline{\text{mod}}\text{-}S_n$.*

It is necessary to define the S_n -morphism $S_n(f)$ for any R_n -morphism $f = \{f_i\} : X = \{X_i, \phi_i\} \rightarrow \{X'_i, \phi'_i\} = X'$, at first. In order to define such a morphism, it is sufficient to define S_n -morphisms $\mathcal{A}(f) : \mathcal{A}(X) \rightarrow \mathcal{A}(X')$ and $\mathcal{B}(f) : \mathcal{B}(X) \rightarrow \mathcal{B}(X')$ such that $u(X') \cdot \mathcal{A}(f) = \mathcal{B}(f) \cdot u(X)$.

Let us put $\mathcal{A}(f)$ and $\mathcal{B}(f)$ as follows :

$$\mathcal{A}(f)_i = f_i \otimes DT : \mathcal{A}(X)_i = X_i \otimes DT \longrightarrow X'_i \otimes DT = \mathcal{A}(X')_i,$$

$$\mathcal{B}(f)_i = \begin{pmatrix} F(f_{i+1}^*) & \\ 0 & F(f_i^*) \otimes DB \end{pmatrix} :$$

$$\mathcal{B}(X)_i = F(V(X_{i+1})) \oplus F(V(X_i)) \otimes DB \longrightarrow F(V(X'_{i+1})) \oplus F(V(X'_i)) \otimes DB = \mathcal{B}(X')_i,$$

where f_i^* is defined by the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & X_i & \xrightarrow{\alpha_{X_i}} & V(X_i) & \xrightarrow{\beta_{X_i}} & T(X_i) \longrightarrow 0 \\ & & f_i \downarrow & & f_i^* \downarrow & & f_i^{**} \downarrow \\ 0 & \longrightarrow & X'_i & \xrightarrow{\alpha_{X'_i}} & V(X'_i) & \xrightarrow{\beta_{X'_i}} & T(X'_i) \longrightarrow 0. \end{array}$$

The fact that $\mathcal{A}(f)$ and $\mathcal{B}(f)$ are S_n -morphisms is clear.

LEMMA 2.5. *The above morphisms $\mathcal{A}(f)$ and $\mathcal{B}(f)$ satisfy $u(X') \cdot \mathcal{A}(f) = \mathcal{B}(f) \cdot u(X)$.*

PROOF. We have to verify the following two equalities :

(a) $\varepsilon_{V(X'_i)} \otimes DT \cdot F(f_i^*) \otimes DB \cdot (\varepsilon_{V(X_i)} \otimes DT)^{-1} \cdot \alpha_{X_i} \otimes DT = \alpha_{X'_i} \otimes DT \cdot f_i \otimes DT$

and

(b) $F(f_{i+1}^* \cdot \alpha_{X_{i+1}} \cdot - \phi_i) \cdot \eta_{X_i \otimes DT} = F(\alpha_{X'_{i+1}} \cdot - \phi'_i) \cdot \eta_{X'_i \otimes DT} \cdot f_i \otimes DT.$

The above two equalities (a) and (b) follow from the naturality of ε and η and the following three equalities : $f_{i+1} \cdot \phi_i = \phi'_i \cdot f_i \otimes DA$, $f_i^* \cdot \alpha_{X_i} = \alpha_{X'_i} \cdot f_i$ and $f_{i+1}^* \cdot \alpha_{X_{i+1}} = \alpha_{X'_{i+1}} \cdot f_{i+1}$

Therefore we have defined the S_n -morphism $S(f)$ by the following commuta-

tive diagram :

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mathcal{A}(X) & \xrightarrow{u(X)} & \mathcal{B}(X) & \longrightarrow & \mathcal{S}_n(X) \longrightarrow 0 \\
 & & \mathcal{A}(f) \downarrow & & \mathcal{B}(f) \downarrow & & \downarrow \mathcal{S}_n(f) \\
 0 & \longrightarrow & \mathcal{A}(X') & \xrightarrow{u(X')} & \mathcal{B}(X') & \longrightarrow & \mathcal{S}_n(X') \longrightarrow 0.
 \end{array}$$

By the definition, $\mathcal{A}(f)$ is uniquely determined by f but $\mathcal{B}(f)$ is not and so $\mathcal{S}_n(f)$ is not uniquely determined by f . However, in the stable category $\underline{\text{mod}}\text{-}\mathcal{S}_n$, we can prove the singleness of the morphism $\mathcal{S}_n(f)$. To show this fact, we shall prove that $\mathcal{S}_n(f)$ factors through projective \mathcal{S}_n -modules if $f=0$.

Since $f_i=0$, there is a morphism $\delta_i : T(X_i) \rightarrow V(X'_i)$ and $f_i^* = \delta_i \cdot \beta_{X_i}$. Let $\mathcal{P}(X)$ be a projective \mathcal{S}_n -module defined by

$$\mathcal{P}(X)_i = F(T(X_{i+1})) \oplus F(T(X_i)) \otimes DB$$

and

$$(\mathcal{P}(X)_i \otimes DB \rightarrow \mathcal{P}(X)_{i+1}) = \begin{pmatrix} 0 & 0 \\ 1_{F(T(X_{i+1})) \otimes DB} & 0 \end{pmatrix} :$$

$$F(T(X_{i+1})) \otimes DB \oplus F(T(X_i)) \otimes DB \otimes DB \longrightarrow F(T(X_{i+2})) \oplus F(T(X_{i+1})) \otimes DB.$$

It is possible to define \mathcal{S}_n -morphisms $\beta(X)$ from $\mathcal{B}(X)$ to $\mathcal{P}(X)$ and Δ from $\mathcal{P}(X)$ to $\mathcal{B}(X')$ so that $\mathcal{B}(f) = \Delta \cdot \beta(X)$ by putting :

$$\beta(X)_i = F(\beta_{X_{i+1}}) \oplus F(\beta_{X_i}) \otimes DB :$$

and

$$F(V(X_{i+1})) \oplus F(V(X_i)) \otimes DB \longrightarrow F(T(X_{i+1})) \oplus F(T(X_i)) \otimes DB$$

$$\Delta_i = F(\delta_{i+1}) \oplus F(\delta_i) \otimes DB :$$

$$F(T(X_{i+1})) \oplus F(T(X_i)) \otimes DB \longrightarrow F(V(X'_{i+1})) \oplus F(V(X'_i)) \otimes DB.$$

It is easy to see that $\beta(X) \cdot u(X) = 0$ and $\mathcal{S}_n(f)$ factors through projective \mathcal{S}_n -module $\mathcal{P}(X)$:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mathcal{A}(X) & \xrightarrow{u(X)} & \mathcal{B}(X) & \longrightarrow & \mathcal{S}_n(X) \longrightarrow 0 \\
 & & & & \beta(X) \downarrow & \swarrow \text{---} & \\
 & & & & \mathcal{P}(X) & & \\
 & & & & \Delta \downarrow & & \\
 0 & \longrightarrow & \mathcal{A}(X') & \xrightarrow{u(X')} & \mathcal{B}(X') & \longrightarrow & \mathcal{S}_n(X') \longrightarrow 0.
 \end{array}$$

Therefore, we have defined the functor $\text{nod-}R_n \rightarrow \underline{\text{mod}}\text{-}\mathcal{S}_n$ and this functor induces the desired stable functor $\tilde{\mathcal{S}}_n : \underline{\text{mod}}\text{-}R_n \rightarrow \underline{\text{mod}}\text{-}\mathcal{S}_n$ by Lemma 2.3. This completes the proof of Proposition 2.4.

From the definition of the functors Φ_n , $\Phi_{m,n}$ and S_n , the commutativity of the diagram in Introduction is now obvious.

3. The functor $Q_n : \underline{\text{mod}}\text{-}S_n \rightarrow \underline{\text{mod}}\text{-}R_n$

The functor Q_n has defined as the composite $\underline{\text{mod}}\text{-}S_n \xrightarrow{D} S_n\text{-}\underline{\text{mod}} \xrightarrow{S_n} R_n\text{-}\underline{\text{mod}} \xrightarrow{D} \underline{\text{mod}}\text{-}R_n$. In this section, we shall show the construction of this functor in an explicit way, for the later use.

In the definition of the functor S_n , we expressed R_n - and S_n -modules as the tensor forms: $\{X_i, \phi_i : X_i \otimes DA \rightarrow X_{i+1}\}$ and $\{Y_i, \psi_i : Y_i \otimes DB \rightarrow Y_{i+1}\}$. But for the definition of the functor Q_n , it is convenient to express the modules as the hom-forms: $\{X_i, \bar{\phi}_i : X_i \rightarrow \text{Hom}_A(DA, X_{i+1})\}$ and $\{Y_i, \bar{\psi}_i : Y_i \rightarrow \text{Hom}_B(DB, Y_{i+1})\}$, where $\bar{\phi}_i$ (resp. $\bar{\psi}_i$) is the adjoint of ϕ_i (resp. ψ_i) which corresponds to ϕ_i (resp. ψ_i) by the canonical isomorphism $\text{Hom}_A(X_i \otimes_A DA, X_{i+1}) \cong \text{Hom}_A(X_i, \text{Hom}_A(DA, X_{i+1}))$ (resp. $\text{Hom}_B(Y_i \otimes_B DB, Y_{i+1}) \cong \text{Hom}_B(Y_i, \text{Hom}_B(DB, Y_{i+1}))$).

In the following we shall sometimes abbreviate $\text{Hom}(?, ?)$ by $[?, ?]$.

For an S_n -module $Y = \{Y_i, \bar{\psi}_i\}$, let us put

$$\begin{aligned} \mathcal{C}(Y) &= \{[DT, Y_i], [DT, -\bar{\psi}_i] : [DT, Y_i] \longrightarrow [DT, [DB, Y_{i+1}]] \\ &= [DA, [DT, Y_{i+1}]]\} \end{aligned}$$

and

$$\begin{aligned} \mathcal{D}(Y) &= \left\{ [DA, G(U(Y_i))] \oplus G(U(Y_{i-1})), \begin{pmatrix} 0 & 0 \\ [DA, G(U(Y_i))] & 0 \end{pmatrix} : \right. \\ & [DA, G(U(Y_i))] \oplus G(U(Y_{i-1})) \longrightarrow [DA, [DA, G(U(Y_{i+1}))]] \\ & \left. \oplus [DA, G(U(Y_i))] \right\} \end{aligned}$$

and define the map $p(Y) : \mathcal{D}(Y) \rightarrow \mathcal{C}(Y)$ as follows:

$$\begin{aligned} p(Y)_i &= ([DT, \gamma_{Y_i} \cdot (\eta_{U(Y_i)})^{-1}, \epsilon_{[DT, Y_i]} \cdot G(-\bar{\psi}_i \cdot \gamma_{Y_{i-1}})] : \\ \mathcal{D}(Y)_i &= [DA, G(U(Y_i))] \oplus G(U(Y_{i-1})) \longrightarrow [DT, Y_i] = \mathcal{C}(Y)_i. \end{aligned}$$

Then $\mathcal{C}(Y)$ and $\mathcal{D}(Y)$ become R_n -modules and $p(Y)$ is an R_n -morphism. The module $Q_n(Y)$ coincides with the kernel $\text{Ker } p(Y)$ of the above morphism $p(Y)$.

For an S_n -morphism $g = \{g_i : Y_i \rightarrow Y'_i\} : Y = \{Y_i, \bar{\psi}_i\} \rightarrow Y' = \{Y'_i, \bar{\psi}'_i\}$, we put $\mathcal{C}(g) : \mathcal{C}(Y) \rightarrow \mathcal{C}(Y')$ and $\mathcal{D}(g) : \mathcal{D}(Y) \rightarrow \mathcal{D}(Y')$ as follows:

$$\begin{aligned} \mathcal{C}(g)_i &= [DT, g_i] : [DT, Y_i] \longrightarrow [DT, Y'_i], \\ \mathcal{D}(g)_i &= \begin{pmatrix} [DA, G(g_i^*)] & 0 \\ 0 & G(g_{i-1}^*) \end{pmatrix} : \\ & [DA, G(U(Y_i))] \oplus G(U(Y_{i-1})) \longrightarrow [DA, G(U(Y'_i))] \oplus G(U(Y'_{i-1})), \end{aligned}$$

where g_i^* is defined by the following commutative diagram :

$$\begin{array}{ccccccc}
 0 & \longrightarrow & W(Y_i) & \xrightarrow{\delta_{Y_i}} & U(Y_i) & \xrightarrow{\gamma_{Y_i}} & Y_i \longrightarrow 0 \\
 & & \downarrow g_i^{**} & & \downarrow g_i^* & & \downarrow g_i \\
 0 & \longrightarrow & W(Y'_i) & \xrightarrow{\delta_{Y'_i}} & U(Y'_i) & \xrightarrow{\gamma_{Y'_i}} & Y'_i \longrightarrow 0.
 \end{array}$$

Then $C(g)$ and $\mathcal{D}(g)$ become R_n -morphisms and satisfy $C(g) \cdot p(Y) = p(Y') \cdot \mathcal{D}(g)$. The R_n -morphism $Q_n(g) : Q_n(Y) \rightarrow Q_n(Y')$ is defined by the following commutative diagram :

$$\begin{array}{ccccccc}
 0 & \longrightarrow & Q_n(Y) & \longrightarrow & \mathcal{D}(Y) & \xrightarrow{p(Y)} & C(Y) \longrightarrow 0 \\
 & & \downarrow Q_n(g) & & \downarrow D(g) & & \downarrow C(g) \\
 0 & \longrightarrow & Q_n(Y') & \longrightarrow & \mathcal{D}(Y') & \xrightarrow{p(Y')} & C(Y') \longrightarrow 0.
 \end{array}$$

Similarly to the functor S_n , Q_n can be considered as a functor $\text{mod-}S_n \rightarrow \text{mod-}R_n$ and it induces $\text{mod-}S_n \rightarrow \text{mod-}R_n$.

4. The proof of the isomorphism $Q_n \cdot S_n \cong I_{\text{mod-}R_n}$

We begin with the survey of the torsion-free resolution of the component of $S_n(X)$, in order to investigate the module $Q_n S_n(X)$.

Let us denote the morphism $\text{cok } u(X)$ by $\theta(X)$:

$$\theta(X)_i = (x_i, y_i) : F(V(X_{i+1})) \oplus F(V(X_i)) \otimes DB \longrightarrow S_n(X)_i.$$

Let $P_0^i \xrightarrow{p_0^i} V(X_i) \rightarrow 0$ be the projective cover, then we have the following commutative diagram with exact rows :

$$\begin{array}{ccccccc}
 0 & \longrightarrow & P_1^i & \xrightarrow{\alpha^i} & P_0^i & \xrightarrow{\beta^i = \beta_{x_i} \cdot p_0^i} & T(X_i) \longrightarrow 0 \\
 & & \downarrow p_1^i & & \downarrow p_0^i & & \parallel \\
 0 & \longrightarrow & X_i & \xrightarrow{\alpha_{x_i}} & V(X_i) & \xrightarrow{\beta_{x_i}} & T(X_i) \longrightarrow 0
 \end{array}$$

Since $\text{proj. dim. } T_A \leq 1$, P_1^i has to be projective. Applying $(? \otimes_A DT)$ to the above diagram, we have the following commutative diagram :

$$\begin{array}{ccccccc}
 0 & \longrightarrow & P_1^i \otimes DT & \xrightarrow{\alpha^i \otimes DT} & P_0^i \otimes DT & \xrightarrow{\beta^i \otimes DT} & T(X_i) \otimes DT \longrightarrow 0 \\
 & & \downarrow p_1^i \otimes DT & & \downarrow p_0^i \otimes DT & & \parallel \\
 0 & \longrightarrow & X_i \otimes DT & \xrightarrow{\alpha_{x_i} \otimes DT} & V(X_i) \otimes DT & \xrightarrow{\beta_{x_i} \otimes DT} & T(X_i) \otimes DT \longrightarrow 0.
 \end{array}$$

Here we used the fact that $\text{Tor}_1^A(T(X_i), DT) \cong D\text{Ext}_A^1(T(X_i), T) = 0$. Hence we know $\text{Ker}(p_1^i \otimes DT)_B \cong \text{Ker}(p_0^i \otimes DT)_B$ by the Snake Lemma.

Consider the following diagram of B -modules:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & P_1^i \otimes DT & \xrightarrow{\zeta} & F(V(X_{i+1})) & & \\
 & & \downarrow p_1^i \otimes DT & & \downarrow \chi & & \\
 0 & \longrightarrow & \mathcal{A}(X)_i & \xrightarrow{u(X)_i} & \mathcal{B}(X)_i & \xrightarrow{\theta(X)_i} & \mathcal{S}_n(X)_i \longrightarrow 0,
 \end{array}$$

where ζ and χ are defined as follows:

$$\zeta = \begin{pmatrix} F(\alpha_{X_i} \cdot -\phi_i) \cdot \eta_{X_i \otimes DT} \cdot p_1^i \otimes DT \\ (\eta_{P_0^i \otimes DT})^{-1} \cdot \alpha^i \otimes DT \end{pmatrix},$$

$$\chi = \begin{pmatrix} 1_{F(V(X_{i+1}))} & 0 \\ 0 & (\varepsilon_{V(X_i)} \otimes DT)^{-1} \cdot p_0^i \otimes DT \cdot (\eta_{P_0^i \otimes DT})^{-1} \end{pmatrix}.$$

From the fact that $\text{Ker}(p_0^i \otimes DT)_B \cong \text{Ker}(p_1^i \otimes DT)_B$, it follows that $\text{Ker} \chi \cong \text{Ker}((\varepsilon_{V(X_i)} \otimes DT)^{-1} \cdot p_0^i \otimes DT \cdot (\eta_{P_0^i \otimes DT})^{-1}) \cong \text{Ker}(p_0^i \otimes DT) \cong \text{Ker}(p_1^i \otimes DT)$. Therefore we have $\text{Cok} \zeta \cong \mathcal{S}_n(X)_i$ and we have a torsion-free resolution of $\mathcal{S}_n(X)_i$.

LEMMA 4.1. *The following exact sequence is a torsion-free resolution of $\mathcal{S}_n(X)_i$:*

$$0 \longrightarrow P_1^i \otimes DT \xrightarrow{\zeta} F(V(X_{i+1})) \oplus F(P_0^i \otimes DA) \longrightarrow \mathcal{S}_n(X)_i \longrightarrow 0.$$

It is clear that $P_1^i \otimes DT \in \text{add}(DT_B)$ and $F(V(X_{i+1})) \oplus F(P_0^i \otimes DA) \in \text{add}(A)$. We shall denote $\text{coker} \zeta$ by $\hat{\theta}_i = (x_i, \hat{y}_i)$.

To define the modules $\mathcal{S}_n(X)$ and $\mathcal{Q}_n(Y)$, we have used the minimal torsion and torsion-free resolutions. But by the remark on torsion and torsion-free resolutions in section one, we may use any such resolutions since we consider modules in the stable categories.

Now, using the torsion-free resolutions given by Lemma 4.1, let us calculate the module $\mathcal{Q}_n \mathcal{S}_n(X)$.

The routine verification shows the following lemma.

LEMMA 4.2. *The map $p(\mathcal{S}_n(X))$ is expressed as follows:*

- (a) $\mathcal{C}\mathcal{S}_n(X)_i = [DT, \mathcal{S}_n(X)_i]$,
- (b) $\mathcal{D}\mathcal{S}_n(X)_i = [DA, GF(V(X_{i+1}))] \oplus [DA, GF(P_0^i \otimes DA)]$
 $\oplus GF(V(X_i)) \oplus GF(P_0^{i-1} \otimes DA)$

$$(c) \quad p(\mathcal{S}_n(X))_i = ([DT, x_i \cdot (\eta_{F(V(X_{i+1}))})^{-1}], [DT, \hat{y}_i \cdot (\eta_{F(P_0^i \otimes DA)})^{-1}], \\ \varepsilon_{[DT, \mathcal{S}_n(X)_i]} \cdot F(-\bar{\varphi}_{i-1} \cdot x_{i-1}), \varepsilon_{[DT, \mathcal{S}_n(X)_i]} \cdot F(-\bar{\varphi}_{i-1} \cdot \hat{y}_{i-1})),$$

where we identify $[DA, ?]$ (resp. $[DB, ?]$) with $[DT, [T, ?]]$ (resp. $[T, [DT, ?]]$) and $\bar{\varphi}_i: \mathcal{S}_n(X)_i \rightarrow [DB, \mathcal{S}_n(X)_{i+1}]$ denotes the i -th structure map of the \mathcal{S}_n -module $\mathcal{S}_n(X)$ in the hom-form.

The remaining part of this section is devoted to prove that $\text{Ker } p(\mathcal{S}_n(X))$ is isomorphic to X as an object in the stable category $\text{mod-}R_n$. In fact, we shall show $\text{Ker } p(\mathcal{S}_n(X)) \cong X \oplus P$ for the projective (= injective) R_n -module P defined as follows:

$$P = \left\{ P_0^i \oplus P_0^{i-1} \otimes DA, \begin{pmatrix} 0 & 0 \\ 1_{P_0^i \otimes DA} & 0 \end{pmatrix} \right\}.$$

LEMMA 4.3. $|\text{Ker } p(\mathcal{S}_n(X))| = |X| + |P|$, where $|*|$ denotes the K -composition length of a module $*$.

PROOF. By Lemma 4.1, we have

$$|\text{Ker } p(\mathcal{S}_n(X))_i| = |[DA, V(X_{i+1})]| + |P_0^i| + |V(X_i)| \\ + |P_0^{i-1} \otimes DA| - |[DT, \mathcal{S}_n(X)_i]|,$$

since $\varepsilon_V: FG(V) \cong V$ for a torsion A -module V and $\eta_U: U \cong GF(U)$ for a torsion-free B -module U by Brenner-Butler's theorem. On the other hand, from the exact sequence

$$0 \longrightarrow P_0^i \otimes DT \longrightarrow F(V(X_{i+1})) \oplus F(P_0^i \otimes DA) \longrightarrow \mathcal{S}_n(X)_i \longrightarrow 0$$

we have the exact sequence

$$0 \longrightarrow [DT, P_0^i \otimes DT] \longrightarrow [DT, F(V(X_{i+1}))] \oplus [DT, F(P_0^i \otimes DA)] \\ \longrightarrow [DT, \mathcal{S}_n(X)_i] \longrightarrow 0$$

and $[DT, F(P_0^i \otimes DA)] \cong [DA, P_0^i \otimes DA] \cong P_0^i$, $[DT, P_0^i \otimes DT] \cong [DT, D[P_0^i, T]] \cong [[P_0^i, T], T] \cong P_0^i$, as well. Therefore, it follows $|[DT, \mathcal{S}_n(X)_i]| = |[DA, V(X_{i+1})]| + |P_0^i| - |P_0^i|$. Hence we have $|\text{Ker } p(\mathcal{S}_n(X))_i| = |V(X_i)| + |P_0^{i-1} \otimes DA| + |P_0^i|$. Further, from the exact commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & P_0^i & \longrightarrow & P_0^i & \longrightarrow & T(X_i) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & X_i & \longrightarrow & V(X_i) & \longrightarrow & T(X_i) \longrightarrow 0, \end{array}$$

we know $|P_0^i| - |P_0^i| = |T(X_i)| = |V(X_i)| - |X_i|$, i.e., $|V(X_i)| + |P_0^i| = |P_0^i| + |X_i|$.

Finally, we have $|\text{Ker } p(\mathcal{S}_n(X))_i| = |P_0^{i-1} \otimes DA| + |P_0^i| + |X_i| = |X_i| + |P_i|$ and this means that $|\text{Ker } p(\mathcal{S}_n(X))| = |X| + |P|$ as desired.

By the above lemma, in order to prove the isomorphism $\text{Ker } p(\mathcal{S}_n(X)) \cong X \oplus P$, it suffices to show the existence of an R_n -monomorphism $(e(X), f(X)): X \oplus P \rightarrow \mathcal{D}\mathcal{S}_n(X)$ such that the composition $p(\mathcal{S}_n(X)) \cdot (e(X), f(X))$ is a zero map. To define such morphisms $e(X)$ and $f(X)$, it is necessary to introduce a notation: For a bimodule ${}_{E_1}M_{E_2}$ over algebras E_1 [and E_2 , we can always consider the adjoint pair of functors $\text{Hom}_{E_2}(M, ?): \text{mod-}E_2 \rightarrow \text{mod-}E_1$ and $(? \otimes_{E_1} M): \text{mod-}E_1 \rightarrow \text{mod-}E_2$. We shall denote the unit and counit of this adjunction by $\eta^M: 1_{\text{mod-}E_1} \rightarrow \text{Hom}_{E_2}(M, ? \otimes_{E_1} M)$ and $\varepsilon^M: \text{Hom}_{E_2}(M, ?) \otimes_{E_1} M \rightarrow 1_{\text{mod-}E_2}$, respectively. Then it is noted that $\eta = \eta^T$ and $\varepsilon = \varepsilon^T$.

Now let us put $e(X): X \rightarrow \mathcal{D}\mathcal{S}_n(X)$ and $f(X): P \rightarrow \mathcal{D}\mathcal{S}_n(X)$ as follows:

$$e(X)_i = \begin{bmatrix} [DA, (\varepsilon_{V(X_{i+1})})^{-1} \cdot \alpha_{X_{i+1}} \cdot \phi_i] \cdot \eta_{X_i}^{DA} \\ 0 \\ (\varepsilon_{V(X_i)})^{-1} \cdot \alpha_{X_i} \\ 0 \end{bmatrix}$$

and

$$f(X)_i = \begin{bmatrix} 0 & 0 \\ [DT, \eta_{[T, P_0^i \otimes DA]}] & 0 \\ (\varepsilon_{V(X_i)})^{-1} \cdot p_0^i & 0 \\ 0 & (\varepsilon_{P_0^{i-1} \otimes DA})^{-1} \end{bmatrix}.$$

In the following, we shall show that $e(X)$ and $f(X)$ are R_n -homomorphisms and $p(\mathcal{S}_n(X)) \cdot e(X) = 0 = p(\mathcal{S}_n(X)) \cdot f(X)$. To do so, it is necessary to provide the following lemma.

LEMMA 4.4. *The following diagrams are commutative for any A -module X and B -morphism $g: Z \otimes DB \rightarrow Y$.*

$$\begin{array}{ccc} & [DT, \eta_{F(X)}] & \\ (a) & [DT, F(X)] \xrightarrow{\quad} & [DT, FGF(X)] \\ & \parallel & \parallel \\ & [DA, X] \xrightarrow{\quad} & [DA, GF(X)] \\ & [DA, \varepsilon_X] & \\ & & \\ & X & \xrightarrow{\eta_X^{DA}} [DA, X \otimes DA] \\ (b) & \eta_X^{DT} \downarrow & \parallel \\ & [DT, X \otimes DT] \xrightarrow{\quad} & [DT, F(X \otimes DA)] \\ & [DT, \eta_{X \otimes DT}] & \end{array}$$

$$(c) \quad \begin{array}{ccc} [DT, Z \otimes DB] & \xrightarrow{[DT, g]} & [DT, Y] \\ \eta_{GF(Z)}^{DT} \uparrow & & \uparrow \varepsilon_{[DT, Y]} \\ G(Z) & & GF([DT, Y]) \\ G(\eta_{Z}^{DB}) \downarrow & & \parallel \\ G([DB, Z \otimes DB]) & \xrightarrow{G([DB, g])} & G([DB, Y]) \end{array}$$

PROOF. Routine verification.

LEMMA 4.5. *The map $e(X)$ is an R_n -morphism.*

PROOF. At first, we have to verify the equality

$$\begin{aligned} & \varepsilon_{GF(V(X_{i+1}))}^{DA} \cdot ([DA, \varepsilon_{V(X_{i+1})}] \otimes DA)^{-1} \cdot [DA, \alpha_{X_{i+1}} \cdot \phi_i] \otimes DA \cdot \eta_{X_i}^{DA} \otimes DA \\ & = (\varepsilon_{V(X_{i+1})})^{-1} \cdot \alpha_{X_{i+1}} \cdot \phi_i. \end{aligned}$$

By the naturality of ε^{DA} and the relation $\varepsilon_{X_i \otimes DA}^{DA} \cdot \eta_{X_i}^{DA} \otimes DA = 1_{X_i}$, we have $\alpha_{X_{i+1}} \cdot \phi_i = \varepsilon_{V(X_{i+1})}^{DA} \cdot [DA, \alpha_{X_{i+1}} \cdot \phi_i] \otimes DA \cdot \eta_{X_i}^{DA} \otimes DA$. Hence it is sufficient to show $(\varepsilon_{V(X_{i+1})})^{-1} \cdot \varepsilon_{GF(V(X_{i+1}))}^{DA} = \varepsilon_{GF(V(X_{i+1}))}^{DA} \cdot ([DA, \varepsilon_{V(X_{i+1})}] \otimes DA)^{-1}$, i.e., $\varepsilon_{V(X_{i+1})} \cdot \varepsilon_{GF(V(X_{i+1}))}^{DA} = \varepsilon_{GF(V(X_{i+1}))}^{DA} \cdot [DA, \varepsilon_{V(X_{i+1})}] \otimes DA$. But this follows again from the naturality of ε^{DA} .

The another necessary condition $([DA, \varepsilon_{V(X_{i+2})}])^{-1} \cdot [DA, \alpha_{X_{i+2}} \cdot \phi_{i+1}] \cdot \eta_{X_{i+1}}^{DA} \cdot \phi_i = 0$ is obvious.

LEMMA 4.6. *The map $f(X)$ is an R_n -morphism, as well.*

PROOF. We have to verify the equality

$$(\varepsilon_{P_0^i \otimes DA})^{-1} = \eta_{GF(P_0^i \otimes DA)}^{DA} \cdot [DT, \eta_{F(P_0^i \otimes DA)}] \otimes DA \cdot \eta_{P_0^i}^{DA}.$$

Since $\varepsilon_{P_0^i \otimes DA}^{DA} \cdot \eta_{P_0^i}^{DA} \otimes DA = 1_{P_0^i \otimes DA}$ and ε^{DA} is a natural transformation, we have

$$(\varepsilon_{P_0^i \otimes DA})^{-1} = \varepsilon_{GF(P_0^i \otimes DA)}^{DA} \cdot ([DA, \varepsilon_{P_0^i \otimes DA}] \otimes DA)^{-1} \cdot \eta_{P_0^i}^{DA} \otimes DA.$$

Hence we have the desired result since $([DA, \varepsilon_{P_0^i \otimes DA}])^{-1} = [DT, \eta_{F(P_0^i \otimes DA)}] \otimes DA$ by Lemma 4.4 (a).

For the proof of $p(X) \cdot (e(X), f(X)) = 0$, we note that the i -th structure map $\phi_i: \mathcal{S}_n(X)_i \otimes DB \rightarrow \mathcal{S}_n(X)_{i+1}$ satisfies $\phi_i \cdot y_i \otimes DB = 0$ and $y_{i+1} = \phi_i \cdot x_i \otimes DB$ and its adjoint $\bar{\phi}_i$ is the same with the composition: $[DB, \phi_i] \cdot \eta_{\mathcal{S}_n(X)_i}^{DB}: \mathcal{S}_n(X)_i \rightarrow [DB, \mathcal{S}_n(X)_i \otimes DB] \rightarrow [DB, \mathcal{S}_n(X)_{i+1}]$. Then it is easy to prove the following lemma, by definition.

LEMMA 4.7. *The following hold.*

$$(a) \quad G(\bar{\phi}_{i-1} \cdot x_{i-1}) = G([DB, y_i] \cdot \eta_{F(V(X_i))}^{DA})$$

$$(b) \quad G(\bar{\phi}_{i-1} \cdot \hat{y}_{i-1}) = 0$$

LEMMA 4.8. $p(X) \cdot e(X) = 0$.

PROOF. By Lemma 4.7 (a), it is sufficient to prove the commutativity of the following diagram :

$$\begin{array}{ccccc}
 & \eta_{X_i}^{DA} & [DA, \phi_i] & [DA, \alpha_{X_{i+1}}] & \\
 X_i & \xrightarrow{[DA, X_i \otimes DA]} & [DA, X_{i+1}] & \xrightarrow{[DA, \alpha_{X_{i+1}}]} & [DT, F(V(X_{i+1}))] \\
 \alpha_{X_i} \downarrow & & & & \downarrow [DT, x_i] \\
 V(X_i) & & & & [DT, S_n(X_i)] \\
 \epsilon_{V(X_i)} \uparrow \wr & & & & \uparrow \epsilon_{[DT, S_n(X)]_i} \\
 GF(V(X_i)) & \xrightarrow{G(\eta_{F(V(X_i))}^{DB})} & G([DB, F(V(X_i)) \otimes DB]) & \xrightarrow{G([DB, y_i])} & G([DB, S_n(X)_i]).
 \end{array}$$

We know $\eta_{X_i}^{DA} = [DT, \eta_{X_i \otimes DT} \cdot \eta_{X_i}^{DT} : X_i \rightarrow [DT, X_i \otimes DT] \rightarrow [DT, F(X_i \otimes DA)] = [DA, X_i \otimes DA]$ by Lemma 4.4 (b) and $\epsilon_{[DT, S_n(X)]_i} \cdot G([DB, y_i] \cdot \eta_{F(V(X_i))}^{DB}) = [DT, y_i] \cdot \eta_{GF(V(X_i))}^{DT}$ by Lemma 4.4 (c). Further, by the definition of the map θ , it holds that

$$y_i \cdot G(\epsilon_{V(X_i)})^{-1} \cdot \alpha_{X_i} \otimes DT = x_i \cdot F(\alpha_{X_{i+1}} \cdot \phi_i) \cdot \eta_{X_i \otimes DT}.$$

Hence we have

$$\begin{aligned}
 & [DT, x_i] \cdot [DA, \alpha_{X_{i+1}} \cdot \phi_i] \cdot \eta_{X_i}^{DA} \\
 &= [DT, x_i \cdot F(\alpha_{X_{i+1}} \cdot \phi_i)] \cdot \eta_{X_i}^{DA} \\
 &= [DT, x_i \cdot F(\alpha_{X_{i+1}} \cdot \phi_i) \cdot \eta_{X_i \otimes DT} \cdot \eta_{X_i}^{DT}] \cdot \eta_{X_i}^{DT} \\
 &= [DT, y_i \cdot (\epsilon_{V(X_i)} \otimes DT)^{-1} \cdot \alpha_{X_i} \otimes DT] \cdot \eta_{X_i}^{DT} \\
 &= [DT, y_i \cdot (\epsilon_{V(X_i)} \otimes DT)^{-1}] \cdot \eta_{V(X_i)}^{DT} \cdot \alpha_{X_i} \\
 &= [DT, y_i] \cdot \eta_{GF(V(X_i))}^{DT} \cdot (\epsilon_{V(X_i)})^{-1} \cdot \alpha_{X_i} \\
 &= \epsilon_{[DT, S_n(X)]_i} \cdot G([DB, y_i] \cdot \eta_{F(V(X_i))}^{DB}) \cdot (\epsilon_{V(X_i)})^{-1} \cdot \alpha_{X_i}.
 \end{aligned}$$

LEMMA 4.9. $p(X) \cdot f(X) = 0$.

PROOF. By Lemma 4.6 (a), it is enough to prove the commutativity of the diagram :

$$\begin{array}{ccc}
 P_0^i \xrightarrow{\eta_{P_0^i}^{DA}} [DA, P_0^i \otimes DA] = [DT, F(P_0^i \otimes DA)] \xrightarrow{[DT, \hat{y}_i]} [DT, \mathcal{S}_n(X)_i] \\
 \begin{array}{c} p_0^i \downarrow \\ V(X_i) \\ \varepsilon_{V(X_i)} \uparrow \end{array} & & \begin{array}{c} \uparrow \varepsilon_{[DT, \mathcal{S}_n(X)_i]} \\ G([DB, \mathcal{S}_n(X)_i]) \end{array} \\
 GF(V(X_i)) \xrightarrow{G(\eta_{F(V(X_i))}^{DB})} G([DB, F(V(X_i)) \otimes DB]) \xrightarrow{G([DB, y_i])} G([DB, \mathcal{S}_n(X)_i])
 \end{array}$$

By Lemma 4.4 (b) and (c), we know $\eta_{P_0^i}^{DA} = [DT, \eta_{P_0^i \otimes DT}^{\#}] \cdot \eta_{P_0^i}^{DT}$ and $\eta_{GF(V(X_i))}^{DB} = \varepsilon_{[DT, F(V(X_i)) \otimes DB]} \cdot G(\eta_{F(V(X_i))}^{DB})$. Hence we have

$$\begin{aligned}
 [DT, \hat{y}_i] \cdot \eta_{P_0^i}^{DA} &= [DT, \hat{y}_i \cdot \eta_{P_0^i \otimes DT}^{\#}] \cdot \eta_{P_0^i}^{DT} \\
 &= [DT, y_i \cdot (\varepsilon_{V(X_i)} \otimes DT)^{-1} \cdot p_0^i \otimes DT] \cdot \eta_{P_0^i}^{DT} \\
 &= [DT, y_i \cdot (\varepsilon_{V(X_i)} \otimes DT)^{-1}] \cdot \eta_{V(X_i)}^{DT} \cdot p_0^i \\
 &= [DT, y_i] \cdot \eta_{GF(V(X_i))}^{DB} \cdot (\varepsilon_{V(X_i)})^{-1} \cdot p_0^i \\
 &= [DT, y_i] \cdot \varepsilon_{[DT, F(V(X_i)) \otimes DB]} \cdot G(\eta_{F(V(X_i))}^{DB}) \cdot (\varepsilon_{V(X_i)})^{-1} \cdot p_0^i \\
 &= \varepsilon_{[DT, \mathcal{S}_n(X)_i]} \cdot GF([DT, y_i]) \cdot G(\eta_{F(V(X_i))}^{DB}) \cdot (\varepsilon_{V(X_i)})^{-1} \cdot p_0^i \\
 &= \varepsilon_{[DT, \mathcal{S}_n(X)_i]} \cdot G([DB, y_i] \cdot \eta_{F(V(X_i))}^{DB}) \cdot (\varepsilon_{V(X_i)})^{-1} \cdot p_0^i.
 \end{aligned}$$


Since $(e(X), f(X))$ is obviously an R_n -monomorphism, we have proved $Q_n \mathcal{S}_n(X) \cong X \oplus P$ as R_n -modules. It is easy to prove that the monomorphism $e(X): X \rightarrow \mathcal{D}\mathcal{S}_n(X)$ has naturality on X . By the duality, we can prove the similar result on $\mathcal{S}_n Q_n$.

Thus we have

THEOREM 4.10. $Q_n \mathcal{S}_n \cong \underline{1}_{\text{mod-}R_n}$ and $\mathcal{S}_n Q_n \cong \underline{1}_{\text{mod-}S_n}$, i.e., the stable categories $\text{mod-}R_n$ and $\text{mod-}S_n$ are always equivalent.

REMARK. D. Happel [15] has proved that $\text{mod-}\hat{A}$ and $\text{mod-}\hat{B}$ are equivalent if $\text{gl. dim. } A < \infty$. And, in the case where $\Phi = \Phi_1: \text{mod-}\hat{A} \rightarrow \text{mod-}R$ is dense, the stable functor \mathcal{S}_n is induced from \mathcal{S}_∞ for each $n \neq \infty$. But, in general, Φ is not dense even if $\text{gl. dim. } A < \infty$, can not be induced from \mathcal{S}_∞ .

EXAMPLE 4.11. Let A be the bound quiver algebra of



Then $\text{soc}(e_4A) \cong \text{top}(e_4A \otimes DA)$ and there is a non-zero map f from $e_4A \otimes DA$ to e_4A such that $\text{Im}(f) = \text{soc}(e_4A)$ and $\text{Ker}(f) = \text{rad}(e_4A \otimes DA)$. Hence we can define an indecomposable R -module $X = (e_4A, f)$. As is easily seen, for any R -module X' , X can not be isomorphic to $\Phi(X')$. Thus $\Phi: \underline{\text{mod-}}\hat{A} \rightarrow \underline{\text{mod-}}R$ is not dense, even though $\text{g.l dim. } A = 2 < \infty$.

References

- [1] Assem, I., Iterated tilted algebras of type B_n and C_n , preprint.
- [2] Assem, I. and Happel, D., Generalized tilted algebras of type A_n , *Comm. Algebra* **9** (1981), 2101-2125.
- [3] Assem, I., Happel, D. and Roldan, O., Representation finite trivial extension algebras, preprint.
- [4] Assem, I. and Iwanaga, Y., Stable equivalence of representation finite trivial extension, preprint.
- [5] Auslander, M., Platzeck, M.I. and Reiten, I., Coxeter functors without diagrams, *Trans. Amer. Math. Soc.* **250** (1979), 1-46.
- [6] Auslander, M. and Reiten, I., Representation theory of artin algebras III, *Comm. Algebra* **5** (1975), 239-294.
- [7] Auslander, M. and Reiten, I., Representation theory of artin algebras V, *Comm. Algebra* **5** (1977), 519-554.
- [8] Bernstein, I., Gelfand, I.M. and Ponomarev, V.A., Coxeter functor and Gabriel's theorem, *Russ. Math. Surveys* **28** (1977), 17-32.
- [9] Bongartz, K., Tilted algebras, *Springer LNM* **903** (1981), 16-38.
- [10] Brenner, S. and Butler, M.C.R., Generalization of Bernstein-Gelfand-Ponomarev reflection functors, *Springer LNM* **832** (1980), 103-169.
- [11] Dlab, V. and Ringel, C.M., Indecomposable representations of graphs and algebras, *Memoirs Amer. Math. Soc.* **173** (1976).
- [12] Fossum, R.M., Griffith, P.A. and Reiten, I., Trivial extensions of abelian categories, *Springer LNM* **456** (1975).
- [13] Gabriel, P. and Riedtmann, C., Group representations without groups, *Comment. Math. Helv.* **54** (1979), 240-287.
- [14] Gabriel, P., Auslander-Reiten sequences and representation finite algebras, *Springer LNM* **831** (1980), 1-71.
- [15] Happel, D., Triangulated categories and trivial extension algebras, to appear in the proceedings of Carleton Univ., 1984.
- [16] Happel, D. and Ringel, C.M., Tilted algebras, *Trans. Amer. Math. Soc.* **247** (1982), 399-443.
- [17] Hoshino, M., Trivial extensions of tilted algebras, *Comm. Algebra* **10** (1982), 1965-1999.
- [18] Hughes, D. and Waschbüsch, J., Trivial extensions of tilted algebras, *Proc. London Math. Soc.* **46** (1983), 427-440.
- [19] Riedtmann, C., Algebren, Darstellungsköcher, Überlagerungen und zurück, *Comment. Math. Helv.* **55** (1980), 199-224.
- [20] Tachikawa, H., Representations of trivial extensions of hereditary algebras, *Springer LNM* **832** (1980), 572-599.
- [21] Tachikawa, H., Reflection functors and Auslander-Reiten translations for trivial extensions of hereditary algebras, *J. Algebra.* **90** (1984), 98-118.
- [22] Tachikawa, H., Self-injective algebras and tilting theory, to appear in the proceed-

- ings of Carleton Univ., 1984.
- [23] Tachikawa H. and Wakamatsu, T., Extensions of tilting functors and QF-3 algebras, to appear in J. Algebra.
 - [24] Tachikawa, H. and Wakamatsu, T., Tilting functors and stable equivalences for self-injective algebras, preprint.
 - [25] Tachikawa, H. and Wakamatsu, T., Applications of reflection functors for self-injective algebras, to appear in the proceedings of Carleton Univ., 1984.
 - [26] Wakamatsu, T., Note on trivial extensions of artin algebras, Comm. Algebra 12 (1984), 33-41.
 - [27] Wakamatsu, T., Partial Coxeter functors and stable equivalences for self-injective algebras, Tsukuba J. Math. 9 (1985), 171-183.
 - [28] Waschbüsch, J., Universal coverings of self-injective algebras, Springer LNM 903 (1981), 331-349.
 - [29] Yamagata, K., On algebras whose trivial extensions are of finite representation type, Springer LNM 903 (1981), 364-371.

Institute of Mathematics,
University of Tsukuba
Sakura-mura, Ibaraki,
305, Japan