

THE LIMITING ABSORPTION PRINCIPLE FOR THE ACOUSTIC WAVE OPERATORS IN TWO UNBOUNDED MEDIA

Dedicated to Professor Mutsuhide Matsumura on his sixtieth birthday

By

Mitsuteru KADOWAKI

1. Introduction.

In the present paper we study the limiting absorption principle for the acoustic wave operators in two unbounded media. We assume that the propagation speed is discontinuous at the interface and the equilibrium density is 1.

Let $n \geq 2$ and $x = (y, z) \in \mathbf{R}^{n-1} \times \mathbf{R}$. The following equation describes the wave propagation here:

$$(1.1) \quad \partial_t^2 u(t, x) - a(x)^2 \Delta u(t, x) = 0, \quad (t, x) \in \mathbf{R} \times \mathbf{R}^n,$$

where $a(x)$ is a propagation speed.

We make the assumptions for the interface separating two media and $a(x)$.

Let $\varphi_0(y) = a|y|$ and $\varphi(y) \in C^1(\mathbf{R}^{n-1} \setminus \{0\})$, where $a \geq 0$. We assume that $\varphi(y)$ describes the interface and satisfies

$$(A.0) \quad \sum_{|\alpha| \leq 1} |y|^{|\alpha|} |\partial^\alpha (\varphi(y) - \varphi_0(y))| = O(|y|^{-\theta}) \quad (|y| \rightarrow \infty),$$

for some $\theta > 0$, and

$$(A.1) \quad \sum_{|\alpha| \leq 1} |y|^{|\alpha|} |\partial^\alpha \varphi(y)| = O(|y|^{-\sigma}) \quad (|y| \rightarrow 0).$$

where $0 < \sigma < 1/2$. For $\varphi(y)$, we use the following notation:

$$\Omega_+ = \{x = (y, z) : z > \varphi(y)\},$$

$$\Omega_- = \{x = (y, z) : z < \varphi(y)\},$$

$$S = \{x = (y, z) : z = \varphi(y)\}.$$

We denote the unit normal vector at the point $x \in S$ by $\nu = (\nu_1, \nu_2, \dots, \nu_z)$ with $\nu_z > 0$.

The propagation speed $a(x) > 0$ is assumed to satisfy the following: for

some $c > 1$,

$$(A.2) \quad 1/c < a(x) < c$$

and there exist $a_{\pm} > 0$, $a_{\pm}^{\sharp}(x) \in B^1(\Omega_{\pm})$ and $a_S(x) \in L^{\infty}(\mathbf{R}^n)$ such that $a(x)$ is decomposed as

$$(A.3) \quad \begin{cases} a(x) = a_{\pm} + a_{\pm}^{\sharp}(x) + a_S(x) \quad (x \in \Omega_{\pm}), \\ \sum_{|\alpha| \leq 1} |x|^{|\alpha|} |\partial^{\alpha} a_{\pm}^{\sharp}(x)| = O(|x|^{-\theta}) \quad (|x| \rightarrow \infty, x \in \Omega_{\pm}), \\ a_S(x) = O(|x|^{-\theta-1}) \quad (|x| \rightarrow \infty) \end{cases}$$

for some $\theta > 0$.

Under (A.0)~(A.3), we show the nonexistence of eigenvalues and the limiting absorption principle of the acoustic operator $-a(x)^2 \Delta$ for (1.1).

There are many works dealing with the acoustic wave propagation problem with the discontinuous propagation speed at the interface separating media. Eidus [5] proved the limiting absorption and amplitude principle for two unbounded media problem with the interface satisfying the following conditions: for any $x \in S$

$$(1.2) \quad \nu_z \geq C_1 > 0,$$

$$(1.3) \quad |x \cdot \nu| \leq C_2,$$

where $C_j > 0$, ($j=1, 2$), are independent of $x \in S$. For example,

$$\varphi(y) \in C^1(\mathbf{R}^{n-1}), \quad \varphi(y) = \frac{\sin |y|}{|y|} \quad (|y| \gg 1), \quad \varphi_0(y) = 0$$

satisfies (1.2) and (1.3), but not satisfies (A.0). We can also deal with the following interface not satisfying (1.2),

$$\varphi(y) = |y|^{-\sigma}, \quad \varphi_0(y) = 0$$

where, $0 < \sigma < 1/2$.

The propagation speed considered in Eidus [5] is a piecewise constant function while we can perturb the propagation speed. Wilcox [17] considered two stratified fluids in a half space and established the eigenfunction expansion theorem. Ben-Artzi [1], Weder [14]~[16] and Dermenjian and Guillot [3] considered perturbed stratified fluids problems. They showed the limiting absorption principle by the approach of Wilcox [17]. Kikuchi and Tamura [8] also proved the limiting amplitude principle for perturbed stratified fluids. On the other hand there are some works dealing with the case where the equilibrium density is discontinuous at the interface separating media, for example,

Debièvre and Pravica [2] and Wilcox [17].

In order to show the limiting absorption principle for our operator, we use Mourre's method. This method was first developed by Mourre [9] to prove the limiting absorption principle for 3-body Schrödinger operators (see also Perry, Sigal and Simon [10] and Tamura [12]). In Froese and Herbst [6], they showed by Mourre's method that N-body Schrödinger operators have no positive eigenvalues. Iwashita [7] and Weder [13] showed the limiting absorption principle for first order symmetric systems. For the acoustic wave operators in perturbed stratified fluids, Debièvre and Pravica [2] obtained the similar results as in [7] and [13]. Tamura [11] used Mourre's method in order to prove the limiting amplitude principle for the acoustic wave operators (see also Kikuchi and Tamura [8]).

We now define the acoustic operator L as

$$(1.4) \quad L = -a(x)^2 \Delta$$

Under the above assumptions, (A.0)~(A.3), L is a symmetric operator in the Hilbert space $L^2(\mathbf{R}^n; E(x)dx)$ with $E(x) = a(x)^{-2}$ and admits a unique self-adjoint realization. We denote by the same notation L this self-adjoint realization. Then L is a positive operator (zero is not an eigenvalue) and the domain $D(L)$ is given by $D(L) = H^s(\mathbf{R}_x^n)$, $H^s(\mathbf{R}_x^n)$ being the Sobolev space of order s over \mathbf{R}_x^n . We also denote by $R(z; L)$ the resolvent $(L - z)^{-1}$ of L for $\text{Im}z \neq 0$.

We need several notations to describe our results. Let L^2 be the usual L^2 space defined on \mathbf{R}^n , with the inner product

$$\langle u, v \rangle = \int u(x) \overline{v(x)} dx$$

and the corresponding norm $|\cdot|_0$. For $\alpha \in \mathbf{R}$ let L_α^2 be the weighted L^2 space defined by

$$L_\alpha^2 = \{u(x) : \langle x \rangle^\alpha u(x) \in L^2(\mathbf{R}_x^n)\}, \quad \langle x \rangle = (1 + |x|^2)^{1/2},$$

with the norm

$$|u|_\alpha^2 = \int \langle x \rangle^{2\alpha} |u(x)|^2 dx.$$

Let $A: L^2 \rightarrow L^2$ be a bounded operator. We denote by $\|A\|$ the operator norm of A .

The main result is

THEOREM 1.1. *Assume that (A.0)~(A.3). Then*

(i) L has no eigenvalues.

(ii) Let $\lambda_0 > 0$ and $\alpha > 1/2$. Then for any compact interval $I \subset \mathbf{R}_+$ containing λ_0 , there exists a positive constant $C = C(I, \alpha)$ such that

$$\| \langle x \rangle^{-\alpha} R(\lambda \pm i\kappa; L) \langle x \rangle^{-\alpha} \| \leq C,$$

for $\lambda \in I, 0 < \kappa < 1$.

(iii) For every $\lambda > 0$ and $\alpha > 1/2$, following two limits

$$R(\lambda \pm i0; L) = \lim_{\kappa \downarrow 0} R(\lambda \pm i\kappa; L),$$

exist in the uniform operator topology of $\mathfrak{B}(L_\alpha^2, L_{-\alpha}^2)$. Moreover $R(\lambda \pm i0; L)$ are locally Hölder continuous.

Acknowledgement. The author would like to express his sincere gratitude to Professors M. Matsumura and K. Mochizuki for their generous advice and kind encouragement.

2. Mourre's estimate.

In this section we shall show Mourre's estimate (Lemma 2.4). First, we prove the following lemma. We need this lemma to estimate an integral on the interface S .

LEMMA 2.2. Let $s > 1/2$. For $u \in \mathcal{S}(\mathbf{R}^n)$ (Schwartz space), we define

$$(T_\varphi u)(y) = u(y, \varphi(y)).$$

T_φ has an extension to a bounded operator from $H^s(\mathbf{R}^n)$ to $L^2(\mathbf{R}^{n-1})$.

PROOF. We show that:

$$(2.1) \quad \|T_\varphi u\|_{L^2(\mathbf{R}^{n-1})} \leq C \|u\|_{H^s(\mathbf{R}^n)}$$

for $u \in \mathcal{S}(\mathbf{R}^n)$. Let $u \in \mathcal{S}(\mathbf{R}^n)$. Then $u(y, \varphi(y))$ is represented as

$$(2.2) \quad u(y, \varphi(y)) = (2\pi)^{-1/2} \int_{-\infty}^{+\infty} e^{i\varphi(y) \cdot \zeta} (\mathfrak{F}_z u)(y, \zeta) d\zeta,$$

where $\xi = (\eta, \zeta) \in \mathbf{R}_\eta^{n-1} \times \mathbf{R}_\zeta = \mathbf{R}_\xi^n$ and \mathfrak{F}_z is the Fourier transform in \mathbf{R}_z . By Schwartz's inequality, we have

$$(2.3) \quad |(T_\varphi u)(y)|^2 \leq (2\pi)^{-1} \int_{-\infty}^{+\infty} (1 + \zeta^2)^{-s} d\zeta \int_{-\infty}^{+\infty} (1 + \zeta^2)^{s/2} |(\mathfrak{F}_z u)(y, \zeta)|^2 d\zeta.$$

By integrating both sides of the above inequality over \mathbf{R}^{n-1} , we obtain (2.1). ■

We consider only the case $1 = a_-^{-2} < a_+^{-2}$. The other cases can be proved similarly.

We define the self-adjoint operator $H(\lambda)$ on L^2 by

$$H(\lambda) = -\Delta - \lambda(E(x) - 1),$$

$$D(H(\lambda)) = H^2(\mathbf{R}^n).$$

Then we have

$$(L - (\lambda \pm i\kappa))^{-1} = (H(\lambda) - \lambda \mp i\kappa E(x))^{-1} E(x).$$

We apply Mourre's commutator method to $H(\lambda)$ on L^2 (see Tamura [11] or Kikuchi and Tamura [8]).

By (A.3), we can decompose $E(x) = E_L^+(x) + E_S(x)$ ($x \in \Omega_\pm$) in such a way that

$$\sum_{|\alpha| \leq 1} |x|^{|\alpha|} |\partial^\alpha (E_L^+(x) - a_\pm^2)| = O(|x|^{-\theta}), \quad (|x| \rightarrow \infty, x \in \Omega_\pm),$$

$$E_S(x) = O(|x|^{-1-\theta}), \quad (|x| \rightarrow \infty).$$

Let A be the generator of the dilation unitary group;

$$A = \frac{1}{2i}(x \cdot \nabla + \nabla \cdot x).$$

We define the commutator $i[H(\lambda), A]$ as a form on $H^2(\mathbf{R}^n) \cap D(A)$ as follows; For $u, v \in H^2(\mathbf{R}^n) \cap D(A)$

$$\begin{aligned} & \langle i[H(\lambda), A]u, v \rangle \\ &= i(\langle Au, H(\lambda)v \rangle - \langle H(\lambda)u, Av \rangle). \end{aligned}$$

LEMMA 2.2. *The form $i[H(\lambda), A]$ defined on $H^2(\mathbf{R}^n) \cap D(A)$ is extended to a bounded operator from $H^1(\mathbf{R}^n)$ to $H^{-1}(\mathbf{R}^n)$ which is denoted by $i[H(\lambda), A]^0$.*

In order to prove Lemma 2.2, we need the following lemma.

LEMMA 2.3 (Sobolev's Theorem). *Suppose that*

$$1/2 - l/m = 1/q, \quad 2 < q < \infty.$$

Then we have the embedding

$$H^l(\mathbf{R}^m) \hookrightarrow L^q(\mathbf{R}^m).$$

PROOF OF LEMMA 2.2. Let $u \in \mathbf{S}(\mathbf{R}^n)$. Then we have

$$(2.4) \quad \langle i[-\Delta, A]u, u \rangle = 2\langle \nabla u, \nabla u \rangle,$$

$$\begin{aligned} (2.5) \quad & \langle i[E-1, A]u, u \rangle \\ &= \langle x \cdot \nabla u, E_L u \rangle + \langle E_L u, x \cdot \nabla u \rangle + n \langle E_L u, u \rangle, \\ & \quad + \langle x \cdot \nabla u, E_S u \rangle + \langle E_S u, x \cdot \nabla u \rangle + n \langle E_S u, u \rangle, \end{aligned}$$

where $E_L = E_L^\pm(x)(x \in \Omega_\pm)$.

We calculate the first and the second terms of right side of (2.5). We set $w_n = z - \varphi(y)$ and $w_j = y_j (j=1, 2, \dots, n-1)$. Then we have $D(y, z)/D(w', w_n) = 1$, where $w' = (w_1, w_2, \dots, w_{n-1})$. Integrating by parts, we have

$$\begin{aligned}
(2.6) \quad & \langle x \cdot \nabla u, E_L^\pm u \rangle_{L^2(\Omega_\pm)} + \langle E_L^\pm u, x \cdot \nabla u \rangle_{L^2(\Omega_\pm)} \\
& = \langle w \cdot \nabla_w \tilde{u}, \tilde{E}_L^\pm \tilde{u} \rangle_{L^2(\mathbb{R}_\pm^n)} + \langle \tilde{E}_L^\pm \tilde{u}, w \cdot \nabla_w \tilde{u} \rangle_{L^2(\mathbb{R}_\pm^n)} \\
& \quad - \langle (w' \cdot \nabla_{w'} \varphi(w') - \varphi(w')) \partial_{w_n} \tilde{u}, \tilde{E}_L^\pm \tilde{u} \rangle_{L^2(\mathbb{R}_\pm^n)} \\
& \quad - \langle \tilde{E}_L^\pm \tilde{u}, (w' \cdot \nabla_{w'} \varphi(w') - \varphi(w')) \partial_{w_n} \tilde{u} \rangle_{L^2(\mathbb{R}_\pm^n)} \\
& = - \langle (w \cdot \nabla_w \tilde{E}_L^\pm) \tilde{u}, \tilde{u} \rangle_{L^2(\mathbb{R}_\pm^n)} - n \langle \tilde{E}_L^\pm \tilde{u}, \tilde{u} \rangle_{L^2(\mathbb{R}_\pm^n)} \\
& \quad + \langle (w' \cdot \nabla_{w'} \varphi(w') - \varphi(w')) (\partial_{w_n} \tilde{E}_L^\pm) \tilde{u}, \tilde{u} \rangle_{L^2(\mathbb{R}_\pm^n)} \\
& \quad \pm \int_{\mathbb{R}^{n-1}} (w' \cdot \nabla_{w'} \varphi(w') - \varphi(w')) \tilde{E}_L^\pm(w', 0) |\tilde{u}(w', 0)|^2 dw' \\
& = - \langle (x \cdot \nabla E_L^\pm) u, u \rangle_{L^2(\Omega_\pm)} - n \langle E_L^\pm u, u \rangle_{L^2(\Omega_\pm)} \\
& \quad \pm \int_{\mathbb{R}^{n-1}} (y \cdot \nabla_y \varphi(y) - \varphi(y)) E_L^\pm(y, \varphi(y)) |u(y, \varphi(y))|^2 dy,
\end{aligned}$$

where $\tilde{E}_L^\pm = E_L^\pm(w', w_n + \varphi(w'))$ and $\tilde{u} = u(w', w_n + \varphi(w'))$. (2.6) implies

$$\begin{aligned}
(2.7) \quad & \langle x \cdot \nabla u, E_L u \rangle + \langle E_L u, x \cdot \nabla u \rangle + n \langle E_L u, u \rangle \\
& = - \langle (x \cdot \nabla E_L) u, u \rangle + \int_{\mathbb{R}^{n-1}} (y \cdot \nabla_y \varphi(y) - \varphi(y)) (E_L^{+0} - E_L^{-0}) |u(y, \varphi(y))|^2 dy,
\end{aligned}$$

where $E_L^{\pm 0} = E_L^\pm(y, \varphi(y))$ and $(x \cdot \nabla E_L) = x \cdot \nabla E_L^\pm(x \in \Omega_\pm)$. By (2.4), (2.5) and (2.7), we have

$$\begin{aligned}
(2.8) \quad & \langle i[H(\lambda), A]u, u \rangle = 2 \langle \nabla u, \nabla u \rangle + \lambda \langle (x \cdot \nabla E_L) u, u \rangle \\
& \quad - \lambda \int_{\mathbb{R}^{n-1}} (y \cdot \nabla_y \varphi(y) - \varphi(y)) (E_L^{+0} - E_L^{-0}) |u(y, \varphi(y))|^2 dy \\
& \quad - n \lambda \langle E_S u, u \rangle - \lambda \langle E_S u, x \cdot \nabla u \rangle - \lambda \langle x \cdot \nabla u, E_S u \rangle.
\end{aligned}$$

For any $r > 0$, we have

$$\begin{aligned}
(2.9) \quad & \lambda \int_{\mathbb{R}^{n-1}} (y \cdot \nabla_y \varphi - \varphi) (E_L^{+0} - E_L^{-0}) |u(y, \varphi(y))|^2 dy \\
& = \lambda \langle T_\varphi^*(y \cdot \nabla_y \varphi - \varphi) (E_L^{+0} - E_L^{-0}) T_\varphi u, u \rangle \\
& = \lambda \langle T_\varphi^*(\chi_{|y| < r} (y \cdot \nabla_y \varphi - \varphi) (E_L^{+0} - E_L^{-0})) T_\varphi u, u \rangle \\
& \quad + \langle T_\varphi^*(\chi_{|y| > r} (y \cdot \nabla_y \varphi - \varphi) (E_L^{+0} - E_L^{-0})) T_\varphi u, u \rangle
\end{aligned}$$

We define an operator R_φ^r as

$$R_\varphi^r u = \chi_{|y| < r}(y)(y \cdot \nabla_y \varphi(y) - \varphi(y))^{1/2} T_\varphi u.$$

We show that R_φ^r is a bounded operator from $H^1(\mathbf{R}^n)$ to $L^2(\mathbf{R}^{n-1})$.

If $0 < \sigma < 1/2$, there exist some $s > 1/2$ and $p > n-1$ such that

$$(2.10) \quad n-1-\sigma p > 0,$$

$$(2.11) \quad \frac{n-1}{2p} + s = 1.$$

By (2.2), (2.3) and Hölder inequality, we have

$$(2.12) \quad \begin{aligned} \|R_\varphi^r u\|_{L^2(\mathbf{R}^{n-1})}^2 &\leq C \int_{-\infty}^{+\infty} (1+\zeta^2)^s \int_{|y| < r} |y \cdot \nabla_y \varphi - \varphi| |(\mathfrak{F}_z u)(y, \zeta)|^2 dy d\zeta \\ &\leq C r^{(n-1-\sigma p)/p} \int_{-\infty}^{+\infty} (1+\zeta^2)^s |(\mathfrak{F}_z u)(\cdot, \zeta)|_{L^{2p/(p-1)}(\mathbf{R}^{n-1})}^2 d\zeta, \end{aligned}$$

where $C > 0$ is independent of r .

Lemma 2.3 implies that

$$(2.13) \quad \|\mathfrak{F}_z u(\cdot, \zeta)\|_{L^{2p/(p-1)}(\mathbf{R}^{n-1})} \leq C \|\mathfrak{F}_z u(\cdot, \zeta)\|_{H^{(n-1)/2p}(\mathbf{R}^{n-1})}.$$

It follows from (2.11), (2.12) and (2.13) that

$$(2.14) \quad \|R_\varphi^r u\|_{L^2(\mathbf{R}^{n-1})} \leq C r^{(n-1-\sigma p)/2p} \|u\|_{H^1(\mathbf{R}^n)},$$

where C is independent of r .

It follows from Lemma 2.1 that T_φ^* is a bounded operator from $L^2(\mathbf{R}^{n-1})$ to $H^{-1}(\mathbf{R}^n)$. Moreover, since $E_S x \cdot \nabla$ is a bounded operator from $H^1(\mathbf{R}^n)$ to $L^2(\mathbf{R}^n)$, the adjoint operator $(\nabla)^* \cdot x E_S$ is a bounded operator from $L^2(\mathbf{R}^n)$ to $H^{-1}(\mathbf{R}^n)$.

Hence, from (2.8), (2.9) and (2.13), the form $i[H(\lambda), A]$ on $\mathbf{S}(\mathbf{R}^n)$ has an extension to a bounded operator $i[H(\lambda), A]^0$ on from $H(\mathbf{R}^n)$ to $H^{-1}(\mathbf{R}^n)$.

Let $u \in H^2(\mathbf{R}^n) \cap D(A)$. Repeating the argument in the proof of Lemma 2.1 of Weder [13], we can show that u satisfies (2.8). ■

LEMMA 2.4. *Let $\lambda_0 > 0$ and $0 < \delta < \min(1, \lambda_0/4)$ and take $f_\delta(p) \in C_0^\infty(\mathbf{R})$, $0 \leq f_\delta \leq 1$ such that f_δ has support in $(\lambda_0 - 3\delta, \lambda_0 + 3\delta)$ and $f_\delta = 1$ on $[\lambda_0 - 2\delta, \lambda_0 + 2\delta]$. Then, there exist a positive constant α and a compact operator K on L^2 which depend on only λ_0 such that*

$$(2.15) \quad \begin{aligned} f_\delta(H(\lambda)) i[H(\lambda), A]^0 f_\delta(H(\lambda)) \\ \geq \alpha f_\delta(H(\lambda))^2 + f_\delta(H(\lambda)) K(\lambda) f_\delta(H(\lambda)) \end{aligned}$$

for $\lambda \in (\lambda_0 - \delta, \lambda_0 + \delta)$, in the form sense.

PROOF. We simply write f_δ instead of $f_\delta(H(\lambda))$. By Lemma 2.2, we have for $u \in L^2$,

$$\begin{aligned}
 (2.16) \quad & \langle f_{\delta i}[H(\lambda), A^0]^0 f_{\delta u}, u \rangle \\
 & = 2\langle -\Delta f_{\delta u}, f_{\delta u} \rangle + \lambda \langle ((x \cdot \nabla E_L) - n E_S) f_{\delta u}, f_{\delta u} \rangle \\
 & \quad - \langle x \cdot \nabla f_{\delta u}, E_S f_{\delta u} \rangle - \langle E_S f_{\delta u}, x \cdot \nabla f_{\delta u} \rangle \\
 & \quad - \langle (T_\varphi)^* \mathcal{X}_{|y| > r} (y \cdot \nabla_y \varphi - \varphi) (E_L^{+0} - E_L^{-0}) T_\varphi f_{\delta u}, f_{\delta u} \rangle \\
 & \quad - \langle (R_\varphi^r)^* (E_L^+ - E_L^-) R_\varphi^r f_{\delta u}, f_{\delta u} \rangle.
 \end{aligned}$$

Let $0 < r \ll 1$. By (2.13), there exists a positive number C independent of r such that,

$$\begin{aligned}
 & |\lambda \langle (R_\varphi^r)^* (E_L^+ - E_L^-) R_\varphi^r f_{\delta u}, f_{\delta u} \rangle| \\
 & = |\lambda \langle (E_L^{+0} - E_L^{-0}) R_\varphi^r f_{\delta u}, R_\varphi^r f_{\delta u} \rangle_{L^2(\mathbb{R}^{n-1})}| \\
 & \leq C r^{(n-1-\sigma p)/p} (|\nabla f_{\delta u}|_0^2 + |f_{\delta u}|_0^2).
 \end{aligned}$$

Let $E_0(x) = a_\pm^{-2}(x \in \Omega_\pm)$. Then we have

$$\begin{aligned}
 (2.17) \quad & 2\langle -\Delta f_{\delta u}, f_{\delta u} \rangle - \lambda \langle (R_\varphi^r)^* (E_L^+ - E_L^-) R_\varphi^r f_{\delta u}, f_{\delta u} \rangle \\
 & \geq (2 - C r^{(n-1-\sigma p)/p}) \langle H(\lambda) f_{\delta u}, f_{\delta u} \rangle - C r^{(n-1-\sigma p)/p} |f_{\delta u}|_0^2 \\
 & \quad + (2 - C r^{(n-1-\sigma p)/p}) \lambda \langle (E - E_0) f_{\delta u}, f_{\delta u} \rangle \\
 & \quad + (2 - C r^{(n-1-\sigma p)/p}) \lambda \langle (E_0 - 1) f_{\delta u}, f_{\delta u} \rangle \\
 & \geq \left(\frac{(2 - C r^{(n-1-\sigma p)/p}) \lambda_0}{4} C r^{(n-1-\sigma p)/p} \right) |f_{\delta u}|_0^2 \\
 & \quad + (2 - C r^{(n-1-\sigma p)/p}) \lambda \langle (E - E_0) f_{\delta u}, f_{\delta u} \rangle.
 \end{aligned}$$

Take $h \in C_0^\infty(\mathbb{R})$, $0 \leq h \leq 1$ such that $h=1$ on $(\lambda_0/4, 7\lambda_0/4)$. Using h , we define an operator $K(\lambda)$ as

$$K(\lambda) = \sum_{j=1}^4 K_j(\lambda)$$

where,

$$K_1(\lambda) = \lambda(h(H(\lambda))((x \cdot \nabla E_L) - n E_S + (2 - C r^{(n-1-\sigma p)/p})(E - E_0))h(H(\lambda)),$$

$$K_2(\lambda) = -\lambda h(H(\lambda)) E_S x \cdot \nabla h(H(\lambda)),$$

$$K_3(\lambda) = -\lambda h(H(\lambda)) \nabla^* \cdot x E_S h(H(\lambda)),$$

$$K_4(\lambda) = -\lambda h(H(\lambda)) T_\varphi^* (E_L^{+0} - E_L^{-0}) \mathcal{X}_{|y| > r} (y \cdot \nabla_y \varphi - \varphi) T_\varphi h(H(\lambda)).$$

For each $\lambda \in (\lambda_0 - \delta, \lambda_0 + \delta)$, we show that $K_j(\lambda)$ ($j = 1, 2, 3, 4$) is a compact

operator on L^2 .

Noting that $h(H(\lambda))$ is a bounded operator from $L^2(\mathbf{R}^n)$ to $H^2(\mathbf{R}^n)$, by Rellich's theorem and (A.3), we have that $K_j(\lambda)$ ($j=1, 2, 3$) is a compact operator on L^2 . Let $u \in L^2$. Noting that

$$\nabla_y T_\varphi h(H(\lambda))u = T_\varphi \nabla_y h(H(\lambda))u + (\nabla_y \varphi(y)) T_\varphi \partial_z h(H(\lambda))u,$$

we have that $T_\varphi h(H(\lambda))$ is a bounded operator from $L^2(\mathbf{R}^n)$ to $H^1(|y| > r)$. Thus, Rellich's theorem and (A.0) imply that $K_4(\lambda)$ is also a compact on L^2 . Thus we obtain (2.15) by (2.16) and (2.17). ■

REMARK. Assuming that $0 < \sigma < 1$, we can also prove that Lemma 2.2 and 2.3. But, we need the condition $0 < \sigma < 1/2$ to prove the limiting absorption principle (see Appendix).

3. The limiting absorption principle.

SKETCH OF THE PROOF OF THEOREM 1.1 (1): Assume that $u \in H^2(\mathbf{R}^n)$ is the eigenfunction associated with eigenvalue $\lambda_0 > 0$ i. e. $Lu = \lambda_0 u$, then we can consider u as the eigenfunction of $H(\lambda_0)$ associated with eigenvalue λ_0 . Using Lemma 2.1 and Lemma 2.2, we can prove

$$\exp(\alpha \langle x \rangle) u \in L^2 (\forall \alpha \geq 0)$$

in the same way as in Froese and Herbst [6]. Moreover, we can also prove that $u=0$ in the same way as in Appendix I of Tamura [11]. ■

If $|\mu| > 1$, $(A+i\mu)^{-1}$ sends $H^k(\mathbf{R}^n)$ into $H^k(\mathbf{R}^n)$ and as an operator on $H^k(\mathbf{R}^n)$ ($k=0, \pm 1, \pm 2$)

$$(3.1) \quad s\text{-}\lim_{|\mu| \rightarrow \infty} i\mu(A+i\mu)^{-1} = I$$

is valid (see Lemma 2.3 of Weder [13]). Put $A(\mu) = i\mu A(A+i\mu)^{-1}$. It follows from $A(\mu) = i\mu + \mu^2(A+i\mu)^{-1}$ that $H(\lambda)A(\mu) - A(\mu)H(\lambda)$ is a bounded operator from $H^1(\mathbf{R}^n)$ to $H^{-1}(\mathbf{R}^n)$. Using (3.1), we can prove the following lemma in the same way as the proof of Lemma 2.4 of Weder [13].

LNMMMA 3.1.

$$i[H(\lambda), A(\mu)] = i\mu(A+i\mu)^{-1}i[H(\lambda), A]^0i\mu(A+i\mu)^{-1}$$

and

$$\begin{aligned} s\text{-}\lim_{\mu \rightarrow \infty} (-\Delta+1)^{-1/2}i[H(\lambda), A(\mu)](-\Delta+1)^{-1/2} \\ = (-\Delta+1)^{-1/2}i[H(\lambda), A]^0(-\Delta+1)^{-1/2} \end{aligned}$$

for all $\lambda \in (\lambda_0 - \delta, \lambda_0 + \delta)$, as an operator on L^2 .

Using Lemma 3.1, we can also prove the following lemma in the same way as the proof of Lemma 2.5 of Weder [13].

LEMMA 3.3. *Let $f \in C_0^\infty(\mathbf{R}^n)$. Then*

(i) *$f(H(\lambda))$ sends $D(A)$ into $D(A)$*

(ii) *$[f(H(\lambda)), A]$ defined as operator on $D(A)$ is extended to a bounded operator on L^2 which is denoted by $[f(H(\lambda)), A]^0$.*

Following Tamura [12], we consider cut off functions, $\chi_n(x) \in C_0^\infty(\mathbf{R}^n)$ such that $\chi_n(x)$ has support in $\{x \in \mathbf{R}^n; |x| < 2\}$ and $\chi_n = 1$ for $|x| \leq 1$. For $\varepsilon > 0$ small enough, we define

$$E_{L,\varepsilon}(x) = E_0(x) + \chi_n(\varepsilon x)(E_L(x) - E_0(x)),$$

$$E_{S,\varepsilon}(x) = \chi_n(\varepsilon x)E_S(x),$$

and

$$V_\varepsilon^r(y) = \chi_{|y|>r}(y)\chi_{n-1}(\varepsilon y)(y \cdot \nabla_y \varphi(y) - \varphi(y)).$$

We further define an operator $B(\varepsilon; \lambda)$ as

$$\begin{aligned} B(\varepsilon; \lambda) = & -2\Delta + \lambda((x \cdot \nabla E_{L,\varepsilon}) - nE_{S,\varepsilon} + \nabla^* \cdot x E_{S,\varepsilon} - E_{S,\varepsilon} x \cdot \nabla \\ & - T_\varphi^* V_\varepsilon^r(y)(E_L^+ - E_L^-)T_\varphi - (R_\varphi^r)^*(E_L^+ - E_L^-)R_\varphi^r). \end{aligned}$$

Note that by the proof of Lemma 2.2, we can consider $B(\varepsilon; \lambda)$ to be a bounded operator from $H^1(\mathbf{R}^n)$ to $H^{-1}(\mathbf{R}^n)$.

LEMMA 3.3. *Let $M(\varepsilon; \lambda) = f_\delta(H(\lambda))B(\varepsilon; \lambda)f_\delta(H(\lambda))$. Then $[M(\varepsilon; \lambda), A]$ defined as a form on $D(A)$ is extended to a bounded operator on L^2 which is denoted by $[M(\varepsilon; \lambda), A]^0$.*

PROOF. We again write f_δ for $f_\delta(H(\lambda))$. Noting that $M(\varepsilon; \lambda)$ is a bounded operator on L^2 , we can see that $[M(\varepsilon; \lambda), A(\mu)]$ is also a bounded operator on L^2 . Let $u \in L^2$. Since $A(\mu)$ sends $H^2(\mathbf{R}^n)$ into $H^2(\mathbf{R}^n)$, we have

$$\begin{aligned} (3.2) \quad & \langle A(\mu)u, M(\varepsilon; \lambda)u \rangle - \langle M(\varepsilon; \lambda)u, A(\mu)^*u \rangle \\ & = \langle A(\mu)f_\delta u, B(\varepsilon; \lambda)f_\delta u \rangle - \langle B(\varepsilon; \lambda)f_\delta u, A(\mu)^*f_\delta u \rangle \\ & \quad + \langle [f_\delta, A(\mu)]u, B(\varepsilon; \lambda)f_\delta u \rangle + \langle B(\varepsilon; \lambda)f_\delta u, [A(\mu)^*, f_\delta]u \rangle \end{aligned}$$

(3.2) implies that

$$\begin{aligned} (3.3) \quad & [M(\varepsilon; \lambda), A(\mu)] \\ & = f_\delta[B(\varepsilon; \lambda), A(\mu)]f_\delta + f_\delta B(\varepsilon; \lambda)[f_\delta, A(\mu)] + [f_\delta, A(\mu)]B(\varepsilon; \lambda)f_\delta. \end{aligned}$$

For $u \in H^2(\mathbf{R}^n) \cap D(A)$ satisfying $Au \in H^1(\mathbf{R}^n)$, we define the commutator $[B(\varepsilon; \lambda), A]$ as follows

$$\begin{aligned} \langle i[B(\varepsilon; \lambda), A]u, u \rangle &= \sum_{j=1}^4 \langle i[B_j(\varepsilon; \lambda), A]u, u \rangle \\ &= \sum_{j=1}^4 i(\langle Au, B_j(\varepsilon; \lambda)u \rangle - \langle B_j(\varepsilon; \lambda)u, Au \rangle), \end{aligned}$$

where

$$\begin{aligned} B_1(\varepsilon; \lambda) &= -2\Delta + \lambda(x \cdot \nabla E_{L,\varepsilon}), \\ B_2(\varepsilon; \lambda) &= -\lambda((E_{S,\varepsilon} x \cdot \nabla)^* + E_{S,\varepsilon} x \cdot \nabla + nE_{S,\varepsilon}) \\ &= -\lambda(\nabla^* \cdot x E_{S,\varepsilon} + E_{S,\varepsilon} x \cdot \nabla + nE_{S,\varepsilon}), \\ B_3(\varepsilon; \lambda) &= -\lambda T_\varphi^* V_\varepsilon^r (E_L^{+0} - E_L^{-0}) T_\varphi, \\ B_4(\varepsilon; \lambda) &= -\lambda (R_\varphi^r)^* (E_L^{+0} - E_L^{-0}) R_\varphi^r. \end{aligned}$$

Then the $i[B(\varepsilon; \lambda), A]$ has an extension $[B(\varepsilon; \lambda), A]^0$ to be a bounded operator from $H^2(\mathbf{R}^n)$ to $H^{-2}(\mathbf{R}^n)$ (see Appendix). Let $u \in L^2$. Again, noting that $A(\mu)H^k(\mathbf{R}^n) \subset H^k(\mathbf{R}^n)$ ($k = \pm 1, \pm 2$), we have

$$\begin{aligned} (3.4) \quad & \langle A(\mu)f_\delta u, B(\varepsilon; \lambda)f_\delta u \rangle - \langle B(\varepsilon; \lambda)f_\delta u, A(\mu)^*f_\delta u \rangle \\ &= \mu^2 \langle B(\varepsilon; \lambda)(A+i\mu)^{-1}f_\delta u, f_\delta u \rangle - \langle f_\delta u, B(\varepsilon; \lambda)(A-i\mu)^{-1}f_\delta u \rangle \\ &= \mu^2 \langle B(\varepsilon; \lambda)(A+i\mu)^{-1}f_\delta u, A(A-i\mu)^{-1}f_\delta u \rangle \\ & \quad - \langle A(A+i\mu)^{-1}f_\delta u, B(\varepsilon; \lambda)(A-i\mu)^{-1}f_\delta u \rangle. \end{aligned}$$

(3.4) implies that

$$(3.5) \quad f_\delta[B(\varepsilon; \lambda), A(\mu)]f_\delta = f_\delta i\mu(A+i\mu)^{-1}[B(\varepsilon; \lambda), A]^0 i\mu(A+i\mu)^{-1}f_\delta$$

as an operator on L^2 . (3.1) and (3.5) imply that

$$(3.6) \quad s - \lim_{|\mu| \rightarrow \infty} f_\delta[B(\varepsilon; \lambda), A(\mu)]f_\delta = f_\delta[B(\varepsilon; \lambda), A]^0 f_\delta,$$

for all $\lambda \in (\lambda_0 - \delta, \lambda_0 + \delta)$, as an operator on L^2 . Using Lemma 3.1, by the same way as in the proof of Lemma 2.4 of Weder [13], we can prove that

$$(3.7) \quad \begin{aligned} s - \lim_{|\mu| \rightarrow \infty} (-\Delta+1)^{1/2}[f_\delta, A(\mu)](-\Delta+1)^{1/2} \\ = (-\Delta+1)^{1/2}[f_\delta, A]^0(-\Delta+1)^{1/2} \end{aligned}$$

for all $\lambda \in (\lambda_0 - \delta, \lambda_0 + \delta)$, as an operator on L^2 . Hence by (3.2), (3.6) and (3.7), we have

$$(3.8) \quad \begin{aligned} s - \lim_{|\mu| \rightarrow \infty} [M(\varepsilon; \lambda), A(\mu)] \\ = f_\delta[B(\varepsilon; \lambda), A]^0 f_\delta + f_\delta B(\varepsilon; \lambda)[f_\delta, A]^0 + [f_\delta, A]^0 B(\varepsilon; \lambda)f_\delta \end{aligned}$$

for all $\lambda \in (\lambda_0 - \delta, \lambda_0 + \delta)$, as an operator on L^2 . We define $[M(\varepsilon; \lambda), A]^0$ by the right side of (3.8).

Finally, for $u, v \in D(A)$, we have

$$\begin{aligned} \langle [M(\varepsilon; \lambda), A]^0 u, v \rangle &= \lim_{|\mu| \rightarrow \infty} (\langle A(\mu)u, M(\varepsilon; \lambda)v \rangle - \langle M(\varepsilon; \lambda)u, A(\mu)v \rangle) \\ &= \langle Au, M(\varepsilon; \lambda)v \rangle - \langle M(\varepsilon; \lambda)u, Av \rangle. \quad \blacksquare \end{aligned}$$

We can prove the following lemma by a straightforward calculation.

LEMMA 3.4. *Let $\lambda_0 - \delta < \lambda < \lambda_0 + \delta$, $0 < \varepsilon < 1$. There exists a positive constant C independent of λ and ε such that*

- (i) $\|(-\Delta + 1)^{-1/2}(B(\lambda) - B(\varepsilon; \lambda))(-\Delta + 1)^{-1/2}\| \leq C\varepsilon^\theta$,
- (ii) $\|(-\Delta + 1)^{-1/2}(d/d\varepsilon)B(\varepsilon; \lambda)(-\Delta + 1)^{-1/2}\| \leq C\varepsilon^{\theta-1}$,
- (iii) $\|(-\Delta + 1)^{-1}[B(\varepsilon; \lambda), A]^0(-\Delta + 1)^{-1}\| \leq C\varepsilon^{\theta-1}$,

where $B(\lambda) = [H(\lambda), A]^0$.

Let $\lambda_0 > 0$. By Theorem 1.1(i), for the compact operator $K(\lambda_0)$ on L^2 , we have

$$\|f_\delta(H(\lambda_0))K(\lambda_0)f_\delta(H(\lambda_0))\| = o(1) \quad (\delta \rightarrow 0).$$

We can also show that

$$\|f_\delta(H(\lambda)) - f_\delta(H(\lambda_0))\| \leq C\delta,$$

where $C > 0$ depends on only λ_0 . Thus by Lemma 2.4, we can take δ so small that

$$\begin{aligned} (3.9) \quad M(\lambda) &\equiv f_\delta(H(\lambda))i[H(\lambda), A]^0f_\delta(H(\lambda)) \\ &\geq (\alpha/2)f_\delta(H(\lambda))^2 \end{aligned}$$

in the form sense.

Moreover, (3.9) together with Lemma 3.4(i) implies that

$$(3.10) \quad M(\varepsilon; \lambda) \geq \gamma f_\delta(H(\lambda))^2$$

for $\varepsilon > 0$ small enough, where $\gamma > 0$.

It follows from (3.10) that $M(\varepsilon; \lambda)$ is non-negative and hence we define an operator, $G_\kappa(\varepsilon; \lambda)$, on L^2 by

$$G_\kappa(\varepsilon; \lambda) = (H(\lambda) - \lambda - i\kappa E(x) - i\varepsilon M(\varepsilon; \lambda))^{-1}$$

for $0 < \kappa < 1$ and $0 < \varepsilon \ll 1$.

LEMMA 3.5. *Let $0 < \kappa < 1$, $\lambda_0 - \delta < \lambda < \lambda_0 + \delta$ and $0 < \varepsilon \ll 1$. There exists a positive constant C independent of κ , λ and ε such that*

$$\|G_\kappa(\varepsilon; \lambda)\| \leq C\varepsilon^{-1}.$$

For a proof of Lemma 3.5, see that of Lemma 5.3 of Kikuchi and Tamura [8] or Lemma 3.2 of Tamura [11].

For $1/2 < \alpha \leq 1$, we write

$$F_\kappa(\varepsilon; \lambda) = A_\alpha(\varepsilon)G_\kappa(\varepsilon; \lambda)A_\alpha(\varepsilon),$$

where

$$A_\alpha(\varepsilon) = (1 + |A|)^{-\alpha}(1 + \varepsilon|A|)^{\alpha-1}.$$

Differentiating $F_\kappa(\varepsilon; \lambda)$ in ε , we have

$$\begin{aligned} (3.11) \quad (d/d\varepsilon)F_\kappa(\varepsilon; \lambda) &= ((d/d\varepsilon)A_\alpha(\varepsilon))G_\kappa A_\alpha(\varepsilon) - iA_\alpha(\varepsilon)G_\kappa M(\varepsilon; \lambda)G_\kappa A_\alpha(\varepsilon) \\ &\quad - i\varepsilon A_\alpha(\varepsilon)G_\kappa((d/d\varepsilon)M(\varepsilon; \lambda))G_\kappa A_\alpha(\varepsilon) \\ &\quad + A_\alpha(\varepsilon)G_\kappa((d/d\varepsilon)A_\alpha(\varepsilon)). \end{aligned}$$

Repeating the argument in the proof of Lemma 2.9 of Weder [13], we can show from Lemma 3.2 and 3.3

$$(3.12) \quad G_\kappa(\varepsilon; \lambda)D(A) \subset D(A) \cap H^2(\mathbf{R}^n).$$

Let $g_\delta(p) = 1 - f_\delta(p)$. We write in brief f_δ and g_δ for $f_\delta(H(\lambda))$ and $g_\delta(H(\lambda))$ respectively.

Using (3.11), (3.12) and Lemma 3.3, we can decompose $(d/d\varepsilon)F_\kappa(\varepsilon; \lambda)$ as a form on L^2

$$(3.13) \quad (d/d\varepsilon)F_\kappa(\varepsilon; \lambda) = \sum_{j=1}^9 Y_\kappa^j(\varepsilon; \lambda),$$

where

$$\begin{aligned} Y_\kappa^1(\varepsilon; \lambda) &= iA_\alpha(\varepsilon)G_\kappa f_\delta B(\varepsilon; \lambda)g_\delta G_\kappa A_\alpha(\varepsilon), \\ Y_\kappa^2(\varepsilon; \lambda) &= iA_\alpha(\varepsilon)G_\kappa g_\delta B(\varepsilon; \lambda)f_\delta G_\kappa A_\alpha(\varepsilon), \\ Y_\kappa^3(\varepsilon; \lambda) &= iA_\alpha(\varepsilon)G_\kappa g_\delta B(\varepsilon; \lambda)g_\delta G_\kappa A_\alpha(\varepsilon), \\ Y_\kappa^4(\varepsilon; \lambda) &= -iA_\alpha(\varepsilon)G_\kappa(B(\varepsilon; \lambda) - B(\lambda))G_\kappa A_\alpha(\varepsilon) \\ Y_\kappa^5(\varepsilon; \lambda) &= -iA_\alpha(\varepsilon)G_\kappa[H(\lambda) - \lambda - i\kappa E(x) - i\varepsilon M(\varepsilon; \lambda), A]G_\kappa A_\alpha(\varepsilon), \\ Y_\kappa^6(\varepsilon; \lambda) &= \kappa A_\alpha(\varepsilon)G_\kappa[E(x), A]G_\kappa A_\alpha(\varepsilon), \\ Y_\kappa^7(\varepsilon; \lambda) &= \varepsilon A_\alpha(\varepsilon)G_\kappa[M(\varepsilon; \lambda), A]G_\kappa A_\alpha(\varepsilon), \\ Y_\kappa^8(\varepsilon; \lambda) &= -i\varepsilon A_\alpha(\varepsilon)G_\kappa((d/d\varepsilon)M(\varepsilon; \lambda))G_\kappa A_\alpha(\varepsilon), \\ Y_\kappa^9(\varepsilon; \lambda) &= ((d/d\varepsilon)A_\alpha(\varepsilon))G_\kappa A_\alpha(\varepsilon) + A_\alpha(\varepsilon)G_\kappa((d/d\varepsilon)A_\alpha(\varepsilon)). \end{aligned}$$

We need the following lemmas to estimate each term of right side of (3.13).

LEMMA 3.6. *Let $0 < \kappa < 1$, $\lambda_0 - \delta < \lambda < \lambda_0 + \delta$, and $0 < \varepsilon \ll 1$. There exists a positive constant independent of κ , λ and ε such that*

$$(i) \quad \|(-\Delta + 1)^{1/2} f_\delta G_\kappa(\varepsilon; \lambda) A_\alpha(\varepsilon)\| \leq C \varepsilon^{-1/2} \|F_\kappa\|^{1/2},$$

$$(ii) \quad \|(-\Delta + 1)^{1/2} g_\delta G_\kappa(\varepsilon; \lambda) A_\alpha(\varepsilon)\| \leq C,$$

$$(iii) \quad \|(-\Delta + 1)^{1/2} G_\kappa(\varepsilon; \lambda) A_\alpha(\varepsilon)\| \leq C \kappa^{-1/2} \|F_\kappa\|^{1/2}.$$

For a proof of Lemma 3.6, see that of Lemma 5.5 of Kikuchi and Tamura [8].

LEMMA 3.7. *Let $\lambda_0 - \delta < \lambda < \lambda_0 + \delta$ and $0 < \varepsilon \ll 1$. There exists a positive constant independent of λ and ε such that*

$$\|[M(\varepsilon; \lambda), A]^0\| \leq C \varepsilon^{\theta-1}.$$

PROOF. By the definition of $[M(\varepsilon; \lambda), A]^0$, we estimate each term of right side of (3.8).

Lemma 3.4(iii) implies that

$$\|f_\delta [B(\varepsilon; \lambda), A]^0 f_\delta\| \leq C \varepsilon^{\theta-1}.$$

Noting the definition of $B(\varepsilon; \lambda)$ and (3.7), we also have that

$$\|f_\delta B(\varepsilon; \lambda) [f_\delta, A]^0\|, \|[f_\delta, A]^0 B(\varepsilon; \lambda) f_\delta\| \leq C,$$

where $C > 0$ is independent of ε . Thus, the proof of Lemma 3.7 is now complete. ■

Using lemma 3.6 and 3.7, we can evaluate the norm of Y_κ^j , $1 \leq j \leq 9$, and obtain the following differential inequality (see Tamura [12])

$$(3.14) \quad \|(d/d\varepsilon)F_\kappa(\varepsilon; \lambda)\| \leq C(\varepsilon^{\alpha-1} + \varepsilon^{\alpha-3/2} \|F_\kappa\|^{1/2} + \varepsilon^{\theta-1} \|F_\kappa\|).$$

Let ε_0 , $0 < \varepsilon_0 \ll 1$. Then by Lemma 3.5, we have

$$(3.15) \quad \|F_\kappa(\varepsilon_0; \lambda)\| \leq C \varepsilon_0^{-1}.$$

By (3.10) and (3.11), we immediately obtain

$$(3.16) \quad \|F_\kappa(\varepsilon; \lambda)\| \leq C,$$

where $C > 0$ independent of $0 < \kappa < 1$, $0 < \varepsilon < \varepsilon_0$ and $\lambda_0 - \delta < \lambda < \lambda_0 + \delta$.

In order to prove (ii) and (iii) of theorem 1.1, we need the following lemma (see Weder [13]).

LEMMA 3.8. *Let $\lambda_0 - \delta < \lambda < \lambda_0 + \delta$ and $0 < \varepsilon \ll 1$. There exists a positive constant*

C independent of λ and ε such that

$$\|A_\alpha(\varepsilon)^{-1}f_\delta X_\alpha(\varepsilon)\| \leq C$$

where

$$X_\alpha(\varepsilon) = (1 + |x|^2)^{-\alpha/2} (1 + \varepsilon^2 |x|^2)^{(\alpha-1)/2}.$$

PROOF. By the same argument as in Weder [13], we can show that $(1 + \varepsilon|A|)f_\delta(1 + \varepsilon^2|x|^2)^{-1/2}$ is a bounded operator on L^2 and

$$(3.17) \quad \|(1 + \varepsilon|A|)f_\delta(1 + \varepsilon^2|x|^2)^{-1/2}\| \leq C,$$

where C independent of ε and λ ($0 < \varepsilon \ll 1$, $\lambda_0 - \delta < \lambda < \lambda_0 + \delta$).

Take $\varepsilon=1$, then we also have

$$(3.18) \quad \|(1 + |A|)f_\delta(1 + |x|^2)^{-1/2}\| \leq C$$

we obtain

$$\|A_\alpha(\varepsilon)^{-1}f_\delta X_\alpha(\varepsilon)\| \leq C$$

by interpolation between (3.17) and (3.18). ■

PROOF OF THEOREM 1.1 (II), (III): Using (3.1) and Lemma 3.8, we have

$$(3.19) \quad \|X_\alpha(\varepsilon)f_\delta G_\varepsilon(\varepsilon; \lambda)f_\delta X_\alpha(\varepsilon)\| \leq C.$$

Moreover, since

$$\|g_\delta G_\varepsilon(\varepsilon; \lambda)\|, \|G_\varepsilon(\varepsilon; \lambda)g_\delta\| \leq C,$$

we have

$$(3.20) \quad \|X_\alpha(\varepsilon)g_\delta G_\varepsilon(\varepsilon; \lambda)f_\delta X_\alpha(\varepsilon)\| \leq C,$$

$$(3.21) \quad \|X_\alpha G_\varepsilon(\varepsilon; \lambda)g_\delta X_\alpha(\varepsilon)\| \leq C.$$

By (3.19), (3.20) and (3.21), we obtain

$$(3.22) \quad \|X_\alpha(\varepsilon)G(\varepsilon; \lambda)X_\alpha(\varepsilon)\| \leq C.$$

We write

$$F_\varepsilon(\varepsilon; \lambda) = X_\alpha(\varepsilon)G_\varepsilon(\varepsilon; \lambda)X_\alpha(\varepsilon).$$

We can prove (3.14) for $F_\varepsilon(\varepsilon; \lambda)$ defined above. Thus using (3.22), we have

$$(3.23) \quad \|(d/d\varepsilon)F_\varepsilon(\varepsilon; \lambda)\| \leq C\varepsilon^{\beta-1}$$

where $0 < \beta < 1$.

Finally, by (3.22) and (3.23), we obtain Theorem 1.1(ii), (iii). ■

4. Appendix.

In this section we show that the commutator $i[B_j(\varepsilon; \lambda), A]$ ($j=1, 2, 3, 4$) is

extended to a bounded operator from $H^2(\mathbf{R}^n)$ to $H^{-2}(\mathbf{R}^n)$. We write $i[B_j(\varepsilon; \lambda), A]^0$ ($j=1, 2, 3, 4$) for the extended operators.

We define an operator P_φ^r as

$$P_\varphi^r u = \chi_{|y| \leq r}(y) ((y \cdot \nabla_y \varphi(y) - \varphi(y)) \varphi(y))^{1/2} T_\varphi u,$$

for $u \in \mathcal{S}(\mathbf{R}^n)$.

LEMMA 4.1. P_φ^r is extended to a bounded operator from $H^1(\mathbf{R}^n)$ to $L^2(\mathbf{R}^{n-1})$.

PROOF. If $0 < \sigma < 1/2$, there exist some $s' > 1/2$ and $p' > n-1$ such that $n-1-2\sigma p' > 0$ and $(n-1)/2p' + s' = 1$. Then we can prove that

$$\|P_\varphi^r u\|_{L^2(\mathbf{R}^{n-1})} \leq C r^{(n-1-2\sigma p')/2p'} \|u\|_{H^2(\mathbf{R}^n)}$$

in the same way as in the proof of (2.14). ■

By a straightforward calculation, the form $i[B_j(\varepsilon; \lambda), A]$ ($j=1, 2$) is extended to a bounded operator from $H^2(\mathbf{R}^n)$ to $H^{-2}(\mathbf{R}^n)$. Moreover, we have that

$$\begin{aligned} i[B_1(\varepsilon; \lambda), A]^0 &= -4\Delta + \lambda((x \cdot \nabla E_{L,\varepsilon})(x \cdot \nabla) + (\nabla^* \cdot x)(x \cdot \nabla E_{L,\varepsilon}) + n(x \cdot \nabla E_{L,\varepsilon})), \\ i[B_2(\varepsilon; \lambda), A]^0 &= -\lambda(E_{S,\varepsilon}(x \cdot \nabla)(x \cdot \nabla) + (E_{S,\varepsilon}(x \cdot \nabla)(x \cdot \nabla))^* + 2(\nabla^* \cdot x)E_{S,\varepsilon}(x \cdot \nabla) \\ &\quad + 2nE_{S,\varepsilon}(x \cdot \nabla) + 2n(\nabla^* \cdot x)E_{S,\varepsilon} + n^2E_{S,\varepsilon}). \end{aligned}$$

Let $u \in H^2(\mathbf{R}^n) \cap \mathcal{D}(A)$ satisfying $Au \in H^1(\mathbf{R}^n)$. Then we have that

$$\begin{aligned} (4.1) \quad \langle i[B_3(\varepsilon; \lambda), A]u, u \rangle &= -\lambda \langle y \cdot T_\varphi \nabla_y u, V_\varepsilon^r(E_L^{+0} - E_L^{-0})T_\varphi u \rangle_{L^2(\mathbf{R}^{n-1})} \\ &\quad + \langle \varphi(y)T_\varphi \partial_z u, V_\varepsilon^r(E_L^{+0} - E_L^{-0})T_\varphi u \rangle_{L^2(\mathbf{R}^{n-1})} \\ &\quad + \langle V_\varepsilon^r(E_L^{+0} - E_L^{-0})T_\varphi u, y \cdot T_\varphi \nabla_y u \rangle_{L^2(\mathbf{R}^{n-1})} \\ &\quad + \langle V_\varepsilon^r(E_L^{+0} - E_L^{-0})T_\varphi u, \varphi(y)T_\varphi \partial_z u \rangle_{L^2(\mathbf{R}^{n-1})} \\ &\quad + n \langle V_\varepsilon^r(E_L^{+0} - E_L^{-0})T_\varphi u, T_\varphi u \rangle_{L^2(\mathbf{R}^{n-1})}, \end{aligned}$$

$$\begin{aligned} (4.2) \quad \langle i[B_4(\varepsilon; \lambda), A]u, u \rangle &= -\lambda \langle y \cdot R_\varphi^r \nabla_y u, (E_L^{+0} - E_L^{-0})R_\varphi^r u \rangle_{L^2(\mathbf{R}^{n-1})} \\ &\quad + \langle P_\varphi^r \partial_z u, (E_L^{+0} - E_L^{-0})P_\varphi^r u \rangle_{L^2(\mathbf{R}^{n-1})} \\ &\quad + \langle (E_L^{+0} - E_L^{-0})R_\varphi^r u, y \cdot R_\varphi^r \nabla_y u \rangle_{L^2(\mathbf{R}^{n-1})} \\ &\quad + \langle (E_L^{+0} - E_L^{-0})P_\varphi^r u, P_\varphi^r \partial_z u \rangle_{L^2(\mathbf{R}^{n-1})} \\ &\quad + n \langle (E_L^{+0} - E_L^{-0})R_\varphi^r u, R_\varphi^r u \rangle_{L^2(\mathbf{R}^{n-1})}. \end{aligned}$$

Lemma 2.1 and (4.1) imply that $i[B_3(\varepsilon; \lambda), A]$ is extended to a bounded operator from $H^2(\mathbf{R}^n)$ to $H^{-2}(\mathbf{R}^n)$. Lemma 4.1, (2.14) and (4.2) imply that

$i[B_3(\varepsilon; \lambda), A]$ is extended to a bounded operator from $H^2(\mathbf{R}^n)$ to $H^{-2}(\mathbf{R}^n)$. We also have that

$$\begin{aligned} i[B_3(\varepsilon; \lambda), A]^0 = & -\lambda(T_\varphi^*(E_L^{+0} - E_L^{-0})V_\varepsilon^r y \cdot T_\varphi \nabla_y + (T_\varphi \nabla_y)^* \cdot y(E_L^{+0} - E_L^{-0})V_\varepsilon^r T_\varphi \\ & + T_\varphi^*(E_L^{+0} - E_L^{-0})\varphi(y)V_\varepsilon^r T_\varphi \partial_z + (T_\varphi \partial_z)^*(E_L^{+0} - E_L^{-0})\varphi(y)V_\varepsilon^r T_\varphi \\ & + nT_\varphi^*(E_L^{+0} - E_L^{-0})V_\varepsilon^r T_\varphi), \end{aligned}$$

$$\begin{aligned} i[B_4(\varepsilon; \lambda), A]^0 = & -\lambda((R_\varphi^r)^*(E_L^{+0} - E_L^{-0})y \cdot R_\varphi^r \nabla_y + (R_\varphi^r \nabla_y)^* \cdot y(E_L^{+0} - E_L^{-0})R_\varphi^r \\ & + (P_\varphi^r)^*(E_L^{+0} - E_L^{-0})P_\varphi^r \partial_z + (P_\varphi^r \partial_z)^*(E_L^{+0} - E_L^{-0})P_\varphi^r \\ & + n(R_\varphi^r)^*(E_L^{+0} - E_L^{-0})R_\varphi^r). \end{aligned}$$

We define an operator $i[B(\varepsilon; \lambda), A]^0$ as

$$i[B(\varepsilon; \lambda), A]^0 = \sum_{j=1}^4 i[B_j(\varepsilon; \lambda), A]^0.$$

Thus, the form $i[B(\varepsilon; \lambda), A]$ is extended to a bounded operator from $H^2(\mathbf{R}^n)$ to $H^{-2}(\mathbf{R}^n)$.

REMARK. We can consider that $i[B_1(\varepsilon; \lambda); A]^0$ belongs to $\mathfrak{B}(H^1, H^{-1})$ and $i[B_3(\varepsilon; \lambda), A]^0$ belongs to $\mathfrak{B}(H^{1+s}, H^{-1-s})$, where $s > 1/2$.

References

- [1] Ben-Artzi. M, Dermenjian. Y and Guillot. J.C, Acoustic waves in perturbed stratified fluids: a spectral theory, *Comm. Partial Differential Equations* 14 (1989), 479-517.
- [2] Debièvre. S and Pravica. W, Spectral analysis for optical fibres and stratified fluids I: the limiting absorption principle, *J. Func. Anal* 98 (1991), 404-436.
- [3] Dermenjian. Y and Guillot. J.C, Théorie spectrale de la propagation des ondes acoustiques dans un milieu stratifié perturbé, *J. Defferential Equations* 62 (1986), 357-409.
- [4] Eastham. M.S and Kalf. H, "Schrödinger-type operators with continuous spectra, Resaech note in Math", Pitman, 1982.
- [5] Eidus. D, The limiting absorption and amplitude principle for the diffraction problem with two unbounded media, *Comm. Math. Phys* 107 (1986), 27-38.
- [6] Froese. R and Herbst. I, Exponential bounds and absence of positive eigenvalues for N -body Schrödinger operator, *Comm. Math. Phys* 87 (1982), 429-447.
- [7] Iwashita. H, Spectral theory for symmetric systems in an exterior domain, *Tsukuba J. Math* 11 (1987), 241-256.
- [8] Kikuchi. K and Tamura. H, Limiting amplitude principle for acoustic propagators in perturbed stratified fluids, *J. Defferential Equations* 93 (1991), 260-282.
- [9] Mourre. E, Absence of singular continuous spectrum for certain self-adjoint operators, *Comm. Math. Phys* 78 (1981), 391-408.
- [10] Perry. P, Sigal. I.M and Simon. B, Spectral analysis of N -body Schrödinger operators, *Ann of Math* 114 (1981), 519-567.

- [11] Tamura. H, Resolvent estimates at low frequencies and limiting ampliting amplitude principle for acoustic propagators, *J. Math. Soc. Japan* **41** (1989), 549-575.
- [12] ———, Principle of limiting absorption for N -body Schrödinger operators, A remark on the commutator method, *Lett. in Math. Phys* **17** (1989), 31-36.
- [13] Weder. R, Spectral analysis of strongly propagator systems, *J. Reine Angew. Math.* **354** (1984), 95-122.
- [14] ———, Spectral and scattering theory in perturbed stratified fluids, *J. Math. Pures et Appl* **64** (1985), 149-173.
- [15] ———, Spectral and scattering theory in perturbed stratified fluids II, transmission and exterior domains, *J. Defferential Equations* **64** (1986), 109-131.
- [16] ———, The limiting absorption principle at thresholds, *J. Math. pures at Appl* **67** (1988), 318-338.
- [17] Wilcox. C. H, Spectral analysis of the Pekeris operator in the theory of acoustic wave propagation in Shallow water, *Arch. Rational Mech. Anal* **60** (1976), 259-300.

University of Tsukuba, Institute of Mathematics,
Ibaraki, 305, Japan