

A NOTE ON MULTIVALENT FUNCTIONS

By

Mamoru NUNOKAWA

The author would like to dedicate this paper to the memory of the late Professor Shigeo Ozaki.

It is well known that if $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is analytic in $E = \{z : |z| < 1\}$ and $\operatorname{Re} f'(z) > 0$ in E , then $f(z)$ is univalent in E .

Ozaki [3, Theorem 2] extended the above result to the following: if $f(z)$ is analytic in a convex domain D and

$$\operatorname{Re}(e^{i\alpha} f^{(p)}(z)) > 0 \quad \text{in } D$$

where α is a real constant, then $f(z)$ is at most p -valent in D .

This shows that if $f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n$ is analytic in E and

$$\operatorname{Re} f^{(p)}(z) > 0 \quad \text{in } E,$$

then $f(z)$ is p -valent in E .

DEFINITION. Let $F(z)$ be analytic and univalent in E and suppose that $F(E) = D$. If $f(z)$ is analytic in E , $f(0) = F(0)$, and $f(E) \subset D$, then we say that $f(z)$ is subordinate to $F(z)$ in E , and we write

$$f(z) \prec F(z).$$

In this paper, we need the following lemmata.

LEMMA 1. If $p(z)$ is analytic in E , with $p(0) = 1$, and

$$p(z) + zp'(z) \prec \left(\frac{1+z}{1-z} \right)^{3/2} \quad \text{in } E,$$

then we have

$$p(z) \prec \frac{1+z}{1-z} \quad \text{in } E.$$

We owe this lemma to [1, Theorem 5 and its remark].

REMARK. From Lemma 1, it is trivial that if $p(0) = 1$ and

Received August 24, 1988.

$$|\arg(p(z) + zp'(z))| < \frac{3}{4}\pi \quad \text{in } E,$$

then we have

$$|\arg p(z)| < \frac{\pi}{2} \quad \text{in } E.$$

LEMMA 2. Let $f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n$ be analytic in E . If there exists a $(p-k+1)$ -valent starlike function $g(z) = z^{p-k+1} + \sum_{n=p-k+2}^{\infty} b_n z^n$ that satisfies

$$\operatorname{Re} \frac{zf^{(k)}(z)}{g(z)} > 0 \quad \text{in } E,$$

then $f(z)$ is p -valent in E .

We owe this lemma to [2, Theorem 8].

MAIN THEOREM. Let $p \geq 2$. If $f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n$ is analytic in E and

$$(1) \quad |\arg f^{(p)}(z)| < \frac{3}{4}\pi \quad \text{in } E,$$

then $f(z)$ is p -valent in E .

PROOF. Let us put

$$p(z) = f^{(p-1)}(z)/(p!z).$$

Then from the assumption (1) and by an easy calculation, we have

$$p(z) + zp'(z) = \frac{f^{(p)}(z)}{p!} < \left(\frac{1+z}{1-z} \right)^{3/2} \quad \text{in } E$$

and $p(0) = 1$. Then, from Lemma 1, we have

$$\frac{f^{(p-1)}(z)}{p!z} < \frac{1+z}{1-z} \quad \text{in } E.$$

This shows that

$$(2) \quad \operatorname{Re} \frac{f^{(p-1)}(z)}{z} = \operatorname{Re} \frac{zf^{(p-1)}(z)}{z^2} > 0 \quad \text{in } E.$$

Moreover, it is trivial that $g(z) = z^2$ is 2-valently starlike in E . Therefore, from Lemma 2 and (2), we have that $f(z)$ is p -valent in E .

The author sincerely appreciate the suggestions which were made by the referee.

References

- [1] Miller, S.S. and Mocanu, P.T., Marx-Strohhäcker differential subordination systems, Proc. Amer. Math. Soc. **99** (1987), 527-534.
- [2] Nunokawa, M., On the theory of multivalent functions, Tsukuba J. Math. **11** (1987), 273-286.
- [3] Ozaki, S., On the theory of multivalent functions, Sci. Rep. Tokyo Bunrika Daigaku A, **2** (1935), 167-188.

Department of Mathematics
Gunma University
Aramaki, Maebashi, 371
Japan