

GENERATORS AND RELATIONS FOR COMPACT LIE ALGEBRAS

By

S. BERMAN¹⁾

1. Introduction.

The main purpose of this paper is to provide a system of generators and relations for each of the nine types of compact simple Lie algebras. Indeed, we are able to give a presentation of each such algebra which depends only on the finite Cartan matrix (A_{ij}) which is attached to the complexification of our compact algebra. One of the main results that lies behind our work is the Theorem of Serre which gives a presentation of the simple Lie algebras over the complex field attached to (A_{ij}) .

Although our main interest is with the compact Lie algebras, we work in the generality of Kac-Moody Lie algebras (see [1], [4], [7], [8]). In this setting we will be able to provide a generalization of the compact simple Lie algebras. We realize these algebras as certain forms of the Kac-Moody algebra. More specifically, if (A_{ij}) is any indecomposable Cartan matrix which is non-Euclidean we let \mathcal{L}_C (resp. $\bar{\mathcal{L}}_C$) be the reduced (resp. standard) Kac-Moody Lie algebra over the complex field C , (see Section 1 for more details). We define a real form \mathcal{L}_C (resp. $\bar{\mathcal{L}}_C$) of \mathcal{L}_C (resp. $\bar{\mathcal{L}}_C$), and show that \mathcal{L}_C is the only simple homomorphic image of $\bar{\mathcal{L}}_C$. We then give generators and relations for $\bar{\mathcal{L}}_C$. The question of when $\bar{\mathcal{L}}_C = \mathcal{L}_C$ is equivalent to the question of when $\mathcal{L}_C = \bar{\mathcal{L}}_C$, and is a major unsolved question about Kac-Moody algebras. However, thanks to Serre's Theorem, we know $\mathcal{L}_C = \bar{\mathcal{L}}_C$, and hence $\mathcal{L}_C = \bar{\mathcal{L}}_C$, when (A_{ij}) is of finite type. This yields a presentation of \mathcal{L}_C in this case.

The content of the paper is as follows. In Section 1 we recall the notation and a few facts about Kac-Moody algebras. In Section 2, the final section, we begin by making a study of the algebras \mathcal{L}_C and $\bar{\mathcal{L}}_C$. We then go on to obtain a presentation of $\bar{\mathcal{L}}_C$, and then use this in dealing with the compact simple Lie algebras. We think it is interesting that analogues of the compact Lie algebras exist in the Kac-Moody setting. Moreover, just as the compact algebras are

1) This work was supported by a National Science and Engineering Research Council of Canada grant.

Received September 8, 1980. Revised December 8, 1980.

important in studying all real forms of the simple Lie algebras over \mathbf{C} (e.g. Cartan and Iwasawa decompositions), no doubt the generalizations studied here will play a similar role. Throughout, we let \mathbf{R} denote the real field and \mathbf{C} the complex field.

Section 1: We use similar notation to [1] and [2] where the reader may find all the necessary facts. Also, one may consult [4], [7], [8]. Thus, (A_{ij}) will denote an $l \times l$ indecomposable Cartan matrix which is not Euclidean. $\tilde{\mathcal{L}}$ denotes the universal Kac-Moody algebra of type (A_{ij}) over \mathbf{R} , so that $\tilde{\mathcal{L}}$ is generated by $3l$ elements $e_j, h_j, f_j, 1 \leq j \leq l$, subject to the relations

- (i) $[e_k, h_j] = A_{jk} e_k,$
- (ii) $[f_k, h_j] = -A_{jk} f_k,$
- (iii) $[h_j, h_k] = 0,$
- (iv) $[e_k, f_j] = \delta_{kj} h_k, 1 \leq j, k \leq l.$

The reduced Kac-Moody algebra is defined to be $\tilde{\mathcal{L}}/\mathcal{R}$ where \mathcal{R} is the radical of $\tilde{\mathcal{L}}$. We let this be denoted by \mathcal{L} and recall that \mathcal{R} is the unique maximal ideal of $\tilde{\mathcal{L}}$. The standard Kac-Moody algebra is denoted $\bar{\mathcal{L}}$ and is $\tilde{\mathcal{L}}$ factored by the ideal \mathcal{E} , where we recall that \mathcal{E} is generated by the elements $e_j(ad e_k)^{-A_{kj}+1}, f_j(ad f_k)^{-A_{kj}+1}, 1 \leq j \neq k \leq l$, and the elements $\{h \in \tilde{H} \mid \alpha_j(h) = 0, 1 \leq j \leq l\}$. Here, as usual, \tilde{H} is the linear span of h_1, \dots, h_l in $\tilde{\mathcal{L}}$. $V_{\mathbf{Z}}$ denotes the free abelian group with basis $\alpha_1, \dots, \alpha_l$ and $V_{\mathbf{Z}}$ acts on \tilde{H} by $\alpha_j(h_k) = A_{kj}$ for $1 \leq j, k \leq l$. Γ_1 denotes the roots of $\tilde{\mathcal{L}}$ and Γ_2 will denote the roots of \mathcal{L} and $\bar{\mathcal{L}}$, (they have the same roots). Thus, we have the decomposition $\mathcal{L} = H \oplus \sum_{\alpha \in \Gamma_2} \mathcal{L}_{\alpha}$; similarly for $\bar{\mathcal{L}}$ and $\tilde{\mathcal{L}}$. As usual, n denotes the automorphism of \mathcal{L} (or $\bar{\mathcal{L}}$ or $\tilde{\mathcal{L}}$) which satisfies $n(e_j) = f_j, n(f_j) = e_j, 1 \leq j \leq l$. Here, we are of course using the convention of letting $e_j, h_j, f_j, 1 \leq j \leq l$, denote the image in \mathcal{L} or $\bar{\mathcal{L}}$ of the corresponding elements of $\tilde{\mathcal{L}}$. Note that $\bar{\mathcal{L}}$ has a unique maximal ideal which is the kernel of the natural map of $\bar{\mathcal{L}}$ onto \mathcal{L} .

If S denotes any one of the algebras $\mathcal{L}, \bar{\mathcal{L}}, \tilde{\mathcal{L}}$ we let $S_{\mathbf{C}}$ denote the complexification of S . Thus, $S_{\mathbf{C}} = \mathbf{C} \otimes_{\mathbf{R}} S$, and we let τ be the map $-\otimes n$ of $S_{\mathbf{C}}$, where the "over bar" denotes complex conjugation. Thus, τ is a semi-linear automorphism of $S_{\mathbf{C}}$ of period 2. We define $S_{\mathbf{C}}$ to be the fixed points of $S_{\mathbf{C}}$ under τ . Clearly $S_{\mathbf{C}}$ is real form of $S_{\mathbf{C}}$. We let $S_+ = \{s \in S \mid n(s) = s\}$ and $S_- = \{s \in S \mid n(s) = -s\}$. Then we have the decompositions

$$(1.1) \quad S = S_+ \oplus S_-, \quad S_{\mathbf{C}} = S_+ \oplus iS_-.$$

Of course, $\mathcal{L}_{\mathbf{C}}$ is our generalization of the simple compact Lie algebras. Indeed, we will see in the next section that $\mathcal{L}_{\mathbf{C}}$ is a finite dimensional simple

compact Lie algebra when (A_{ij}) is of finite type. For the present we remark, since \mathcal{L}_C is simple, and \mathcal{L}_C is a real form of \mathcal{L}_C , that \mathcal{L}_C is simple. Moreover, it is clear, and follows from the corresponding fact for $\tilde{\mathcal{L}}_C$ (or $\tilde{\mathcal{L}}_C$), that $\tilde{\mathcal{L}}_C$ (or $\tilde{\mathcal{L}}_C$) has a unique maximal ideal which is the kernel of the obvious natural homomorphism onto \mathcal{L}_C .

Section 2: As in the previous section we let S denote any one of the algebras \mathcal{L} , $\tilde{\mathcal{L}}$, or $\tilde{\mathcal{L}}$. Letting H_S denote the image of \tilde{H} in S we have the decomposition $S=H_S\oplus\sum_{\alpha\in\Gamma}S_\alpha$, where Γ is the root system of S . Let $n_\alpha=\dim S_\alpha$, and for $\alpha\in\Gamma^+$ we let $x_{\alpha,1}, \dots, x_{\alpha,n_\alpha}$ be a basis of the space S_α chosen from among the elements $[e_{j_1}, \dots, e_{j_l}]$ where $\alpha_{j_1}+\dots+\alpha_{j_l}=\alpha$. It's worth noting that the n_α 's are known when $S=\mathcal{L}$ or $S=\tilde{\mathcal{L}}$, (see [3]). Thus, for example, $x_{\alpha,1}=e_j$ for $1\leq j\leq l$. Let $x_{-\alpha,j}=n(x_{\alpha,j})$ for $\alpha\in\Gamma^+$, $1\leq j\leq n_\alpha$, so that $x_{-\alpha,1}, \dots, x_{-\alpha,n_\alpha}$ is a basis of $S_{-\alpha}$. Then we have

LEMMA 2.1. $\{x_{\alpha,j}+x_{-\alpha,j}|\alpha\in\Gamma^+, 1\leq j\leq n_\alpha\}$ is a basis of S_+ .

PROOF. Let $x=h+\sum_{\alpha\in\Gamma^+}\sum_{j=1}^{n_\alpha}(a_{\alpha,j}x_{\alpha,j}+b_{\alpha,j}x_{-\alpha,j})$ be an element in S_+ where $h\in H_S$. Then $n(x)=x$ implies that $h=0$ and $a_{\alpha,j}=b_{\alpha,j}$ for all $\alpha\in\Gamma^+$, $1\leq j\leq n_\alpha$. Thus, x is in the linear span of the set $\{x_{\alpha,j}+x_{-\alpha,j}|\alpha\in\Gamma^+, 1\leq j\leq n_\alpha\}$. The rest is clear. □

We now define some elements of interest to us. Let $x_j=e_j+f_j$, $y_j=i(e_j-f_j)$, and $z_j=ih_j$ for $1\leq j\leq l$. Clearly $x_j, y_j, z_j\in S_C$ for $1\leq j\leq l$.

LEMMA 2.2. S_+ is generated by the elements x_1, \dots, x_l .

PROOF. Let M be the subalgebra of S_+ generated by the elements x_1, \dots, x_l . By Lemma 2.1 it is enough to show that for all $\alpha\in\Gamma^+$ and $j\in\{1, \dots, n_\alpha\}$ that $x_{\alpha,j}+x_{-\alpha,t}\in M$. We do this by induction on $l(\alpha)$, where $l(\alpha)$ is defined to be $\sum C_j$, when $\alpha=\sum C_j\alpha_j$; the case when $l(\alpha)=1$ being clear because $\dim S_{\alpha_k}=1$ for $1\leq k\leq l$, so that $x_{\alpha_k,1}+x_{-\alpha_k,1}=x_k$ for $1\leq k\leq l$. Assume $\alpha\in\Gamma^+$ and $l(\alpha)=n+1$ where $n\geq 1$ and that if $\beta\in\Gamma^+$ and $l(\beta)\leq n$ that $x_{\alpha,j}+x_{-\beta,j}\in M$ for $1\leq j\leq n_\beta$. Now, since $l(\alpha)\geq 2$ then for any $j\in\{1, \dots, n_\alpha\}$ we may assume that there exists some $\beta\in\Gamma^+$, $k\in\{1, \dots, l\}$ such that $l(\beta)=n$, and $x_{\alpha,j}=[x_{\beta,t}, e_k]$, for some $t\in\{1, \dots, n_\alpha\}$. (This is because $x_{\alpha,j}=[e_{j_1}, \dots, e_{j_n}, e_k]$ for some k and $x_{\alpha_k,1}=e_k$). Thus, $x_{\alpha,j}+x_{-\alpha,j}=x_{\alpha,j}+n(x_{\alpha,j})=[x_{\beta,t}, e_k]+[x_{-\beta,t}, f_k]$.

Next, note that $x_{\beta,t}+x_{-\beta,t}\in M$ and $e_k+f_k\in M$, so that $[x_{\beta,t}+x_{-\beta,t}, e_k+f_k]=x_{\alpha,j}+x_{-\alpha,j}+[x_{\beta,t}, f_k]+[x_{-\beta,t}, e_k]\in M$. But the element $[x_{\beta,t}, f_k]+[x_{-\beta,t}, e_k]$ is in $S_+\cap(S_{\beta-\alpha_k}\oplus S_{-\langle\beta-\alpha_k\rangle})$, and hence by Lemma 2.1 is an \mathbf{R} -linear combination

of the elements $x_{\gamma,s} + x_{-\gamma,s}$ for $s \in \{1, \dots, n_\gamma\}$ and $\gamma = \beta - \alpha_k$. Thus, by induction $[x_{\beta,l}, f_k] + [x_{-\beta,l}, e_k]$ is in M . It now follows that $x_{\alpha,j} + x_{-\alpha,j} \in M$, as desired. \square

LEMMA 2.3. *The elements ih_j , $1 \leq j \leq l$, together with the elements $i(x_{\alpha,k} - x_{-\alpha,k})$ for $\alpha \in \Gamma^+$, $k \in \{1, \dots, n_\alpha\}$ form a basis of iS .*

PROOF. It is enough to show that the elements h_j , $1 \leq j \leq l$, together with the elements $x_{\alpha,k} - x_{-\alpha,k}$, for $\alpha \in \Gamma^+$, $k \in \{1, \dots, n_\alpha\}$, span S_- . Let M be the subspace spanned by these elements and note that h_j , $1 \leq j \leq l$, $x_{\alpha,k} + x_{-\alpha,k}$, $x_{\alpha,k} - x_{-\alpha,k}$, for $\alpha \in \Gamma^+$, $k \in \{1, \dots, n_\alpha\}$ form a basis of S , so that $S = S_+ \oplus M$. Since we also have $S = S_+ \oplus S_-$ and $M \subseteq S_-$, it follows that $M = S_-$. \square

PROPOSITION 2.4. S_C is generated by the $3l$ elements x_j, y_j, z_j , $1 \leq j \leq l$.

PROOF. $1/2[x_j, z_j] = 1/2[e_j + f_j, ih_j] = (i/2)(2e_j - 2f_j) = y_j$, $1 \leq j \leq l$. Thus, letting M be the subalgebra of S_C generated by x_j, z_j , $1 \leq j \leq l$, we have that $y_j \in M$, $1 \leq j \leq l$. By Lemma 2.1 and Lemma 2.3 we know that S_C has basis ih_j , $1 \leq j \leq l$, $(x_{\alpha,k} + x_{-\alpha,k})$, $i(x_{\alpha,k} - x_{-\alpha,k})$, for $\alpha \in \Gamma^+$, $k \in \{1, \dots, n_\alpha\}$; hence it is enough to show that these elements are in M . By Lemma 2.2 we have that $(x_{\alpha,k} + x_{-\alpha,k}) \in M$ for $\alpha \in \Gamma^+$, $k \in \{1, \dots, n_\alpha\}$. It is clear that $iH_S \subseteq M$. Next, let $\alpha \in \Gamma^+$, $k \in \{1, \dots, n_\alpha\}$ and choose $h \in H_S$ such that $\alpha(h) \neq 0$, (this is possible since (A_{ij}) is not Euclidean). Then we get $ih \in M$ and $x_{\alpha,k} + x_{-\alpha,k} \in M$ so that $[x_{\alpha,k} + x_{-\alpha,k}, ih] = \alpha(h)(i(x_{\alpha,k} - x_{-\alpha,k})) \in M$. It follows that $i(x_{\alpha,k} - x_{-\alpha,k}) \in M$. \square

DEFINITION 2.5. Let $j, k \in \{1, \dots, l\}$, $j \neq k$ and let $s, t \in \mathbf{Z}$.

We define the integer $C_{s,t}^{(j,k)}$ as follows:

$C_{0,0}^{(j,k)} = 1$, $C_{s,t}^{(j,k)} = 0$ if either $s < 0$, $t < 0$, or if $t > s$. Otherwise $C_{s,t}^{(j,k)}$ is defined inductively by $C_{s,t}^{(j,k)} = C_{s-1,t-1}^{(j,k)} + (s-1)[A_{kj} + (s-2)]C_{s-2,t}^{(j,k)}$. Note that $C_{s,s}^{(j,k)} = 1$ for all $s \geq 0$, and that $C_{s,t}^{(j,k)} = 0$ if $(-1)^s \neq (-1)^t$.

PROPOSITION 2.6. *The elements x_j, y_i, z_j for $1 \leq j \leq l$ satisfy the following relations:*

$$F_1 \quad y_j = 1/2[x_j, z_j], \quad 1 \leq j \leq l,$$

$$F_2 \quad [x_j, z_k] = A_{kj}y_j, \quad 1 \leq j, k \leq l,$$

$$F_3 \quad [y_j, z_k] = -A_{kj}x_j, \quad 1 \leq j, k \leq l,$$

$$F_4 \quad [z_j, z_k] = 0, \quad 1 \leq j, k \leq l,$$

$$F_5 \quad [x_j, x_k] + [y_j, y_k] = 0, \quad 1 \leq j, k \leq l,$$

$$F_6 \quad [x_j, y_k] + [x_k, y_j] = -4\delta_{jk}z_j, \quad 1 \leq j, k \leq l,$$

$$F_7 \quad e_j(ad e_k)^{2n} + f_j(ad f_k)^{2n} = \sum_{t=0}^n (-1)^{n-t} C_{2n, 2t}^{(j, k)} x_j(ad x_k)^{2t},$$

$$\text{for } n \geq 0, 1 \leq j, k \leq l, j \neq k,$$

$$F_8 \quad e_j(ad e_k)^{2n+1} + f_j(ad f_k)^{2n+1} = \sum_{t=0}^n (-1)^{n-t} C_{2n+1, 2t+1}^{(j, k)} x_j(ad x_k)^{2t+1},$$

$$\text{for } n \geq 0, 1 \leq j, k \leq l, j \neq k,$$

$$F_9 \quad i(e_j(ad e_k)^{2n} - f_j(ad f_k)^{2n}) = \sum_{t=0}^n (-1)^{n-t} C_{2n, 2t}^{(j, k)} y_j(ad x_k)^{2t},$$

$$\text{for } n \geq 0, 1 \leq j, k \leq l, j \neq k,$$

$$F_{10} \quad i(e_j(ad e_k)^{2n+1} - f_j(ad f_k)^{2n+1}) = \sum_{t=0}^n (-1)^{n-t} C_{2n+1, 2t+1}^{(j, k)} y_j(ad x_k)^{2t+1},$$

$$\text{for } n \geq 0, 1 \leq j, k \leq l, j \neq k.$$

In particular, these relations are a consequence of the definitions and the relations (i)-(iv) of Section 1.

PROOF. The relations F_1 through F_6 are easy to establish so we do F_7 and F_8 together, by induction on n . In doing this we write $C_{s, t}$ for $C_{s, t}^{(j, k)}$. Also, we will use the following well known formulas.

$$e_j(ad e_k)^t(ad f_k) = (tA_{kj} + t(t-1))e_j(ad e_k)^{t-1} \quad \text{for } t \geq 1,$$

$$f_j(ad f_k)^t(ad e_k) = (tA_{kj} + t(t-1))f_j(ad f_k)^{t-1} \quad \text{for } t \geq 1.$$

When $n=0$, $e_j(ad e_k)^{2n} + f_j(ad f_k)^{2n} = e_j + f_j = x_j$ while

$$\sum_{t=0}^n (-1)^{n-t} C_{2n, 2t} x_j(ad x_k)^{2t} = C_{0, 0} x_j = x_j,$$

so F_7 holds when $n=0$. By definition, $[e_j, e_k] + [f_j, f_k] = [x_j, x_k]$ for $j \neq k$, so F_8 holds when $n=0$. Also F_7 holds when $n=1$, since we have

$$\begin{aligned} & e_j(ad e_k)^2 + f_j(ad f_k)^2 \\ &= [e_j(ad e_k) + f_j(ad f_k), e_k + f_k] - [f_j(ad f_k), e_k] - [e_j(ad e_k), f_k] \\ &= x_j(ad x_k)^2 - A_{kj}f_j - A_{kj}e_j \\ &= C_{2, 2} x_j(ad x_k)^2 - C_{2, 0} x_j. \end{aligned}$$

Next, assume that $m \geq 1$ and that F_7 holds when $n \leq m$, and that F_8 holds when $n \leq m-1$. We show F_8 holds when $n=m$. We have that

$$\begin{aligned}
& e_j(ad e_k)^{2m+1} + f_j(ad f_k)^{2m+1} \\
&= [e_j(ad e_k)^{2m} + f_j(ad f_k)^{2m}, e_k + f_k] \\
&\quad - [e_j(ad e_k)^{2m}, f_k] - [f_j(ad f_k)^{2m}, e_k] \\
&= \left[\sum_{t=0}^m (-1)^{m-t} C_{2m, 2t} x_j (ad x_k)^{2t}, x_k \right] \\
&\quad - (2mA_{kj} + 2m(2m-1))e_j(ad e_k)^{2m-1} - (2mA_{kj} + 2m(2m-1))f_j(ad f_k)^{2m-1} \\
&= \sum_{t=0}^m (-1)^{m-t} C_{2m, 2t} x_j (ad x_k)^{2t+1} \\
&\quad - (2mA_{kj} + 2m(2m-1)) \sum_{t=0}^{m-1} (-1)^{m-1-t} C_{2m-1, 2t+1} x_j (ad x_k)^{2t+1}.
\end{aligned}$$

This equals

$$\begin{aligned}
& \sum_{t=0}^{m-1} \{ (-1)^{m-t} C_{2m, 2t} - (2mA_{kj} + 2m(2m-1))(-1)^{m-1-t} C_{2m-1, 2t+1} \} x_j (ad x_k)^{2t+1} \\
& \quad + C_{2m, 2m} x_j (ad x_k)^{2m+1}.
\end{aligned}$$

Now

$$C_{2m, 2t} + 2m(A_{kj} + (2m-1))C_{2m-1, 2t-1} = C_{2m+1, 2t+1}$$

and

$$\begin{aligned}
& (-1)^{m-t} C_{2m, 2t} - (2mA_{kj} + 2m(2m-1))(-1)^{m-1-t} C_{2m-1, 2t+1} \\
&= (-1)^{m-t} (C_{2m, 2t} + (2mA_{kj} + 2m(2m-1))C_{2m-1, 2t+1}) \\
&= (-1)^{m-t} (C_{2m, 2t} + 2m(A_{kj} + (2m-1))C_{2m-1, 2t+1}) \\
&= (-1)^{m-t} C_{2m+1, 2t+1}.
\end{aligned}$$

Thus, we have that

$$e_j(ad e_k)^{2m+1} + f_j(ad f_k)^{2m+1} = \sum_{t=0}^m (-1)^{m-t} C_{2m+1, 2t+1} x_j (ad x_k)^{2t+1},$$

as desired.

Next, one lets $m \geq 1$ and assumes F_8 holds when $n \leq m$ and that F_7 holds when $n \leq m$ and shows that F_7 holds when $n = m+1$. This is similar to the above and so is omitted. In the same way F_9 and F_{10} can be shown to hold. \square

DEFINITION 2.7. Let $F\mathcal{L} = F\mathcal{L}(X_j, Y_j, Z_j | 1 \leq j \leq l)$ be the free Lie algebra over \mathbf{R} generated by the $3l$ -symbols $X_j, Y_j, Z_j, 1 \leq j \leq l$. Recall that (A_{ij}) is a fixed $l \times l$ indecomposable Cartan matrix which is not of Euclidean type. Let J denote the ideal of $F\mathcal{L}$ generated by the following elements: $(R_1 - R_{10})$

$$R_1: Y_j - 1/2[X_j, Z_j], \quad 1 \leq j \leq l,$$

$$R_2: [X_j, Z_k] - A_{kj}Y_j, \quad 1 \leq j, k \leq l,$$

$$R_3: [Y_j, Z_k] + A_{kj}X_j, \quad 1 \leq j, k \leq l,$$

$$R_4: [Z_j, Z_k], \quad 1 \leq j, k \leq l,$$

$$R_5: [X_j, X_k] + [Y_j, Y_k], \quad 1 \leq j, k \leq l,$$

$$R_6: [X_j, X_k] + [X_k, Y_j] + 4\delta_{jk}Z_j, \quad 1 \leq j, k \leq l.$$

Next, let $j, k \in \{1, \dots, l\}$, $j \neq k$. Let $m = -A_{kj} + 1$. If m is even we put $n = m/2$. Then

$$R_7: \sum_{t=0}^n (-1)^{n-t} C_{2n, 2t}^{(j,k)} X_j (ad X_k)^{2t}, \text{ and}$$

$$R_9: \sum_{t=0}^n (-1)^{n-t} C_{2n, 2t}^{(j,k)} Y_j (ad X_k)^{2t}.$$

If m is odd we put $n = \frac{m-1}{2}$. Then

$$R_8: \sum_{t=0}^n (-1)^{n-t} C_{2n+1, 2t+1}^{(j,k)} X_j (ad X_k)^{2t+1}, \text{ and}$$

$$R_{10}: \sum_{t=0}^n (-1)^{n-t} C_{2n+1, 2t+1}^{(j,k)} Y_j (ad X_k)^{2t+1}.$$

Finally, we let $\bar{L}_C = \frac{F\mathcal{L}}{J}$ and let \bar{L}_C be the complexification of \bar{L}_C . Let $E_j, F_j, H_j \in \bar{L}_C$ be defined by

$$E_j = 1/2(X_j - iY_j), \quad F_j = 1/2(X_j + iY_j),$$

and

$$H_j = -iZ_j, \quad 1 \leq j \leq l.$$

PROPOSITION 2.8. *The algebras \bar{L}_C and \bar{L}_C are isomorphic. In particular, R_1 - R_{10} provides a presentation of \bar{L}_C . Moreover, \bar{L}_C has a unique maximal ideal and the corresponding simple factor is isomorphic to \mathcal{L}_C .*

PROOF. We first note that formulas (i)-(iv) of section 1 hold in \bar{L}_C . Indeed, R_4 implies that $[H_j, H_k] = 0$ for $1 \leq j, k \leq l$. Now

$$\begin{aligned} [E_k, H_j] &= 1/2[X_k - iY_k, -iZ_j] \\ &= -i/2[X_k, Z_j] - 1/2[Y_k, Z_j] \\ &= -i/2 A_{jk} Y_k + 1/2 A_{jk} X_k \quad (\text{by } R_2 \text{ and } R_3) \\ &= A_{jk}(1/2(X_k - iY_k)) = A_{jk} E_k, \end{aligned}$$

as desired. Similarly, one finds that $[F_k, H_j] = -A_{jk}F_k$ and that $[E_j, F_k] = \delta_{jk}H_j$.

By Proposition 2.6 we obtain a Lie algebra homomorphism ϕ from \bar{L}_C onto the subalgebra of \bar{L}_C generated by the elements $x_j, y_j, z_j, 1 \leq j \leq l$; and by Proposition 2.4 this is the algebra \bar{L}_C . Thus, ϕ is a surjective homomorphism of \bar{L}_C onto \bar{L}_C .

Since the relations (i)-(iv) of Section 1 hold in \bar{L}_C we get a Lie algebra homomorphism $\tilde{\Psi}$ from the universal Kac-Moody algebra \bar{L}_C to \bar{L}_C such that $\tilde{\Psi}(e_j) = E_j, \tilde{\Psi}(f_j) = F_j, \text{ and } \tilde{\Psi}(h_j) = H_j, 1 \leq j \leq l$. Now formulas F_1 - F_{10} hold in \bar{L}_C thanks to Proposition 2.6. Thus, since R_7 - R_{10} hold in \bar{L}_C we see that $E_j(ad E_k)^{-A_{kj}+1} = 0 = F_j(ad F_k)^{-A_{kj}+1}$. It follows that $\tilde{\Psi}$ induces a homomorphism Ψ of \bar{L}_C to \bar{L}_C . Clearly, $\Psi(x_j) = X_j, \Psi(y_j) = Y_j, \text{ and } \Psi(z_j) = Z_j, 1 \leq j \leq l$, so that $\Psi(\bar{L}_C) = \bar{L}_C$. Finally, it is clear that $\phi \circ \tilde{\Psi} = id_{\bar{L}_C}$ and $\tilde{\Psi} \circ \phi = id_{\bar{L}_C}$, so that \bar{L}_C and \bar{L}_C are isomorphic. As in Section 1 it is clear that \bar{L}_C has a unique maximal ideal with the corresponding simple factor being isomorphic to \mathcal{L}_C . \square

Assume now that (A_{ij}) is one of the 9 types of $l \times l$ indecomposable finite Cartan matrices. Then by Serre's Theorem $\mathcal{L}_C = \bar{L}_C$ is the split simple Lie algebra of type (A_{ij}) over \mathbf{C} . Let (\cdot, \cdot) denote the Killing form \mathcal{L}_C and let n be as in Section 1. As in [6 pg. 147-149] a compact subalgebra C of \mathcal{L}_C has basis $i h_j, 1 \leq j \leq l, e_\alpha + e_{-\alpha}, i(e_\alpha - e_{-\alpha})$; for $\alpha \in \mathcal{A}$, (the root system of \mathcal{L}_C) and $e_\alpha \in \mathcal{L}_\alpha$ is chosen such that $n(e_\alpha) = e_{-\alpha}$ and $(e_\alpha, e_{-\alpha}) = -1$, for all $\alpha \in \mathcal{A}$.

We are going to show that $x_j, y_j \in C, 1 \leq j \leq l$. As usual, h_α denotes the element in H satisfying $\alpha(h) = (h_\alpha, h)$ for all $h \in H, \alpha \in \mathcal{A}$. Then $[e_j, f_j] = h_j$ and $\alpha_j(h_j) = 2$ imply that

$$(e_j, f_j) = \frac{-2}{(\alpha_j, \alpha_j)}, \quad 1 \leq j \leq l,$$

since

$$\begin{aligned} 2 &= (h_j, h_{\alpha_j}) = ([e_j, f_j], h_{\alpha_j}) = (e_j, [f_j, h_{\alpha_j}]) \\ &= -\alpha_j(h_{\alpha_j})(e_j, f_j) = (-\alpha_j, \alpha_j)(e_j, f_j). \end{aligned}$$

Let

$$\lambda_j = \left(\frac{(\alpha_j, \alpha_j)}{2} \right)^{1/2} \in \mathbf{R}, \quad 1 \leq j \leq l.$$

Then $(\lambda_j e_j, \lambda_j f_j) = -1$ so that, as part of our basis of C , we can take $e_{\alpha_j} = \lambda_j e_j, e_{-\alpha_j} = \lambda_j f_j$. Then $e_{\alpha_j} + e_{-\alpha_j} = \lambda_j(e_j + f_j) \in C$, hence $x_j = \lambda_j^{-1}(e_{\alpha_j} + e_{-\alpha_j}) \in C$, and similarly $y_j \in C$ for $1 \leq j \leq l$. It follows that $\mathcal{L}_C \subseteq C$. But $(\mathcal{L}_C)_C = \mathcal{L}_C = C_C$ so that $\mathcal{L}_C = C$. This completes the proof of the following result.

THEOREM 2.9. *Let (A_{ij}) be an $l \times l$ indecomposable Cartan matrix of finite type. Then the Lie algebra generated by the $3l$ elements $X_j, Y_j, Z_j, 1 \leq j \leq l$, satisfying the relations R_1 - R_{10} is the compact simple Lie algebra of type (A_{ij}) .*

One consequence of this result is the following Corollary.

COROLLARY 2.10. *Let (A_{ij}) be an $l \times l$ indecomposable Cartan matrix of finite type. Then there is one and only one simple Lie algebra generated by $3l$ elements $X_j, Y_j, Z_j, 1 \leq j \leq l$, satisfying the relations R_1 - R_6 . Moreover, this algebra is compact.*

PROOF. The algebra $\tilde{\mathcal{L}}_c$ satisfies R_1 - R_6 , and has a unique simple factor. This factor is compact. □

References

- [1] Berman, S., On the construction of simple Lie algebras, *J. Algebra*, **27** (1973), 158-183.
- [2] Berman, S., Isomorphisms and Automorphisms of universal Heffalump Lie algebras, *Proc. A. M. S.*, **65** (1977), 29-34.
- [3] Berman, S. and Moody, R. V., Lie algebra multiplicities, *Proc. A. M. S.*, **76** (1979), 223-228.
- [4] Garland, H. and Lepowsky, J., Lie algebra homology and the Macdonald-Kac formulas, *Invent. Math.*, **34** (1976), 37-76.
- [5] Humphreys, J. E., *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag, (1972), New York.
- [6] Jacobson, N., *Lie Algebras*, Wiley Interscience, (1962), New York.
- [7] Kac, V. G., Simple irreducible graded Lie algebras of finite growth, *Math. U.S.S.R. —Izv.*, **2** (1968), 1271-1311.
- [8] Moody, R. V., A new class of Lie algebras, *J. Algebra*, **10** (1968), 211-230.
- [9] Serre, J. P., *Algèbres de Lie semi-simples complexes*, W. A. Benjamin, (1966), New York.

Department of Mathematics
University of Saskatchewan
Saskatoon, Saskatchewan
CANADA S7N 0W0