

## ON $q$ -PSEUDOCONVEX OPEN SETS IN A COMPLEX SPACE

By

Edoardo BALLICO

In a series of (perhaps not widely known) papers T. Kiyosawa ([1], [2], [3], [4], [5]) introduced and developed the notion of Levi  $q$ -convexity. Here we show how to use this notion to improve one of his results ([2] Th. 2) (for a different extension, see [7]). To state and prove our results, we recall few definitions.

Let  $M$  be a complex manifold of dimension  $n$ ; a real  $C^2$  function  $u$  on  $M$  is said to be  $q$ -convex at a point  $P$  of  $M$  if the hermitian form  $L(u)(P) = \sum_{i,j} \left( \frac{\partial^2 u}{\partial z_i \partial \bar{z}_j} \right) \times (P) a_i \bar{a}_j$ ,  $z_1, \dots, z_n$  local coordinates around  $P$ , has at least  $n-q+1$  strictly positive eigenvalues; we say that  $u$  is Levi  $q$ -convex at  $P$  if either  $(du)_P = 0$  and  $u$  is  $q$ -convex at  $P$  or  $(du)_P \neq 0$  and the restriction of  $L(u)(P)$  to the hyperplane  $\left\{ \sum_i \left( \frac{\partial u}{\partial z_i} \right) (P) a_i = 0 \right\}$  has at least  $n-q$  strictly positive eigenvalues. Let  $X$  be a complex space,  $A \in X$ , and  $f: X \rightarrow \mathbf{R}$  a  $C^2$  function; we say that  $f$  is  $q$ -convex (or Levi  $q$ -convex) at  $A$  if there is a neighborhood  $V$  of  $A$  in  $X$ , a closed embedding  $p: V \rightarrow U$  with  $U$  open subset of an euclidean space, a  $C^2$  function  $u$  on  $U$  such that  $f|V = u \circ p$  and  $u$  is  $q$ -convex (or respectively Levi  $q$ -convex) at  $P = p(A)$ . It is well-known that a  $q$  convex function is Levi  $q$  convex and that both notions do not depend upon the choice of charts and local coordinates; for any fixed choice of charts and local coordinates we will call  $L(u)(P)$  the Levi form of  $u$  at  $P$  and of  $f$  at  $A$ .

An open subset  $D$  of a complex space  $X$  is said to have regular Levi  $q$ -convex boundary if we can take a covering  $\{V_i\}$  of a neighbourhood of the boundary  $bD$  of  $D$  with closed embeddings  $p_i: V_i \rightarrow U_i$ ,  $U_i$  open in an euclidean space and  $C^2$  functions  $f_i$  on  $U_i$  with  $V_i \cap D = \{x \in V: f_i \circ p_i(x) < 0\}$  and such that if  $x \in V_i \cap V_j$ , there is a neighborhood  $A$  of  $x$  in  $V_i \cap V_j$  such that on  $A(f_i \circ p_i)|_A = f_{ij}(f_j \circ p_j)|_A$  with  $f_{ij} > 0$ ,  $f_{ij} \in C^2$  on  $A$ . The last condition is always satisfied for a domain  $D$  defined locally by Levi  $q$ -convex functions  $s_i$  if the set of points of  $bD$  at which either  $ds_i$  vanishes or  $X$  is singular is discrete.

A complex space  $X$  is called  $q$ -complete if it has a  $C^2$   $q$ -convex exhausting function  $f$ ; if  $f$  is both  $q$ -convex and weakly plurisubharmonic,  $X$  is called very

strongly  $q$ -convex (in the sense of T. Ohsawa [6]).

Now we can state our results.

**THEOREM.** *Let  $D$  be a regular Levi  $q$ -convex open subset of a complex space  $X$ . Then there exist a neighbourhood  $V$  of the boundary  $bD$  and a  $q$ -convex real function  $t$  such that  $D \cap V = \{x \in V : t(x) < 0\}$ .*

**COROLLARY.** *Let  $X$  be a very strongly  $q$ -convex space and  $D$  an open subset of  $X$  with regular Levi  $q$ -convex boundary. Then  $D$  is  $q$ -complete.*

Compare the corollary with the main result in [7].

**PROOF** of the theorem. Note that the proof of [2] Theorem 2 goes on verbatim even if  $D$  is not relatively compact in  $X$ . The quoted result gives a neighbourhood  $W$  of  $bD$  and a Levi  $q$ -convex function  $g$  in  $W$  such that  $D \cap W = \{x \in W : g(x) < 0\}$ . Consider a strictly positive real function  $v$  on  $W$ . Set  $t = ge^{vg}$ . Since  $g$  vanishes on  $bD$ , the Levi form of  $t$  at a point  $y$  in  $bD$  is proportional to the Levi form at  $y$  of  $e^{cg}$ , with  $c = g(y)$ . Hence if  $g(y)$  is sufficiently high,  $t$  is  $q$ -convex at  $y \in bD$  ([3] Prop. 2 or [5] Lemma 2); how big must be  $g(y)$  depend only from the eigenvalues of the Levi form of  $g$  at  $y$ ; hence the same constant works also in a neighbourhood of  $y$ . Let  $\{V_n\}$ ,  $\{U_n\}$  be locally finite coverings of  $W$  with  $V_n$  relatively compact in  $U_n$ ,  $\{U_n\}$  fine enough (in particular with local charts on which  $g$  may be found constants  $c_n > 0$  such that if  $u < c_n$  on  $V_n$ ,  $t = ge^{ug}$  is  $q$ -convex at every point of  $bD$ , hence in a neighbourhood  $V$  of  $bD$ . Q. E. D.

**PROOF** of the corollary. By the theorem we may find an open neighbourhood  $V$  of  $bD$  and a real  $C^2$   $q$ -convex function  $f$  on  $V$  such that  $V \cap D = \{x \in V : f(x) < 0\}$ . Let  $W$  be an open neighbourhood of  $bD$  with closure contained in  $V$ . Note that the function  $s := -f^{-1}$  is  $q$ -convex on  $V \cap D$  and goes to infinity near  $bD$ . Let  $u$  be a real non-negative  $C^2$  function on  $U$  with support contained in  $V \cap D$ ,  $u = 1$  in  $W \cap D$ . We may consider  $us$  as a function on  $D$  setting  $(us)(x) = 0$  if  $x \notin V$ . Take an exhaustive, positive,  $q$ -convex function  $h$  on  $X$ . Take an increasing sequence  $\{K_n\}$  or compact subset of  $X$ , with union  $X$  and a sequence  $\{c_n\}$  of strictly positive real numbers. Take a  $C^2$  function  $b : \mathbf{R} \rightarrow \mathbf{R}$  with  $b(t) = 0$  for  $t \leq -1$ ,  $b(t) \geq c_j$  for  $j \leq t \leq j+1$  and  $b'(t) > 0$  for  $t > -1$ . Set

$$g(t) = \int_{-\infty}^t b(x) dx$$

and set  $F = g \circ h$ . For every  $P \in X$  and any choice of local coordinates, we have  $L(F)(P) \geq b(h(P))L(g)(P)$ . Hence we may choose the constants  $c_j$  with  $c \geq j$  and such that  $F + s$  is  $q$ -convex on  $(D \setminus W) \cap K_j$  for every  $j$ . Since  $F$  is plurisubharmonic,  $F + s$  is  $q$ -convex on  $D$ . If  $\{x_n\}$  is a sequence in  $D$  without accumula-

tion points in  $X$ , then  $\{F(x_n)\}$  and  $\{F(x_n)+s(x_n)\}$  are unbounded on  $\{x_n\}$ . The function  $s$  is unbounded on every sequence of points in  $D$  converging to a point in  $\partial D$ , hence  $F+s$  is an exhaustion function on  $D$ . Q. E. D.

### References

- [1] Kiyosawa, T., On the extension of holomorphic functions for a complex manifold, Sci. Rep. Tokyo Kyoiku Daigaku, Sec. A **9** (1968), 193-198.
- [2] Kiyosawa, T., On the Levi  $s$ -convex set in complex spaces, Sci. Rep. Tokyo Kyoiku Daigaku, Sec. A **10** (1969-1970), 168-177.
- [3] Kiyosawa, T., On the dimension of the cohomology for Levi  $q$ -convex spaces, Sci. Rep. Tokyo Kyoiku Daigaku, Sec. A **10** (1969-1970), 223-233.
- [4] Kiyosawa, T., On the Levi  $q$ -convex spaces, Sci. Rep. Tokyo Kyoiku Daigaku, Sec. A **10** (1969-1970), 241-244.
- [5] Kiyosawa, T., Some equivalent relations of  $q$ -convex domains in  $\mathbb{C}^n$ , Sci. Rep. Tokyo Kyoiku Daigaku, Sec. A **10** (1969-1970), 245-249.
- [6] Ohsawa, T., A reduction theorem for cohomology groups of very strongly  $q$ -convex Kähler manifolds, Invent. Math. **63** (1981), 335-354.
- [7] Vigna Suria, G.,  $q$ -pseudoconvex and  $q$ -complete domains, Compositio Math. **53** (1984), 105-111.

Scuola Normale Superiore  
56100 Pisa  
Italy