

1 **Title:**

2 Transcription regulation of rRNA gene by a multi-functional nucleolar protein,  
3 B23/nucleophosmin through its histone chaperone activity

4  
5 **Authors:**

6 Kensaku Murano,<sup>1</sup> Mitsuru Okuwaki,<sup>1,2</sup> Miharuru Hisaoka,<sup>1</sup> and Kyosuke Nagata<sup>1\*</sup>

7  
8 **Authors' affiliations:** <sup>1</sup>Department of Infection Biology, Graduate School of Comprehensive  
9 Human Sciences and Institute of Basic Medical Sciences, University of Tsukuba, 1-1-1  
10 Tennodai, Tsukuba 305-8575, <sup>2</sup>PRESTO, Japan Science and Technology Agency, 4-1-8  
11 Honcho, Kawaguchi, 322-0012, *Japan*

12  
13 **\*Corresponding author:** Department of Infection Biology, Graduate School of  
14 Comprehensive Human Sciences and Institute of Basic Medical Sciences, University of  
15 Tsukuba, 1-1-1 Tennodai, Tsukuba 305-8575, *Japan*; Phone and Fax +81-29-853-3233;  
16 E-mail: knagata@md.tsukuba.ac.jp

17  
18 **Running title:** B23-mediated transcription regulation of rRNA gene

19  
20 **Word count for the Materials and Methods section:** 1309 words

21 **Word count for the Introduction, Results, and Discussion sections:** 5731 words

1 **ABSTRACT**

2

3 **It is well established that the transcription rate of the rRNA gene is closely**

4 **associated with profound alterations in the cell growth rate. Regulation of the rRNA**

5 **gene transcription is likely to be dependent on the dynamic conversion of the chromatin**

6 **structure. Previously, we identified B23/nucleophosmin, a multi-functional nucleolar**

7 **phosphoprotein, as a component of Template Activating Factor-III that remodels the**

8 **chromatin-like structure of the adenovirus genome complexed with viral basic proteins.**

9 **It is also shown that B23 has the histone chaperone activity. Here, we examined the**

10 **effect of B23 on the rRNA gene transcription. B23 was found to be associated with the**

11 **rRNA gene chromatin. siRNA-mediated down-regulation of the B23 expression level**

12 **resulted in reduction of the transcription rate of the rRNA gene. We constructed a B23**

13 **mutant termed B23 $\Delta$ C, which lacks the domain essential for the histone chaperone**

14 **activity and inhibited the histone binding activity of B23 in a dominant-negative manner.**

15 **Expression of B23 $\Delta$ C decreased the rRNA gene transcription and the rate of cell**

16 **proliferation. These suggest that B23 is involved in the transcription regulation of the**

17 **rRNA gene as a nucleolar histone chaperone.**

18

19

# 1 INTRODUCTION

2

3 To ensure the protein synthesis level, growing cells require continuous ribosome  
4 supply by synthesis of ribosomal proteins and ribosomal RNA (rRNA). The transcription  
5 rate of the rRNA gene by RNA polymerase I (Pol I) is closely correlated with the cell growth  
6 rate (19). In eukaryotic cells, rRNA genes are found in multiple tandem-arrayed copies in  
7 the nucleolus, known as nucleolar organizer region (NOR). The tandem rRNA gene repeats  
8 exist in two distinct types of chromatin, an 'open' one permissive for transcription and a  
9 'closed' one under the transcriptionally repressive state (6). Several epigenetic  
10 characteristics including differential DNA methylation and specific histone modification also  
11 differ between potentially active and inactive rRNA genes (18), so that the epigenetic control  
12 mechanism is important for the transcription regulation of the rRNA gene. For instance, it is  
13 reported that the nucleolar remodeling complex (NoRC) is a member of ATP-dependent  
14 chromatin remodeling factors, and consists of ATPase Snf2h and TIP5 localized in nucleoli  
15 (54). It is suggested that NoRC plays an active role in nucleosome dynamics at the rRNA  
16 gene promoter in an ATP-dependent manner, and specific nucleosome positions determine the  
17 transcriptional readout of the rRNA gene (33). NoRC interacts with histone deacetylases,  
18 histone methyltransferases, and DNA methyltransferases, and targets these enzymatic  
19 activities to the rRNA gene promoter to form heterochromatin (18). In addition to histone  
20 modification enzymes and ATP-dependent chromatin remodeling factors, it is shown that  
21 nucleolin, one of histone chaperones localized in the nucleolus, enhances the activity of  
22 chromatin remodeling machineries, SWI/SNF and ACF, and promotes the transcription from  
23 nucleosomal template *in vitro* (1). Experiments using siRNA for nucleolin showed that

1 nucleolin is involved in the transcription of the rRNA gene *in vivo*. Moreover, chromatin  
2 immunoprecipitation (ChIP) experiments showed that nucleolin is associated with the  
3 chromatin around the 18S rRNA gene (48).

4 To identify a factor involved in structural change of chromatin, we have taken  
5 advantage of the adenovirus (Ad) DNA complexed with viral basic core proteins, called Ad  
6 core (28, 36, 44). The viral genome is packed around core protein VII hexamer, and a unit  
7 of the viral nucleosome is bridged by core protein V (8, 55). Because of the chromatin-like  
8 structure of Ad core, the access to the viral genome of the factors required for the replication  
9 and transcription is restricted. Using a *cell-free* DNA replication system using Ad core as a  
10 template, we have identified three host factors termed Template Activating Factor (TAF)-I,  
11 TAF-II, and TAF-III. TAF-I and TAF-II are found to be identical to the leukemia-associated  
12 gene product SET (58) and Nucleosome Assembly Protein-1 (NAP-1) (28), respectively.  
13 The major component of TAF-III is B23/nucleophosmin, a nucleolar phosphoprotein (44).  
14 Recombinant B23 induces the structural alteration of Ad core, thereby stimulating the Ad core  
15 DNA replication (44, 49). TAF-I and NAP-1 bind to core histones directly and transfers  
16 them to naked DNA (25, 35). TAF-I is not only involved in the replication and transcription  
17 of Ad core (20, 21, 37), but also suggested to be involved in regulation of chromatin structure  
18 in eukaryotic cells (14, 27). In addition, TAF-I promotes vitamin D<sub>3</sub> receptor- and  
19 Gal4-VP16-dependent transcription from chromatin template *in vitro* (14), and stimulates the  
20 transcription of a sub-set of genes in HeLa cells in a histone acetylation-independent manner  
21 (27). TAF-I also plays an important role in transcription regulated by transcription factors  
22 such as MLL (mixed lineage leukemia), Sp1, and KLF5 (39, 53, 56).

23 We have demonstrated that B23 binds to core histones directly and transfers them to

1 naked DNA (45) as do TAF-I (35) and NAP-1 (25), and the acidic regions of B23 are  
2 important for its activity (45). The amino-terminal domain of B23 shows significant  
3 similarity to nucleoplasmin, which is the first identified histone chaperone (32). Recently, it  
4 is indicated that B23 is acetylated, and acetylated B23 has the enhanced histone chaperone  
5 activity and stimulates *in vitro* transcription from chromatin templates by RNA polymerase II  
6 in the presence of p300 (57). The crystal structure of the N-terminal domain of NO38, a  
7 *Xenopus* homolog of human B23, indicates that NO38 forms a pentamer and each pentamer  
8 further assembles into a decamer as does nucleoplasmin (10, 41). These studies suggest that  
9 oligomer formation of nucleoplasmin family proteins may be important for their histone  
10 chaperone activity. Based on its nucleolar localization and biochemical properties, it is  
11 reasonably assumed that B23 is involved in regulation of the chromatin structure around the  
12 rRNA gene through its histone chaperone activity.

13 B23 is also termed as nucleophosmin, numatrin, or NPM1, and was initially  
14 identified as an abundant nucleolar phosphoprotein (47). These properties are similar to  
15 those of nucleolin, alternatively termed C23 (47). The expression level of B23 is  
16 up-regulated in association with growth factor-induced mitogenesis (12). A positive  
17 correlation has been reported among the quantity of B23 and nucleolin, the rate of Pol I  
18 transcription, and the cell proliferation rate (7). There exist two subtypes of B23 termed  
19 B23.1 and B23.2, the latter of which lacks the carboxyl-terminal region present in B23.1 (4).  
20 B23 is a multi-functional protein: It is suggested that B23 is involved in rRNA maturation  
21 through its nucleic acid binding and ribonuclease activities (9, 22). B23 is shown to play a  
22 critical role in centrosome duplication (43). Inactivation of B23 in mouse embryonic  
23 fibroblasts (MEFs) leads to unrestricted centrosome duplication and severe genomic

1 instability in addition to reduction of the cell proliferation rate (17). B23 is also involved in  
2 regulation of the cell cycle progression. In normal cells, B23 targets ARF in the nucleolus  
3 by binding to ARF through the Mdm2 binding domain of ARF, thus allowing Mdm2 to  
4 interact with p53 for degradation of p53. Upon exposure to genotoxic stresses, the  
5 interaction between B23 and ARF is abrogated, so that the stability of p53 is increased (2, 5,  
6 29). These indicate that B23 controls the cell cycle through the p53 pathway.

7           Here, we examined the effect of the histone chaperone activity of B23 on the rRNA  
8 gene transcription. We showed that down-regulation of B23 by siRNA reduces the  
9 transcription rate of the rRNA gene. Nucleolar chromatin immunoprecipitation analyses  
10 demonstrated that B23 binds to the chromatin around the rRNA gene promoter. In order to  
11 characterize the function of B23 as a histone chaperone in the nucleolus, we constructed  
12 deletion mutants lacking the histone binding activity. B23 $\Delta$ C, devoid of the functional  
13 acidic regions, formed a hetero-oligomer with B23 and inhibited the histone binding activity  
14 in a dominant-negative manner. Over-expression of B23 $\Delta$ C reduced the transcription level  
15 from the rRNA gene. These results altogether suggest that B23 regulates the transcription  
16 from the rRNA gene chromatin as a nucleolar histone chaperone.

17

# 1 MATERIALS AND METHODS

2

## 3 Cell culture and transfection

4 293T and *p53*<sup>-/-</sup> MEF cells were cultured at 37°C in DMEM supplemented with 10%  
5 fetal bovine serum (FBS). HeLa cells were cultured at 37°C in MEM supplemented with  
6 10% FBS. Transient transfection was performed using *TransIT*-293 (Mirus) for 293T cells,  
7 *TransIT*-LT1 (Mirus) for *p53*<sup>-/-</sup> MEF, and GeneJuice (Novagen) for HeLa cells. HeLa cells  
8 were transfected with the siRNA using Lipofectamine 2000 (Invitrogen).

9

## 10 Preparation of plasmids

11 pCHA-B23.1 was prepared as previously described (44). To generate pCHA-B23ΔC,  
12 a fragment of B23ΔC3 cDNA was prepared by digestion of pET14b-hB23ΔC3 (44) with *Nde*  
13 I and *Bam*H I. The fragment was then cloned in-frame to *Bst*E II- and *Bgl* II-digested pCHA  
14 (44). To generate pUC119-HurDNA, the human rRNA gene (-192~+703 relative to the  
15 transcription start site (+1)) was amplified by PCR using genomic DNA purified from HeLa  
16 cells as template. To amplify the human rRNA gene, two oligonucleotides,  
17 5'-CGCGGATCCTGTCCTTGGGTTGACCAGAG-3' and  
18 5'-CCGGAATTCGCAAGTCGACAACCACTGGA-3', were used as primers. The  
19 amplified human rRNA gene fragment was cloned into the *Bam*H I- and *Eco*R I-digested  
20 pUC119. For expression of Flag-tagged B23.1, B23.1 cDNA attached by the Flag-tag was  
21 excised from pBS-Flag-B23.1 (46) by *Bam*H I, and subcloned into *Bam*H I-digested pcDNA3  
22 vector (CLONTECH). To create pCAGGS-Flag-B23.1, a DNA fragment encoding  
23 Flag-tagged B23.1 was excised from pBS-Flag-B23.1 by *Bam*H I, and subcloned into *Bgl*

1 II-digested pCAGGS vector. To generate pET-14b-B23ΔA1, a DNA fragment of B23ΔA1  
2 was amplified by two-step PCR using four oligonucleotides,  
3 5'-GCTAGTTATTGCTCAGCGG-3' and 5'-AGTAGCTGTGGTGAAACTCTTAAGTATA-3',  
4 and 5'-TAATACGACTCACTATAGG-3' and  
5 5'-AGAGTTTCACCACAGCTACTAAGTGCTG-3', and pET-14b-B23.1 as a template, and  
6 cloned into the *Nde* I and *Bam*H I-digested pET-14b. To generate pET-14b-B23ΔA, a DNA  
7 fragment of B23ΔA was amplified by two-step PCR using four oligonucleotides,  
8 5'-GCTAGTTATTGCTCAGCGG-3' and  
9 5'-ACTTGCTGCTAAAGCGCCAGTGAAGAAA-3', and  
10 5'-TAATACGACTCACTATAGG-3' and 5'-CTGGCGCTTTAGCAGCAAGTTTTACTTT-3',  
11 and pET-14b-B23ΔA1 as a template, and cloned into the *Nde* I and *Bam*H I-digested pET-14b.  
12 To generate pBS-Flag-B23ΔA, a DNA fragment of B23ΔA was prepared by digestion of  
13 pET-14b-B23ΔA with *Nde* I and *Hind* III and cloned into the *Nde* I and *Hind* III-digested  
14 pBS-Flag. To generate pCAGGS-Flag- B23ΔA, a DNA fragment of Flag-B23ΔA was  
15 prepared by digestion of pBS-Flag-B23ΔA with *Bam*H I and cloned into *Bgl* II-digested  
16 pCAGGS.

17

## 18 **Antibodies**

19 Antibodies used in this study were as follows: Mouse monoclonal antibody for B23  
20 (Zymed) recognizes both endogenous B23.1 and B23ΔA proteins, but not B23ΔC protein.  
21 For immunofluorescence assays of B23, goat polyclonal anti-B23 antibody (Santa Cruz  
22 Biotechnology) was used; for histone H3, rabbit polyclonal antibody (Abcam); for acetylated  
23 histone H3, rabbit polyclonal antibody (Upstate); for TBP, rabbit polyclonal antibody (Santa



1 Cruz Biotechnology); for UBF, rabbit polyclonal antibody (F-9, Santa Cruz Biotechnology);  
2 for nucleolin, mouse monoclonal antibody (Santa Cruz Biotechnology); for fibrillarin, rabbit  
3 polyclonal antibody (Santa Cruz Biotechnology); for HA-tag, rabbit polyclonal antibody  
4 (MBL) or rat monoclonal antibody (3F10, Roche); for Flag-tag, mouse monoclonal antibody  
5 (M2, Sigma); for His-tag, mouse monoclonal antibody (Sigma); for  $\beta$ -actin, mouse  
6 monoclonal antibody (Sigma).

7

### 8 **Immunoprecipitation**

9           Cells were lysed and sonicated in buffer A (50 mM Tris-HCl [pH 7.4], 0.1% NP-40)  
10 containing 100 mM NaCl. Antibodies, BSA (1 mg/ml at the final concentration), and  
11 protein A-Sepharose beads (GE Healthcare) were added to the lysate, and incubated at 4°C for  
12 2-4 h. The beads were washed extensively with buffer A containing 300 mM NaCl, and  
13 proteins bound to the beads were eluted with an SDS sample buffer. Proteins were separated  
14 through 10% SDS-PAGE and detected by immunoblotting using anti-HA, anti-B23, anti-Flag,  
15 or anti-histone H3 antibodies.

16

### 17 **Pulse labeling of B23**

18           293T cells ( $1 \times 10^5$ ) were suspended in 200  $\mu$ l per assay of methionine free-MEM  
19 (Invitrogen). Met  $^{35}$ S label (American Radiolabeled Chemicals) containing [ $^{35}$ S]methionine  
20 (20  $\mu$ Ci per assay) was added, and cells were incubated at 37°C for 1 h. Cell lysates were  
21 prepared and B23 was immuno-purified using anti-B23 antibody. Labeled proteins were  
22 separated by a 10% SDS-PAGE and visualized by autoradiography.

23

## 1 **Short interference RNA (siRNA) for B23**

2 B23 Stealth RNAi select containing siB23-A (NPM1-HSS143152), siB23-B  
3 (NPM1-HSS143153), and siB23-C (NPM1-HSS143154), and control Stealth RNAi Negative  
4 control (Cat. 12935-200, Invitrogen) were introduced into HeLa cells with Lipofectamine  
5 2000 (Invitrogen) according to the manufacturer's protocol. At 24 h post transfection, the  
6 medium was replaced, and cells were harvested at 72 h post siRNA introduction. Proteins in  
7 HeLa cell lysates was separated by a 10% SDS-PAGE and detected by immunoblotting.

8 Total RNA was prepared by RNeasy mini kit (Qiagen) and RNase-free DNase I  
9 (Qiagen). For analysis of 45S pre-rRNA, total RNA was subjected to reverse transcription  
10 with a primer, 5'-CCATAACGGAGGCAGAGACA-3', corresponding to the 5' external  
11 transcribed spacer (ETS) of the rRNA gene, and the synthesized cDNA was used for  
12 quantitative PCR using FullVelocity SYBR Green Q-PCR Master Mix (STRATAGENE) in  
13 the presence of a primer set, 5'-GCCTTCTCTAGCGATCTGAGAG-3' and  
14 5'-CCATAACGGAGGCAGAGACA-3' corresponding to the 5' external transcribed spacer of  
15 rRNA as described (48). For analysis of  $\beta$ -actin, total RNA was subjected to reverse  
16 transcription with oligo dT as a primer, and the synthesized cDNA was used as a template for  
17 PCR amplification for  $\beta$ -actin in the presence of a primer set,  
18 ATGGGTCAGAAGGATTCCTATGT-3' and 5'-GGTCATCTTCTCGCGGTT-3'.

19 For immunofluorescence, HeLa cells transfected with siRNA on glass slides were  
20 permeablized with 0.1% Triton X-100, and then subjected to cross-linking with 4%  
21 formaldehyde. Then, B23 was detected with goat anti-B23 polyclonal antibody. DNA was  
22 stained with 4',6-diamidino-2-phenylindole (DAPI).

23

## 1 **Nucleolar chromatin immunoprecipitation**

2 Preparation of nucleoli and nucleolar ChIP were carried out according to the  
3 protocol described previously (42). Nucleoli were prepared from  $6 \times 10^6$  cells pre-fixed by  
4 formaldehyde (0.25% for 10 min at 37°C). Nucleolar ChIP assays were carried out  
5 according to the manual of the ChIP assay kit (Upstate). Quantitative PCR was carried out  
6 using FullVelocity SYBR Green QPCR Master Mix (STRATAGENE). PCR reactions were  
7 carried out using primer sets corresponding to the rRNA gene indicated in Table 1.

8

## 9 **Run-on assay**

10 Transcriptionally active nuclei were prepared from 293T cells ( $1 \times 10^6$ ) by  
11 treatment with NP-40 buffer (10 mM Tris-HCl [pH7.9], 10 mM NaCl, 3 mM  $MgCl_2$ , and  
12 0.1% NP-40) and resuspended in 30  $\mu$ l of glycerol stock buffer (10 mM Tris-HCl [pH7.9], 0.1  
13 mM EDTA, 3 mM  $MgCl_2$ , and 40% glycerol), and stored at  $-80^\circ C$  until use. The isolated  
14 nuclei (30  $\mu$ l) were mixed with 30  $\mu$ l of a reaction buffer containing 10 mM Tris-HCl [pH7.9],  
15 300 mM KCl, 5 mM  $MgCl_2$ , 0.5 mM each A, C, UTP, 50  $\mu$ Ci [ $\alpha$ - $^{32}P$ ]GTP ( $\sim 400$  Ci/mmol)  
16 (Amersham), and 40 units RNase inhibitor (TOYOBO), and incubated at  $30^\circ C$  for 30 min.  
17 Genomic DNA was digested with 200 units of DNase I at  $30^\circ C$  for 10 min followed by  
18 incubation with 100  $\mu$ g proteinase K at  $42^\circ C$  for 30 min. RNA was extracted with  
19 phenol/chloroform, and the newly transcribed RNA was monitored by hybridization with 1  $\mu$ g  
20 of denatured pUC119-HurDNA or pUC119 immobilized on Hybond- $N^+$  membranes. The  
21 intensity of hybridization dots was measured using the Image Gauge Program (Fuji-film).

22

## 23 ***In vitro* histone binding assay**

1 His-tagged B23.1, B23 $\Delta$ A, and B23 $\Delta$ C proteins were prepared as described (44).  
2 His-B23.1 and His-B23 $\Delta$ C proteins were mixed at appropriate ratios and precipitated by cold  
3 acetone. The precipitated proteins were denatured in a guanidine buffer (50 mM  
4 HEPES-NaOH [pH7.9], 150 mM NaCl, 0.1 mM EDTA, 0.5 mM DTT, 0.1% NP-40, and 6 M  
5 Guanidine-HCl) at room temperature for 15 min and renatured by dialysis against buffer  
6 containing 20 mM HEPES-NaOH [pH7.9], 50 mM NaCl, 0.5 mM EDTA, 1 mM DTT, 0.1  
7 mM PMSF, and 10% Glycerol. Renatured B23 proteins were mixed and incubated at 4°C  
8 for 30 min with core histones purified from HeLa cells in 50  $\mu$ l of 50 mM Tris-HCl [pH7.9],  
9 270 mM NaCl, 1 mM MgCl<sub>2</sub>, 0.1 mM PMSF, 0.1 mg/ml BSA, and 0.1% NP-40.  
10 Immunoprecipitation was carried out using anti-His or anti-B23 antibodies. Precipitated  
11 proteins were analyzed by a 12.5% SDS-PAGE and visualized by silver staining and  
12 immunoblotting using anti-His and anti-histone H3 antibodies.

13

#### 14 **Nucleosome assembly assay**

15 Supercoiled plasmid DNA (pCAGGS) was relaxed by incubation with  
16 topoisomerase I (TaKaRa) at 37°C for 30 min. The supercoiling assay was performed  
17 essentially as described (24).

## 1 RESULTS

2

### 3 Relationship between the cell growth rate and the rRNA gene transcription level

4 The relationship among the cell growth rate, the transcription rate of the rRNA gene,  
5 and the expression level of B23 was examined using 293T cells maintained in the presence of  
6 either 1% or 10% of serum. The expression level of B23 was examined by pulse-labeling  
7 with [<sup>35</sup>S]methionine. The pulse-labeled B23 was immunoprecipitated with an anti-B23  
8 antibody and analyzed through a 10% SDS-PAGE followed by autoradiography. The  
9 expression level of B23 was approximately four fold higher in the presence of 10% serum  
10 than in the presence of 1% serum at 48 h (Fig. 1A, lanes 3 and 4). This result is consistent  
11 with the report that the expression of B23 is associated with growth factor-induced  
12 mitogenesis in Swiss 3T3 cells (12). The expression level of nucleolus-related proteins was  
13 examined in cells maintained in the presence of 1% or 10% serum by immunoblotting using  
14 anti-UBF, anti-nucleolin, anti-fibrillarin, and anti- $\beta$ -actin antibodies. The protein expression  
15 level of UBF and nucleolin, both of which are known to be involved in the transcription of the  
16 rRNA gene, was reduced in the presence of 1% serum compared with that of 10% serum at 48  
17 h (Fig. 1B) as was the case for B23. On the other hand, the expression level of fibrillarin  
18 involved in the processing of pre-rRNA and  $\beta$ -actin was almost equal between cells  
19 maintained in 1% and 10% serum (Fig. 1B). The reduction level of mRNAs of *UBF*,  
20 *nucleolin*, *TIF-1A*, *fibrillarin*, and  *$\beta$ -actin* by low serum was virtually equal (see Fig. S1 in the  
21 supplemental material). These results suggest that the transcription by RNA polymerase II is  
22 evenly affected by serum concentrations and the expression level of proteins involved in the  
23 rRNA gene transcription is coordinately regulated at the level of translation and/or protein

1 degradation. In parallel, the transcription rate of the rRNA gene was monitored by run-on  
2 assays. The transcription level of the rRNA gene in cells maintained in the presence of 1%  
3 serum was lower by 0.7 fold than that in the presence of 10% serum (Fig. 1C). The amount  
4 of pre-rRNA was also monitored by quantitative RT-PCR using a primer set corresponding to  
5 the 5' external transcribed spacer (5' ETS) of pre-rRNA (48) as described in Materials and  
6 Methods in detail. The results were normalized as a ratio to that of  $\beta$ -actin. The amount of  
7 pre-rRNA in cells maintained in the presence of 1% serum was lower by 0.6 fold than that in  
8 the presence of 10% serum (Fig. 1D). The number of rRNA transcripts in cells under the  
9 growing condition is about 80% of the total cellular RNA quantity (40). Thus, this  
10 difference in the rRNA gene transcription rate between cells in 1% and 10% serum affect the  
11 cellular metabolism significantly. These results indicate that the rate of the rRNA gene  
12 transcription and the cell growth are positively correlated with the expression level of genes  
13 involved in the rRNA gene transcription including B23. Hereafter, we focused functional  
14 studies on B23.

15

### 16 **Knock-down of B23 decreases the transcription rate of the rRNA gene**

17 To investigate the role of B23.1 in transcription of the rRNA gene, knock-down  
18 (KD) of the B23.1 expression was performed using short interfering RNA (siRNA) specific  
19 for B23.1, siB23-A, -B, and -C. Introduction of the siRNA decreased the protein level of  
20 B23.1, but not  $\beta$ -actin, whereas a control siRNA had no effect (Fig. 2A). The expression  
21 level of other nucleolus-related genes was also examined in cells treated with siB23-C, and  
22 siB23-C was found to have no effect on the expression of other nucleolus-related genes (Fig.  
23 2B). Therefore, we could examine the function of B23.1 without other possible indirect

1 effect of siB23-C. KD of B23.1 was also confirmed in nucleoli of HeLa cells by  
2 immunofluorescence using anti-B23 antibody (Fig. 2C). The amount of pre-rRNA in HeLa  
3 cells transfected with siB23 or control siRNA was examined by quantitative RT-PCR using 5'  
4 ETS-specific primers for the rRNA gene. The amount of pre-rRNA was normalized as a  
5 ratio to that of  $\beta$ -actin mRNA. KD of the B23 expression level resulted in decreased  
6 transcription of the rRNA gene (Fig. 2D, right panel). The transcription level of the rRNA  
7 gene in cells treated by siB23-C was lower by 0.5 fold than that of control siRNA. These  
8 results indicate that B23 plays an important role in transcription of the rRNA gene by Pol I.

9

#### 10 **B23 is associated with the chromatin around the rRNA gene**

11 The amino-terminal domain of B23 shows significant similarity to nucleoplasmin,  
12 which is the first identified histone chaperone (32). B23 has been shown to bind with  
13 histones (45) and localized in nucleoli. Thus, it is reasonably hypothesized that B23 binds to  
14 chromatin around the rRNA gene. To examine this, we performed nucleolar chromatin  
15 immunoprecipitation (ChIP) as described in Materials and Methods. Nucleoli isolated from  
16 HeLa cells cross-linked with formaldehyde were sonicated to release chromatin fragments.  
17 DNA purified from chromatin fragments ranged in size less than 1 Kbp (Fig. S3 in  
18 supplemental material). Sonicated chromatin was subjected to immunoprecipitation with  
19 anti-Flag antibody as a negative control and anti-B23 antibody, followed by quantitative PCR  
20 using a primer set specific for the rRNA gene promoter. The antibody against TBP was also  
21 used, since TBP is one of landmarks of the active promoter. Figure 3A indicates that B23 as  
22 well as TBP is associated with the chromatin around the rRNA gene promoter. Next,  
23 quantitative PCR was performed using primer sets corresponding to the entire rRNA gene

1 (Table 1) and immunoprecipitated chromatin DNA as a template. It is indicated that B23  
2 binds the overall rRNA gene unit, but is concentrated at the transcription initiation site and the  
3 region encoding 5.8S rRNA (Fig. 3B). Binding of B23 at the region encoding 5.8S rRNA  
4 could correspond to the function of B23 for the processing of internal transcribed spacer 2  
5 (ITS2) of pre-rRNA (22, 26, 51). The binding activity of B23 at the transcription initiation  
6 site was examined in cells maintained in the presence of 1% or 10% of serum. The binding  
7 activity of B23 to the rRNA gene in cells maintained in the presence of 10% serum is higher  
8 than that in the presence of 1% (Fig. 3C). Moreover, we examined the chromatin structure  
9 of the rRNA gene promoter region in cells treated with siB23-C. The amount of B23  
10 associated with the rRNA gene promoter region was decreased in cells treated with siB23-C,  
11 while that of fibrillarin or transcription factors of the rRNA gene, that are TBP and UBF, were  
12 not influenced upon treatment with siB23-C (Fig. 3D, upper and middle panels). In contrast,  
13 the amount of histone H3 associated with the rRNA gene increased upon treatment with  
14 siB23-C (Fig. 3D, lower panel). The amount of histone H3 associated with the rRNA gene  
15 chromatin was 2 fold higher in cells treated with siB23-C than that of control siRNA.  
16 Histone distribution over the rRNA gene locus was examined (Fig. 3E, upper panel). The  
17 amount of histone H3 associated with the rRNA gene in the cells treated with siB23-C was  
18 higher than that of control cells over the rRNA gene locus. The ratio of the histone density  
19 on the rRNA gene in cells treated with siB23-C relative to that in control cells is plotted (Fig.  
20 3E, middle panel). This plot shows that the histone density around the transcription start site  
21 and non-coding region between +15 Kbp and +20 Kbp relative to the transcription start site is  
22 increased more than other regions. It is possible that the increase of the histone density on  
23 the promoter region may decrease the transcription efficiency of the rRNA gene, although it is



1 presently unclear how the high histone density of the non-coding region affects the rRNA  
2 gene transcription. These results suggest that B23 is involved in control of the histone  
3 density over the rRNA gene.

4

#### 5 **B23 $\Delta$ C inhibits the histone binding activity of B23.1 *in vitro***

6 To understand the function of B23 as a histone chaperone, we tried to prepare  
7 deletion mutants of B23.1 aiming construction of a dominant-negative type mutant to block  
8 the histone chaperone function of B23 specifically. The acidic regions of B23 are crucial for  
9 both histone binding and histone chaperone activities (44, 45). The crystal structure of the  
10 N-terminal domain of NO38, a *Xenopus* homolog of human B23, indicates that NO38 forms a  
11 pentamer and each pentamer further assembles into a decamer as does nucleoplasmin (10, 41).  
12 These studies suggest that oligomer formation of nucleoplasmin family proteins is important  
13 for their histone chaperone activity. Thus, we hypothesized that a B23 protein containing the  
14 oligomeric domain but lacking domains responsible for its histone chaperone activity is to be  
15 a dominant-negative type protein. To test this, B23 mutants termed B23 $\Delta$ A and B23 $\Delta$ C  
16 lacking either the acidic region or both the acidic and the carboxyl-terminal region,  
17 respectively, was prepared (Fig. 4A). Recombinant hexa histidine-tagged (His-) B23.1,  
18 B23 $\Delta$ A, and B23 $\Delta$ C proteins were expressed in *E. coli* and purified (Fig. 4B).

19 To examine the histone binding activity of B23 mutants, histone binding assays  
20 were carried out *in vitro* using recombinant B23 mutant proteins and core histones purified  
21 from HeLa cells. B23 proteins and core histones were mixed and incubated, followed by  
22 immunoprecipitation with anti-His antibody and immunoblotting with anti-His and  
23 anti-histone H3 antibodies. His-B23 $\Delta$ C did not bind to core histones, whereas the histone

1 binding activity of His-B23 $\Delta$ A was essentially the same as that of wild type His-B23.1 (Fig.  
2 4C). The fact that B23 $\Delta$ A binds to histone H3 indicates that the other domain except for  
3 acidic domains is important for histone binding. Nucleosome assembly activity of B23 $\Delta$ A  
4 was examined using supercoiling assay (Fig. 4D). His-B23.1 assembled nucleosomes in a  
5 dose-dependent manner, and His-B23 $\Delta$ C did not as previously described (45). Although  
6 His-B23 $\Delta$ A bound with core histones, it could not transfer core histones to plasmid DNA (Fig.  
7 4D). This result indicates that acidic domains of B23.1 are crucial for the histone chaperone  
8 activity.

9 We confirmed that B23 $\Delta$ C does not have histone binding and nucleosome assembly  
10 activities (Figs. 4C and 4D). Then, we examined whether B23 $\Delta$ C is a dominant-negative  
11 type protein for B23.1. When mixed directly, His-B23 $\Delta$ C did not form a hetero-oligomer  
12 with His-B23.1 *in vitro*, suggesting that B23 forms a stable oligomer (data not shown).  
13 However, when a mixture of His-B23.1 and His-B23 $\Delta$ C was subjected to the  
14 denature-renature protocol as described in Materials and Methods, two proteins were found to  
15 form a hetero-oligomer successfully. The interaction between B23 oligomers and core  
16 histones was examined by immunoprecipitation using the anti-B23 antibody that can not  
17 recognize B23 $\Delta$ C (data not shown). Immunoprecipitated proteins were separated by a  
18 12.5% SDS-PAGE, followed by silver staining (Fig. 4E, left panel) and immunoblotting (Fig.  
19 4E, right panel) using anti-His antibody. B23.1 formed a hetero-oligomer with His-B23 $\Delta$ C  
20 (Fig. 4E, right panel, lanes 3 and 4). B23.1 interacted efficiently with core histones (Fig. 4E,  
21 left panel, lane 2) as shown in Fig. 4C and a previous report (45). However, the histone  
22 binding activity of B23.1 was decreased by oligomer formation with increasing amounts of  
23 His-B23 $\Delta$ C (Fig. 4E, left panel, lanes 3 and 4). The molar ratio between His-B23.1 and

1 His-B23 $\Delta$ C used here was 1:2.5 and 1:5 in lanes 3 and 4, respectively. Since B23 was  
2 shown to form a pentamer, it is likely that a His-B23.1 pentamer including four His-B23 $\Delta$ C  
3 molecules lost the histone binding activity completely. We concluded that oligomer  
4 formation of B23 is critical for its histone binding activity and B23 $\Delta$ C inhibits the histone  
5 binding activity of B23.1 in a dominant-negative manner.

6

### 7 **B23 $\Delta$ C inhibits the histone binding activity of B23.1 *in vivo***

8 To use the B23 mutant proteins as tools for functional analyses of B23 *in vivo*, we  
9 examined whether B23 mutants inhibit the histone binding activity of B23.1. First, we  
10 confirmed the expression of each B23 protein by immunoblotting using lysates of 293T cells  
11 transfected with plasmids encoding HA-B23.1, HA-B23 $\Delta$ C, Flag-B23.1, or Flag-B23 $\Delta$ A  
12 proteins (Fig. 5A). To examine the histone binding activity of B23 mutants,  
13 immunoprecipitation with anti-HA and anti-Flag antibodies was performed using lysates  
14 prepared from 293T cells expressing HA-B23.1, HA-B23 $\Delta$ C, Flag-B23.1, or Flag-B23 $\Delta$ A,  
15 followed by immunoblotting with anti-Histone H3 antibody (Fig. 5B). B23 $\Delta$ C did not bind  
16 to histone H3, whereas B23 $\Delta$ A had the histone binding activity as did wild type B23.1. This  
17 result is consistent with the result of histone binding assays *in vitro* (Fig. 4C).

18 We investigated whether B23 $\Delta$ A and B23 $\Delta$ C inhibit the histone binding activity of  
19 wild type B23.1 *in vivo* (Figs. 5C and 5D). HA-B23.1 was expressed in 293T cells without  
20 or with Flag-B23 $\Delta$ A, and immunoprecipitation was carried out using anti-HA antibody,  
21 followed by immunoblotting with anti-HA, anti-Flag, and anti-histone H3 antibodies. As  
22 expected, Flag-B23 $\Delta$ A did not influence the histone binding activity of HA-B23.1, but rather  
23 the amount of co-precipitated histone H3 with B23 $\Delta$ A was slightly increased (Fig. 5C, lanes 8,

1 9, and 10). Then, Flag-B23.1 was expressed in 293T cells without or with HA-B23 $\Delta$ C, and  
2 immunoprecipitation was carried out with anti-Flag antibody, followed by immunoblotting  
3 with anti-Flag, anti-HA, and anti-histone H3 antibodies (Fig. 5D). In contrast to  
4 Flag-B23 $\Delta$ A, HA-B23 $\Delta$ C inhibited the histone binding activity of Flag-B23.1 in a  
5 dose-dependent manner through interacting with Flag-B23.1 (Fig. 5D, lanes 7 and 8). This  
6 result indicates that B23 $\Delta$ C forms a hetero-oligomer with B23.1 and inhibits the histone  
7 binding activity of B23.1 *in vivo*.

8

### 9 **Inhibition of the rRNA gene transcription by B23 $\Delta$ C**

10 We have shown that B23 $\Delta$ C inhibits the histone binding activity of B23.1 in a  
11 dominant-negative manner. Therefore, B23 $\Delta$ C could be a powerful tool to clarify the  
12 cellular function of B23 as a histone chaperone. Next, we examined the effect of B23 $\Delta$ C on  
13 the rRNA gene transcription. 293T cells expressing HA-B23 $\Delta$ C were lysed, and HA-B23 $\Delta$ C  
14 was immunoprecipitated with anti-HA antibody (Fig. 6A). Approximately 50% of  
15 endogenous B23.1 were co-precipitated with HA-B23 $\Delta$ C under the condition employed here.  
16 Therefore, it is quite likely that HA-B23 $\Delta$ C can quench the function of endogenous B23 in  
17 cells. In order to examine the effect of HA-B23 $\Delta$ C on the chromatin binding activity of B23,  
18 nucleolar ChIP with anti-B23 antibody was carried out using nucleoli purified from 293T  
19 cells transfected with plasmids encoding HA-B23.1 and HA-B23 $\Delta$ C. The association of  
20 B23 with the rRNA gene promoter region was examined by quantitative PCR using  
21 immunoprecipitated DNA and a specific primer set (Fig. 6B). We found that B23 binds to  
22 the promoter region of the rRNA gene, and over-expression of HA-B23 $\Delta$ C decreased this  
23 association by 25%. Next, we examined the effect of HA-B23 $\Delta$ C on the chromatin structure

1 of the rRNA gene promoter region. Binding of transcription factors of the rRNA gene, UBF  
2 and TBP, was not affected or slightly up-regulated by expression of HA-B23 $\Delta$ C (Fig. 6C),  
3 while the amount of histone H3 or acetylated histone H3 associated with the rRNA gene  
4 increased by expression of HA-B23 $\Delta$ C (Fig. 6D). Histone H3 distribution over the rRNA  
5 gene locus was examined (Fig. 6E, upper panel). The amount of histone in cells expressing  
6 B23 $\Delta$ C was higher than that in mock-treated cells over the rRNA gene locus.  
7 Over-expression of B23 $\Delta$ C increased the histone density of the coding region of 18S rRNA  
8 gene (Fig. 6E, lower panel). The histone density on the rRNA gene except for the region  
9 encoding 18S rRNA is 1.3 ~ 1.8 fold higher in cells expressing B23 $\Delta$ C compared with that of  
10 mock-treated cells, consisting with the result using siB23 (Fig. 3E), whereas the histone  
11 density on the region coding 18S rRNA was 2.9 fold higher in cells expressing B23 $\Delta$ C than  
12 that in mock-treated cells. The reason for this discrepancy between the results in Fig. 3E and  
13 Fig. 6E is presently unclear, but assumed to be due to the difference of cell types used in  
14 experiments. The results shown in Fig. 3D and Fig. 6D indicate that the amount ratio of  
15 acetylated histone H3 to histone H3 on the rRNA gene in HeLa cells was lower than that of  
16 293T cells. In other words, this could be interpreted as that the ratio of the number of fully  
17 active rRNA genes to the total number of the rRNA genes is higher in 293T cells than that of  
18 HeLa cells. Nevertheless, it is suggested that B23 controls the histone density on the rRNA  
19 gene.

20           Next, we analyzed the transcription activity of Pol I in cells expressing HA-B23 $\Delta$ C.  
21 Nuclei were isolated from 293T cells expressing HA-B23 $\Delta$ C, and run-on assays were carried  
22 out (Fig. 6F). Run-on assays demonstrated that the transcription level of the rRNA gene in  
23 293T cells expressing HA-B23 $\Delta$ C was lower by 0.6 fold than that of control cells, suggesting

1 that over-expression of HA-B23 $\Delta$ C down-regulates the activity of Pol I possibly by blocking  
2 the function of endogenous B23. Moreover, we examined the effect of B23 $\Delta$ A on the  
3 transcription of the rRNA gene, which binds to histones but lacks the histone chaperone  
4 activity (Fig. 4D). Quantitative RT-PCR for pre-rRNA demonstrated that the amount of  
5 pre-rRNA in cells expressing B23 $\Delta$ A was lower than that of control cells (Fig. 6G). This  
6 result may be interpreted as that B23 $\Delta$ A forms oligomer with endogenous B23.1 and inhibits  
7 the rRNA transcription as does B23 $\Delta$ C. Thus, it is quite possible that B23 controls the  
8 histone density on the rRNA gene, and thereby regulates the transcription of the rRNA gene as  
9 a nucleolar histone chaperone.

10 We tried to confirm the above notion by complementation experiments. By  
11 monitoring the transcription of the rRNA gene, we examined the rescue ability of B23 mutant  
12 proteins for HeLa cells treated with siRNA specific for B23.1. After treatment of siB23-C,  
13 HeLa cells were super-transfected with plasmids encoding various B23 proteins with  
14 pEF321-T encoding SV40 T antigen in order to amplify the expression level of B23 proteins  
15 supplied by pCAGGS and/or pCHA, which contain the replication origin of SV40. siB23-C  
16 recognizes the 3'UTR of endogenous mRNA of B23.1, and we constructed pCAGGS-based  
17 plasmids encoding B23 mutant proteins without the siB23-C target sequence. Therefore,  
18 siB23-C does not affect the expression of exogenous B23 mutant proteins. The expression  
19 level of B23 proteins was detected by immunoblotting using anti-B23 or anti-HA antibodies  
20 (Fig. 7A). The amount of 45S pre-rRNA in HeLa cells was examined by quantitative  
21 RT-PCR using the 5'ETS-specific primer set for the rRNA gene. The reduction in synthesis  
22 of 45S pre-rRNA by siB23-C was recovered by the expression of Flag-B23.1 in a  
23 dose-dependent manner (Fig. 7B, lanes 2-4). In contrast, Flag-B23 $\Delta$ A and HA-B23 $\Delta$ C could

1 not rescue the synthesis of 45S pre-rRNA reduced by siB23-C, but rather B23 $\Delta$ C decreased  
2 the synthesis of 45S pre-rRNA (Fig. 7B, lanes 2, and 5-8). These results indicate that  
3 Flag-B23.1 promotes the transcription of the rRNA gene, but Flag-B23 $\Delta$ A and HA-B23 $\Delta$ C do  
4 not. B23 $\Delta$ A was shown to be capable of binding to histones (Fig. 4C), but it does not have  
5 the histone chaperone activity (Fig. 4D), indicating that the acidic domain of B23.1 is critical  
6 for the histone chaperone activity and the promoting activity of the rRNA gene transcription.  
7 Thus, it is concluded that the histone binding activity of B23 is not sufficient for the rRNA  
8 gene transcription promotion, and not only the histone binding activity but also the histone  
9 chaperone activity is required (27).

10

#### 11 **Inhibition of cell growth by over-expression of B23 $\Delta$ C**

12 We have shown the positive correlation between the B23 histone chaperone activity  
13 and the rRNA gene transcription. Next, we investigated the function of B23 in the cell  
14 growth. To this end, we took advantage of B23 $\Delta$ C, a dominant-negative type mutant for the  
15 histone chaperone activity of B23. 293T cells were co-transfected with plasmids encoding  
16 B23 $\Delta$ C and EGFP. The number of EGFP-positive cells, thus quite possibly B23 $\Delta$ C-positive  
17 cells, was counted every 24 hrs post transfection (Fig. 8A). The cell growth of 293T cells  
18 expressing B23 $\Delta$ C seemed to be stopped. The cell growth rate was decreased in proportion  
19 to the expression level of B23 $\Delta$ C (Fig. 8B). In cells expressing B23 $\Delta$ C3, the rRNA  
20 synthesis was inhibited (Fig. 6). Moreover, KD of the expression of B23.1 by siB23-C  
21 decreased the cell growth rate (data not shown). Thus, it is possible that B23 $\Delta$ C suppresses  
22 the synthesis of the rRNA gene, and thereby affects the cell growth rate.

23 Alternatively, B23 $\Delta$ C may affect the cell growth through other pathway(s), since

1 B23 is a multifunctional protein. We examined the cell cycle profile of cells expressing  
2 B23 $\Delta$ C. It is suspected that the growth inhibition by expression of B23 $\Delta$ C is due to  
3 apoptosis. However, the population of apoptotic cells was not increased by expression of  
4 B23 $\Delta$ C (Fig. 8C, inset table). As shown in Fig. 8C, G2/M population was slightly increased  
5 by expression of B23 $\Delta$ C. This result may be consistent with the observation with B23<sup>-/-</sup>  
6 MEFs (17, 31). Tetraploid cells contain 4n DNA as do cells in G2/M phase. On this line,  
7 the p53 pathway involved in centrosome duplication that is essential for ploidy maintenance  
8 is one of the candidates for the cell growth defect by expressing B23 $\Delta$ C. It was reported that  
9 B23 regulates the stability and the transcriptional activity of p53 directly or indirectly (5, 29).  
10 To exclude the possibility that B23 $\Delta$ C inhibited the cell growth through affecting the p53  
11 pathway, we examined the effect of over-expression of B23 $\Delta$ C on the cell growth rate of  
12 p53<sup>-/-</sup> MEF (mouse embryonic fibroblast). As shown in Fig. 8D, the cell growth rate of  
13 p53<sup>-/-</sup> MEF was similarly decreased by over-expression of B23 $\Delta$ C. The effect of B23 $\Delta$ C on  
14 the cell growth of p53<sup>-/-</sup> MEF seems lower than that of 293T cells. This could be due to the  
15 fact that a pCAGGS-based expression vector is amplified in 293T cells expressing SV40 T  
16 antigen but not in p53<sup>-/-</sup> MEF, so that the expression level of B23 $\Delta$ C is higher in 293T cells  
17 than p53<sup>-/-</sup> MEF. Thus, the cell growth inhibition activity of B23 $\Delta$ C is independent of the  
18 p53 pathway. From these results, we concluded that over-expression of B23 $\Delta$ C leads to the  
19 growth inhibition most possibly through down-regulation of the rRNA gene transcription.



## 1 DISCUSSION

2

3 In this study, we examined the function of B23 as a nucleolar histone chaperone.

4 Here, we have confirmed that the expression level of B23 is correlated with the rate of the

5 rRNA gene transcription (Fig. 1). KD of B23 by siRNA decreased the transcription rate of

6 rRNA gene (Fig. 2). It was shown using nucleolar ChIP assays that B23 is associated with

7 the rRNA gene in cells (Fig. 3). To test the histone chaperone activity of B23 in cells, we

8 constructed B23 $\Delta$ C, a dominant-negative mutant, which inhibited the histone binding activity

9 of B23.1 (Figs. 4 and 5). B23 $\Delta$ C over-expressed in 293T cells bound to endogenous B23.1

10 and decreased the transcription rate of the rRNA gene *in vivo* (Fig. 6). Expression of

11 exogenous B23.1 but neither B23 $\Delta$ A nor B23 $\Delta$ C could rescue the rRNA synthesis reduced by

12 siRNA (Fig. 7). Finally, over-expression of B23 $\Delta$ C results in inhibition of the cell growth

13 (Fig. 8). These results altogether suggest that B23 regulates the transcription of rRNA gene

14 as a nucleolar histone chaperone and thus affects the cell growth rate.

15 The inhibitory effect of siB23-C (Fig. 2D) and B23 $\Delta$ C (Fig. 6F and G) on the rRNA

16 transcription seemed incomplete. However, these changes in the transcription rate of the

17 rRNA genes are ranged in almost the same level as the difference between the rRNA synthesis

18 levels of cells maintained in 10% and 1% serum (Fig. 1C and D). The tandem rRNA gene

19 repeats exist in two distinct types of chromatin, an 'open' one permissive for transcription and

20 a 'closed' one under the transcriptionally repressive state. It is reported that the number of

21 the rRNA genes existing as open chromatin is independent of the cell growth rate (6). Based

22 on these, it is possible that B23 may not be involved in remodeling of chromatin around the

23 rRNA gene between open and closed states. Then, we would consider the possibility that

1 B23 is required for maintenance of an open chromatin. The results in this study show that  
2 the rRNA gene transcription is proportional to the amount of B23. B23 remodels the  
3 chromatin structure and stimulates the transcription from chromatin template (57, unpublished  
4 results, and see below). We assume that the amount of B23 is a switch for the rRNA  
5 transcription fluctuation in response to a variety of continuous stresses/stimuli from outside  
6 and in side of cells.

7 In eukaryotic cells, the chromatin structure plays an important role in regulation of  
8 gene expression. In the nucleolus, it was shown that NoRC, a nucleolar ATP-dependent  
9 chromatin remodeling complex, recruits histone deacetylase and DNA methyltransferase  
10 activities on the rRNA genes to induce formation of heterochromatin on the rRNA genes (38,  
11 50). B23 has been shown to be acetylated by p300 and stimulate transcription preferentially  
12 from a chromatin template containing acetylated histones, although the stimulation by  
13 acetylated B23 occurs in transcription assay using RNA polymerase II in the presence of  
14 transactivator Gal4-VP16 *in vitro* (57). This observation suggests that B23 facilitates  
15 disassembly of histones over the rRNA gene in combination with p300 to stimulate the  
16 transcription of the rRNA gene. Non-acetylatable B23 mutants would be powerful tools for  
17 further analysis of the coordination mechanism between B23 and p300 on the rRNA gene  
18 transcription. In addition, B23 was shown to interact with nucleolin, which is one of  
19 nucleolar histone chaperones and enhances the activity of the chromatin remodeling  
20 machineries, SWI/SNF and ACF (1, 34, 48). B23 and nucleolin as nucleolar histone  
21 chaperones may regulate the transcription of the rRNA gene through the structural change of  
22 chromatin with ATP-dependent chromatin remodelers and histone modification enzymes in  
23 coordinated manner.

1 Nucleoplasmin 3 (NPM3) was identified as a B23 binding protein from mouse liver  
2 cDNA library using yeast two hybrid system (23). NPM3 shows the significant structural  
3 similarity to B23 $\Delta$ C and is found to interact with endogenous B23. Over-expression of  
4 NPM3 decreased the rate of pre-rRNA synthesis and processing. It is possible that NPM3  
5 also functions as an inhibitor for the histone binding activity of endogenous B23 as does  
6 B23 $\Delta$ C. We have also observed retardation of rRNA processing by overexpression of  
7 B23 $\Delta$ C (data not shown). Since B23.2 lacking the C-terminal RNA binding domain was  
8 found to form an oligomer with B23.1 and inhibit the RNA binding activity of B23.1 (46), it  
9 is likely that B23 $\Delta$ C also inhibits not only the histone chaperone activity but also the RNA  
10 binding activity of B23.1. The RNA binding activity of B23.1 could be important to target  
11 B23.1 to pre-rRNA to be processed. This could be one of the reasons why over-expression  
12 of B23 $\Delta$ C and NPM3 inhibits the pre-rRNA processing.

13 From the observation that B23, NPM3, and nucleolin are involved in pre-rRNA  
14 synthesis and processing (1, 15, 23, 48, 51), it is likely that the rRNA gene transcription and  
15 pre-rRNA processing are coupled and co-regulated. This idea is supported by the result that  
16 B23 associates with not only the promoter region but also the region encoding 5.8S rRNA  
17 next to ITS2 cleaved by B23 (Fig. 3B). In yeast cells, Pol I transcription and pre-rRNA  
18 processing were shown to be linked by pre-rRNA processing factors (13, 16, 52) and the  
19 complex containing Pol I drives both transcription and pre-rRNA modifications *in vitro* (11).  
20 In mammalian cells, B23 could be one of the molecules to link between the rRNA gene  
21 transcription and pre-rRNA processing.

22 We demonstrated that B23 $\Delta$ C drastically decreased the cell proliferation rate of  
23 293T cells (Fig. 8A). B23<sup>+/-</sup>, B23<sup>hy/hy</sup> (homozygous hypomorphic mutants) and B23<sup>-/-</sup> MEFs

1 show the decreased cell proliferation rate compared with wild type MEF (17). It was  
2 previously reported that B23 binds to p53 directly and enhances its stability and the  
3 transcriptional activity of p53, so that over-expression of B23.1 induces inhibition of MEF  
4 proliferation (5). B23 interacts with ARF and Mdm2, those of which are involved in cell  
5 cycle regulation in concert with p53 (3, 29, 30). We had suspected that B23 $\Delta$ C inhibits the  
6 cell proliferation through the p53 pathway. However, this possibility was ruled out by  
7 experiments using p53<sup>-/-</sup> MEF (Fig. 8D). This is supported by the report that a deletion  
8 mutant of B23 similar to B23 $\Delta$ C does not bind to p53 in vitro, and the expression of B23.1  
9 increases the proliferation rate of p53<sup>-/-</sup> MEF slightly (5).

## 1 **ACKNOWLEDGMENTS**

2           We thank Dr. K Fukasawa for the gift of p53<sup>-/-</sup>MEF and Dr. H Ariga for the gift of  
3 pEF321-T. This work was supported in part by grants-in-aid from the Ministry of Education,  
4 Culture, Sports, Science, and Technology of Japan (to K.N. and M.O.), and by grants from the  
5 Bioarchitect Research Program from RIKEN and for the project of Tsukuba Advanced  
6 Research Alliance (to K.N.).

1   **REFERENCES**

2

- 3   1.    **Angelov, D., V. A. Bondarenko, S. Almagro, H. Menoni, F. Mongelard, F. Hans, F.**  
4       **Mietton, V. M. Studitsky, A. Hamiche, S. Dimitrov, and P. Bouvet.** 2006. Nucleolin  
5       is a histone chaperone with FACT-like activity and assists remodeling of nucleosomes.  
6       EMBO J **25**:1669-1679.
- 7   2.    **Bertwistle, D., M. Sugimoto, and C. J. Sherr.** 2004. Physical and functional  
8       interactions of the Arf tumor suppressor protein with nucleophosmin/B23. Mol Cell  
9       Biol **24**:985-996.
- 10  3.    **Brady, S. N., Y. Yu, L. B. Maggi, Jr., and J. D. Weber.** 2004. ARF impedes  
11       NPM/B23 shuttling in an Mdm2-sensitive tumor suppressor pathway. Mol Cell Biol  
12       **24**:9327-9338.
- 13  4.    **Chang, J. H., and M. O. Olson.** 1990. Structure of the gene for rat nucleolar protein  
14       B23. J Biol Chem **265**:18227-18233.
- 15  5.    **Colombo, E., J. C. Marine, D. Danovi, B. Falini, and P. G. Pelicci.** 2002.  
16       Nucleophosmin regulates the stability and transcriptional activity of p53. Nat Cell Biol  
17       **4**:529-533.
- 18  6.    **Conconi, A., R. M. Widmer, T. Koller, and J. M. Sogo.** 1989. Two different  
19       chromatin structures coexist in ribosomal RNA genes throughout the cell cycle. Cell  
20       **57**:753-761.
- 21  7.    **Derenzini, M.** 2000. The AgNORs. Micron **31**:117-120.
- 22  8.    **Dery, C. V., M. Toth, M. Brown, J. Horvath, S. Allaire, and J. M. Weber.** 1985.  
23       The structure of adenovirus chromatin in infected cells. J Gen Virol **66**:2671-2684.

- 1 9. **Dumbar, T. S., G. A. Gentry, and M. O. Olson.** 1989. Interaction of nucleolar  
2 phosphoprotein B23 with nucleic acids. *Biochemistry* **28**:9495-9501.
- 3 10. **Dutta, S., I. V. Akey, C. Dingwall, K. L. Hartman, T. Laue, R. T. Nolte, J. F. Head,**  
4 **and C. W. Akey.** 2001. The crystal structure of nucleoplasmin-core: implications for  
5 histone binding and nucleosome assembly. *Mol Cell* **8**:841-853.
- 6 11. **Fath, S., P. Milkereit, A. V. Podtelejnikov, N. Bischler, P. Schultz, M. Bier, M.**  
7 **Mann, and H. Tschochner.** 2000. Association of yeast RNA polymerase I with a  
8 nucleolar substructure active in rRNA synthesis and processing. *J Cell Biol.*  
9 **149**:575-590.
- 10 12. **Feuerstein, N., S. Spiegel, and J. J. Mond.** 1988. The nuclear matrix protein,  
11 numatrin (B23), is associated with growth factor-induced mitogenesis in Swiss 3T3  
12 fibroblasts and with T lymphocyte proliferation stimulated by lectins and anti-T cell  
13 antigen receptor antibody. *J Cell Biol* **107**:1629-1642.
- 14 13. **Gallagher, J. E., D. A. Dunbar, S. Granneman, B. M. Mitchell, Y. Osheim, A. L.**  
15 **Beyer, and S. J. Baserga.** 2004. RNA polymerase I transcription and pre-rRNA  
16 processing are linked by specific SSU processome components. *Genes Dev.*  
17 **18**:2506-2517.
- 18 14. **Gamble, M. J., H. Erdjument-Bromage, P. Tempst, L. P. Freedman, and R. P.**  
19 **Fisher.** 2005. The histone chaperone TAF-I/SET/INHAT is required for transcription  
20 in vitro of chromatin templates. *Mol Cell Biol* **25**:797-807.
- 21 15. **Ginisty, H., F. Amalric, and P. Bouvet.** 1998. Nucleolin functions in the first step of  
22 ribosomal RNA processing. *EMBO J* **17**:1476-1486.
- 23 16. **Granneman, S., and S. J. Baserga.** 2005. Crosstalk in gene expression: coupling and

- 1 co-regulation of rDNA transcription, pre-ribosome assembly and pre-rRNA processing.  
2 Curr Opin Cell Biol. **17**:281-286.
- 3 17. **Grisendi, S., R. Bernardi, M. Rossi, K. Cheng, L. Khandker, K. Manova, and P. P.**  
4 **Pandolfi.** 2005. Role of nucleophosmin in embryonic development and tumorigenesis.  
5 Nature **437**:147-153.
- 6 18. **Grummt, I.** 2007. Different epigenetic layers engage in complex crosstalk to define  
7 the epigenetic state of mammalian rRNA genes. Hum Mol Genet **16 Spec No**  
8 **1**:R21-27.
- 9 19. **Grummt, I.** 1999. Regulation of mammalian ribosomal gene transcription by RNA  
10 polymerase I. Prog Nucleic Acid Res Mol Biol **62**:109-154.
- 11 20. **Haruki, H., B. Gyurcsik, M. Okuwaki, and K. Nagata.** 2003. Ternary complex  
12 formation between DNA-adenovirus core protein VII and TAF-Ibeta/SET, an acidic  
13 molecular chaperone. FEBS Lett **555**:521-527.
- 14 21. **Haruki, H., M. Okuwaki, M. Miyagishi, K. Taira, and K. Nagata.** 2006.  
15 Involvement of template-activating factor I/SET in transcription of adenovirus early  
16 genes as a positive-acting factor. J Virol **80**:794-801.
- 17 22. **Herrera, J. E., R. Savkur, and M. O. Olson.** 1995. The ribonuclease activity of  
18 nucleolar protein B23. Nucleic Acids Res **23**:3974-3979.
- 19 23. **Huang, N., S. Negi, A. Szebeni, and M. O. Olson.** 2005. Protein NPM3 interacts  
20 with the multifunctional nucleolar protein B23/nucleophosmin and inhibits ribosome  
21 biogenesis. J Biol Chem **280**:5496-5502.
- 22 24. **Ishimi, Y., J. Hirosumi, W. Sato, K. Sugasawa, S. Yokota, F. Hanaoka, and M.**  
23 **Yamada.** 1984. Purification and initial characterization of a protein which facilitates



- 1 assembly of nucleosome-like structure from mammalian cells. *Eur J Biochem*  
2 **142**:431-439.
- 3 25. **Ishimi, Y., M. Kojima, M. Yamada, and F. Hanaoka.** 1987. Binding mode of  
4 nucleosome-assembly protein (AP-I) and histones. *Eur J Biochem* **162**:19-24.
- 5 26. **Itahana, K., K. P. Bhat, A. Jin, Y. Itahana, D. Hawke, R. Kobayashi, and Y.**  
6 **Zhang.** 2003. Tumor suppressor ARF degrades B23, a nucleolar protein Involved in  
7 ribosome biogenesis and cell proliferation. *Mol Cell* **12**:1151-1164.
- 8 27. **Kato, K., M. Miyaji-Yamaguchi, M. Okuwaki, and K. Nagata.** 2007. Histone  
9 acetylation-independent transcription stimulation by a histone chaperone. *Nucleic*  
10 *Acids Res* **35**:705-715.
- 11 28. **Kawase, H., M. Okuwaki, M. Miyaji, R. Ohba, H. Handa, Y. Ishimi, T.**  
12 **Fujii-Nakata, A. Kikuchi, and K. Nagata.** 1996. NAP-I is a functional homologue of  
13 TAF-I that is required for replication and transcription of the adenovirus genome in a  
14 chromatin-like structure. *Genes Cells* **1**:1045-1056.
- 15 29. **Korgaonkar, C., J. Hagen, V. Tompkins, A. A. Frazier, C. Allamargot, F. W.**  
16 **Quelle, and D. E. Quelle.** 2005. Nucleophosmin (B23) targets ARF to nucleoli and  
17 inhibits its function. *Mol Cell Biol* **25**:1258-1271.
- 18 30. **Kurki, S., K. Peltonen, and M. Laiho.** 2004. Nucleophosmin, HDM2 and p53:  
19 players in UV damage incited nucleolar stress response. *Cell Cycle* **3**:976-979.
- 20 31. **Lanni, J. S., and T. Jacks.** 1998. Characterization of the p53-dependent postmitotic  
21 checkpoint following spindle disruption. *Mol Cell Biol* **18**:1055-1064.
- 22 32. **Laskey, R. A., B. M. Honda, A. D. Mills, and J. T. Finch.** 1978. Nucleosomes are  
23 assembled by an acidic protein which binds histones and transfers them to DNA.

- 1 Nature **275**:416-420.
- 2 33. **Li, J., G. Langst, and I. Grummt.** 2006. NoRC-dependent nucleosome positioning  
3 silences rRNA genes. EMBO J **25**:5735-5741.
- 4 34. **Liu, H. T., and B. Y. Yung.** 1999. In vivo interaction of nucleophosmin/B23 and  
5 protein C23 during cell cycle progression in HeLa cells. Cancer Lett **144**:45-54.
- 6 35. **Matsumoto, K., and O. M. Nagata K, Tsujimoto M.** 1999. Histone- and  
7 chromatin-binding activity of template activating factor-I. FEBS Lett **463**:285-288.
- 8 36. **Matsumoto, K., K. Nagata, M. Ui, and F. Hanaoka.** 1993. Template activating  
9 factor I, a novel host factor required to stimulate the adenovirus core DNA replication.  
10 J Biol Chem **268**:10582-10587.
- 11 37. **Matsumoto, K., M. Okuwaki, H. Kawase, H. Handa, F. Hanaoka, and K. Nagata.**  
12 1995. Stimulation of DNA transcription by the replication factor from the adenovirus  
13 genome in a chromatin-like structure. J Biol Chem **270**:9645-9650.
- 14 38. **Mayer, C., K. M. Schmitz, J. Li, I. Grummt, and R. Santoro.** 2006. Intergenic  
15 Transcripts Regulate the Epigenetic State of rRNA Genes. Mol Cell **22**:351-361.
- 16 39. **Miyamoto, S., T. Suzuki, S. Muto, K. Aizawa, A. Kimura, Y. Mizuno, T. Nagino, Y.**  
17 **Imai, N. Adachi, M. Horikoshi, and R. Nagai.** 2003. Positive and negative  
18 regulation of the cardiovascular transcription factor KLF5 by p300 and the oncogenic  
19 regulator SET through interaction and acetylation on the DNA-binding domain. Mol  
20 Cell Biol **23**:8528-8541.
- 21 40. **Moss, T., F. Langlois, T. Gagnon-Kugler, and V. Stefanovsky.** 2007. A housekeeper  
22 with power of attorney: the rRNA genes in ribosome biogenesis. Cell Mol Life Sci  
23 **64**:29-49.

- 1 41. **Namboodiri, V. M., I. V. Akey, M. S. Schmidt-Zachmann, J. F. Head, and C. W.**  
2 **Akey.** 2004. The structure and function of *Xenopus* NO38-core, a histone chaperone  
3 in the nucleolus. *Structure (Camb)* **12**:2149-2160.
- 4 42. **O'Sullivan, A. C., G. J. Sullivan, and B. McStay.** 2002. UBF binding in vivo is not  
5 restricted to regulatory sequences within the vertebrate ribosomal DNA repeat. *Mol*  
6 *Cell Biol* **22**:657-668.
- 7 43. **Okuda, M., H. F. Horn, P. Tarapore, Y. Tokuyama, A. G. Smulian, P. K. Chan, E.**  
8 **S. Knudsen, I. A. Hofmann, J. D. Snyder, K. E. Bove, and K. Fukasawa.** 2000.  
9 Nucleophosmin/B23 is a target of CDK2/cyclin E in centrosome duplication. *Cell*  
10 **103**:127-140.
- 11 44. **Okuwaki, M., A. Iwamatsu, M. Tsujimoto, and K. Nagata.** 2001. Identification of  
12 nucleophosmin/B23, an acidic nucleolar protein, as a stimulatory factor for in vitro  
13 replication of adenovirus DNA complexed with viral basic core proteins. *J Mol Biol*  
14 **311**:41-55.
- 15 45. **Okuwaki, M., K. Matsumoto, M. Tsujimoto, and K. Nagata.** 2001. Function of  
16 nucleophosmin/B23, a nucleolar acidic protein, as a histone chaperone. *FEBS Lett*  
17 **506**:272-276.
- 18 46. **Okuwaki, M., M. Tsujimoto, and K. Nagata.** 2002. The RNA Binding Activity of a  
19 Ribosome Biogenesis Factor, Nucleophosmin/B23, Is Modulated by Phosphorylation  
20 with a Cell Cycle-dependent Kinase and by Association with Its Subtype. *Mol Biol*  
21 *Cell* **13**:2016-2030.
- 22 47. **Orrick, L. R., M. O. Olson, and H. Busch.** 1973. Comparison of nucleolar proteins  
23 of normal rat liver and Novikoff hepatoma ascites cells by two-dimensional

- 1 polyacrylamide gel electrophoresis. Proc Natl Acad Sci U S A. **70**:1316-1320.
- 2 48. **Rickards, B., S. J. Flint, M. D. Cole, and G. LeRoy.** 2007. Nucleolin is required for  
3 RNA polymerase I transcription in vivo. Mol Cell Biol **27**:937-948.
- 4 49. **Samad, M. A., M. Okuwaki, H. Haruki, and K. Nagata.** 2007. Physical and  
5 functional interaction between a nucleolar protein nucleophosmin/B23 and adenovirus  
6 basic core proteins. FEBS Lett **581**:3283-3288.
- 7 50. **Santoro, R., and I. Grummt.** 2005. Epigenetic mechanism of rRNA gene silencing:  
8 temporal order of NoRC-mediated histone modification, chromatin remodeling, and  
9 DNA methylation. Mol Cell Biol **25**:2539-2546.
- 10 51. **Savkur, R. S., and M. O. Olson.** 1998. Preferential cleavage in pre-ribosomal RNA  
11 by protein B23 endoribonuclease. Nucleic Acids Res. **26**:4508-4515.
- 12 52. **Schneider, D. A., A. Michel, M. L. Sikes, L. Vu, J. A. Dodd, S. Salgia, Y. N.**  
13 **Osheim, A. L. Beyer, and M. Nomura.** 2007. Transcription elongation by RNA  
14 polymerase I is linked to efficient rRNA processing and ribosome assembly. Mol Cell  
15 **26**:217-229.
- 16 53. **Shimoyama, T., K. Kato, M. Miyaji-Yamaguchi, and K. Nagata.** 2005. Synergistic  
17 action of MLL, a TRX protein with template activating factor-I, a histone chaperone.  
18 FEBS Lett **579**:757-762.
- 19 54. **Strohner, R., A. Nemeth, P. Jansa, U. Hofmann-Rohrer, R. Santoro, G. Langst,**  
20 **and I. Grummt.** 2001. NoRC--a novel member of mammalian ISWI-containing  
21 chromatin remodeling machines. EMBO J **20**:4892-4900.
- 22 55. **Sung, M. T., M. A. Lischwe, J. C. Richards, and K. Hosokawa.** 1977. Adenovirus  
23 chromatin I. Isolation and characterization of the major core protein VII and precursor

- 1 Pro-VII. J Biol Chem **252**:4981-4987.
- 2 56. **Suzuki, T., S. Muto, S. Miyamoto, K. Aizawa, M. Horikoshi, and R. Nagai.** 2003.
- 3 Functional interaction of the DNA-binding transcription factor Sp1 through its
- 4 DNA-binding domain with the histone chaperone TAF-I. J Biol Chem
- 5 **278**:28758-28764.
- 6 57. **Swaminathan, V., A. H. Kishore, K. K. Febitha, and T. K. Kundu.** 2005. Human
- 7 histone chaperone nucleophosmin enhances acetylation-dependent chromatin
- 8 transcription. Mol Cell Biol **25**:7534-7545.
- 9 58. **von Lindern, M., S. van Baal, J. Wiegant, A. Raap, A. Hagemeijer, and G.**
- 10 **Grosveld.** 1992. Can, a putative oncogene associated with myeloid leukemogenesis,
- 11 may be activated by fusion of its 3' half to different genes: characterization of the set
- 12 gene. Mol Cell Biol **12**:3346-3355.
- 13
- 14

1 **FIGURE LEGENDS**

2

3 **Figure 1** Relationship between the cell growth and the level of rRNA gene transcription.

4 (A) Expression level of B23. 293T cells were maintained in DMEM in the  
5 presence of 1% (lanes 1 and 3) or 10% (lanes 2 and 4) of FBS for 24 h (lanes 1 and 2) or 48 h  
6 (lanes 3 and 4), and then pulse-labeled with [<sup>35</sup>S]methionine for 1 h. [<sup>35</sup>S]-labeled B23  
7 proteins were immunopurified using anti-B23 antibody and analyzed by electrophoresis on a  
8 10% SDS-PAGE followed by autoradiography. As a loading control, 10% of input lysates  
9 was separated on a 10% SDS-PAGE, and  $\beta$ -actin was detected by immunoblotting. (B)  
10 Expression level of nucleolus-related genes. 293T cells were maintained in DMEM in the  
11 presence of 1% (lanes 1 and 3) or 10% (lanes 2 and 4) of FBS for 24 h (lanes 1 and 2) or 48 h  
12 (lanes 3 and 4). Expression level of each indicated proteins were analyzed by  
13 immunoblotting. (C) Transcription rate of the rRNA gene. 293T cells were maintained in  
14 DMEM in the presence of 1% or 10% of FBS for 72 h. The expression level of the rRNA  
15 gene was detected by run-on assays using [ $\alpha$ -<sup>32</sup>P]GTP. Labeled rRNA was used as a probe  
16 for hybridization to membrane-blotted plasmid DNAs containing the rRNA gene  
17 (pUC-119-HurDNA) and pUC119 as a control. Results are represented as mean values  $\pm$  SD  
18 from three independent experiments. (D) Accumulation level of pre-rRNA. 293T cells  
19 were maintained in DMEM in the presence of 1% or 10% of FBS for 72 h. The expression  
20 level of pre-rRNA and  $\beta$ -actin mRNA as a loading control was detected by quantitative  
21 RT-PCR (left and middle panels). The amount of pre-rRNA was normalized with that of  
22  $\beta$ -actin mRNA in the right panel. Results are represented as mean values  $\pm$  SD from three  
23 independent experiments.

1

2 **Figure 2** Repression of the rRNA gene transcription by siB23

3 (A) Expression level of B23.1. In the left panel, immunoblotting was performed  
4 using lysates prepared from HeLa cells treated with 40 pmol of control siRNA (lane 1) and  
5 siB23-A (lane 2), B (lane 3), and C (lane 4) (siRNA for B23.1). The mRNA level of B23.1  
6 was also decreased by transfection of siB23-C compared with control siRNA (Fig. S2 in  
7 supplemental material). (B) Expression level of nucleolus-related proteins. The expression  
8 level of indicated each protein was determined by immunoblotting using HeLa cells treated  
9 with control siRNA and siB23-C. (C) Expression of B23 in nucleoli. Cellular B23 was  
10 stained by indirect immunofluorescence method using anti-B23 antibody in HeLa cells treated  
11 with control siRNA or siB23-C. Nuclei were visualized with DAPI. Arrows indicate the  
12 position of nucleoli. (D) Quantitative determination of pre-rRNA by RT-PCR. RNA  
13 isolated from HeLa cells treated with control siRNA or siB23-C was examined by RT-PCR  
14 using 5' ETS-specific primers for pre-rRNA (left panel) and primers for  $\beta$ -actin (middle  
15 panel). The amount of pre-rRNA was normalized with that of  $\beta$ -actin mRNA (right panel).  
16 Results are represented as mean values  $\pm$  SD from three independent experiments.

17

18 **Figure 3** Association of B23 with the chromatin around the rRNA gene

19 (A) Nucleolar ChIP analysis using quantitative PCR (Q-PCR). DNA was extracted  
20 from soluble nucleolar chromatin immunoprecipitated using anti-Flag, anti-B23, and anti-TBP  
21 antibodies. Q-PCR was performed using promoter-specific primers for the rRNA gene and  
22 ChIPed DNA as templates. Results are represented as mean values  $\pm$  SD from three  
23 independent experiments. (B) ChIP analysis using Q-PCR through whole rRNA gene unit.

1 The structure of the human rRNA gene unit is shown under the graph. Grey boxes indicate  
2 18S, 5.8S, and 28S rRNA encoding sequences. Horizontal arrow indicates the transcription  
3 initiation site (+1) of the rRNA gene transcribed by Pol I. DNA was extracted from soluble  
4 chromatin immunoprecipitated using anti-Flag and anti-B23 antibodies. Q-PCR was  
5 performed using ChIPed DNA as templates and primer sets indicated in Table 1. Results  
6 are represented as mean values  $\pm$  SD from three independent experiments. (C) ChIP  
7 analysis using Q-PCR. 293T cells were maintained in DMEM in the presence of 1% or 10%  
8 of FBS for 72 h. DNA was extracted from chromatin that was immunoprecipitated using  
9 anti-Flag, and anti-B23 antibodies. Q-PCR was performed using promoter specific primers  
10 for the rRNA gene and ChIPed DNA as templates. Results are represented as mean values  $\pm$   
11 SD from three independent experiments. (D) Histone density on the rRNA gene promoter.  
12 Soluble chromatin fraction from HeLa cells treated with control siRNA or siB23-C was  
13 examined by ChIP analyses using anti-Flag, anti-B23, anti-UBF, anti-TBP, anti-acetylated  
14 histone H3, and anti-histone H3 antibodies. Q-PCR was performed using promoter-specific  
15 primers for the rRNA gene and ChIPed DNA as templates. Results are represented as mean  
16 values  $\pm$  SD from three independent experiments. (E) Histone distribution over the rRNA  
17 gene. At the same time in (D), Q-PCR was carried out over the rRNA gene (upper panel).  
18 The ratio of the histone density over the rRNA gene in cells treated with siB23-C relative to  
19 that in cells treated with control siRNA is plotted (middle panel). Results are represented as  
20 mean values  $\pm$  SD from three independent experiments.

21

22 **Figure 4** Inhibition of the histone binding activity of B23.1 by B23 $\Delta$ C *in vitro*

23 (A) Schematic representation of the structure of B23.1, B23 $\Delta$ A, and B23 $\Delta$ C. Grey,



1 black, and striped boxes indicate oligomeric domain, acidic domain, and RNA binding  
2 domain, respectively. **(B)** Purification of recombinant proteins. Recombinant His-B23.1,  
3 His-B23 $\Delta$ A, and His-B23 $\Delta$ C were purified, and proteins (400 ng) were separated with a  
4 12.5% SDS-PAGE followed by CBB staining. **(C)** Histone binding activity of B23 mutants.  
5 Recombinant B23 proteins (500 ng) were mixed with 300 ng core histones purified from  
6 HeLa cells. Immunoprecipitation was carried out using an anti-His antibody. Precipitated  
7 proteins were analyzed by a 12.5% SDS-PAGE followed by immunoblotting with anti-His  
8 and anti-histone H3 ( $\alpha$ -H3) antibodies. **(D)** Nucleosome assembly activity of B23 mutants.  
9 The nucleosome assembly activity of B23 mutants was examined by supercoiling assay.  
10 Core histones (200 ng) preincubated without (lane 1) or with His-B23.1 (167 ng for lane 2,  
11 500 ng for lane 3, and 1500 ng for lane 4), His-B23 $\Delta$ A (142 ng for lane 5, 425 ng for lane 6,  
12 and 1275 ng for lane 7), and His-B23 $\Delta$ C (66.8 ng for lane 8, 200 ng for lane 9, and 600 ng for  
13 lane 10), were mixed with closed circular DNA relaxed by topoisomerase I and further  
14 incubated. The amounts of B23 mutant proteins were adjusted according to their sizes so as  
15 to be added at the same molecular numbers. The DNA was purified and separated by  
16 electrophoresis on a 1% agarose gel, and visualized by staining with ethidium bromide.  
17 Positions of relaxed (R), supercoiled (S), or nicked (N) circular plasmid DNA are indicated.  
18 **(E)** Inhibition of the histone binding activity of B23.1 by B23 $\Delta$ C. His-B23.1 (200 ng) was  
19 mixed without (lane 2) or with 200 ng (lane 3) or 400 ng (lane 4) of B23 $\Delta$ C and subjected to  
20 the denature-renature protocol. Renatured B23 complexes were mixed with 300 ng core  
21 histones purified from HeLa cells. Immunoprecipitation of His-B23.1 was carried out using  
22 an anti-B23 antibody that can not recognize B23 $\Delta$ C protein. Precipitated proteins were  
23 analyzed by a 12.5% SDS-PAGE followed by silver staining (left panel) and immunoblotting

1 (IB) with an anti-His antibody (right panel). H and L indicate heavy and light chains of  
2 immunoglobulin, respectively. The position of core histones are indicated beside the left  
3 panel.

4  
5 **Figure 5** Inhibition of the histone binding activity of B23.1 by B23 $\Delta$ C *in vivo*

6 (A) Expression of B23 proteins in 293T cells. Immunoblotting was carried out  
7 using lysate prepared from 293T cells transfected with each plasmids indicated above each  
8 lane. Left and right panels indicate immunoblotting by anti-HA and anti-Flag antibodies,  
9 respectively. (B) Histone binding activity of B23 mutants. Immunoprecipitation (IP) with  
10 anti-HA and anti-Flag antibodies were performed using lysate of 293T cells transfected with  
11 plasmids expressing B23 proteins. Immunoprecipitated proteins were analyzed by  
12 immunoblotting using anti-Flag, anti-HA, and anti-histone H3 antibodies, respectively. (C)  
13 No effect of B23 $\Delta$ A on the histone binding activity of B23.1. Lysates of 293T cells (24 h  
14 post transfection) transfected with pCHA-B23.1 (160 ng for lanes 2-5, and 7-10) and  
15 pCAGGS-Flag-B23 $\Delta$ A (160 ng for lanes 3 and 8, 480 ng for lanes 4 and 9, and 1440 ng for  
16 lanes 5 and 10), were subjected to immunoprecipitation with an anti-HA antibody.  
17 Immunoprecipitated proteins were analyzed by immunoblotting using anti-Flag, anti-HA, and  
18 anti-histone H3 antibodies, respectively. (D) Inhibition of the histone binding activity of  
19 B23.1 by B23 $\Delta$ C *in vivo*. Lysates of 293T cells (24 h post transfection) transfected with  
20 pcDNA3-Flag-B23.1 (100 ng for lanes 2-4, and 6-8) and pCHA-B23 $\Delta$ C (300 ng for lanes 3  
21 and 7, and 1400 ng for lanes 4 and 8), were subjected to immunoprecipitation with anti-Flag  
22 antibody. Immunoprecipitated proteins were analyzed by immunoblotting using anti-Flag,  
23 anti-HA, and anti-histone H3 antibodies, respectively.

1

2 **Figure 6** Inhibition of the rRNA gene transcription by B23ΔC

3 (A) Interaction of B23ΔC with endogenous B23. 293T cells were transiently  
4 transfected with pCHA (control) and pCHA-B23ΔC, and cultured for 72 h post transfection.  
5 Lysates prepared from 293T cells were used for immunoprecipitation with anti-HA antibody  
6 (lanes 3, 4, 7, and 8). Immunoprecipitates (P) and flow-through fraction (F) were resolved  
7 by a 12.5% SDS-PAGE, transferred to a membrane, and probed with anti-B23 (upper panel)  
8 and anti-HA (lower panel) antibodies. (B) Inhibition of association of B23 with the rRNA  
9 gene by B23ΔC. 293T cells were co-transfected with pBabe-puro and indicated plasmids  
10 (pCHA, pCHA-B23.1, and pCHA-B23ΔC), and 0.5 μg/ml puromycin was added after 24 h  
11 post transfection. After puromycin selection for 30 h, 293T cells were maintained in fresh  
12 medium for 18 h. Nucleolar ChIP was carried out using anti-B23 and anti-Flag antibodies.  
13 Quantitative PCR was performed with primers around the promoter region of the rRNA gene.  
14 Results are represented as mean values ± SD from three independent experiments. A paired  
15 Student's *t*-test gave  $P=0.02$  between pCHA- and pCHA-B23ΔC-transfected cells. (C)  
16 Association of transcription factors on the rRNA gene promoter. ChIP was carried out using  
17 anti-UBF and anti-TBP antibodies and lysates prepared for mock-treated cells or cells  
18 expressing B23ΔC. Q-PCR was performed with primers around the promoter region of the  
19 rRNA gene. Results are represented as mean values ± SD from three independent  
20 experiments. (D) Histone density on the rRNA gene promoter. ChIP was carried out using  
21 anti-B23, anti-histone H3, and anti-acetylated histone H3 (AcH3) antibodies in the presence  
22 or absence of B23ΔC. Q-PCR was performed with primers around the promoter region of  
23 the rRNA gene. Results are represented as mean values ± SD from three independent

1 experiments. (E) Histone distribution over the rRNA gene. ChIP was performed with  
2 anti-histone H3 antibody using cells expressing HA-B23 $\Delta$ C or mock-treated cells. Amount  
3 of ChIPed DNA was determined by Q-PCR using primer sets over the rRNA gene (upper  
4 panel). The ratio of the histone density over the rRNA gene in cells expressing B23 $\Delta$ C  
5 relative to that in mock-treated cells is plotted (middle panel). Results are represented as  
6 mean values  $\pm$  SD from three independent experiments. (F) Inhibition of the rRNA gene  
7 transcription by B23 $\Delta$ C. 293T cells were transiently transfected with pCHA (control) and  
8 pCHA-B23 $\Delta$ C, and cultured for 72 h post transfection. Nuclei isolated from these 293T  
9 cells were used for run-on transcription assays. Radiolabeled nuclear RNA was isolated and  
10 hybridized with 1  $\mu$ g of pUC119 (control) and pUC119-HurDNA (rRNA) on a membrane.  
11 This assay was done in triplicate, and a typical pattern is shown. The right panel shows a  
12 densitometry quantitation of the hybridization signals. Results are represented as mean  
13 values  $\pm$  SD from three independent experiments. (G) Inhibition of the rRNA gene  
14 transcription by B23 $\Delta$ A. 293T cells were co-transfected with pBabe-puro and either one of  
15 indicated plasmids (pCHA, pCHA-B23 $\Delta$ C, and pCHA-B23 $\Delta$ A), and 0.5  $\mu$ g/ml puromycin  
16 was added after 24 h post transfection. After puromycin selection for 30 h, 293T cells were  
17 maintained in fresh medium for 18 h. RNA isolated from 293T cells was examined by  
18 RT-PCR using 5' ETS-specific primers for pre-rRNA (left panel) and primers for  *$\beta$ -actin*  
19 mRNA (middle panel). Results are represented as mean values  $\pm$  SD from three independent  
20 experiments.

21

22 **Figure 7** Involvement of acidic domains of B23 in the rRNA gene transcription

23 (A) Expression of B23 proteins in HeLa cells transfected with siB23. HeLa cells

1 were transfected with 40 pmol of control siRNA (lane 1) and 40 pmol of siB23-C (lanes 2-8).  
2 At 12 h after siRNAs transfection, HeLa cells were super-transfected with  
3 pCAGGS-Flag-B23.1 (180 ng for lane 3, and 360 ng for lane 4), pCAGGS-Flag B23ΔA (180  
4 ng for lane 5, and 360 ng for lane 6), pCHA-B23ΔC (180 ng for lane 7, and 360 ng for lane 8),  
5 and pEF321-T (40 ng for lanes 1-8) encoding SV40 T antigen for amplification of transfected  
6 plasmid. After 72 h post siRNAs transfection, lysate of  $2.5 \times 10^3$  cells were analyzed by  
7 immunoblotting using anti-B23, anti-HA, and anti-β-actin antibodies, respectively. **(B)**  
8 Quantitative determination of pre-rRNA by Q-PCR. At the same time with (A), total RNA  
9 extracted from HeLa cells were examined by RT-PCR using 5'ETS-specific primers for the  
10 rRNA gene. The amount of pre-rRNA was normalized with that of β-actin mRNA. Results  
11 are represented as mean values ± SD from three independent experiments.

12

### 13 **Figure 8** Inhibition of cell proliferation by B23ΔC

14 **(A)** Proliferation inhibition of 293T cells by expression of B23ΔC. 293T cells  
15 were co-transfected with 1400 ng of pCHA and 200 ng of pEGFP-N1 or with 1400 ng of  
16 pCHA- B23ΔC and 200 ng of pEGFP-N1. The number of EGFP-positive cells was counted  
17 every 24 hrs as a function of culture time. Results are represented as mean values ± SD from  
18 three independent experiments. **(B)** B23 dose-dependent inhibition of the cell proliferation  
19 rate. The number of EGFP-positive cells was counted at 72 h after transfection of 200 ng of  
20 pEGFP-N1 with pCHA-B23ΔC (350 ng (lane 2), 700 ng (lane 3), and 1400 ng (lane 4)), and  
21 pCHA (1400 ng (lane 1), 700 ng (lane 2), and 350 ng (lane 3)). Results are represented as  
22 mean values ± SD from three independent experiments. **(C)** Cell cycle population of cells  
23 expressing B23ΔC. 293T cells were co-transfected with 1400 ng of pCHA and 200 ng of

1 pEGFP-N1 or with 1400 ng of pCHA-B23ΔC and 200 ng of pEGFP-N1, and then incubated  
2 for 72 h. Cell population of 293T cells expressing EGFP was determined by FACS analysis  
3 (left panel). Sub-G1 fraction was quantitatively determined as a population of apoptotic  
4 cells and shown in the right inset table. **(D)** p53-independent effect of B23ΔC on the cell  
5 proliferation. The number of EGFP-positive p53<sup>-/-</sup> MEF cells was counted at 72 h post  
6 transfection of pEGFP-N1 (125 ng) with pCHA-B23.1 (875 ng) or pCHA-B23ΔC (875 ng).  
7 Results are represented as mean values ± SD from three independent experiments.

8