

ON FUNCTION SPACES FOR GENERAL TOPOLOGICAL SPACES

By

Mitsuaki KAKIMI and Toshio YAGI

1. Introduction.

In this paper we mean by a space a topological space with no separation axiom unless otherwise specified, and we denote by \mathbf{R} and \mathbf{I} the real line and the closed unit interval respectively.

Given two spaces X and Y , let $F(X, Y)$ denote the set of all maps from X into Y , $C(X, Y)$ the set of all continuous maps from X into Y . In case Y is the real line \mathbf{R} , $C(X, \mathbf{R})$ is denoted more simply by $C(X)$. The map $\rho: F(X \times Y, T) \rightarrow F(Y, F(X, T))$ defined by the formula $[\rho(f)(y)](x) = f(x, y)$ for $f \in F(X \times Y, T)$ is bijective; this correspondence is called the *exponential map*.

A topology on $C(X, T)$ is called *proper* if for every space Y and any $f \in C(X \times Y, T)$ the map $\rho(f)$ belongs to $C(Y, C(X, T))$. Similarly, a topology on $C(X, T)$ is called *admissible* if for every space Y and any $g \in C(Y, C(X, T))$ the map $\rho^{-1}(g)$ belongs to $C(X \times Y, T)$. A topology on $C(X, T)$ that is both proper and admissible is called an *acceptable topology* (see [1], [2] and [3]).

As is well known, the compact-open topology on $C(X, T)$ is acceptable for any space T when X is locally compact Hausdorff (see [4]). Furthermore, the following theorem was proved by R. Arens [1].

THEOREM 1.1. *Let X be a Tychonoff space. Then the following conditions are equivalent.*

- (1) X is locally compact.
- (2) There exists an acceptable topology on $C(X)$.

In the case that X is not necessarily Tychonoff, Professor T. Ishii raised the following problem: Characterize a space X such that there exists an acceptable topology on $C(X)$.

The main purpose of this paper is to give the solution for this problem by proving the following theorem.

THEOREM 1.2. *For a space X , the following conditions are equivalent.*

- (1) *X is locally relatively w -compact (that is, every point of X has a relatively w -compact nbd (=neighborhood)).*
- (2) *There exists an acceptable topology on $C(X, T)$ for any Tychonoff space T .*
- (3) *There exists an acceptable topology on $C(X)$.*

The definition of relatively w -compact subsets is given in section 2. Section 3 is devoted to a study of a new topology on $C(X, T)$ which is acceptable when X is locally relatively w -compact. Theorem 1.2 is proved in section 4.

The authors wish to thank Professor T. Ishii for his valuable comments.

2. Properties of relatively w -compact subsets.

A subset P of a space X is called τ -open if P is a union of cozero-sets of X (see [6]). For any subset A of X , we call the intersection of all zero-sets of X containing A the τ -closure of A , and we denote it by \bar{A}^τ . A subset A of X is said to be τ -closed if $A = \bar{A}^\tau$ holds.

DEFINITION 2.1. A subset A of a space X is *relatively w -compact* if for any family $\{P_\lambda | \lambda \in A\}$ of τ -open sets of X such that $\{A \cap P_\lambda | \lambda \in A\}$ has the f.i.p. (=finite intersection property), we have $\bigcap \{cl P_\lambda | \lambda \in A\} \neq \emptyset$.

T. Ishii introduced the notion of w -compact spaces, in connection with the problem concerning a product formula for the Tychonoff functor ([6]). A space X is called *w -compact* if for any family $\{P_\lambda | \lambda \in A\}$ of τ -open sets of X with the f.i.p., we have $\bigcap \{cl P_\lambda | \lambda \in A\} \neq \emptyset$. Clearly every w -compact subset of a space X is relatively w -compact.

PROPOSITION 2.2. *Let A be a subset of a space X . Then the following conditions are equivalent.*

- (1) *A is relatively w -compact.*
- (2) *For any collection $\{A_\alpha | \alpha \in \Omega\}$ of closed sets of X such that it is closed under the finite intersection and each A_α contains a cozero-set of X which meets A , we have $\bigcap \{A_\alpha | \alpha \in \Omega\} \neq \emptyset$.*
- (3) *For every open cover $\{U_\alpha | \alpha \in \Omega\}$ of X there exists a finite set $\{\alpha(1), \dots, \alpha(n)\} \subset \Omega$ such that $A \subset \bigcup \{\bar{U}_{\alpha(i)}^\tau | i=1, \dots, n\}$.*
- (4) *For every family $\{U_\alpha | \alpha \in \Omega\}$ of open sets of X such that $\bar{A}^\tau \subset \bigcup \{U_\alpha | \alpha \in \Omega\}$ there exists a finite set $\{\alpha(1), \dots, \alpha(n)\} \subset \Omega$ such that $A \subset \bigcup \{\bar{U}_{\alpha(i)}^\tau | i=1, \dots, n\}$.*

PROOF. The equivalences of (1) and (2) and of (1) and (3) are clear. And

the implication (4) \Rightarrow (3) is obvious. We prove only (3) \Rightarrow (4).

Let $\{U_\alpha | \alpha \in \mathcal{Q}\}$ be a family of open sets of X with $\bar{A}^\tau \subset \cup \{U_\alpha | \alpha \in \mathcal{Q}\}$. For each $x \in X - \bar{A}^\tau$ we can take a zero-set nbd of x which misses \bar{A}^τ since $X - \bar{A}^\tau$ is a union of cozero-sets of X . Thus we obtain an open nbd V_x of x with $\bar{V}_x \cap \bar{A}^\tau = \emptyset$. Then $\{U_\alpha | \alpha \in \mathcal{Q}\} \cup \{V_x | x \in X - \bar{A}^\tau\}$ is an open cover of X . From this fact (4) follows from (3).

As is easily seen, a relatively w -compact subset of a Tychonoff space is relatively compact. Hence, if a Tychonoff space is locally relatively w -compact, then it is locally compact. However, there exists a regular T_1 -space on which every continuous real-valued function is constant (for instance, see [5]). This example shows that there exists a locally relatively w -compact regular T_1 -space that is not locally compact. Hence, Theorem 1.2 shows that in case X is not Tychonoff, the local compactness of X is unessential in Theorem 1.1.

Let X and Y be two spaces and A a subset of X . The projection $\pi_Y: A \times Y \rightarrow Y$ is called a *relative Z -map* if $\pi_Y((A \times Y) \cap Z)$ is closed in Y for any zero-set Z of $X \times Y$ (see [10]).

The following proposition is a generalization of [6, Proposition 2.7].

PROPOSITION 2.3. *Let A be a subset of a space X . Then the following conditions are equivalent.*

- (1) *A is relatively w -compact.*
- (2) *For any space Y the set $\pi_Y((\bar{A}^\tau \times Y) \cap F)$ is closed in Y for every τ -closed subset F of $X \times Y$, where π_Y is the projection: $\bar{A}^\tau \times Y \rightarrow Y$.*
- (3) *The projection $\pi_Y: \bar{A}^\tau \times Y \rightarrow Y$ is a relative Z -map for any space Y .*
- (4) *The projection $\pi_Y: \bar{A}^\tau \times Y \rightarrow Y$ is a relative Z -map for any paracompact Hausdorff space Y .*

PROOF. (1) \Rightarrow (2). Let A be a relatively w -compact subset of X and F a τ -closed set of $X \times Y$. Take a point $y_0 \in Y - \pi_Y((\bar{A}^\tau \times Y) \cap F)$. Since $(\bar{A}^\tau \times \{y_0\}) \cap F = \emptyset$ and $X \times Y - F$ is τ -open, for each $x \in \bar{A}^\tau$ the point (x, y_0) has an open nbd of the form $U_x \times V_x$ such that $(\bar{U}_x \times V_x) \cap F = \emptyset$. Clearly $\bar{A}^\tau \subset \cup \{U_x | x \in \bar{A}^\tau\}$, so that by Proposition 2.2, there exists a finite set $\{x(1), \dots, x(n)\} \subset \bar{A}^\tau$ such that $A \subset \cup \{\bar{U}_{x(i)} | i=1, \dots, n\}$. Put $V = \cap \{V_{x(i)} | i=1, \dots, n\}$. Then V is an open nbd of y_0 and $V \cap \pi_Y((\bar{A}^\tau \times Y) \cap F) = \emptyset$. It follows that $\pi_Y((\bar{A}^\tau \times Y) \cap F)$ is closed in Y .

(2) \Rightarrow (3) and (3) \Rightarrow (4) are obvious and (4) \Rightarrow (1) is easily verified by making use of the method of [6, Proposition 2.6]. Thus the proof is completed.

Let A be a relatively w -compact subset of a space X . Then, by Proposition

2.2, we clearly have the following facts:

- (1) \bar{A}^τ is also relatively w -compact.
- (2) If $B \subset A$, then B is also relatively w -compact.

Let X and Y be two spaces. Then $\overline{A \times B}^\tau = \bar{A}^\tau \times \bar{B}^\tau$ holds for $A \subset X$ and $B \subset Y$. Thus, the following proposition is easy to prove.

PROPOSITION 2.4. *Let X and Y be two spaces. If A and B are relatively w -compact subsets of X and Y respectively, then $A \times B$ is also a relatively w -compact subset of $X \times Y$.*

The following proposition is also clear.

PROPOSITION 2.5. *Let X and Y be two spaces and f a map in $C(X, Y)$. If A is a relatively w -compact subset of X , then $f(A)$ is a relatively w -compact subset of Y .*

3. A topology on function spaces.

In this section, we consider a new topology on function spaces. Let X be a space and T a Tychonoff space. For $A \subset X$ and $B \subset T$ we denote by $M(A, B)$ the totality of maps f in $C(X, T)$ for which $f(A) \subset B$. We consider the topology on $C(X, T)$ generated by the base consisting of all sets $\bigcap \{M(\bar{A}_i^\tau, U_i) \mid i=1, \dots, n\}$, where A_i is a relatively w -compact subset of X and U_i is an open subset of T for $i=1, \dots, n$. Throughout this section, a topology on function spaces is assumed to be the topology defined above.

The following proposition easily follows from Proposition 2.3.

PROPOSITION 3.1. *Let X be a space and T a Tychonoff space. Then the topology on $C(X, T)$ is proper.*

The following lemma is also clear.

LEMMA 3.2. *Let X be a locally relatively w -compact space. Then for each point x of X and for any τ -open nbd G of x there exists a relatively w -compact nbd U of x such that $\bar{U}^\tau \subset G$.*

PROPOSITION 3.3. *Let X be a locally relatively w -compact space and T a Tychonoff space. Then the topology on $C(X, T)$ is acceptable.*

PROOF. By Proposition 3.1, it suffices to prove that the topology on $C(X, T)$

is admissible. Let Y be a space and g a map in $C(Y, C(X, T))$. We shall show that $\rho^{-1}(g)$ is continuous. Take a point (x_0, y_0) in $X \times Y$ and an open nbd U of $\rho^{-1}(g)(x_0, y_0)$ in T . Here, notice $\rho^{-1}(g)(x_0, y_0) = [g(y_0)](x_0)$. Since T is a Tychonoff space and $g(y_0)$ is continuous, $g(y_0)^{-1}(U)$ is a τ -open nbd of x_0 . By Lemma 3.2, there exists a relatively w -compact nbd V of x_0 such that $\bar{V}^\tau \subset g(y_0)^{-1}(U)$. Hence we have $g(y_0) \in M(\bar{V}^\tau, U)$. Because of the continuity of g there exists a nbd W of y_0 such that $g(W) \subset M(\bar{V}^\tau, U)$. Therefore we obtain a nbd $V \times W$ of (x_0, y_0) such that $\rho^{-1}(g)(V \times W) \subset U$. Hence $\rho^{-1}(g)$ is continuous. This completes the proof.

Let X be a space, T a Tychonoff space, and f a map in $C(X, T)$. Then it is easily shown that $f(\bar{A}^\tau)$ is compact for any relatively w -compact subset A of X . From this fact, one can easily prove the following lemma and proposition (see [3, 3.4.14 and 3.4.15]).

LEMMA 3.4. *Let X be a space and A a relatively w -compact subset of X . Assigning to each $f \in C(X, I)$ the number $\xi(f) = \sup\{f(x) \mid x \in \bar{A}^\tau\}$ defines a continuous function $\xi: C(X, I) \rightarrow I$.*

PROPOSITION 3.5 *Let X be a space and T a Tychonoff space. Then $C(X, T)$ is also a Tychonoff space.*

LEMMA 3.6. *Let X be a space, T a Tychonoff space, and \mathcal{B} a subbase for T . Then the sets $M(\bar{A}^\tau, U)$, where A is a relatively w -compact subset of X and $U \in \mathcal{B}$, form a subbase for the space $C(X, T)$.*

PROOF. Let A be a relatively w -compact subset of X , U an open set of T , and f a map in $M(\bar{A}^\tau, U)$. For each $x \in \bar{A}^\tau$ we can take sets $U_1^x, \dots, U_{n(x)}^x \in \mathcal{B}$ with $x \in W_x = \bigcap \{f^{-1}(U_j^x) \mid j=1, \dots, n(x)\}$ and $\bigcap \{U_j^x \mid j=1, \dots, n(x)\} \subset U$. Since W_x is a τ -open nbd of x , we can take an open nbd V_x of x such that $\bar{V}_x^\tau \subset W_x$. By Proposition 2.2, there exists a finite set $\{x(1), \dots, x(k)\} \subset \bar{A}^\tau$ such that $\bar{A}^\tau \subset \bigcup \{\bar{V}_{x(i)}^\tau \mid i=1, \dots, k\}$. Put $A_i = \bar{A}^\tau \cap \bar{V}_{x(i)}^\tau$. Clearly, A_i is relatively w -compact, and we have $\bar{A}^\tau = \bigcup \{\bar{A}_i^\tau \mid i=1, \dots, k\}$ and $\bar{A}_i^\tau \subset \bigcap \{f^{-1}(U_j^i) \mid j=1, \dots, n(i)\}$, where $U_j^i = U_{x(i)}^j$ and $n(i) = n(x(i))$. Therefore, $f \in \bigcap \{M(\bar{A}_i^\tau, U_j^i) \mid j=1, \dots, n(i) \mid i=1, \dots, k\} \subset M(\bar{A}^\tau, U)$. Thus the proof is completed.

THEOREM 3.7. *Let X and Y be two spaces and T a Tychonoff space. Then the exponential map $\rho: C(X \times Y, T) \rightarrow C(Y, C(X, T))$ is a homeomorphic embedding.*

PROOF. We first notice that $\rho(C(X \times Y, T)) \subset C(Y, C(X, T))$ by Proposition 3.1.

Let A and B be relatively w -compact subsets of X and Y respectively, and U an open set of T . Then we clearly have $\rho^{-1}[M(\bar{B}^\tau, M(\bar{A}^\tau, U))] = M(\bar{A}^\tau \times \bar{B}^\tau, U)$.

Since $\bar{A}^\tau \times \bar{B}^\tau$ is relatively w -compact by Proposition 2.4, the last lemma implies that ρ is continuous.

The above equality implies that

$$\rho(M(\bar{A}^\tau \times \bar{B}^\tau, U)) = M(\bar{B}^\tau, M(\bar{A}^\tau, U)) \cap \rho(C(X \times Y, T));$$

hence— ρ being a one-to-one map—it suffices to show that the sets $M(\bar{A}^\tau \times \bar{B}^\tau, U)$, where A and B are relatively w -compact subsets of X and Y respectively and U is open in T , form a subbase for $C(X \times Y, T)$.

Take a relatively w -compact subset C of $X \times Y$, an open set U of T and a map $f \in M(\bar{C}^\tau, U)$. Since C is relatively w -compact and $f^{-1}(U)$ is τ -open, there exist open sets $V_1, \dots, V_n \subset X$ and $W_1, \dots, W_n \subset Y$ with $\bar{C}^\tau \subset \bigcup \{\bar{V}_i^\tau \times \bar{W}_i^\tau \mid i=1, \dots, n\} \subset f^{-1}(U)$. Let $A_i = \pi_X(\bar{C}^\tau) \cap \bar{V}_i^\tau$ and $B_i = \pi_Y(\bar{C}^\tau) \cap \bar{W}_i^\tau$ for $i=1, \dots, n$, where $\pi_X: X \times Y \rightarrow X$ and $\pi_Y: X \times Y \rightarrow Y$ are the projections. Then A_i and B_i are relatively w -compact subsets of X and Y respectively for $i=1, \dots, n$. Moreover, we have $\bar{A}_i^\tau \subset \bar{V}_i^\tau$, $\bar{B}_i^\tau \subset \bar{W}_i^\tau$ and $\bar{C}^\tau \subset \bigcup \{\bar{A}_i^\tau \times \bar{B}_i^\tau \mid i=1, \dots, n\}$. Hence, we have $f \in \bigcap \{M(\bar{A}_i^\tau \times \bar{B}_i^\tau, U) \mid i=1, \dots, n\} \subset M(\bar{C}^\tau, U)$, and this completes the proof.

Proposition 3.3 and Theorem 3.7 imply

THEOREM 3.8. *Let X be a locally relatively w -compact space, Y a space and T a Tychonoff space. Then the exponential map $\rho: C(X \times Y, T) \rightarrow C(Y, C(X, T))$ is a homeomorphism.*

4. Proof of Theorem 1.2.

The following two lemmas are due to R. Arens and J. Dugundji [2].

LEMMA 4.1. *For spaces X and T , a topology on $C(X, T)$ is admissible if and only if the evaluation mapping $\omega(f, x) = f(x)$ of $C(X, T) \times X$ into T is continuous.*

LEMMA 4.2. *Let X and T be spaces and let \mathcal{T}_1 and \mathcal{T}_2 be topologies on $C(X, T)$. If \mathcal{T}_1 is proper and \mathcal{T}_2 is admissible, then $\mathcal{T}_1 \subset \mathcal{T}_2$.*

We are now in a position to prove the main theorem. We modify the proof by R. Arens ([1, Theorem 3]).

PROOF OF THEOREM 1.2. Since the implication (1) \Rightarrow (2) is a direct consequence of Proposition 3.3 and (2) \Rightarrow (3) is obvious, we prove only (3) \Rightarrow (1).

Suppose that there exists an acceptable topology on $C(X)$ and denote it by \mathcal{T}_{ac} . Let g be the element of $C(X)$ such that $g(x)=0$ for each $x \in X$. Since $\omega: C(X) \times X \rightarrow \mathbf{R}$ is continuous with respect to the topology \mathcal{T}_{ac} on $C(X)$ by Lemma 4.1, for any $x_0 \in X$ there exist an open nbd V of x_0 and an element W of \mathcal{T}_{ac} such that $g \in W$ and $W \times V \subset \omega^{-1}((-1, 1))$.

Now we shall prove that V is relatively w -compact. Let $\{G_\lambda | \lambda \in A\}$ be a family of open sets of X such that $\bar{V}^\tau \subset \bigcup \{G_\lambda | \lambda \in A\}$ and \mathcal{T} the topology on $C(X)$ with its subbase consisting of the sets of the form $M(\bar{A}^\tau, U)$, where A is a subset of X such that A is contained in some element of $\{G_\lambda | \lambda \in A\}$ or $\bar{A}^\tau \cap \bar{V}^\tau = \emptyset$, and U is open in \mathbf{R} . Then \mathcal{T} is admissible. To see this, it suffices to show that $\omega: C(X) \times X \rightarrow \mathbf{R}$ is continuous with respect to \mathcal{T} by Lemma 4.1. Let U be an open set of \mathbf{R} , and take a point $(f, x) \in \omega^{-1}(U)$. Then $f^{-1}(U)$ is a cozero-set nbd of x in X . Consequently, there exists an open nbd V_1 of x such that $f(\bar{V}_1^\tau) \subset U$.

Case (a). If $x \in \bar{V}^\tau$, then there exists $\lambda \in A$ such that $x \in G_\lambda$. Put $A = V_1 \cap G_\lambda$, then A is an open nbd of x such that $A \subset G_\lambda$.

Case (b). If $x \notin \bar{V}^\tau$, then there is an open nbd V_2 of x such that $\bar{V}_2^\tau \cap \bar{V}^\tau = \emptyset$. Put $A = V_1 \cap V_2$, then $\bar{A}^\tau \cap \bar{V}^\tau = \emptyset$.

Hence each of the two cases above implies that there exists an open nbd A of x such that $f(\bar{A}^\tau) \subset U$ and $M(\bar{A}^\tau, U) \in \mathcal{T}$. Furthermore $(f, x) \in M(\bar{A}^\tau, U) \times A \subset \omega^{-1}(U)$. Thus ω is continuous.

This implies $\mathcal{T}_{ac} \subset \mathcal{T}$ by Lemma 4.2, so that there exist subsets A_1, \dots, A_n of X and open subsets U_1, \dots, U_n of \mathbf{R} such that $g \in \bigcap \{M(\bar{A}_i^\tau, U_i) | i=1, \dots, n\} \subset W$. Here, notice $0 \in U_i$ for each i .

Finally, we shall show that $V \subset \bigcup \{\bar{A}_i^\tau | i=1, \dots, n\}$. Assume that there exists a point $x_1 \in V$ such that $x_1 \notin \bigcup \{\bar{A}_i^\tau | i=1, \dots, n\}$. Then there exists a continuous map $h: X \rightarrow \mathbf{I}$ such that $h(x_1)=1$ and $h(x)=0$ for $x \in \bigcup \{\bar{A}_i^\tau | i=1, \dots, n\}$, and so $h \in \bigcap \{M(\bar{A}_i^\tau, U_i) | i=1, \dots, n\}$. Hence $h \in W$. Since $x_1 \in V$ and $W \times V \subset \omega^{-1}((-1, 1))$, we have $h(x_1) \in (-1, 1)$. This is a contradiction. This implies $V \subset \bigcup \{\bar{A}_i^\tau | i=1, \dots, n\}$. Thus V is contained in the union of finitely many members of $\{\bar{G}_\lambda^\tau | \lambda \in A\}$. Hence V is relatively w -compact by Proposition 2.2, and this completes the proof of Theorem 1.2.

5. An example.

A space X is called *locally cozero-set w -compact* if for each point x of X there exists a cozero-set nbd G of x such that clG is w -compact ([6] and [7]). The following theorem, where we denote by τ the Tychonoff functor, was proved

by T. Ishii in [6].

THEOREM 5.1. *A space X is locally cozero-set w -compact if and only if $\tau(X \times Y) = \tau(X) \times \tau(Y)$ for any space Y .*

Clearly every locally cozero-set w -compact space is locally relatively w -compact. But the converse is false. We construct such an example. A space X is called τ -compact if $\tau(X)$ is compact ([7]). The following lemma is easy to prove.

LEMMA 5.2 *Every τ -compact locally cozero-set w -compact space is w -compact.*

EXAMPLE 5.3. Let ω_1 be the first uncountable ordinal and let us put $S = W(\omega_1 + 1) \times W(\omega_1 + 1) - \{(\omega_1, \omega_1)\}$, where $W(\omega_1 + 1)$ is the space of all ordinals less than $\omega_1 + 1$ with the usual interval topology. Now let X be a space obtained by adding a new point ξ to S and introducing the topology in X as follows: the base at ξ is given by the totality of the sets $U_\beta(\xi) = \{(\alpha, \alpha) \mid \alpha \text{ is a non-limit ordinal and } \beta < \alpha\} \cup \{\xi\}$, $\beta < \omega_1$ and the base at $x \neq \xi$ is the same as in S . Then X has the following properties:

- (1) X is Hausdorff but not regular.
- (2) X is τ -compact.
- (3) X is not w -compact.
- (4) X is locally relatively w -compact.

Indeed (1) is obvious and (2) follows from the fact that any cozero-set of X containing ξ has to contain a set of the form $\{\xi\} \cup T_\alpha$ for some $\alpha < \omega_1$, where $T_\alpha = \{(\lambda, \mu) \mid \lambda, \mu > \alpha\} \subset S$, and (3) follows from the fact that $\{T_\alpha - U_\alpha(\xi) \mid \alpha < \omega_1\}$ is a family of closed sets of X such that $T_\alpha - U_\alpha(\xi)$ contains isolated points of X and $\bigcap \{T_\alpha - U_\alpha(\xi) \mid \alpha < \omega_1\} = \emptyset$ (see [6, p. 175]). Furthermore, each point $x \in S$ has a compact nbd, and $A_\beta = \{(\alpha, \alpha) \mid \beta < \alpha\} \cup \{\xi\}$ for $\beta < \omega_1$ is a relatively w -compact nbd of ξ . To show this, take an open cover $\{G_\lambda \mid \lambda \in A\}$ of X and a $\theta \in A$ such that $\xi \in G_\theta$. Then there exists a $\gamma < \omega_1$ such that $U_\gamma(\xi) \subset G_\theta$. It is easily seen that $cl U_\gamma(\xi)$ contains the set A_γ and $A_\beta - A_\gamma$ is compact. Hence, A_β is contained in the union of finitely many members of $\{cl G_\lambda \mid \lambda \in A\}$, which implies that A_β is relatively w -compact. Thus, (4) is proved.

Lemma 5.2 and properties (2) and (3) imply that X is not locally cozero-set w -compact.

REMARK. Recently, T. Ishii has obtained the following result ([8]): Let X be a space. Then $\tau(X \times Y) = \tau(X) \times \tau(Y)$ holds for any Tychonoff space Y if and only if X satisfies the following property (*):

(*) Let $\{P_\lambda | \lambda \in A\}$ be a family of τ -open sets of X such that there exists a point x_0 of X such that $\{P_\lambda | \lambda \in A\} \cup \{U\}$ has the f.i.p. for any cozero-set nbd U of x_0 . Then we have $\bigcap \{cl P_\lambda | \lambda \in A\} \neq \emptyset$.

He also showed in [8] that if a space X satisfies the property (*) and $\tau(X)$ is locally compact, then X is locally cozero-set w -compact.

From this fact and Example 5.3, it follows that there exists a locally relatively w -compact space which does not satisfy the property (*).

After submitting this paper, the authors observed the following fact: If a space X is locally relatively w -compact and the equality $\tau(X \times Y) = \tau(X) \times \tau(Y)$ holds for any Tychonoff space Y , then X is locally cozero-set w -compact. See the proof of Lemma 1.4 in: K. Morita, Čech cohomology and covering dimension for topological spaces, Fund. Math. 87 (1975), 31-52.

References

- [1] Arens, R., A topology for spaces of transformations, Ann. of Math. 47 (1946), 480-495.
- [2] Arens, R. and Dugundji, J., Topologies for function spaces, Pacific J. of Math. 1 (1951), 5-31.
- [3] Engelking, R., General topology, PWN. Warsaw, 1977.
- [4] Fox, R.H., On topologies for function spaces, Bull. Amer. Math. Soc. 51 (1945), 429-432.
- [5] Herrlich, H., Wann sind alle stetigen Abbildungen in Y konstant?, Math. Zeitschr. 90 (1965), 152-154.
- [6] Ishii, T., On the Tychonoff functor and w -compactness, Topology and Appl. 11 (1980), 173-187.
- [7] Ishii, T., Some results on w -compact spaces, Uspehi Mat. Nauk. 35 (1980), no. 3 (213), 61-66=Russian Math. Surveys 35 (1980), 71-77.
- [8] Ishii, T., On w -paracompactness and its applications, to appear.
- [9] Jackson, J.R., Spaces of mappings on topological products with applications to homotopy theory, Proc. Amer. Math. Soc. 3 (1952), 327-333.
- [10] Noble, N., A note on z -closed projections, Proc. Amer. Math. Soc. 23 (1969), 73-76.

Shizuoka Gakuen High School
226-3, Hijiriishiki, Shizuoka
422 Japan

Department of Mathematics
Faculty of Science
Shizuoka University
Ohya, Shizuoka
422 Japan