

ON UNICOHERENCE AT SUBCONTINUA*

By

Zhou YOUCHENG

Abstract. In this paper an Eilenberg-type characterization of unicoherence at subcontinua and a mapping property about this unicoherence are given.

In [5], a localization of the notion of unicoherence, i.e., unicoherence at subcontinua was introduced. Several mapping properties about unicoherence at subcontinua are studied in [1]. This property is related to other properties of unicoherence closely.

The main purpose of this paper is to establish an Eilenberg-type characterization of unicoherence at continua and to show that local homeomorphism preserves unicoherence at subcontinua for locally connected continua. The latter partially answers a question raised by J.J. Charatonik in [1].

1. Preliminary

A continuum is a compact connected metric space. A continuum is unicoherent if the intersection of every two subcontinua having union X is connected; a continuum X is hereditarily unicoherent if every subcontinuum of X is unicoherent. Let Y be a subcontinuum of X ; X is unicoherent at Y , denoted $U_n(Y)$, if for each pair of proper subcontinua A and B of X such that $X = A \cup B$ the set $A \cap B \cap Y$ is connected.

Let S^1 denote the unit circle. The mapping $f \in S^{1X}$ is said to be inessential ($f \sim 1$) if there exists a mapping $\phi \in R^X$ such that $f(x) = e^{i\phi(x)}$ for every $x \in X$.

The mapping $f \in S^{1X}$ is said to be inessential on the subspace Y of X ($f \sim 1$ on Y), if there exists a mapping $\phi \in R^Y$ such that $f(x) = e^{i\phi(x)}$ for every $x \in Y$.

S. Eilenberg introduced the property (b) for studying unicoherence. A continuum X is said to have property (b) if for each mapping $f \in S^{1X}$, there is

*The Project is supported by National Science Foundation of China
AMS (MOS) Sub. Class 54F20, 54F55.

Key words and phrases: Unicoherence at a subcontinuum, property (b), property (b) on a subcontinuum, local homeomorphism, monotone, light.

Received June 10, 1994

$f \sim 1$ ([3] p.63).

We say that a continuum X has property (b) on a subcontinuum Y of X if for each mapping $f \in S^{1^X}$, there is $f \sim 1$ on Y .

It is clear to have

LEMMA 1. *Suppose that a continuum X has property (b) on a subcontinuum Y of X and Z is a subcontinuum of Y . Then X has property (b) on Z and Y also has property (b) on Z .*

PROPOSITION 2 ([3]). *Any continuum which has property (b) is unicoherent.*

PROPOSITION 3 ([3]). *Let continuum X be locally connected. The following conditions are equivalent:*

- (1) X is unicoherent;
- (2) X has property (b).

PROPOSITION 4 ([5] corollary 1.5). *Let Y_1, Y_2, \dots, Y_n be a finite collection of subcontinua of a continuum X such that X is $U_n(Y_i)$ for every i , and suppose that for each $i > 1$*

$$Y_i \cap \bigcup \{Y_j : j < i\} \neq \emptyset.$$

Then X is $U_n(\bigcup_{i=1}^n Y_i)$.

PROPOSITION 5 ([5] Theorem 1.6). *Let Y be a subcontinuum of a continuum X . If X is $U_n(Y)$ and A and B are proper subcontinua of X such that $X = A \cup B$, then the sets $A \cap Y$ and $B \cap Y$ are connected.*

PROPOSITION 6. *If Y is a subcontinuum of a continuum X and Z is a subcontinuum of Y . Suppose that X is $U_n(Y)$ and Y is $U_n(Z)$. Then X is $U_n(Z)$.*

PROOF. Assuming that the conclusion is false, then there is a pair of proper subcontinua A and B of X such that $X = A \cup B$ and $A \cap B \cap Z$ is not connected. Suppose $A \cap B \cap Z = H \cup K$ is a separation. Because X is $U_n(Y)$, by proposition 5, $A \cap Y$ and $B \cap Y$ are all connected. One can assume that both of $A \cap Y$ and $B \cap Y$ are a nonempty proper subcontinuum of Y (Otherwise the case is simple). Then $[(A \cap Y) \cap (B \cap Y)] \cap Z = (A \cap B \cap Y) \cap Z = (A \cap B \cap Z) \cap Y = (H \cup K) \cap Y = H \cup K$. Since $H \cup K \subset Z \subset Y$, this contradicts to that Y is $U_n(Z)$.

PROPOSITION 7 (Corollary 7 of [1]). *Monotone mappings preserve uni-*

coherence at subcontinua.

2. An Eilenberg-type characterization of unicoherence at subcontinua

In this section we give a characterization of unicoherence at subcontinua which is similar with the characterization of unicoherence given by S. Eilenberg.

THEOREM 8. *Suppose that Y is a subcontinuum of a continuum X and for each pair of proper subcontinua A and B of X has property (b) on $A \cap B \cap Y$. Then X is unicoherent at Y .*

PROOF. Suppose X does not be unicoherent at Y . Then there are subcontinua A and B of X such that $X = A \cup B$ and $A \cap B \cap Y$ is not connected. Write $A \cap B \cap Y$ as a disjoint union of nonempty closed subsets C and D . One can assume that $A \cap Y \neq \emptyset \neq B \cap Y$. Otherwise $A \cap B \cap Y$ must be connected. Define a function $\phi: X \rightarrow \mathbb{R}$ by

$$\phi(x) = \pi \frac{d(x, C)}{d(x, C) + d(x, D)},$$

for each $x \in X$, and mapping $f: X \rightarrow S^1$ by

$$f(x) = \begin{cases} e^{i\phi(x)}, & \text{if } x \in A, \\ e^{-i\phi(x)}, & \text{if } x \in B. \end{cases}$$

Thus the mapping f is well defined and continuous. By hypothesis of property (b) on $A \cap B \cap Y$, one have that $f \sim 1$ on $A \cap B \cap Y$. Then there is a $\xi \in R^{A \cap B \cap Y}$ such that $f(x) = e^{i\xi(x)}$ for each $x \in A \cap B \cap Y$. According to Proposition 5 both of $A \cap Y$ and $B \cap Y$ are connected. There exist integers m and n such that

$$\phi(x) = \xi(x) + 2m\pi, \quad \text{if } x \in A \cap Y$$

and

$$-\phi(x) = \xi(x) + 2n\pi, \quad \text{if } x \in B \cap Y.$$

However, if $x \in C \subset A \cap B \cap Y$, then $\xi(x) = -2m\pi = -2n\pi$ and hence $m=n$. On the other hand, if $x \in D \subset A \cap B \cap Y$, then $\xi(x) = \pi - 2m\pi = -\pi - 2n\pi$ and get $\pi = -\pi$. This contradiction establishes the Theorem.

THEOREM 9. *Let X be a locally connected continuum and Y is its subcontinuum. The following conditions are equivalent:*

- (1) X is unicoherent at Y ;

(2) X has property (b) on Y .

PROOF. Theorem 7 has established $(2) \Rightarrow (1)$. We prove $(1) \Rightarrow (2)$. For every $f: X \rightarrow S^1$ one can let $f = f_2 \circ f_1$ by the monotone-light factorization of f , where $f_1 X \rightarrow X'$ is monotone mapping and $f_2: X' \rightarrow S^1$ is light mapping. By Proposition 7, X' is unicoherent at $Y' = f_1(Y)$. Since f_2 is light, X' must be at most 1-dimensional. Hence Y' must contain no any simple closed curve since X' is $U_n(Y)$. Y' is a locally connected continuum that contains no simple closed curve, i.e., it is a dendrite. Thus Y' is unicoherent and locally connected continuum. By Eilenberg's characterization, Y' has property (b). It is not difficult to see that X' has property (b) on Y' . This means that there is a $\psi \in S^{1Y'}$ such that $f_2(x') = e^{i\psi(x')}$, for each $x' \in Y'$. Let $\phi = \psi \circ f_1 \in S^{1Y}$. Then $f(x) = f_2 f_1(x) = e^{i\phi(x)}$, for each $x \in Y$. This is $f \sim 1$ on Y .

3. Unicoherence at continua under local homeomorphism

It is known that a surjective mapping on a continuum is a local homeomorphism if and only if it is open and n -to-1 for some fixed $n \geq 1$. It is proved that open finite-to-one mapping do not preserve unicoherence at subcontinua, even if the domain space is a linear graph in [1]. In the paper J.J. Charatonik raised a question: Do local homeomorphism perserve unicoherence at continua?

THEOREM 10. Suppose X be a locally connected continuum, Y is a subcontinuum of X and X is $U_n(Y)$, $f: X \rightarrow X'$ is a local homeomorphism. Then X' is $U_n(Y')$, here $Y' = f(Y)$.

PROOF. Whole proof consists of three steps.

CLAIM 1. Y can be covered by finite subcontinua Y_1, Y_2, \dots, Y_m such that X is $U_n(Y_i)$ and $f|_{Y_i}$ is a homeomorphis, for $i = 1, \dots, m$.

Since X is locally connected, for any $x \in Y$ there is a connected open neighborhood V_x of x in Y . Moreover one can assume that $f|_{\bar{V}_x}$ a homeomorphism because f is a local homeomorphism.

The local connectedness of X and its unicoherence at Y imply property (b) on Y by Theorem 9, i.e., for each $f: X \rightarrow S^1 f|_Y \sim 1$. Thus, by Proposition 1, $f|_{\bar{V}_x} \sim 1$ and this means that X is $U_n(\bar{V}_x)$. By compactness of Y , finite subcontinua Y_1, \dots, Y_m as required above can be found. Denote $Y'_i = f(Y_i)$.

CLAIM 2. X' is $U_n(Y'_i)$.

For any pair of proper subcontinua A' and B' of X' such that $A' \cup B' = X'$ we'll show that $A' \cap B' \cap Y'_i$ is connected. Local homeomorphism between continua is exactly a n -to-one open continuous mapping ([1]) and it is a confluent mapping ([4]). Thus one can get disjoint unions of subcontinua of X

$$f^{-1}(A') = A_1 \cup \dots \cup A_k \quad \text{and} \quad f^{-1}(B') = B_1 \cup \dots \cup B_s,$$

here $k, s \leq n$ and each of A_j and B_j is mapped onto A' and B' by f respectively. Since $f|_{Y_i}$ is a homeomorphism, it is not difficult to see that only one of A_j and B_j intersects Y_i respectively. Assume that A_1 and B_1 are they. One can consider two subcontinua

$$A_1 \cup \{B_j : B_j \cap A_1 \neq \emptyset\} \quad \text{and} \quad B_1 \cup \{A_j : A_j \cap B_1 \neq \emptyset\}.$$

Similarly, consider the rest of A_j and B_j which meet $A_1 \cup \{B_j : B_j \cap A_1 \neq \emptyset\}$ or $B_1 \cup \{A_j : A_j \cap B_1 \neq \emptyset\}$ and continue to form unions. Because numbers of A_j and B_j are finite, so finally get two continua

$$A = A_1 \cup \{B_j : j \in J' \subset \{2, \dots, s\}\} \cup \{A_j : j \in J \subset \{2, \dots, k\}\}$$

and

$$B = B_1 \cup \{A_j : j \in \{2, \dots, k\} \setminus J\} \cup \{B_j : j \in \{2, \dots, s\} \setminus J'\}.$$

Then $A \cup B = X$ and $A_1 \cap B_1 \cap Y_i = A \cap B \cap Y_i$ is connected by unicoherence at Y_i . Therefore $A' \cap B' \cap Y'_i = f(A_1 \cap B_1 \cap Y_i)$ is also connected.

CLAIM 3. It is from Claim 2 that X' is $U_n(Y')$.

$Y' = f(Y) = \bigcup_{i=1}^m Y'_i$. The finite collection $\{Y'_1, \dots, Y'_m\}$ can be selected such that for each $i > 1$, $Y'_i \cap \bigcup\{Y'_j : j < i\} \neq \emptyset$. By Corollary 1.5 of [5], the final conclusion yields.

A unicoherent continuum X is strong unicoherent if for every pair of proper subcontinua A and B such that $X = A \cup B$ both A and B are unicoherent. By Theorem 10 and Theorem 2.1 of [5] we have.

COROLLARY 11. *An image of a locally connected strongly unicoherent continuum under a local homeomorphism is locally connected strongly unicoherent.*

References

- [1] J. J. Charatonik, "Monotone mappings and unicoherence at subcontinua", *Topology and its Appl.* **33** (1989), 209–215.

- [2] S. T. Czuba, "Some property of unicoherence", *Bll. Acad Polon. Sci. Ser. Sci. Math. Astronom. Phys.* **27** (1979), 711–716.
- [3] S. Eilenberg, "Transformations continues en la topologie du plan", *Fund. Math.* **26** (1936), 61–112.
- [4] T. Mackowiak, "Continuous mappings on continua", *Dissertations Math. (Rozprawy Mat.)* **158** (1979), 1–91.
- [5] M. A. Owens, "Unicoherence at subcontinua", *Topology and its Appl.* **22** (1986), 145–155.
- [6] G. T. Whyburn, "Analytic Topology", *Amer. Math. Soc. Collq. Publ.* **28** (Amer. Math. Soc. Providence, RI, 1942).

Department of Mathematical
Zhejiang University
Hangzhou, 310027
P.R. China