



# SYNTHETIC VECTOR ANALYSIS III FROM VECTOR ANALYSIS TO DIFFERENTIAL FORMS

**HIROKAZU NISHIMURA**

Institute of Mathematics

University of Tsukuba

Tsukuba, Ibaraki, 305-8571, Japan

## Abstract

In our previous paper [Synthetic vector analysis, *Internat. J. Theor. Phys.* 41 (2002), 1165-1190], we have shown, following the tradition of synthetic differential geometry, that  $\text{div}$  and  $\text{rot}$  are uniquely determined, so long as we require that the divergence theorem and the Stokes theorem should hold on the infinitesimal level. In this paper, we simplify the discussion considerably in terms of differential forms, leading to the natural derivation of exterior differentiation in the usual form.

## 1. Introduction

Vector analysis presupposes dogmatically that every physical quantity is either a scalar or a vector, excluding the possibility of tensors as natural physical quantities. In vector analysis, the force and the flux are equally vector fields, but, to tell the truth, the former is a field of tensors of degree 1, while the latter is a field of skew-symmetric tensors of degree 2. In electromagnetism,  $\mathbf{E}$  and  $\mathbf{B}$  are fields of tensors of degree

---

2000 Mathematics Subject Classification: 51K10.

Keywords and phrases: vector analysis, differential form, synthetic differential geometry, nilpotent infinitesimal, divergence theorem, Stokes' theorem, exterior differentiation.

Received December 17, 2008

1, while  $\mathbf{D}$  and  $\mathbf{H}$  are fields of skew-symmetric tensors of degree 2. It is not desirable to apply  $\text{div}$  to  $\mathbf{E}$  or  $\mathbf{B}$ , though  $\text{curl}$  is indeed applicable to both of them. It is not desirable to apply  $\text{curl}$  to  $\mathbf{D}$  or  $\mathbf{H}$ , while  $\text{div}$  is indeed applicable to both of them. Since  $\mathbf{E}$  and  $\mathbf{D}$  as well as  $\mathbf{B}$  and  $\mathbf{H}$  are proportional in the vacuum, the confusion is apt to occur and develop! Some physicists even insist wrongly that the CGS system of units, in which  $\epsilon_0 = \mu_0 = 1$  holds, is superior to the MKSA system of units.

Nowadays the number of textbooks on elementary physics (elementary electromagnetism in particular) using differential forms in place of vector analysis is increasing, though there are still only a few. It is easy to give a dictionary of vector analysis into the framework of differential forms, so that vector analysis is really to be absorbed into the calculus of differential forms. Nevertheless vector analysis is still popular among physicists and students of physics, mainly because vector analysis is highly intuitive, while the calculus of differential forms is not. The exterior differentiation in the calculus of differential forms is usually given as a decree without taking care of its intuitive or physical foundations at all.

What is easily forgotten, such geniuses as Newton and Leibniz discussed advanced calculus in terms of nilpotent infinitesimals without using limits at all. It was in the 19th century, in the midst of the industrial revolution, that advanced calculus was reformulated in terms of limits, while nilpotent infinitesimals were intentionally neglected as anathema. Synthetic differential geometry, born in the middle of the 20th century, succeeded in reviving nilpotent infinitesimals in advanced calculus and differential geometry without hurting mathematical rigor at all. Newton and Leibniz saw nilpotent infinitesimals not in this world but in another world, and the 20th century witnessed powerful gadgets, such as seen in forcing techniques of set theory, sheaf theory and topos theory, by which many other mathematically meaningful worlds can be coherently constructed. Synthetic differential geometry regained the natural meaning of exterior differentiation in differential forms. The principal objective in this paper is to convince physicists that the exterior

differentiation is well motivated, just as div and curl in vector analysis. The paper is more expository than anything else. We have tried to help physicists understand how naturally vector analysis develops into the calculus of differential forms.

## 2. Preliminaries

We assume that the reader is familiar with Chapter 1 of Lavendhomme [5]. The set  $\mathbb{R}$  of (extended) real numbers is required to abide by the Kock-Lawvere axiom (cf. p. 2 of [5]). We denote by  $D$  the set of real numbers whose squares vanish. The Kock-Lawvere axiom implies that, given a mapping  $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $\mathbf{x}, \mathbf{a} \in \mathbb{R}^n$ , there exists a unique  $\varphi'(\mathbf{x})(\mathbf{a}) \in \mathbb{R}^n$  such that

$$\varphi(\mathbf{x} + \mathbf{a}d) - \varphi(\mathbf{x}) = \varphi'(\mathbf{x})(\mathbf{a})d$$

for any  $d \in D$ . It can be shown easily that the mapping  $\mathbf{a} \in \mathbb{R}^n \mapsto \varphi'(\mathbf{x})(\mathbf{a}) \in \mathbb{R}^n$ , which is to be regarded as the derivative of  $\varphi$  at  $\mathbf{x}$ , is linear. The mapping  $\varphi'(\mathbf{x})$  goes as follows:

$$\varphi'(\mathbf{x}) = \frac{\partial \varphi}{\partial x_1}(\mathbf{x}) \mathbf{d}x_1 + \cdots + \frac{\partial \varphi}{\partial x_n}(\mathbf{x}) \mathbf{d}x_n.$$

We denote by  $\mathbf{e}_i$

$$\begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

( $1 \leq i \leq n$ ), where 1 is positioned at the  $i$ th place. Given  $\gamma : D^{m+1} \rightarrow \mathbb{R}^n$ ,  $e \in D$  and a natural number  $i$  with  $1 \leq i \leq m+1$ , we write  $\gamma_e^i$  for the mapping  $(d_1, \dots, d_m) \in D^m \mapsto \gamma(d_1, \dots, d_{i-1}, e, d_i, \dots, d_m) \in \mathbb{R}^n$ .

Let us consider the usual three-dimensional space  $\mathbb{R}^3$ , which is the favorite space of vector analysis. Viewing the force  $\mathbf{f}(\mathbf{x})$  at  $\mathbf{x} \in \mathbb{R}^3$  as a vector in the usual way should be called an *idealistic* or *Platonic* view of force. Our *pragmatic* or *operational* view of force is to consider how to measure  $\mathbf{f}(\mathbf{x})$  experimentally. If we move from  $\mathbf{x}$  to  $\mathbf{x} + \mathbf{a}d$  infinitesimally with  $\mathbf{a} \in \mathbb{R}^3$  and  $d \in D$ , then we get the power  $\mathbf{f}(\mathbf{x}) \cdot \mathbf{a}d$ , where  $\cdot$  denotes the inner product of vectors. Our pragmatic view of force recommends that the force at  $\mathbf{x}$  should not be  $\mathbf{f}(\mathbf{x})$  but the linear mapping  $\mathbf{a} \in \mathbb{R}^3 \mapsto \mathbf{f}(\mathbf{x}) \cdot \mathbf{a} \in \mathbb{R}$ . We stress that  $\mathbf{f}(\mathbf{x})$  is recognized via the linear mapping  $\mathbf{a} \in \mathbb{R}^3 \mapsto \mathbf{f}(\mathbf{x}) \cdot \mathbf{a} \in \mathbb{R}$ . This is our view of force as a tensor of degree 1. Therefore a field of forces is no other than a differential 1-form from a mathematical viewpoint.

Let us consider a flow of air in  $\mathbb{R}^3$ , which is very often represented by a field  $\mathbf{f}$  of vectors. Our pragmatic view of flow recommends that we should measure how much air passes in a unit time through the infinitesimal parallelogram whose four vertices are  $\mathbf{x}$ ,  $\mathbf{x} + \mathbf{a}d_1$ ,  $\mathbf{x} + \mathbf{b}d_2$  and  $\mathbf{x} + \mathbf{a}d_1 + \mathbf{b}d_2$  with  $\mathbf{x}, \mathbf{a}, \mathbf{b} \in \mathbb{R}^3$  and  $d_1, d_2 \in D$ . The result is surely  $\mathbf{f}(\mathbf{x}) \cdot (\mathbf{a} \times \mathbf{b})d_1d_2$ , where  $\times$  stands for the vector product. We would like to consider pragmatically that the skew-symmetric bilinear mapping  $(\mathbf{a}, \mathbf{b}) \in \mathbb{R}^3 \times \mathbb{R}^3 \mapsto \mathbf{f}(\mathbf{x}) \cdot (\mathbf{a} \times \mathbf{b}) \in \mathbb{R}$  is no other than the mathematical representation of the flow at  $\mathbf{x}$ . In this sense, the flow is represented by a field of skew-symmetric tensors of degree 2, namely, by a differential 2-form.

We know well that every linear mapping from  $\mathbb{R}^3$  to  $\mathbb{R}$  is of the form  $\alpha_1 \mathbf{d}x + \alpha_2 \mathbf{d}y + \alpha_3 \mathbf{d}z$  with  $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ , while every skew-symmetric bilinear mapping from  $\mathbb{R}^3 \times \mathbb{R}^3$  to  $\mathbb{R}$  is of the form  $\alpha_1 \mathbf{d}y \wedge \mathbf{d}z + \alpha_2 \mathbf{d}z \wedge \mathbf{d}x + \alpha_3 \mathbf{d}x \wedge \mathbf{d}y$ . We know well that every skew-symmetric trilinear mapping from  $\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3$  to  $\mathbb{R}$  is of the form  $\alpha \mathbf{d}x \wedge \mathbf{d}y \wedge \mathbf{d}z$  with  $\alpha \in \mathbb{R}$ . More generally, every skew-symmetric  $k$ -linear mapping from

$\underbrace{\mathbb{R}^n \times \cdots \times \mathbb{R}^n}_k$  to  $\mathbb{R}$  is of the form

$$\sum_{1 \leq i_1 < \cdots < i_k \leq n} \alpha_{i_1, \dots, i_k} \mathbf{d}x_{i_1} \wedge \cdots \wedge \mathbf{d}x_{i_k}.$$

In vector analysis, the operators *div* and *rot* are determined uniquely so that the divergence theorem and the Stokes' theorem should hold on the infinitesimal level, respectively. In the same way, the exterior differentiation from a differential  $k$ -form to a differential  $(k + 1)$ -form is determined uniquely so that Stokes' theorem should hold on the infinitesimal level. The principal objective in this paper is to give a lucid explanation on these facts as elementarily as possible from the standpoint of synthetic differential geometry, while avoiding the utmost generality, which would usually be liable to defy ordinary physicists.

### 3. The Fundamental Theorem for Gradient

**Theorem 1.** *Let  $\varphi$  be a scalar field on  $\mathbb{R}^3$ . Let  $t : d \in D \mapsto \mathbf{x} + \mathbf{a}d$  be a tangent vector at  $\mathbf{x}$  on  $\mathbb{R}^3$ . Let  $e \in D$ . Then we have*

$$\int_{\partial(t; e)} \varphi = \int_{(t; e)} \mathbf{d}\varphi,$$

where

$$\partial(t; e) = (\mathbf{x} + \mathbf{a}e) - (\mathbf{x}).$$

**Proof.** This is no other than the definition of  $\mathbf{d}\varphi = \varphi'$ , namely,

$$\varphi(\mathbf{x} + \mathbf{a}e) - \varphi(\mathbf{x}) = \varphi'(\mathbf{x})(\mathbf{a})e. \quad \square$$

### 4. The Fundamental Theorem for Rotation

**Theorem 2.** *Let  $\omega = f\mathbf{d}x + g\mathbf{d}y + h\mathbf{d}z$  be a differential 1-form on  $\mathbb{R}^3$ . Let  $\gamma : (d_1, d_2) \in D^2 \mapsto \mathbf{x} + \mathbf{a}d_1 + \mathbf{b}d_2$  be an infinitesimal parallelogram at  $\mathbf{x}$  on  $\mathbb{R}^3$ . Let  $(e_1, e_2) \in D^2$ . Then we have*

$$\int_{\partial(\gamma; e_1, e_2)} \omega = \int_{(\gamma; e_1, e_2)} \mathbf{d}\omega,$$

where

$$\partial(\gamma; e_1, e_2) = (\gamma_0^2; e_1) + (\gamma_{e_1}^1; e_2) - (\gamma_{e_2}^2; e_1) - (\gamma_0^1; e_2)$$

and

$$\mathbf{d}\omega = \left( \frac{\partial h}{\partial y} - \frac{\partial g}{\partial z} \right) \mathbf{d}y \wedge \mathbf{d}z + \left( \frac{\partial f}{\partial z} - \frac{\partial h}{\partial x} \right) \mathbf{d}z \wedge \mathbf{d}x + \left( \frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \right) \mathbf{d}x \wedge \mathbf{d}y.$$

**Proof.** We have

$$\begin{aligned} & \int_{\partial(\gamma; e_1, e_2)} \omega \\ &= \int_{(\gamma_0^2; e_1)} \omega + \int_{(\gamma_{e_1}^1; e_2)} \omega - \int_{(\gamma_{e_2}^2; e_1)} \omega - \int_{(\gamma_0^1; e_2)} \omega \\ &= \{f(\mathbf{x})a_1 + g(\mathbf{x})a_2 + h(\mathbf{x})a_3\}e_1 \\ & \quad + \{f(\mathbf{x} + \mathbf{a}e_1)b_1 + g(\mathbf{x} + \mathbf{a}e_1)b_2 + h(\mathbf{x} + \mathbf{a}e_1)b_3\}e_2 \\ & \quad - \{f(\mathbf{x} + \mathbf{b}e_2)a_1 + g(\mathbf{x} + \mathbf{b}e_2)a_2 + h(\mathbf{x} + \mathbf{b}e_2)a_3\}e_1 \\ & \quad - \{f(\mathbf{x})b_1 + g(\mathbf{x})b_2 + h(\mathbf{x})b_3\}e_2 \\ &= \{f'(\mathbf{x})(\mathbf{a})b_1 + g'(\mathbf{x})(\mathbf{a})b_2 + h'(\mathbf{x})(\mathbf{a})b_3\}e_1e_2 \\ & \quad - \{f'(\mathbf{x})(\mathbf{b})a_1 + g'(\mathbf{x})(\mathbf{b})a_2 + h'(\mathbf{x})(\mathbf{b})a_3\}e_1e_2 \end{aligned}$$

The first term delineated by  $\{ \}$  and followed by  $e_1e_2$  is obtained by combining the second term and the fourth of the preceding formula, while the second term delineated by  $\{ \}$  and followed by  $e_1e_2$  is obtained by combining the first term and the third of the preceding formula.

$$= \left\{ \begin{array}{l} f'(\mathbf{x})(\mathbf{a}) \\ g'(\mathbf{x})(\mathbf{a}) \cdot \mathbf{b} - g'(\mathbf{x})(\mathbf{b}) \cdot \mathbf{a} \\ h'(\mathbf{x})(\mathbf{a}) \end{array} \right\} e_1e_2.$$

Let  $\varphi : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}$  be the mapping

$$\varphi(\mathbf{a}, \mathbf{b}) = \begin{pmatrix} f'(\mathbf{x})(\mathbf{a}) \\ g'(\mathbf{x})(\mathbf{a}) \cdot \mathbf{b} - g'(\mathbf{x})(\mathbf{b}) \cdot \mathbf{a} \\ h'(\mathbf{x})(\mathbf{a}) \end{pmatrix}$$

for any  $(\mathbf{a}, \mathbf{b}) \in \mathbb{R}^3 \times \mathbb{R}^3$ , so that

$$\int_{\partial(\gamma; e_1, e_2)} \omega = \varphi(\mathbf{a}, \mathbf{b})e_1e_2.$$

Then it is easy to see that  $\varphi$  is a skew-symmetric bilinear mapping, so that  $\varphi$  is of the form

$$\varphi = \alpha_1 \mathbf{d}y \wedge \mathbf{d}z + \alpha_2 \mathbf{d}z \wedge \mathbf{d}x + \alpha_3 \mathbf{d}x \wedge \mathbf{d}y$$

with  $\alpha_i \in \mathbb{R}$  ( $i = 1, 2, 3$ ). By taking

1.  $\mathbf{a} = \mathbf{e}_2$  and  $\mathbf{b} = \mathbf{e}_3$ ,
2.  $\mathbf{a} = \mathbf{e}_3$  and  $\mathbf{b} = \mathbf{e}_1$ , or
3.  $\mathbf{a} = \mathbf{e}_1$  and  $\mathbf{b} = \mathbf{e}_2$ ,

we get

$$\alpha_1 = \frac{\partial h}{\partial y}(\mathbf{x}) - \frac{\partial g}{\partial z}(\mathbf{x}),$$

$$\alpha_2 = \frac{\partial f}{\partial z}(\mathbf{x}) - \frac{\partial h}{\partial x}(\mathbf{x}),$$

$$\alpha_3 = \frac{\partial g}{\partial x}(\mathbf{x}) - \frac{\partial f}{\partial y}(\mathbf{x})$$

easily. This completes the proof.  $\square$

### 5. The Fundamental Theorem for Divergence

**Theorem 3.** Let  $\omega = f\mathbf{d}y \wedge \mathbf{d}z + g\mathbf{d}z \wedge \mathbf{d}x + h\mathbf{d}x \wedge \mathbf{d}y$  be a differential 2-form on  $\mathbb{R}^3$ . Let  $\gamma : (d_1, d_2, d_3) \in D^3 \mapsto \mathbf{x} + \mathbf{a}d_1 + \mathbf{b}d_2 + \mathbf{c}d_3$  be an infinitesimal parallelepiped at  $\mathbf{x}$  on  $\mathbb{R}^3$ . Let  $(e_1, e_2, e_3) \in D^3$ . Then we have

$$\int_{\partial(\gamma; e_1, e_2, e_3)} \omega = \int_{(\gamma; e_1, e_2, e_3)} \mathbf{d}\omega,$$

where

$$\begin{aligned} \partial(\gamma; e_1, e_2, e_3) = & -(\gamma_0^1; e_2, e_3) + (\gamma_{e_1}^1; e_2, e_3) + (\gamma_0^2; e_1, e_3) - (\gamma_{e_2}^2; e_1, e_3) \\ & - (\gamma_0^3; e_1, e_2) + (\gamma_{e_3}^3; e_1, e_2) \end{aligned}$$

and

$$\mathbf{d}\omega = \left( \frac{\partial f}{\partial x} - \frac{\partial g}{\partial y} + \frac{\partial h}{\partial z} \right) \mathbf{d}x \wedge \mathbf{d}y \wedge \mathbf{d}z.$$

**Proof.** We have

$$\begin{aligned} & \int_{\partial(\gamma; e_1, e_2, e_3)} \omega \\ &= - \int_{(\gamma_0^1; e_2, e_3)} \omega + \int_{(\gamma_{e_1}^1; e_2, e_3)} \omega + \int_{(\gamma_0^2; e_1, e_3)} \omega \\ & \quad - \int_{(\gamma_{e_2}^2; e_1, e_3)} \omega - \int_{(\gamma_0^3; e_1, e_2)} \omega + \int_{(\gamma_{e_3}^3; e_1, e_2)} \omega \\ &= - \left\{ f(\mathbf{x}) \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix} + g(\mathbf{x}) \begin{vmatrix} b_3 & c_3 \\ b_1 & c_1 \end{vmatrix} + h(\mathbf{x}) \begin{vmatrix} b_1 & c_1 \\ b_2 & c_2 \end{vmatrix} \right\} e_2 e_3 \\ & \quad + \left\{ f(\mathbf{x} + \mathbf{a}e_1) \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix} + g(\mathbf{x} + \mathbf{a}e_1) \begin{vmatrix} b_3 & c_3 \\ b_1 & c_1 \end{vmatrix} + h(\mathbf{x} + \mathbf{a}e_1) \begin{vmatrix} b_1 & c_1 \\ b_2 & c_2 \end{vmatrix} \right\} e_2 e_3 \\ & \quad + \left\{ f(\mathbf{x}) \begin{vmatrix} a_2 & c_2 \\ a_3 & c_3 \end{vmatrix} + g(\mathbf{x}) \begin{vmatrix} a_3 & c_3 \\ a_1 & c_1 \end{vmatrix} + h(\mathbf{x}) \begin{vmatrix} a_1 & c_1 \\ a_2 & c_2 \end{vmatrix} \right\} e_1 e_3 \\ & \quad - \left\{ f(\mathbf{x} + \mathbf{b}e_1) \begin{vmatrix} a_2 & c_2 \\ a_3 & c_3 \end{vmatrix} + g(\mathbf{x} + \mathbf{b}e_1) \begin{vmatrix} a_3 & c_3 \\ a_1 & c_1 \end{vmatrix} + h(\mathbf{x} + \mathbf{b}e_1) \begin{vmatrix} a_1 & c_1 \\ a_2 & c_2 \end{vmatrix} \right\} e_1 e_3 \\ & \quad - \left\{ f(\mathbf{x}) \begin{vmatrix} a_2 & b_2 \\ a_3 & b_3 \end{vmatrix} + g(\mathbf{x}) \begin{vmatrix} a_3 & b_3 \\ a_1 & b_1 \end{vmatrix} + h(\mathbf{x}) \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} \right\} e_1 e_2 \\ & \quad + \left\{ f(\mathbf{x} + \mathbf{c}e_3) \begin{vmatrix} a_2 & b_2 \\ a_3 & b_3 \end{vmatrix} + g(\mathbf{x} + \mathbf{c}e_3) \begin{vmatrix} a_3 & b_3 \\ a_1 & b_1 \end{vmatrix} + h(\mathbf{x} + \mathbf{c}e_3) \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} \right\} e_1 e_2 \\ &= \left\{ f'(\mathbf{x})(\mathbf{a}) \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix} + g'(\mathbf{x})(\mathbf{a}) \begin{vmatrix} b_3 & c_3 \\ b_1 & c_1 \end{vmatrix} + h'(\mathbf{x})(\mathbf{a}) \begin{vmatrix} b_1 & c_1 \\ b_2 & c_2 \end{vmatrix} \right\} e_1 e_2 e_3 \\ & \quad - \left\{ f'(\mathbf{x})(\mathbf{b}) \begin{vmatrix} a_2 & c_2 \\ a_3 & c_3 \end{vmatrix} + g'(\mathbf{x})(\mathbf{b}) \begin{vmatrix} a_3 & c_3 \\ a_1 & c_1 \end{vmatrix} + h'(\mathbf{x})(\mathbf{b}) \begin{vmatrix} a_1 & c_1 \\ a_2 & c_2 \end{vmatrix} \right\} e_1 e_2 e_3 \\ & \quad + \left\{ f'(\mathbf{x})(\mathbf{c}) \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix} + g'(\mathbf{x})(\mathbf{c}) \begin{vmatrix} b_3 & c_3 \\ b_1 & c_1 \end{vmatrix} + h'(\mathbf{x})(\mathbf{c}) \begin{vmatrix} b_1 & c_1 \\ b_2 & c_2 \end{vmatrix} \right\} e_1 e_2 e_3 \end{aligned}$$



$$= \left\{ \begin{array}{c} \left| \begin{array}{ccc} f'(\mathbf{x})(\mathbf{a}) & & \\ g'(\mathbf{x})(\mathbf{a}) & \mathbf{b} & \mathbf{c} \\ h'(\mathbf{x})(\mathbf{a}) & & \end{array} \right| + \left| \begin{array}{ccc} f'(\mathbf{x})(\mathbf{b}) & & \\ g'(\mathbf{x})(\mathbf{b}) & \mathbf{c} & \\ h'(\mathbf{x})(\mathbf{b}) & & \end{array} \right| + \left| \begin{array}{ccc} f'(\mathbf{x})(\mathbf{c}) & & \\ g'(\mathbf{x})(\mathbf{c}) & \mathbf{a} & \mathbf{b} \\ h'(\mathbf{x})(\mathbf{c}) & & \end{array} \right| \end{array} \right\} e_1 e_2 e_3.$$

Let  $\varphi : \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}$  be the mapping

$$\varphi(\mathbf{a}, \mathbf{b}, \mathbf{c}) = \left| \begin{array}{ccc} f'(\mathbf{x})(\mathbf{a}) & & \\ g'(\mathbf{x})(\mathbf{a}) & \mathbf{b} & \mathbf{c} \\ h'(\mathbf{x})(\mathbf{a}) & & \end{array} \right| + \left| \begin{array}{ccc} f'(\mathbf{x})(\mathbf{b}) & & \\ g'(\mathbf{x})(\mathbf{b}) & \mathbf{c} & \\ h'(\mathbf{x})(\mathbf{b}) & & \end{array} \right| + \left| \begin{array}{ccc} f'(\mathbf{x})(\mathbf{c}) & & \\ g'(\mathbf{x})(\mathbf{c}) & \mathbf{a} & \mathbf{b} \\ h'(\mathbf{x})(\mathbf{c}) & & \end{array} \right|$$

for any  $(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3$ , so that

$$\int_{\partial(\gamma; e_1, e_2, e_3)} \omega = \varphi(\mathbf{a}, \mathbf{b}, \mathbf{c}) e_1 e_2 e_3.$$

Then it is easy to see that  $\varphi$  is a skew-symmetric trilinear mapping, so that  $\varphi$  is of the form

$$\varphi = \alpha \mathbf{d}\mathbf{x} \wedge \mathbf{d}\mathbf{y} \wedge \mathbf{d}\mathbf{z}$$

with  $\alpha \in \mathbb{R}$ . By taking  $\mathbf{a} = \mathbf{e}_1$ ,  $\mathbf{b} = \mathbf{e}_2$  and  $\mathbf{c} = \mathbf{e}_3$ , we get

$$\alpha = \frac{\partial f}{\partial x}(\mathbf{x}) + \frac{\partial g}{\partial y}(\mathbf{x}) + \frac{\partial h}{\partial z}(\mathbf{x})$$

easily. This completes the proof.  $\square$

## 6. The Fundamental Theorem for Exterior Differentiation

**Theorem 4.** Let  $\omega = \sum_{1 \leq i_1 < \dots < i_k \leq n} f_{i_1, \dots, i_k} \mathbf{d}x_{i_1} \wedge \dots \wedge \mathbf{d}x_{i_k}$  be a differential  $k$ -form on  $\mathbb{R}^n$ . Let  $\gamma : (d_1, \dots, d_{k+1}) \in D^{k+1} \mapsto \mathbf{a}^1 d_1 + \dots + \mathbf{a}^{k+1} d_{k+1}$  be an infinitesimal  $(k+1)$ -parallelepiped at  $\mathbf{x}$  on  $\mathbb{R}^n$ . Let  $(e_1, \dots, e_{k+1}) \in D^{k+1}$ . Then we have

$$\int_{\partial(\gamma; e_1, \dots, e_{k+1})} \omega = \int_{(\gamma; e_1, \dots, e_{k+1})} \mathbf{d}\omega,$$

where

$$\begin{aligned} & \partial(\gamma; e_1, \dots, e_{k+1}) \\ &= \sum_{i=1}^{k+1} (-1)^i \{(\gamma_0^i; e_1, \dots, \widehat{e}_i, \dots, e_{k+1}) - (\gamma_{e_i}^i; e_1, \dots, \widehat{e}_i, \dots, e_{k+1})\} \end{aligned}$$

and

$$\mathbf{d}\omega = \sum_{1 \leq i_1 < \dots < i_{k+1} \leq n} \left( \sum_{j=1}^{k+1} (-1)^{j+1} \frac{\partial f_{i_1, \dots, i_j, \dots, i_{k+1}}}{\partial x_{i_j}} \right) \mathbf{d}x_{i_1} \wedge \dots \wedge \mathbf{d}x_{i_{k+1}}.$$

**Proof.** We have

$$\begin{aligned} & \int_{\partial(\gamma; e_1, \dots, e_{k+1})} \omega \\ &= \sum_{i=1}^{k+1} (-1)^i \left\{ \int_{(\gamma_0^i; e_1, \dots, \widehat{e}_i, \dots, e_{k+1})} \omega - \int_{(\gamma_{e_i}^i; e_1, \dots, \widehat{e}_i, \dots, e_{k+1})} \omega \right\} \\ &= \sum_{1 \leq i_1 < \dots < i_k \leq n} \sum_{i=1}^{k+1} (-1)^i \left\{ \begin{array}{l} f_{i_1, \dots, i_k}(\mathbf{x}) \begin{vmatrix} a_{i_1}^1 & \dots & a_{i_1}^{i-1} & a_{i_1}^{i+1} & \dots & a_{i_1}^{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{i_k}^1 & \dots & a_{i_k}^{i-1} & a_{i_k}^{i+1} & \dots & a_{i_k}^{k+1} \end{vmatrix} e_1 \cdots \widehat{e}_i \cdots e_{k+1} - \\ f_{i_1, \dots, i_k}(\mathbf{x} + \mathbf{a}^i e_i) \begin{vmatrix} a_{i_1}^1 & \dots & a_{i_1}^{i-1} & a_{i_1}^{i+1} & \dots & a_{i_1}^{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{i_k}^1 & \dots & a_{i_k}^{i-1} & a_{i_k}^{i+1} & \dots & a_{i_k}^{k+1} \end{vmatrix} e_1 \cdots \widehat{e}_i \cdots e_{k+1} \end{array} \right\} \\ &= \sum_{1 \leq i_1 < \dots < i_k \leq n} \sum_{i=1}^{k+1} (-1)^{i+1} f'_{i_1, \dots, i_k}(\mathbf{x})(\mathbf{a}^i) \begin{vmatrix} a_{i_1}^1 & \dots & a_{i_1}^{i-1} & a_{i_1}^{i+1} & \dots & a_{i_1}^{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{i_k}^1 & \dots & a_{i_k}^{i-1} & a_{i_k}^{i+1} & \dots & a_{i_k}^{k+1} \end{vmatrix} e_1 \cdots e_{k+1} \end{aligned}$$

$$= \sum_{1 \leq i_1 < \dots < i_k \leq n} \begin{vmatrix} f'_{i_1, \dots, i_k}(\mathbf{x})(\mathbf{a}^1) & \dots & f'_{i_1, \dots, i_k}(\mathbf{x})(\mathbf{a}^{k+1}) \\ a_{i_1}^1 & \dots & a_{i_1}^{k+1} \\ \vdots & \dots & \vdots \\ a_{i_k}^1 & \dots & a_{i_k}^{k+1} \end{vmatrix} e_1 \cdots e_{k+1}.$$

Let  $\varphi : \underbrace{\mathbb{R}^n \times \dots \times \mathbb{R}^n}_{k+1} \rightarrow \mathbb{R}$  be the mapping

$$\varphi(\mathbf{a}^1, \dots, \mathbf{a}^{k+1}) = \sum_{1 \leq i_1 < \dots < i_k \leq n} \begin{vmatrix} f'_{i_1, \dots, i_k}(\mathbf{x})(\mathbf{a}^1) & \dots & f'_{i_1, \dots, i_k}(\mathbf{x})(\mathbf{a}^{k+1}) \\ a_{i_1}^1 & \dots & a_{i_1}^{k+1} \\ \vdots & \dots & \vdots \\ a_{i_k}^1 & \dots & a_{i_k}^{k+1} \end{vmatrix}$$

for any  $(\mathbf{a}^1, \dots, \mathbf{a}^{k+1}) \in \underbrace{\mathbb{R}^n \times \dots \times \mathbb{R}^n}_{k+1}$ , so that

$$\int_{\partial(\gamma; e_1, \dots, e_{k+1})} \omega = \varphi(\mathbf{a}^1, \dots, \mathbf{a}^{k+1}) e_1 \cdots e_{k+1}.$$

Then it is easy to see that  $\varphi$  is a skew-symmetric  $(k+1)$ -linear mapping, so that  $\varphi$  is of the form

$$\varphi = \sum_{1 \leq i_1 < \dots < i_{k+1} \leq n} \alpha_{i_1, \dots, i_{k+1}} \mathbf{d}x_{i_1} \wedge \dots \wedge \mathbf{d}x_{i_{k+1}}$$

with

$$\alpha_{i_1, \dots, i_{k+1}} \in \mathbb{R} \quad (1 \leq i_1 < \dots < i_{k+1} \leq n).$$

By taking  $\mathbf{a}^1 = \mathbf{e}_{i_1}$ ,  $\mathbf{a}^2 = \mathbf{e}_{i_2}$ , ...,  $\mathbf{a}^{k+1} = \mathbf{e}_{i_{k+1}}$ , we get

$$\alpha_{i_1, \dots, i_{k+1}} = \sum_{j=1}^{k+1} (-1)^{j+1} \frac{\partial f_{i_1, \dots, \hat{i}_j, \dots, i_{k+1}}}{\partial x_{i_j}}$$

easily. This completes the proof.  $\square$

**References**

- [1] John L. Bell, *A Primer of Infinitesimal Analysis*, Cambridge University Press, Cambridge, 1998.
- [2] Frierich W. Hehl and Yuri N. Obukhov, *Foundations of Classical Electrodynamics*, Birkhäuser, Boston, 2003.
- [3] Darryl D. Holm, *Geometric Mechanics, 2 Volumes*, Imperial College Press, London, 2008.
- [4] M. Kitano, *Maxwell Equations*, Science, Tokyo, 2005 (in Japanese).
- [5] R. Lavendhomme, *Basic Concepts of Synthetic Differential Geometry*, Kluwer, Dordrecht, 1996.
- [6] H. Nishimura, Synthetic vector analysis, *Internat. J. Theor. Phys.* 41 (2002), 1165-1190.
- [7] H. Nishimura, Synthetic vector analysis II, *Internat. J. Theor. Phys.* 43 (2004), 505-517.
- [8] Waldyr Muniz Oliva, *Geometric Mechanics*, Springer, Berlin, Heidelberg, 2002.