

ON THE GROUPS WITH HOMOGENEOUS THEORY

By

Kikyo HIROTAKA

1. Introduction

D. MacPherson [5] proved that no infinite groups are interpretable in any finitely homogeneous structure. A countable structure M is called *finitely homogeneous* if its language is finite, its domain is countable, and every isomorphism between finite tuples in M extends to an automorphism of M .

We shall consider a similar condition which applies to general structures.

DEFINITION 1.1. Let $2 \leq m < n$. We say that a structure M is (m, n) -homogeneous if for any two n -tuples \bar{a}, \bar{b} from M , $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$ if and only if corresponding m -tuples from \bar{a} and \bar{b} have the same type. A complete theory T is (m, n) -homogeneous if every model of T is (m, n) -homogeneous.

Note that the additive group of integers $(\mathbb{Z}, +)$ is $(2, n)$ -homogeneous for any $n > 2$. But it turns out that its theory is not (n, n) -homogeneous for any m, n by the stability and Theorem 2.3 below.

In this paper, we treat the following conjecture:

CONJECTURE 1.2. *If (M, \cdot) is a group (it may have other structures) then the theory of (M, \cdot) is not (m, n) -homogeneous for any m, n such that $2 \leq m < n$.*

We call a theory (m, ∞) -homogeneous if it is (m, n) -homogeneous for any $n > m$. Handa [2] studied (m, ∞) -homogeneous theories and proved that no infinite Abelian p -groups are interpretable in a model of such a theory, and if the theory is ω -stable in addition then no infinite groups are interpretable.

If the above conjecture is true then no groups are interpretable in a model of (m, ∞) -homogeneous theories. However, we cannot claim that no groups are interpretable in a model of an (m, n) -homogeneous theory. The following

example suggested by Ehud Hrushovski is ω -stable, $(2, 3)$ -homogeneous, not $(3, 4)$ -homogeneous, and interprets an infinite group.

EXAMPLE 1.3. Consider the projective line \mathbf{P}^1 over an algebraically closed field K and the action of $\mathrm{PGL}(2, K)$ on it. This group acts sharply 3-transitively on \mathbf{P}^1 . Define a relation $R(z_1, z_2, z_3, z_4, w_1, w_2, w_3, w_4)$ on \mathbf{P}^1 as follows: There is a regular linear map A in $\mathrm{PGL}(2, K)$ such that $Az_i = w_i$ for each $i = 1, 2, 3$ and 4.

R is invariant under the action of $\mathrm{PGL}(2, K)$. Since this group acts sharply 3-transitively on \mathbf{P}^1 , given two sets of three points $\{p, q, r\}$ and $\{p', q', r'\}$ for \mathbf{P}^1 , the relation $R(z, p, q, r, w, p', q', r')$ between z and w represents an automorphism of (\mathbf{P}^1, R) which belongs to $\mathrm{PGL}(2, K)$.

Now we can easily see that $\mathrm{Th}(\mathbf{P}^1, R)$ is $(2, 3)$ -homogeneous but (\mathbf{P}^1, R) interprets the infinite group $\mathrm{PGL}(2, K)$. As we can interpret (\mathbf{P}^1, R) in the field K , $\mathrm{Th}(\mathbf{P}^1, R)$ is ω -stable.

Moreover, the theory is not $(3, 4)$ -homogeneous. Choose three distinct points, a, b, c from \mathbf{P}^1 and a linear map A from $\mathrm{PGL}(2, K)$ sending a, b, c to b, c, a respectively. Since K is algebraically closed, A has a fixed point d in \mathbf{P}^1 . Note that d is different from a, b and c . Choose a new point d' from \mathbf{P}^1 that is not fixed by A . Then $R(d, a, b, c, d, b, c, a)$ holds but $R(d', a, b, c, d', b, c, a)$ does not hold. Since there is only one 3-type realized by three distinct points, this shows that the theory is not $(3, 4)$ -homogeneous.

Also, we cannot claim that no groups are definable in a model of an (m, n) -homogeneous theory. The following example is due to Akito Tsuboi. This example is ω -categorical, ω -stable, $(2, 3)$ -homogeneous, not $(2, 4)$ -homogeneous, but some infinite groups are definable with three parameters.

EXAMPLE 1.4. Let V_1, V_2, V_3, V_4 be four copies of $\mathbf{Z}_2^{(\omega)}$ where \mathbf{Z}_2 is the Abelian group of order 2. Let M be the disjoint union of these four sets, and define the relation $R(x_1, x_2, x_3, x_4)$ by $x_i \in V_i$ and $x_1 + x_2 + x_3 + x_4 = 0$. Then $\mathrm{Th}(M, R)$ is $(2, 3)$ -homogeneous but $\mathbf{Z}_2^{(\omega)}$ is definable in it.

First, we can recover a group structure on each V_i . Fix three elements a, b, c one from each V_2, V_3 and V_4 . The formula

$$\exists x_2, x_3 [R(u_2, x_2, b, c) \wedge R(u_3, a, x_3, c) \wedge R(u_1, x_2, x_3, c)]$$

is equivalent to $u_1 + u_2 + u_3 + a + b + c = 0$ for u_1, u_2, u_3 in V_1 which gives a group structure on V_1 . The same argument works for each V_i .

To show that there is only one 3-type realized by three distinct elements from V_1 is the most essential in the proof of (2, 3)-homogeneity of the theory. Consider each V_i as a vector space over the prime field of characteristic 2. Let $\{a_1, a_2, a_3\}$ and $\{b_1, b_2, b_3\}$ be two sets of three distinct elements from V_1 . Whether each set is dependent or not, we can choose c from V_1 so that $\{a_1 - c, a_2 - c, a_3 - c\}$ and $\{b_1 - c, b_2 - c, b_3 - c\}$ are both linearly independent sets. Let s be a linear automorphism on V_1 sending each $a_i - c$ to $b_i - c$. Then $\sigma(x) = s(x - c) + c$ is an automorphism of V_1 which sends a_i to b_i for $i = 1, 2, 3$. Extend σ to V_2, V_3 and V_4 by $\sigma(x) = s(x + c) - c$ on V_2 , and $\sigma(x) = s(x)$ on V_3 and V_4 . Then σ is an automorphism of (M, R) .

We prove Conjecture 1.2 with various additional conditions such as ω -categoricity, o-minimality, stability and simplicity (in Shelah's sense), but it seems very hard to prove it in general. In the simple case, we only prove that the theory is not (2, 3)-homogeneous. Also, we have not found a pure group with the (m, n) -homogeneous theory for some m and n .

In this paper, the language is countable and the notation follows Pillay's book [7].

2. (m, n) -Homogeneous Theory

In this section, we prove that Conjecture 1.2 holds if $Th(M)$ is ω -categorical, stable, or o-minimal.

THEOREM 2.1 *If (M, \cdot) has an infinite Abelian p -subgroup then $Th(M, \cdot)$ is not (m, n) -homogeneous for any m and n .*

PROOF. The proof is much the same as Handa's proof in [2] which is a modification of Macpherson's argument [5]. We give the proof for reader's convenience.

We work in an Abelian subgroup and write the group operation additively.

It is enough to show that the theory is not $(m, m + 1)$ -homogeneous for any m . We find elements a_1, \dots, a_{m+1} that are linearly independent over a finite prime field F_p with the p elements, and the corresponding m -tuples from $(a_1, \dots, a_m, a_1 + \dots + a_m + a_{m+1})$ and $(a_1, \dots, a_m, a_1 + \dots + a_m)$ have the same type. Note that we can describe this condition by a set of elementary formulas. We show that for given finite set Δ of m -formulas, we can find elements a_1, \dots, a_{m+1} satisfying the above condition except that the phrase "have the same type"

changed to “have the same Δ -type”. Then by compactness, we get the desired tuple.

Let V be an infinite Abelian p -subgroup of (M, \cdot) . Consider V as a vector space over F_p . We can assume that V has the countable dimension over F_p . Choose a basis $(v_i : i < \omega)$ of V . Now we give a rule for coloring the m -dimensional subspaces of V .

First, we give a rule for ordering the elements of a such subspace. If U is an m -dimensional subspace of V , then the cardinality of U is p^m . Since U is a finite dimensional subspace of V , U is covered by the F_p -span of $(v_i : i < n)$ for some natural number n . Every element of U can be written as a linear combination of $(v_i : i < n)$ over F_p . If we list all of them, we naturally get a $|U| \times n$ matrix with entries in F_p . Then we can find a unique row reduced echelon form of the matrix. It has $m (= \dim U)$ nonzero rows, and the tuple of elements of U represented by those rows is an ordered basis of U . We call it the *canonical basis* of U . Order the elements of U lexicographically according to their coordinates with respect to the canonical basis.

Now, if U and U' are m -dimensional subspaces of V , we say that U and U' have the same color if every corresponding m -tuples with respect to the above ordering have the same Δ -type. Note that the number of the colors is finite.

By the affine version of Ramsey’s theorem [1], V has an $(m + 1)$ -dimensional subspace W all of whose m -dimensional subspaces have the same color. Let $(a_1, \dots, a_m, a_{m+1})$ be the canonical basis of W .

All we have to show is that the corresponding m -tuples from $(a_1, \dots, a_m, a_1 + \dots + a_m + a_{m+1})$ and $(a_1, \dots, a_m, a_1 + \dots + a_m)$ have the same Δ -type. Let U_1 be the F_p -span of $\{a_1, \dots, a_m, a_1 + \dots + a_m + a_{m+1}\} \setminus \{a_i\}$ and U_2 the F_p -span of $\{a_1, \dots, a_m, a_1 + \dots + a_m\} \setminus \{a_i\}$. Then their dimensions are both m . Since U_1 has the canonical basis $(a_1, \dots, a_i + a_{m+1}, \dots, a_m)$ and U_2 has the canonical basis $(a_1, \dots, a_i, \dots, a_m)$, $a_1 + \dots + a_m + a_{m+1}$ in U_1 and $a_1 + \dots + a_m$ in U_2 have the same coordinate $(1, \dots, 1)$. Thus, we get the desired result. ■

As there exists an infinite Abelian p -subgroup in a ω -categorical group (see [5]), we have the following.

COROLLARY 2.2. *If $Th(M, \cdot)$ is countably categorical then it is not (m, n) -homogeneous for any m and n .*

We now turn to the stable case. In this case, Conjecture 1.2 holds by the existence of stationary generic types.

THEOREM 2.3. *If $Th(M, \cdot)$ is stable then it is not (m, n) -homogeneous for any m and n .*

PROOF. It is enough to show that the theory is not $(m, m + 1)$ -homogeneous for any m . Let p be a stationary generic type over a model N , and a_1, \dots, a_m independent (over N) realizations of p . Let $b = a_1 \cdots a_m$. Since p is generic, $tp(b/N)$ is also a stationary generic type, and any m elements from a_1, \dots, a_m, b are independent over N .

Now choose c such that $tp(c/a_1 \dots a_m N)$ is a nonforking extension of $tp(b/N)$ and consider the two $(m + 1)$ -tuples (a_1, \dots, a_m, b) and (a_1, \dots, a_m, c) . They do not have the same type since b is algebraic (definable) over $\{a_1, \dots, a_m\}$ and c is independent of $\{a_1, \dots, a_m\}$. But the corresponding m -tuples from both tuples have the same type by the stationarity of types over a model. This shows that the theory is not $(m, m + 1)$ -homogeneous. ■

To finish this section, we consider the o-minimal case.

THEOREM 2.4. *If $Th(M, \cdot, <)$ is o-minimal then it is not (m, n) -homogeneous for any m and n .*

PROOF. Choose algebraically independent elements a_1, \dots, a_m (in the big model). If we cannot choose such elements, then by compactness, there are formulas $\psi_i(x; y_1, \dots, y_{m-1})$ ($i = 1, \dots, m$) such that any m -tuple satisfies one of ψ_i 's (by permuting if necessary) and if x, y_1, \dots, y_{m-1} satisfies ψ_i then x is algebraic over y_1, \dots, y_{m-1} . But if we choose an infinite indiscernible sequence $\langle a_i \mid i < \omega \rangle$, we get a contradiction by considering $a_k, a_{2k}, \dots, a_{mk}$ for sufficiently large k .

Let $b = a_1 \cdots a_m$ and consider the types

$$tp(b/A_i) \text{ where } A_i = \{a_1, \dots, a_m\} \setminus \{a_i\}.$$

Note that they are non-algebraic types. If a formula $\varphi_i(x)$ belongs to $tp(b/A_i)$ then it is a finite union of intervals by o-minimality. Without loss of generality, we can assume that $\varphi_i(x)$ represents a single interval $[c_i, d_i]$ where c_i and d_i are definable elements over A_i (this may not be a closed interval, but the argument will be the same in any case). Since b is not algebraic over A_i , b belongs to the open interval (c_i, d_i) . As this is true for each $i = 1, \dots, m$, the type

$$tp(b/A_1) \cup \dots \cup tp(b/A_m)$$

is non-algebraic by compactness. Choose $b' \neq b$ satisfying this type. Considering the tuples (a_1, \dots, a_m, b) and (a_1, \dots, a_m, b') , we see that the theory is not $(m, m+1)$ -homogeneous. ■

3. (2,3)-Homogeneous Theory

If the theory is simple then we can still find a generic type, but it is not necessarily stationary. Instead, we can use the Independence Theorem due to B. Kim and A. Pillay to prove the conjecture in a special form. But we could not prove the conjecture in the general form.

We use the following definition and facts from [4] and [6].

DEFINITION 3.1. A 1-type $p(x)$ over A is called *generic* if for any a realizing p and b such that a is independent from b over A , $a \cdot b$ is independent from Ab over \emptyset and so is $b \cdot a$.

FACT 3.2. *If $\text{Th}(M, \cdot)$ is simple then there is a generic type.*

FACT 3.3 (Independence Theorem). *Suppose the theory is simple. If A and B are independent over a model M and a type p_1 over A and a type q_2 over B are both nonforking extensions of a type p over M , then there is a type q over $A \cup B$ such that q extends both p_1 and p_2 , and q does not fork over M .*

THEOREM 3.4. *If $\text{Th}(M, \cdot)$ is simple then it is not (2,3)-homogeneous.*

PROOF. Let p be a generic type over some model N , and a_1, a_2 independent realizations of p . Let $b = a_1 \cdot a_2$. Then both $\text{tp}(b/a_1N)$ and $\text{tp}(b/a_2N)$ do not fork over N . By the Independence Theorem, we can choose c such that $\text{tp}(c/a_1a_2N)$ does not fork over N and $\text{tp}(c/a_1a_2N)$ extends both $\text{tp}(b/a_1N)$ and $\text{tp}(b/a_2N)$. This implies that corresponding pairs from (a_1, a_2, b) and (a_1, a_2, c) have the same type. On the other hand, (a_1, a_2, b) and (a_1, a_2, c) have different types over \emptyset since $b = a_1 \cdot a_2$ but c is non-algebraic over $\{a_1, a_2\} \cup N$. ■

References

- [1] R. L. Graham, K. Leeb, and B. L. Rothschild, Ramsey's theorem for a class of categories, *Advances in Math.* **8** (1972), 417–433.
- [2] M. Handa, Master's Thesis (in Japanese), University of Tsukuba, 1994.
- [3] B. Kim, Forking in simple unstable theories, *J. London Math. Soc.*, to appear.
- [4] B. Kim and A. Pillay, Simple theories, preprint.

- [5] D. MacPherson, Interpreting groups in ω -categorical structures, *J. Symb. Logic* **56** (1991), 1317–1324.
- [6] A. Pillay, Definability and definable groups in simple theories, preprint, July, 1996.
- [7] A. Pillay, *An Introduction to Stability Theory*, Clarendon Press, Oxford, 1983.
- [8] B. Poizat, Sous-groupes définissables d'un groupe stable, *J. Symb. Logic* **46** (1981), 137–146.
- [9] S. Shelah, *Classification Theory (revised)*, North-Holland, Amsterdam, 1990.

Department of Mathematical Sciences
Tokai University
Hiratsuka, Kanagawa 259-1292
Japan