

GENERALIZED HELICAL IMMERSIONS OF A
RIEMANNIAN MANIFOLD ALL OF WHOSE
GEODESICS ARE CLOSED INTO
A EUCLIDEAN SPACE

By

Naoyuki KOIKE

Abstract. In this paper, we investigate an isometric immersion of a compact connected Riemannian manifold M into a Euclidean space and a sphere such that every geodesic in M is closed and viewed as a helix (of general order) in the ambient space.

Introduction

Let f be an isometric immersion of a Riemannian manifold M into a Riemannian manifold \tilde{M} . If geodesics in M are viewed as specific curves in \tilde{M} , what are the shape of $f(M)$? Several geometers studied this problem. K. Sakamoto investigated an isometric immersion f of a complete connected Riemannian manifold M into a Euclidean space and a sphere such that every geodesic in M is viewed as a helix in the ambient space and that the order and the Frenet curvatures of the helix are independent of the choice of the geodesic (cf. [13], [14]). Such an immersion is called a *helical immersion*. On the other hand, we recently investigated an isometric immersion of a compact connected Riemannian manifold M into a Euclidean space and a sphere such that every geodesic in M is viewed as a helix in the ambient space, where the order and the Frenet curvatures of the helix may depend on the choice of the geodesic. We called such an immersion a *generalized helical immersion* and the maximal order of those helices the *order of f* . It is easy to show that f is of even order if M is compact and the ambient space is a Euclidean space. In [9], we obtained the following characterizing theorem:

Mathematics Subject Classification (1991):53C42.
Received December 19, 1997.

Let f be a generalized helical immersion of order $2d$ of a compact connected Riemannian manifold M into a Euclidean space. Assume that the following condition holds: (*) for each $p \in M$, there is at least one geodesic σ in M through p such that $f \circ \sigma$ is a generic helix of order $2d$ in the ambient space. Then the second fundamental form of f is parallel and hence f is congruent to the standard isometric embedding of a symmetric R -space of rank d .

Here a generic helix of order $2d$ is a helix in a Euclidean space whose closure is a d -dimensional Clifford torus. In case of $d \geq 2$, the above condition (*) assures the existence of a non-closed geodesic in M because a generic helix of order $2d$ ($d \geq 2$) is non-closed. In this paper, we investigate a generalized helical immersion f of a compact connected Riemannian manifold M all of whose geodesics are closed into a Euclidean space or a sphere. Concretely, we show that, if such an immersion f is an embedding, then M is a SC-manifold (see Theorem 3.2) and, under certain additional conditions, f is helical, where f may not be an embedding (see Theorem 3.6). Here a SC-manifold is a Riemannian manifold all of whose geodesics are simply closed geodesics with the same length.

In Sect. 1 and 2, we prepare basic notations, definitions and lemmas. In Sect. 3 and 4, we prove main results in terms of basic lemmas prepared in Sect. 2.

Throughout this paper, unless otherwise mentioned, we assume that all geometric objects are of class C^∞ and all manifolds are connected ones without boundary.

1. Notations and definitions

In this section, we shall state basic notations and definitions. Let $\sigma : I \rightarrow M$ be a curve in a Riemannian manifold M parametrized by the arclength s , where I is an open interval of the real line \mathbb{R} . Denote by v_0 the velocity vector field $\dot{\sigma}$ of σ . Let ∇ be the Levi-Civita connection of M . If there exist an orthonormal system field (v_1, \dots, v_{d-1}) along σ and positive constants $\lambda_1, \dots, \lambda_{d-1}$ satisfying the following relations

$$(1.1) \quad \left\{ \begin{array}{l} \nabla_{v_0} v_0 = \lambda_1 v_1 \\ \nabla_{v_0} v_1 = -\lambda_1 v_0 + \lambda_2 v_2 \\ \vdots \\ \nabla_{v_0} v_{d-2} = -\lambda_{d-2} v_{d-3} + \lambda_{d-1} v_{d-1} \\ \nabla_{v_0} v_{d-1} = -\lambda_{d-1} v_{d-2}, \end{array} \right.$$

then σ is called a *helix of order d* . The relation (1.1), λ_i , v_i ($1 \leq i \leq d-1$) and (v_0, \dots, v_{d-1}) are called the Frenet formula, the i -th Frenet curvature, the i -th

Frenet normal vector field and the Frenet frame field of σ , respectively. In particular, a helix σ of order $2d$ in an m -dimensional Euclidean space \mathbf{R}^m is expressed as follows:

$$(1.2) \quad \sigma(s) = c_0 + \sum_{i=1}^d r_i (\cos(a_i s) e_{2i-1} + \sin(a_i s) e_{2i}),$$

where c_0 is a constant vector of \mathbf{R}^m , (e_1, \dots, e_{2d}) is an orthonormal system of \mathbf{R}^m , r_i ($1 \leq i \leq d$) are positive constants and a_i ($1 \leq i \leq d$) are mutually distinct positive constants. Note that the image $\text{Im } \sigma$ of σ is contained in the d -dimensional Clifford torus

$$T := \left\{ c_0 + \sum_{i=1}^d r_i (\cos \theta_i \cdot e_{2i-1} + \sin \theta_i \cdot e_{2i}) \mid 0 \leq \theta_i < 2\pi (i = 1, \dots, d) \right\}.$$

Also, helices in an m -dimensional sphere S^m are as follows. Let σ be a helix in S^m and ι the totally umbilic embedding of S^m into \mathbf{R}^{m+1} . Then we see that $\iota \circ \sigma$ is a helix of even order in \mathbf{R}^{m+1} . Let $2d$ be the order of $\iota \circ \sigma$. It is shown that the order of σ is $2d - 1$ (resp. $2d$) if the centroid of the d -dimensional Clifford torus containing $\text{Im}(\iota \circ \sigma)$ coincides (resp. does not coincide) with the center of S^m .

Let f be an isometric immersion of an n -dimensional Riemannian manifold M^n into an m -dimensional Riemannian manifold \tilde{M}^m . Denote by $T_p M$ (resp. $S_p M$) the tangent space (resp. the unit tangent sphere) of M at p and SM the unit tangent bundle of M . We shall identify $T_p M$ with $f_*(T_p M)$, where f_* is the differential of f . Denote by ∇ (resp. $\tilde{\nabla}$) the Levi-Civita connection on M (resp. \tilde{M}) and A, h and ∇^\perp the shape operator, the second fundamental form and the normal connection of f , respectively. Denote by the same symbol $\bar{\nabla}$ both $\nabla^* \otimes \cdots \otimes \nabla^* \otimes \nabla^\perp$ and $\nabla^\perp \otimes \nabla^* \otimes \cdots \otimes \nabla^* \otimes \nabla$, where ∇^* is the dual connection of ∇ . Also, we shall denote the i -th order derivative of h (resp. A) with respect to $\bar{\nabla}$ by $\bar{\nabla}^i h$ (resp. $\bar{\nabla}^i A$). If, for every geodesic σ in M , $f \circ \sigma$ is a helix of order d and the Frenet curvatures of $f \circ \sigma$ do not depend on the choice of σ , then f is called a *helical immersion of order d* . Also, if, for every geodesic σ in M , $f \circ \sigma$ is a helix of order at most d and there is at least one geodesic σ_0 in M such that $f \circ \sigma_0$ is a helix of order d , then we shall call f a *generalized helical immersion of order d* .

2. Basic lemmas

In this section, we prepare basic lemmas which will be used in the next section. Let f be a generalized helical immersion of an n -dimensional complete

Riemannian manifold M^n into an m -dimensional Euclidean space \mathbf{R}^m . For each $v \in SM$, denote by σ_v the maximal geodesic in M parametrized by the arc-length s whose velocity vector at $s = 0$ is equal to v . For $p \in M$, we set $V_{p,i} := \{v \in S_p M \mid f \circ \sigma_v : \text{helix of order } i\}$ ($i \geq 1$) and define a function $\hat{\lambda}_i$ ($i \geq 1$) on SM by

$$\hat{\lambda}_i(v) := \begin{cases} \lambda_i & (v \in \bigcup_{i+1 \leq j} \bigcup_{p \in M} V_{p,j}) \\ 0 & (v \in \bigcup_{1 \leq j \leq i} \bigcup_{p \in M} V_{p,j}), \end{cases}$$

where λ_i is the i -th Frenet curvature of $f \circ \sigma_v$. It is easy to show that $\hat{\lambda}_i$ is continuous on $\bigcup_{i \leq j} \bigcup_{p \in M} V_{p,j}$ ($i \geq 1$). In [9], we proved the following lemma.

LEMMA 2.1. *Assume that $V_{p,d} \neq \emptyset$ and $V_{p,i} = \emptyset$ ($i \geq d+1$) for $p \in M$. Then the set $V_{p,i}$ ($1 \leq i \leq d-1$) are closed sets of measure zero in $S_p M$ and $V_{p,d}$ is a dense open set in $S_p M$.*

In the sequel, assume that $V_{p,2d} \neq \emptyset$ and $V_{p,i} = \emptyset$ ($i \geq 2d+1$) for some $p \in M$. For each $v \in V_{p,2d}$, $f \circ \sigma_v$ is uniquely expressed as

$$(f \circ \sigma_v)(s) = c(v) + \sum_{i=1}^d r_i(v) (\cos(a_i(v)s) e_{2i-1}(v) + \sin(a_i(v)s) e_{2i}(v)),$$

where $c(v)$ is a constant vector of \mathbf{R}^m , $(e_1(v), \dots, e_{2d}(v))$ is an orthonormal system of \mathbf{R}^m , $r_i(v)$ ($1 \leq i \leq d$) are positive constants and $a_i(v)$ ($1 \leq i \leq d$) are positive constants with $a_1(v) < \dots < a_d(v)$. We regard r_i and a_i ($1 \leq i \leq d$) as functions on $V_{p,2d}$. In [9], we proved the following lemma.

LEMMA 2.2. *The functions a_i ($1 \leq i \leq d$) are analytic.*

Also, we prepare the following lemma.

LEMMA 2.3 *On $V_{p,2d}$, the following relation holds:*

$$\begin{pmatrix} a_1^2 & \cdots & a_d^2 \\ a_1^4 & \cdots & a_d^4 \\ \vdots & & \vdots \\ a_1^{2d} & \cdots & a_d^{2d} \end{pmatrix} \begin{pmatrix} r_1^2 \\ \vdots \\ r_d^2 \end{pmatrix} = \begin{pmatrix} 1 \\ F_1(\hat{\lambda}_1) \\ \vdots \\ F_{d-1}(\hat{\lambda}_1, \dots, \hat{\lambda}_{d-1}) \end{pmatrix},$$

where F_i is a polynomial of i -variables ($1 \leq i \leq d-1$).

PROOF. Fix $v \in V_{p,2d}$. Let $(v_0, v_1, \dots, v_{2d-1})$ be the Frenet frame field of $f \circ \sigma_v$. Then we have

$$(2.1) \quad v_0 = \sum_{i=1}^d r_i(v) a_i(v) (-\sin(a_i(v)s)e_{2i-1}(v) + \cos(a_i(v)s)e_{2i}(v))$$

and hence $\sum_{i=1}^d r_i(v)^2 a_i(v)^2 = 1$. Thus, if $d = 1$, then the proof is completed. In the sequel, assume $d \geq 2$. By operating $\tilde{\nabla}_{v_0}$ to (2.1), we have

$$(2.2) \quad \hat{\lambda}_1(v)v_1 = - \sum_{i=1}^d r_i(v) a_i(v)^2 (\cos(a_i(v)s)e_{2i-1}(v) + \sin(a_i(v)s)e_{2i}(v))$$

and hence $\sum_{i=1}^d r_i(v)^2 a_i(v)^4 = \hat{\lambda}_1(v)^2$. Thus, if $d = 2$, then the proof is completed.

In the sequel, assume $d \geq 3$. Furthermore, by operating $\tilde{\nabla}_{v_0}$ to (2.2), we have

$$\begin{aligned} & \hat{\lambda}_1(v)(-\hat{\lambda}_1(v)v_0 + \hat{\lambda}_2(v)v_2) \\ &= \sum_{i=1}^d r_i(v) a_i(v)^3 (\sin(a_i(v)s)e_{2i-1}(v) - \cos(a_i(v)s)e_{2i}(v)) \end{aligned}$$

and hence $\sum_{i=1}^d r_i(v)^2 a_i(v)^6 = \hat{\lambda}_1(v)^4 + \hat{\lambda}_1(v)^2 \hat{\lambda}_2(v)^2$. Thus, if $d = 3$, then the proof is completed. In case of $d \geq 4$, by repeating the same process, we can obtain

$$\sum_{i=1}^d r_i(v)^2 a_i(v)^{2j} = F_{j-1}(\hat{\lambda}_1(v), \dots, \hat{\lambda}_{j-1}(v)) \quad (4 \leq j \leq d),$$

where F_{j-1} is a polynomial of $(j-1)$ -variables ($4 \leq j \leq d$). This completes the proof. \square

3. Generalized helical immersions into a Euclidean space

In this section, we shall investigate a generalized helical immersion f of an n -dimensional compact Riemannian manifold M all of whose geodesics are closed into an m -dimensional Euclidean space \mathbb{R}^m . Since M is compact, f is of even order. Let $2d$ be the order of f . Take $p \in M$ with $V_{p,2d} \neq \emptyset$. Since all of geodesics in M are closed, they admit a common period by Lemma 7.11 of [1, P182]. Let $\mu := \max_{v \in S_p M} l(\sigma_v)$, where $l(\sigma_v)$ is the length of σ_v (i.e., the minimal period of σ_v). Let $W := \{v \in V_{p,2d} \mid l(\sigma_v) = \mu\}$. Since $l(\sigma_v)$ ($v \in V_{p,2d}$) are divisors of the common period, $\{l(\sigma_v) \mid v \in V_{p,2d}\}$ is a discrete set. The function ϕ on $V_{p,2d}$ defined by $\phi(v) = l(\sigma_v)$ is lower semi-continuous. These facts deduce that $W = \phi^{-1}(\mu)$ is an open set in $V_{p,2d}$. Let a_i ($1 \leq i \leq d$) be functions on $V_{p,2d}$ stated in Sect. 2. First we shall show the following lemma.

LEMMA 3.1. *The set $V_{p,2d}$ coincides with $S_p M$ and a_i ($1 \leq i \leq d$) are constant on $S_p M$.*

PROOF. (Step I) First we shall show that the functions a_i ($1 \leq i \leq d$) are constant on each component of $V_{p,2d}$ which intersects with W . Let W_0 be a component of W . For each $v \in W_0$, set $s_v := \min\{s \mid a_i(v)s \in N \ (1 \leq i \leq d)\}$. Clearly $l(f \circ \sigma_v) = 2\pi s_v$ holds. Also, we can show $l(\sigma_v)/l(f \circ \sigma_v) \in N$. Hence we have $\mu/2\pi s_v \in N$, which together with $a_i(v)s_v \in N$ implies $a_i(v)\mu/2\pi \in N$. Therefore, it follows from the continuity of a_i that a_i is constant on W_0 . Thus a_i is constant on each component of W . This together with the analyticity of a_i (by Lemma 2.2) implies that a_i is constant on each component of $V_{p,2d}$ which intersects with W .

(Step II) Next we shall show $V_{p,2d} = S_p M$. Let V_0 be a component of $V_{p,2d}$ which intersects with W . We showed that a_i ($1 \leq i \leq d$) are constant on V_0 . Denote by \bar{V}_0 the closure of V_0 in $S_p M$. Take $v \in \bar{V}_0$ and a sequence $\{w_k\}_{k=1}^\infty$ in V_0 with $\lim_{k \rightarrow \infty} w_k = v$. The helix $f \circ \sigma_{w_k}$ is uniquely expressed as

(3.1)

$$(f \circ \sigma_{w_k})(s) = c(w_k) + \sum_{i=1}^d r_i(w_k)(\cos(a_i(w_k)s)e_{2i-1}(w_k) + \sin(a_i(w_k)s)e_{2i}(w_k)).$$

Since helices $f \circ \sigma_{w_k}$ ($k \in N$) are contained in a compact set $f(M)$, we have $\sup_k \|c(w_k)\| < \infty$ and $\sup_k r_i(w_k) < \infty$ ($1 \leq i \leq d$). Set $C := \sup_k \|c(w_k)\|$ and $R_i := \sup_k r_i(w_k)$ ($1 \leq i \leq d$). Since $\{(e_1(w_k), \dots, e_{2d}(w_k), c(w_k), r_1(w_k), \dots, r_d(w_k))\}_{k=1}^\infty$ is a sequence in a compact set $S_{m,2d} \times B^m(C) \times [0, R_1] \times \dots \times [0, R_d]$, its convergent subsequence $\{(e_1(w_{\alpha(k)}), \dots, e_{2d}(w_{\alpha(k)}), c(w_{\alpha(k)}), r_1(w_{\alpha(k)}), \dots, r_d(w_{\alpha(k)}))\}_{k=1}^\infty$ exists, where $S_{m,2d}$ is the Stiefel manifold of all orthonormal $2d$ -frames in \mathbb{R}^m , $B^m(C)$ is the m -dimensional ball of center C in \mathbb{R}^m and $[0, R_i]$ ($1 \leq i \leq d$) are closed intervals. Let $(e_1^0, \dots, e_{2d}^0, c^0, r_1^0, \dots, r_d^0) := \lim_{k \rightarrow \infty} (e_1(w_{\alpha(k)}), \dots, e_{2d}(w_{\alpha(k)}), c(w_{\alpha(k)}), r_1(w_{\alpha(k)}), \dots, r_d(w_{\alpha(k)}))$. From (3.1) and the constancy of a_i on V_0 , we have

$$\begin{aligned} \lim_{k \rightarrow \infty} (f \circ \sigma_{w_{\alpha(k)}})(s) &= \lim_{k \rightarrow \infty} \left\{ c(w_{\alpha(k)}) + \sum_{i=1}^d r_i(w_{\alpha(k)})(\cos(a_i(w_{\alpha(k)})s)e_{2i-1}(w_{\alpha(k)}) \right. \\ &\quad \left. + \sin(a_i(w_{\alpha(k)})s)e_{2i}(w_{\alpha(k)})) \right\} \\ &= c_0 + \sum_{i=1}^d r_i^0(\cos(a_i(w_1)s)e_{2i-1}^0 + \sin(a_i(w_1)s)e_{2i}^0). \end{aligned}$$

On the other hand, we have

$$\begin{aligned}\lim_{k \rightarrow \infty} (f \circ \sigma_{w_{\alpha(k)}})(s) &= \lim_{k \rightarrow \infty} (f \circ \exp_p)(sw_{\alpha(k)}) = (f \circ \exp_p)(sv) \\ &= (f \circ \sigma_v)(s),\end{aligned}$$

where \exp_p is the exponential map of M at p . Thus we can obtain

$$(f \circ \sigma_v)(s) = c_0 + \sum_{i=1}^d r_i^0 (\cos(a_i(w_1)s)e_{2i-1}^0 + \sin(a_i(w_1)s)e_{2i}^0),$$

which implies that $f \circ \sigma_v$ is a helix of order $2d$, that is, $v \in V_{p,2d}$. Clearly, this implies $v \in V_0$. Therefore, we have $\bar{V}_0 = V_0$, that is, V_0 is closed in $S_p M$. On the other hand, since $V_{p,2d}$ is open in $S_p M$ by Lemma 2.1, so is also V_0 . Hence, it follows from the connectedness of $S_p M$ that $V_0 = S_p M$, that is, $V_{p,2d} = S_p M$. This completes the proof. \square

From this lemma, we can prove the following result.

THEOREM 3.2. *Let f be a generalized helical immersion of an n -dimensional compact Riemannian manifold M all of whose geodesics are closed into an m -dimensional Euclidean space \mathbb{R}^m . Then the following statements (i) and (ii) hold:*

- (i) *all geodesics in M are viewed as closed helices of the same order with the same length in \mathbb{R}^m ,*
- (ii) *if f is an embedding, then M is a SC-manifold.*

PROOF. Let $2d$ be the order of f . Take $p \in M$ with $V_{p,2d} \neq \emptyset$. From Lemma 3.1, it follows that $V_{p,2d} = S_p M$ and that a_i ($1 \leq i \leq d$) are constant on $S_p M$. This implies that all geodesics in M through p are viewed as closed helices of order $2d$ with the same length in \mathbb{R}^m . Take an arbitrary $q \in M$. Since M is compact and hence complete, there is a geodesic in M through p and q . This implies $V_{q,2d} \neq \emptyset$. Hence, we see that all geodesics in M through p or q are viewed as closed helices of order $2d$ with the same length in \mathbb{R}^m . Thus the statement (i) is deduced from the arbitrariness of q . Assume that f is an embedding. Since a closed helix in \mathbb{R}^m are simply closed, all geodesics in M are simply closed geodesics. Also, $l(\sigma) = l(f \circ \sigma)$ holds for each geodesic σ in M . From the statement (i), $l(f \circ \sigma)$ is independent of the choice of σ . Therefore, so is also $l(\sigma)$, that is, all geodesics in M have the same length. Thus M is a SC-manifold. \square

From the statement (ii) of this theorem, we can obtain the following corollary.

COROLLARY 3.3. *Let $f : M \hookrightarrow \mathbb{R}^m$ be an immersion as in Theorem 3.2. If f is an embedding and M is a Riemannian homogeneous space, then M is isometric to a compact symmetric space of rank one.*

PROOF. By the statement (ii) of Theorem 3.2, M is a SC-manifold. Hence, since M is a Riemannian homogeneous space, M is isometric to a compact symmetric space of rank one by Theorem 7.55 of [1, P196]. \square

Next we shall investigate in what case an immersion as in Theorem 3.2 is helical. First we shall show the following lemma.

LEMMA 3.4. *Let f be an immersion as in Theorem 3.2 and $2d$ the order of f . In case of $d \geq 2$, assume that $\hat{\lambda}_i$ ($1 \leq i \leq d-1$) are constant on SM , where $\hat{\lambda}_i$ ($1 \leq i \leq d-1$) are functions defined in Sect. 2. Then f is helical.*

PROOF. Fix $p \in M$. By Theorem 3.2, $V_{p,2d} = S_p M$ holds. For each $v \in S_p M$, $f \circ \sigma_v$ is uniquely expressed as

$$(3.2) \quad (f \circ \sigma_v)(s) = c(v) + \sum_{i=1}^d r_i(v) (\cos(a_i(v)s)e_{2i-1}(v) + \sin(a_i(v)s)e_{2i}(v)).$$

It follows from Lemma 3.1 that a_i ($1 \leq i \leq d$) are constant on $S_p M$. Hence, since $\hat{\lambda}_i$ ($1 \leq i \leq d-1$) are constant on $S_p M$ by the assumption, so are also r_i ($1 \leq i \leq d$) by Lemma 2.3. Therefore, by (3.2), $f \circ \sigma_v$ ($v \in S_p M$) are mutually congruent, that is, they have the same Frenet curvatures. This together with the arbitrariness of p and the completeness of M implies that f is helical. \square

Define functions F_{ij} ($i \geq 0, j \geq 0$) on SM by

$$F_{ij}(v) := \langle (\bar{\nabla}^i h)(v, \dots, v), (\bar{\nabla}^j h)(v, \dots, v) \rangle \quad (v \in SM)$$

and functions G_{ij} on the Stiefel bundle $V_2(M)$ of M of all orthonormal 2-frames of M by

$$G_{ij}(v, w) := \langle (\bar{\nabla}^i h)(v, \dots, v), (\bar{\nabla}^j h)(v, \dots, v, w) \rangle \quad ((v, w) \in V_2(M)).$$

Now we shall prepare another lemma.

LEMMA 3.5. *Let f be an immersion as in Theorem 3.2 and $2d$ the order of f . Assume that F_{kk} is constant on SM and $\bar{\nabla}^i h$ ($i \geq 2$) are symmetric, where k is a fixed non-negative integer. Then the following relations hold:*

$$\begin{aligned} G_{kk} &= 0, \quad \sum_{i=0}^j \binom{j}{i} F_{k+j-i, k+i+1} = 0, \\ \sum_{i=0}^j \binom{j}{i} G_{k+i+1, k+j-i} &= 0, \quad \sum_{i=0}^j \binom{j}{i} G_{k+i, k+j-i+1} = 0 \quad (j \geq 0). \end{aligned}$$

PROOF. Take an arbitrary point p of M and furthermore, take an arbitrary orthonormal 2-frame (v, w) of M at p . Let \tilde{v} be the velocity vector field of the geodesic σ_v . By operating d/ds to the constant function $F_{kk}(\tilde{v})$, we have $F_{k,k+1}(\tilde{v}) = 0$, where s is the arclength of σ_v . Furthermore, by operating $(d/ds)^j$ to $F_{k,k+1}(\tilde{v}) = 0$ and substituting $s = 0$, we can obtain $\sum_{i=0}^j \binom{j}{i} F_{k+j-i, k+i+1}(v) = 0$. Hence, by the arbitrariness of v and p , $\sum_{i=0}^j \binom{j}{i} F_{k+j-i, k+i+1} = 0$ holds on SM . By differentiating $F_{kk}|_{S_p M}$ in the direction $w (\in T_v(S_p M))$, we have $G_{kk}(v, w) = 0$. By the arbitrariness of (v, w) and p , $G_{kk} = 0$ holds on $V_2(M)$. Let \tilde{w} be the parallel vector field along σ_v with $\tilde{w}(0) = w$. By operating d/ds to $G_{kk}(\tilde{v}, \tilde{w})$ and substituting $s = 0$, we have $G_{k+1,k}(v, w) + G_{k,k+1}(v, w) = 0$. Also, by differentiating $F_{k,k+1}|_{S_p M} = 0$ in the direction $w (\in T_v(S_p M))$, we have

$$(k+2)G_{k+1,k}(v, w) + (k+3)G_{k,k+1}(v, w) = 0.$$

Therefore, we can obtain $G_{k+1,k}(v, w) = G_{k,k+1}(v, w) = 0$ and hence, by the arbitrariness of (v, w) and p , $G_{k+1,k} = G_{k,k+1} = 0$ holds on $V_2(M)$. Furthermore, by operating $(d/ds)^j$ to $G_{k+1,k}(\tilde{v}, \tilde{w}) = G_{k,k+1}(\tilde{v}, \tilde{w}) = 0$ and substituting $s = 0$, we can obtain

$$\sum_{i=0}^j \binom{j}{i} G_{k+i+1, k+j-i}(v, w) = \sum_{i=0}^j \binom{j}{i} G_{k+i, k+j-i+1}(v, w) = 0.$$

Hence, by the arbitrariness of (v, w) and p ,

$$\sum_{i=0}^j \binom{j}{i} G_{k+i+1, k+j-i} = \sum_{i=0}^j \binom{j}{i} G_{k+i, k+j-i+1} = 0$$

holds on $V_2(M)$. □

From these lemmas, we can show the following result.

THEOREM 3.6. *Let f be a generalized helical immersion of order $2d$ of an n -dimensional compact Riemannian manifold M all of whose geodesics are closed into an m -dimensional Euclidean space \mathbb{R}^m . In case of $d \geq 2$, assume that, for each $p \in M$, $\|(\bar{\nabla}^i h)(v, \dots, v)\|$ is independent of the choice of $v \in S_p M$ ($0 \leq i \leq d-2$) and furthermore, in case of $d \geq 6$, assume that $\bar{\nabla}^i h$ is symmetric ($2 \leq i \leq [d/2]-1$), where $[]$ is the Gauss's symbol. Then f is helical.*

PROOF. If $d = 1$, then f is a planar geodesic immersion and hence a helical immersion of order 2. In the sequel, assume that $d \geq 2$. Take an arbitrary point p_0 of M and furthermore take an arbitrary orthonormal 2-frame (v, w) of M at p_0 . Let (v_0, \dots, v_{2d-1}) (resp. λ_i ($1 \leq i \leq 2d-1$)) be the Frenet frame (resp. i -th Frenet curvature) of $f \circ \sigma_v$, where we note that $f \circ \sigma_v$ is of order $2d$ by Theorem 3.2. From the Gauss formula and the Frenet formula, we have

$$(3.3) \quad \lambda_1 v_1 = h(v_0, v_0)$$

and hence $\hat{\lambda}_1(v)^2 = F_{00}(v)$. By the arbitrariness of v and p_0 , we see that $\hat{\lambda}_1^2 = F_{00}$ holds on SM . By the assumption, $\hat{\lambda}_1$ is constant on $S_p M$ for each $p \in M$. Furthermore, since f is generalized helical and there exists a geodesic through arbitrary two points of M by the compactness of M , $\hat{\lambda}_1 (= F_{00})$ is constant on SM . Thus, if $d = 2$, then f is helical by Lemma 3.4. In the sequel, assume $d \geq 3$. By operating $\tilde{\nabla}_{v_0}$ to (3.3), we have

$$-\lambda_1^2 v_0 + \lambda_1 \lambda_2 v_2 = -A_{h(v_0, v_0)} v_0 + (\bar{\nabla} h)(v_0, v_0, v_0).$$

Also, since F_{00} is constant on SM , it follows from Lemma 3.5 that $G_{00} = 0$ and hence $A_{h(v_0, v_0)} v_0 = F_{00}(v_0) v_0$, where we note that the symmetricness of h is used. So we can obtain

$$(3.4) \quad -\lambda_1^2 v_0 + \lambda_1 \lambda_2 v_2 = -F_{00}(v_0) v_0 + (\bar{\nabla} h)(v_0, v_0, v_0)$$

and hence

$$\hat{\lambda}_1(v)^4 + \hat{\lambda}_1(v)^2 \hat{\lambda}_2(v)^2 = F_{00}(v)^2 + F_{11}(v).$$

By the arbitrariness of v and p_0 , we see that

$$(3.5) \quad \hat{\lambda}_1^4 + \hat{\lambda}_1^2 \hat{\lambda}_2^2 = F_{00}^2 + F_{11}$$

holds on SM . Since $\hat{\lambda}_1$ and F_{00} are constant on SM and F_{11} is constant on S_pM for each $p \in M$, it follows from (3.5) that $\hat{\lambda}_2$ is constant on S_pM for each $p \in M$. Furthermore, since f is a generalized helical and there exists a geodesic through arbitrary two points of M , $\hat{\lambda}_2$ is constant on SM . Thus, if $d = 3$, then f is helical by Lemma 3.4. In the sequel, assume $d \geq 4$. By operating $\tilde{\nabla}_{v_0}$ to (3.4), we have

$$\begin{aligned} & -\lambda_1(\lambda_1^2 + \lambda_2^2)v_1 + \lambda_1\lambda_2\lambda_3v_3 \\ &= -2F_{10}(v_0)v_0 + F_{00}(v_0)h(v_0, v_0) \\ & \quad - A_{(\bar{\nabla}h)(v_0, v_0, v_0)}v_0 + (\bar{\nabla}^2h)(v_0, \dots, v_0). \end{aligned}$$

Also, since F_{00} is constant on SM , we have $F_{10} = 0$ and $G_{10} = 0$ by Lemma 3.5, where we note that the symmetricness of $\bar{\nabla}h$ is used. So we have

$$(3.6) \quad -\lambda_1(\lambda_1^2 + \lambda_2^2)v_1 + \lambda_1\lambda_2\lambda_3v_3 = F_{00}(v_0)h(v_0, v_0) + (\bar{\nabla}^2h)(v_0, \dots, v_0).$$

and hence

$$\begin{aligned} & \hat{\lambda}_1(v)^2(\hat{\lambda}_1(v)^2 + \hat{\lambda}_2(v)^2)^2 + \hat{\lambda}_1(v)^2\hat{\lambda}_2(v)^2\hat{\lambda}_3(v)^2 \\ &= F_{00}(v)^3 + 2F_{00}(v)F_{20}(v) + F_{22}(v). \end{aligned}$$

By the arbitrariness of v and p_0 , we see that

$$(3.7) \quad \hat{\lambda}_1^2(\hat{\lambda}_1^2 + \hat{\lambda}_2^2)^2 + \hat{\lambda}_1^2\hat{\lambda}_2^2\hat{\lambda}_3^2 = F_{00}^3 + 2F_{00}F_{20} + F_{22}$$

holds on SM . Since $\hat{\lambda}_1$, $\hat{\lambda}_2$ and F_{00} are constant on SM , so is also F_{11} by (3.5). Furthermore, since F_{00} and F_{11} are constant on SM , so is also F_{20} by Lemma 3.5. Therefore, since $\hat{\lambda}_1$, $\hat{\lambda}_2$, F_{00} and F_{20} are constant on SM and F_{22} is constant on S_pM for each $p \in M$, it follows from (3.7) that $\hat{\lambda}_3$ is constant on S_pM for each $p \in M$. Moreover, since f is generalized helical and there exists a geodesic through arbitrary two points of M , $\hat{\lambda}_3$ is constant on SM . Thus, if $d = 4$, then f is helical by Lemma 3.4. In case of $d \geq 5$, by repeating the same process, we can show that $\hat{\lambda}_i$ ($4 \leq i \leq d-1$) are constant on SM . Hence f is helical by Lemma 3.4. \square

Now we shall recall examples of a helical immersion into a sphere (or a Euclidean space) given by K. Tsukada in [18]. Let M be an n -dimensional compact symmetric space of rank one. Let V_k be the eigenspace for k -th eigenvalue λ_k of the Laplace operator on M and let $\dim V_k = m(k) + 1$. We define an inner product \langle , \rangle on V_k by $\langle \phi, \psi \rangle := \int_M \phi \psi dV$, where dV is the volume element of M . We define a map $\Phi_k : M \rightarrow \mathbf{R}^{m(k)+1}$ by $\Phi_k(p) := \sqrt{n/\lambda_k}(\phi_0(p), \dots, \phi_{m(k)}(p))$, where $(\phi_0, \dots, \phi_{m(k)})$ is an orthonormal base of V_k .

Then Φ_k becomes a helical immersion. Furthermore, it is shown that $\Phi_k(M)$ is contained in a hypersphere $S^{m(k)}$ of $\mathbf{R}^{m(k)+1}$ and that $\Phi_k : M \hookrightarrow S^{m(k)}$ is minimal and helical. The isometric immersion Φ_k is called the k -th *standard minimal immersion* into $S^{m(k)}$. K. Tsukada defined an isometric immersion $\Phi_{k_1 \dots k_r}$ of M into $\mathbf{R}^{m(k_1) + \dots + m(k_r) + r}$ by

$$\Phi_{k_1 \dots k_r}(p) := (c_1 \Phi_{k_1}(p), \dots, c_r \Phi_{k_r}(p)),$$

where k_1, \dots, k_r are positive integers and c_1, \dots, c_r are positive numbers with $c_1^2 + \dots + c_r^2 = 1$. He showed that it is a helical immersion into a hypersphere $S^{m(k_1) + \dots + m(k_r) + r - 1}$ of $\mathbf{R}^{m(k_1) + \dots + m(k_r) + r}$ (cf. [18]). Now we can obtain the following result in terms of Theorem 3.6 and Theorem 4.7 of [14].

COROLLARY 3.7. *Under the hypothesis in Theorem 3.6, assume that f is a full embedding and $\dim M = 2$ or odd integer. Then M is isometric to a sphere or a real projective space and f is congruent to the above immersion $\Phi_{k_1 \dots k_r}$.*

4. Generalized helical immersions into a sphere

In this section, we shall deduce some results for a generalized helical immersion into a sphere in terms of results in the previous section. First we can deduce the following result from Theorem 3.2.

THEOREM 4.1. *Let f be a generalized helical immersion of an n -dimensional compact Riemannian manifold M all of whose geodesics are closed into an m -dimensional sphere S^m . Then the following statements (i) and (ii) hold:*

- (i) *if f is of odd (resp. even) order d , then all geodesics in M are viewed as closed helices of order d (resp. d or $d - 1$) with the same length in S^m ,*
- (ii) *if f is an embedding, then M is a SC-manifold.*

PROOF. Let ι be the totally umbilical embedding of S^m into \mathbf{R}^{m+1} and set $\tilde{f} := \iota \circ f$. It is clear that \tilde{f} is generalized helical. Hence, it follows from Theorem 3.2 that all geodesics in M are viewed as closed helices of the same order with the same length in \mathbf{R}^{m+1} . This deduces the statement (i) because a helix of order d in S^m is viewed as a helix of order $2[(d + 1)/2]$ in \mathbf{R}^{m+1} . If f is an embedding, then so is also \tilde{f} . Hence, the statement (ii) is deduced from Theorem 3.2. \square

From the statement (ii) of this theorem, we can obtain the following corollary.

COROLLARY 4.2. *Let $f : M \hookrightarrow S^m$ be an immersion as in Theorem 4.1. If f is an embedding and M is a Riemannian homogeneous space, then M is isometric to a compact symmetric space of rank one.*

Also, we can deduce the following result from Theorem 3.6.

THEOREM 4.3. *Let f be a generalized helical immersion of order $2d - 1$ or $2d$ of an n -dimensional compact Riemannian manifold M all of whose geodesics are closed into an m -dimensional sphere S^m . In case of $d \geq 2$, assume that, for each $p \in M$, $\|(\bar{\nabla}^i h)(v, \dots, v)\|$ is independent of $v \in S_p M$ ($0 \leq i \leq d - 2$) and furthermore, in case of $d \geq 6$, assume that $\bar{\nabla}^i h$ is symmetric ($2 \leq i \leq [d/2] - 1$). Then f is helical.*

PROOF. Let ι be the totally umbilical embedding of S^m into \mathbb{R}^{m+1} and set $\tilde{f} := \iota \circ f$. From the assumptions, we can show that \tilde{f} satisfies the conditions of Theorem 3.6. Hence \tilde{f} is helical by Theorem 3.6. This implies that so is also f . \square

Also, we can obtain the following result from Corollary 3.7.

COROLLARY 4.4. *Under the hypothesis in Theorem 4.3, assume that f is a full embedding and $\dim M = 2$ or odd integer. Then M is isometric to a sphere or a real projective space and f is congruent to the immersion $\Phi_{k_1 \dots k_r}$ stated in Sect. 3.*

References

- [1] Besse, A. L.: Manifolds all of whose geodesics are closed. Berlin, Heidelberg, New York: Springer 1978.
- [2] Ferus, D.: Immersionen mit paralleler zweiter Fundamentalform. Beispiele und Nicht-Beispiele. Manuscripta Math. **12**, 153–162 (1974).
- [3] Ferus, D.: Immersions with parallel second fundamental form. Math. Z. **140**, 87–93 (1974).
- [4] Ferus, D., Schirrmacher, S.: Submanifolds in Euclidean space with simple geodesics. Math. Ann. **260**, 57–62 (1982).
- [5] Helgason, S.: Differential geometry, Lie groups and symmetric spaces, Academic Press, New York, 1978.
- [6] Hong, S. L.: Isometric immersions of manifolds with plane geodesics into a Euclidean space. J. Differential Geometry **8**, 259–278 (1973).
- [7] Ki, U. H., Kim, Y. H.: Surfaces with simple geodesics through a point. J. of Geometry **51**, 67–78 (1994).
- [8] Kobayashi, S.: Isometric imbeddings of compact symmetric spaces. Tôhoku Math. J. **20**, 21–25 (1968).
- [9] Koike, N.: Generalized Helical Immersions. Tsukuba J. of Math. **22**, 407–425 (1998).
- [10] Little, J. A.: Manifolds with planar geodesics. J. Differential Geometry **11**, 265–285 (1976).

- [11] Nakagawa, H.: On a certain minimal immersion of a Riemannian manifold into a sphere. *Kōdai Math. J.* **3**, 321–340 (1980).
- [12] Sakamoto, K.: Planar geodesic immersions. *Tôhoku Math. J.* **29**, 25–56 (1977).
- [13] Sakamoto, K.: Helical immersions into a unit sphere. *Math. Ann.* **261**, 63–80 (1982).
- [14] Sakamoto, K.: Helical immersions into a Euclidean space. *Michigan Math. J.* **33**, 353–364 (1986).
- [15] Song, H. H., Kimura, T., Koike, N.: On proper helices and extrinsic spheres in pseudo-Riemannian geometry. *Tsukuba J. of Math.* **20**, 263–280 (1996).
- [16] Takeuchi, M.: Parallel submanifolds of space forms. *Manifolds and Lie groups: Papers in honor of Yôzô Matsushima*, Birkhäuser **1981**, 429–447.
- [17] Takeuchi, M., Kobayashi, S.: Minimal imbeddings of R-spaces. *J. Differential Geometry* **2**, 203–215 (1968).
- [18] Tsukada, K.: Helical geodesic immersions of compact rank one symmetric spaces into spheres. *Tokyo J. of Math.* **6**, 267–285 (1983).

Department of Mathematics,
Faculty of Science,
Science University of Tokyo, Tokyo 162-0827,
Japan