

# TOTALLY COMPLEX SUBMANIFOLDS OF THE CAYLEY PROJECTIVE PLANE

By

Liu XIMIN

**Abstract.** Let  $K$  be the sectional curvature of a compact submanifold  $M$  of the Cayley projective plane  $CaP^2$ . In this paper, we prove that the compact totally complex submanifold  $M$  of complex dimension 2 in  $CaP^2$  satisfying  $K > (1/8)$  is totally geodesic and  $M = CP^2$ .

## 1. Introduction

Let  $M$  be an  $n$ -dimensional compact Kaehler submanifold of complex projective space  $CP^n(1)$ . Denote by  $K$  the sectional curvature of  $M$ . In [6], Ros and Verstraelen showed that if  $K > (1/8)$ , then  $M$  is totally geodesic. The analogous result in the case of totally complex submanifolds of quaternion projective space  $HP^n(1)$  was obtained by Xia [7]. In the present paper, we prove the following same type result for totally complex submanifolds of the Cayley projective plane  $CaP^2$ .

**THEOREM.** *Let  $M$  be a compact totally complex submanifold of complex dimension 2, immersed in the Cayley projective plane  $CaP^2$ . If the sectional curvature  $K$  of  $M$  satisfying  $K > (1/8)$ , then  $M$  is totally geodesic in  $CaP^2$  and  $M^2 = CP^2$ .*

## 2. Cayley projective plane

In this section, we review simply the fundamental results about the Cayley projective plane, for details see [4].

---

**Key words and phrases.** totally complex submanifold, Cayley projective plane, sectional curvature.  
1991 Mathematics Subject Classifications. 53C42, 53C40.

Received July 3, 1996

Revised October 14, 1996

Let us denote by  $Ca$  the Cayley number, it possesses a multiplicative identity 1 and a positive definite bilinear form  $\langle \cdot, \cdot \rangle$  with norm  $\|a\| = \langle a, a \rangle$ , satisfying  $\|ab\| = \|a\| \|b\|$ , for  $a, b \in Ca$ . Every element  $a \in Ca$  can be expressed in the form  $a = a_0 1 + a_1$  for  $a_0 \in R$  and  $\langle a_1, 1 \rangle = 0$ . The conjugation map  $a \rightarrow a^* = a_0 1 - a_1$  is an anti-automorphism  $(ab)^* = b^* a^*$ .

A canonical basis for  $Ca$  is any basis of the form  $\{1, e_0, e_1, \dots, e_6\}$  satisfying: (i)  $\langle e_i, 1 \rangle = 0$ ; (ii)  $\langle e_i, e_j \rangle = \{0 \text{ for } i \neq j, \text{ and } 1 \text{ otherwise}\}$ ; (iii)  $e_i^2 = -1$ ;  $e_i e_j + e_j e_i = 0$  ( $i \neq j$ ); (iv)  $e_i e_{i+1} = e_{i+3}$  for  $i \in Z_7$ .

Let  $V$  be a vector space of real dimension 16 with automorphism group  $Spin(9)$ . the splitting

$$V = Ca \oplus Ca$$

together with the above canonical basis on each summand, endows  $V$  with what we may refer to as a Cayley structure. We know that the Cayley projective plane  $CaP^2$  is the 16-dimensional Riemannian symmetric space whose tangent space admits the Cayley structure pointwise. In the following, let  $\{I_0, \dots, I_6\}$  be the Cayley structure on  $CaP^2$ .

The curvature tensor  $\bar{R}$  of  $CaP^2$  is given in [2] as follows

$$(2.1) \quad \begin{aligned} \bar{R}((a, b), (c, d))(e, f) = & \frac{1}{4}((4\langle c, e \rangle a - 4\langle a, e \rangle c + (ed)b^* - (eb)d^* \\ & + (ad - cb)f^*), (4\langle d, f \rangle b - 4\langle b, f \rangle d + a^*(cf) \\ & - c^*(af) - e^*(ad - cb)). \end{aligned}$$

On  $Ca \oplus Ca$  we have the positive definite bilinear form  $\langle \cdot, \cdot \rangle$  given by

$$(2.2) \quad \langle (a, b), (c, d) \rangle = \langle a, c \rangle + \langle b, d \rangle.$$

### 3. Totally complex submanifolds

Let  $V \subset T_x CaP^2$  be a real vector subspace, we say that  $V$  is a totally complex subspace if there exists an  $I$  such that there exists a basis with  $I = I_0$  and (i)  $I_0 V \subset V$ , and (ii)  $I_k V$  is perpendicular to  $V$  for  $1 \leq k \leq 6$ . Clearly, if  $V$  is a maximal subspace of this kind then  $\dim_R V = 4$ .

Let  $M$  be a compact Riemannian manifold isometrically immersed in  $CaP^2$  by  $j : M \rightarrow CaP^2$ . Denote by  $h$  and  $A$  the second fundamental form of  $j$  and the Weingarten endomorphism respectively. Then we have

$$(3.1) \quad \langle h(X, Y), N \rangle = \langle X, A_N(Y) \rangle \quad X, Y \in TM, N \in TM^\perp$$

We take  $\bar{\nabla}$ ,  $\nabla$  and  $\nabla^\perp$  to be the Riemannian connections on  $CaP^2$ ,  $M$  and the normal connection on  $M$  respectively. The corresponding curvature tensors are denoted by  $\bar{R}$ ,  $R$ , and  $R^\perp$  respectively. The first and second covariant derivatives of  $h$  are given by

$$(3.2) \quad (\bar{\nabla}h)(X, Y; Z) = \nabla_z^\perp(h(X, Y)) - h(\nabla_z X, Y) - h(X, \nabla_z Y)$$

$$(3.3) \quad (\bar{\nabla}^2h)(X, Y; Z; W) = \nabla_w^\perp(\bar{\nabla}h)(X, Y; Z) - (\bar{\nabla}h)(\nabla_w X, Y; Z) \\ - (\bar{\nabla}h)(X, \nabla_w Y; Z) - (\bar{\nabla}h)(X, Y; \nabla_w Z)$$

$$X, Y, Z, W \in TM.$$

The Codazzi equation takes the following form

$$(3.4) \quad (\bar{\nabla}h)(X_{\tau(1)}, X_{\tau(2)}; X_{\tau(3)}) = (\bar{\nabla}h)(X_1, X_2; X_3),$$

where  $\tau(i) \in S_3$  the permutation group and the arguments are in the tangent space of  $M$ . Recalling that  $h$  and  $(\bar{\nabla}h)$  are symmetric, we have the Ricci identity

$$(3.5) \quad (\bar{\nabla}^2h)(X, Y; Z; W) - (\bar{\nabla}^2h)(X, Y; W; Z) \\ = -R^\perp(Z, W)h(X, Y) + h(R(Z, W)X, Y) + h(X, R(Z, W)Y).$$

We say that  $j : M \rightarrow CaP^2$  is a totally complex immersion if  $W = j_*(TM)$  is a totally complex subspace for each point of  $M$ . Observe that every totally complex submanifold of  $CaP^2$  has a Kaehler structure. We set  $I = I_0$ , and consequently we have

$$(3.6) \quad \begin{aligned} (a) \quad & \bar{\nabla}_X I = 0 \\ (b) \quad & h(IX, Y) = Ih(X, Y) \\ (c) \quad & A_{IN} = IA_N = -A_N I \\ (d) \quad & IR(X, IX)X = R(X, IX)IX \end{aligned}$$

where  $X, Y \in T_x M$  and  $N \in T_x M^\perp$ .

Define  $f(u) = |h(u, u)|^2$ , where  $u \in UM$ , the unit tangent bundle over  $M$ . Assume  $f$  attains its maximum at some vector  $v \in UM_p$ ,  $p \in M$ , then ([5]):

$$(3.7) \quad A_{h(v,v)}v = |h(v, v)|^2 v.$$

LEMMA 3.1. *Let  $M^n$  be a compact totally complex submanifold in  $CaP^2$ . Assume  $f$  attains its maximum at  $v \in UM_p$ , then*

$$(3.8) \quad 3|h(v, v)|^2(1 - 4|h(v, v)|^2) + \sum_{i=1}^6 \langle h(v, v), I_i v \rangle^2 + 4|(\bar{\nabla}h)(v, v; v)|^2 \leq 0.$$

PROOF. Fix  $v$  in  $UM_p$ . For any  $u \in UM_p$ , let  $r_u(t)$  be the geodesic in  $M$  satisfying the initial conditions  $r_u(0) = p$ ,  $r'_u(0) = u$ . Parallel translating along  $r_u(t)$  gives rise to a vector field  $V_u(t)$ . Put  $f_u(t) = f(V_u(t))$ , then

$$(3.9) \quad \frac{d^2}{dt^2} f_u(0) = 2\langle (\bar{\nabla}^2 h)(v, v; u; u), h(v, v) \rangle + 2|(\bar{\nabla}h)(u, v; v)|^2.$$

Using (3.4), (3.5) and (3.6), we have

$$(3.10) \quad \begin{aligned} \langle (\bar{\nabla}^2 h)(v, v; Iv; Iv), h(v, v) \rangle &= \langle \bar{\nabla}^2 h)(v, Iv; v; Iv), h(v, v) \rangle \\ &= -\langle (\bar{\nabla}^2 h)(v, v; v; v), h(v, v) \rangle \\ &\quad + \langle R^\perp(Iv, v)h(Iv, v), h(v, v) \rangle \\ &\quad - 2\langle R(Iv, v)Iv, A_{h(v, v)}v \rangle. \end{aligned}$$

From the Ricci equation, (2.1), (2.2) and (3.6), we obtain

$$(3.11) \quad \begin{aligned} \langle R^\perp(Iv, v)h(Iv, v), h(v, v) \rangle &= \langle \bar{R}(Iv, v)h(Iv, v), h(v, v) \rangle + \langle [A_{h(Iv, v)}, A_{h(v, v)}]Iv, v \rangle \\ &= -\frac{1}{2}|h(v, v)|^2 - 2|A_{h(v, v)}v|^2 + \frac{1}{2} \sum_{i=1}^6 \langle h(v, v), I_i v \rangle^2 \end{aligned}$$

Now, by the Gauss equation and using (2.1), (2.2) and (3.6) we have

$$(3.12) \quad \langle R(Iv, v)Iv, A_{h(v, v)}v \rangle = -|h(v, v)|^2 + 2|A_{h(v, v)}v|^2.$$

Since  $f$  attains its maximum at  $v$ , we have

$$(3.13) \quad \frac{d^2}{dt^2} f_v(0) + \frac{d^2}{dt^2} f_{Iv}(0) \leq 0.$$

Combining (3.9)–(3.13) and noticing (3.7), we get (3.8).

#### 4. Proof of the Theorem

We will prove the Theorem by showing that under its assumptions the hypothesis that  $M$  is not totally geodesic leads to a contradiction.

From Lemma (3.1) it follows that, by the hypothesis  $h \neq 0$ .

$$(4.1) \quad |h(v, v)|^2 \geq \frac{1}{4}$$

For any  $u \in UM_p$ , let  $r_u(t)$  be the geodesic in  $M$  determined by the initial conditions  $r_u(0) = p$  and  $r'_u(0) = u$ . Parallel translation of  $v$  along  $r_u(t)$  yields a vector field  $V_u(t)$ . Then we know that the function  $f_u$  defined by  $f_u(t) = f(V_u(t))$  attains a maximum at  $t = 0$ . This implies that

$$(4.2) \quad \frac{d^2}{dt^2} f_u(0) + \frac{d^2}{dt^2} f_{Iu}(0) \leq 0.$$

for all  $u \in UM_p$ .

By direct computations we have

$$(4.3) \quad \frac{d^2}{dt^2} f_u(0) = 2\langle (\bar{\nabla}^2 h)(v, v; u, u), h(v, v) \rangle + 2|(\bar{\nabla} h)(u, v; v)|^2$$

Using (3.4), (3.5) and (3.6), we have

$$(4.4) \quad \begin{aligned} \langle (\bar{\nabla}^2 h)(v, v; Iu, Iu), h(v, v) \rangle &= \langle (\bar{\nabla}^2 h)(v, Iv; u, Iu), h(v, v) \rangle \\ &= -\langle (\bar{\nabla}^2 h)(v, Iv; Iu, u), h(v, v) \rangle \\ &\quad + \langle R^\perp(Iu, u)Ih(v, v), h(v, v) \rangle \\ &\quad - 2\langle R(Iu, u)Iv, A_{h(v,v)}v \rangle. \end{aligned}$$

From the Ricci equation, (2.1), (2.2), and (3.6), we obtain

$$(4.5) \quad \begin{aligned} \langle R^\perp(Iu, u)Ih(v, v), h(v, v) \rangle &= \langle \bar{R}(Iu, u)Ih(v, v), h(v, v) \rangle + \langle [A_{h(Iv,v)}, A_{h(v,v)}]Iu, u \rangle \\ &= -\frac{1}{2}|h(v, v)|^2 - 2|A_{h(v,v)}u|^2 + \frac{1}{2} \sum_{i=1}^6 \langle h(v, v), I_i u \rangle^2 \end{aligned}$$

By the Gauss equation, we get

$$(4.6) \quad \langle R(Iu, u)Iv, A_{h(v,v)}v \rangle = -|h(v, v)|^2 \langle R(u, Iu)Iv, v \rangle.$$

From (4.2)–(4.6), we obtain

$$(4.7) \quad 2|h(v, v)|^2 \langle R(u, Iu)Iv, v \rangle - \frac{1}{2}|h(v, v)|^2 - 2|A_{h(v,v)}u|^2 + \sum_{i=1}^6 \langle h(v, v), I_i u \rangle^2 \leq 0.$$

Since  $n = 2$ , we can always choose a unit eigenvector  $u$  of  $A_{h(v,v)}$  such that  $\langle u, v \rangle = \langle u, Iv \rangle = 0$ , using the equation of Gauss which implies that

$$(4.8) \quad R(u, v)v = \frac{1}{4}u + A_{h(v,v)}u - A_{h(u,u)}v$$

$$(4.9) \quad R(u, Iv)Iv = \frac{1}{4}u - A_{h(v,v)}u - A_{h(u,u)}v$$

we have

$$(4.10) \quad A_{h(v,v)}u = \frac{1}{2}(R(u, v)v - R(u, Iv)Iv) = \frac{1}{2}(K(u, v) - K(u, Iv))u$$

where  $K(r, s)$  is the sectional curvature of  $M$  at  $p$  for the plane spanned by  $r, s \in T_pM$ . The Bianchi identity shows that

$$(4.11) \quad \langle R(u, Iv)Iv, v \rangle = K(u, v) + K(u, Iv)$$

From (4.7), (4.10) and (4.11) we obtain

$$(4.12) \quad 2|h(v, v)|^2(K(u, v) + K(u, Iv)) - \frac{1}{2}|h(v, v)|^2 - \frac{1}{2}(K(u, v)^2 + K(u, Iv)^2) - 2K(u, v)K(u, Iv) + \sum_{i=1}^6 \langle h(v, v), I_i u \rangle^2 \leq 0.$$

or equivalently,

$$(4.13) \quad aK(u, v) + bK(u, Iv) - \frac{1}{2}|h(v, v)|^2 + \sum_{i=1}^6 \langle h(v, v), I_i u \rangle^2 \leq 0.$$

where

$$(4.14) \quad a = 2|h(v, v)|^2 - \frac{1}{2}K(u, v) + \frac{1}{2}K(u, Iv)$$

$$(4.15) \quad b = 2|h(v, v)|^2 - \frac{1}{2}K(u, Iv) + \frac{1}{2}K(u, v)$$

Now, we prove that  $a, b > 0$ . From the equation of Gauss it follows that

$$(4.16) \quad K(u, v) + K(u, Iv) = \frac{1}{2} - 2|h(v, v)|^2 \leq \frac{1}{2}$$

By (4.1) and (4.14), we have

$$(4.17) \quad 1 - K(u, v) + K(u, Iv) \leq 2a$$

From (4.16) and (4.17), we know

$$(4.18) \quad 1 + 2K(u, Iv) \leq 2a + \frac{1}{2}$$

Which by the assumption  $K > (1/8)$  implies that  $a > 0$ . In the same way it follows that also  $b > 0$ . Since  $a$  and  $b$  are strictly positive and  $K > (1/8)$ , by (4.13) we get the strictly inequality

$$(4.19) \quad \frac{1}{8}(a + b) - \frac{1}{2}|h(v, v)|^2 + \sum_{i=1}^6 \langle h(v, v), I_i u \rangle^2 < 0$$

But from (4.14) and (4.15) it follows that

$$(4.20) \quad a + b = 4|h(v, v)|^2$$

Which combines with (4.19) yields the desired contradiction. So  $M$  is totally geodesic, by the Theorem 2.2 in [4], we known that  $M = CP^2$ .

ACKNOWLEDGEMENTS. The author is grateful to the referee for a careful reading and very helpful comments on earlier version of the manuscript.

### References

- [1] Besse, A. L., manifolds all of whose geodesic are closed, Springer-Verlag, Berlin, 1978.
- [2] Brown, R. and Gray, A., Riemannian manifolds with holonomy group  $Spin(9)$ , in Diff. Geom. in honor of K. Yano, Tokyo, 1972, pp. 42–59.
- [3] Coulton, P. and Gauchman, H., Submanifold of quaternion projective space with bounded second fundamental forms, Kodai Math. J. **12** (1989), 296–307.
- [4] Goulton, P. and Glazebrook, J., Submanifolds of the Cayley projective plane with bounded second fundamental form, Geom. Dedi. **33** (1990), 265–275.
- [5] Ros, A., Positively curved Kaehler submanifolds, Proc. Amer. Math. Soc. **93** (1985), 329–331.
- [6] Ros, A. and Verstraelen, L., On a conjecture of K. Ogiue, J. Diff. Geom. **19** (1984), 561–565.
- [7] Xia, C. Y., Totally complex submanifolds in  $HP^m(1)$ , Geom. Dedi., **54** (1995), 103–112.

Department of Applied Mathematics  
Dalian University of Technology  
Dalian 116024  
China  
E-mail: xmliu@dlut.edu.cn