

# SINGULAR SETS OF IDEAL INSTANTONS AND POINCARÉ DUALITY

By

Yasuyuki NAGATOMO

**Abstract.** A point in the boundary of a moduli space of quaternion ASD connections can be regarded as a singular ASD connection with a particular singular set. In the case of generalized 1 instantons on  $HP^n$  and  $Gr_2(\mathbf{C}^{n+2})$ , it is proved that Poincaré dual of the singular set is the Chern class of the vector bundle.

## 1. Introduction

The purpose of the present paper is to show that Poincaré dual of the homology class represented by the singular set of an ideal instanton is the even degree Chern class of the vector bundle on which the ideal instanton is defined (Theorem 2.3 and the Table).

It is well known that the moduli space of 1 instantons on the 4 dimensional sphere  $S^4$  is identified with the 5 dimensional open ball. From the viewpoint of Uhlenbeck compactification, a point in the boundary of the moduli space can be considered as an ASD connection with a point singularity. Since the Chern class of the bundle is equal to 1, Poincaré dual of the singular set is the second Chern class  $c_2$ .

The quaternion projective space  $HP^n$  and the complex Grassmannian manifold  $Gr_2(\mathbf{C}^{n+2})$  are quaternion-Kähler manifolds and in particular, in the case  $n = 1$ , these manifolds are  $S^4$  and the complex projective plane  $CP^2$ , respectively. By definition, a quaternion-Kähler manifold is a  $4n$ -dimensional Riemannian manifold for which the linear holonomy group can be reduced to  $Sp(1) \cdot Sp(n)$ . Anti-self-duality can be defined over quaternion-Kähler manifolds in the same way as in the 4-dimensional case (see, for example, Mamone Capria and Salamon [3]).

From this point of view, 1 instantons on  $S^4$  and  $CP^2$  are generalized to objects on  $HP^n$  and  $Gr_2(\mathbf{C}^{n+2})$  (Definition 2.1). The moduli spaces of these

generalized 1 instantons can be described by the theory of monads on the Salamon twistor space (see §2). This theory indicates a natural compactification of moduli spaces and a point in the boundary of the moduli can also be regarded as an ASD connection which is defined only on the complementary set of a closed subset of the base manifold. We call such a closed subset the singular set. In the case of generalized 1 instantons, the singular sets are described in Theorem 2.2, and all of them are quaternion submanifolds.

As for the method of our proof, see §3. In particular, we use holomorphic vector bundles and sections of them on the twistor spaces. All the holomorphic vector bundles stated here have common properties in terms of ASD bundles and line bundles on the twistor spaces. These properties will be formulated in the forthcoming paper.

The author would like to express his sincere gratitude to Professor Simon Salamon for many valuable comments, including computation of the Chern classes. The author is a Monbusho-sponsored Japanese Overseas Reserch Fellow and would also like to thank the Ministry of Education and Culture for the support.

## 2. Preliminaries

The symmetric spaces  $HP^n$  and  $Gr_2(\mathbf{C}^{n+2})$  are quaternion-Kähler manifolds with positive scalar curvature. A connection on a vector bundle on a quaternion-Kähler manifold is called an *ASD connection* if its curvature 2-form is invariant under the action of  $I, J, K$  (see, for example, [3]). A vector bundle with an ASD connection is also called *instanton* (bundle). On the other hand, every quaternion-Kähler manifold has a twistor space with a natural complex structure [9]. The twistor spaces of  $HP^n$  and  $Gr_2(\mathbf{C}^{n+2})$  are the odd dimensional complex projective spaces  $CP^{2n+1}$  and the generalized flag manifolds  $F^{2n+1} = SU(n+2)/S(U(1) \times U(n) \times U(1))$ , respectively. The pull-back bundle with ASD connection on the twistor space has a holomorphic structure induced by the pull-back connection. Hence we do not distinguish ASD bundles on quaternion-Kähler manifolds from the pull-back bundles on the twistor space, and we use the same symbol for both.

To specify vector bundles, we need to describe the cohomology rings on the twistor spaces. As for the complex projective spaces,  $x$  is defined as the standard positive generator of  $H^2(CP^{2n+1}, \mathbf{Z})$ . Since the twistor space  $F^{2n+1}$  can be expressed as the projective bundle of the holomorphic cotangent bundle of  $CP^{n+1}$ , the Leray-Hirsch theorem implies that the cohomology ring  $H^*(F^{2n+1}, \mathbf{Z})$  is isomorphic to the quotient ring of  $\mathbf{Z}[x, y]$  by the ideal generated by  $x^{n+2}$  and

$x^{n+1} - x^n y + \dots + (-1)^{n+1} y^{n+1}$ , where  $x$  is the pull-back of the standard positive generator of  $H^2(\mathbf{C}P^{n+1}, \mathbf{Z})$ . The twistor space  $F^{2n+1}$  has another fibration over the dual complex projective space  $\mathbf{C}P^{n+1*}$ . Then  $y$  is the pull-back of the standard positive generator of  $H^2(\mathbf{C}P^{n+1*}, \mathbf{Z})$ .

We now define the ASD bundles which we wish to consider in the present paper.

**DEFINITION 2.1.** (0) Let  $E_0$  be an ASD bundle on  $\mathbf{H}P^n$  with structure group  $Sp(n)$ . The total Chern class  $c(E_0)$  of  $E_0$  is assumed to be equal to  $(1 - x^2)^{-1}$ . More precisely, the Chern classes are expressed as  $c_2(E_0) = x^2$ ,  $c_4(E_0) = x^4, \dots, c_{2n}(E_0) = x^{2n}$ .

(1) Let  $E_1$  be an ASD bundle on  $Gr_2(\mathbf{C}^{n+2})$  with structure group  $Sp(n)$ . The total Chern class  $c(E_1)$  of  $E_1$  is assumed to be equal to  $(1 - x^2)^{-1}(1 - y^2)^{-1}$ . More precisely, the Chern classes are expressed as  $c_{2i}(E_1) = \sum_{k=0}^i x^{2i-2k} y^{2k}$ , where  $i = 1, 2, \dots, n$ .

(2) Let  $E_2$  be an ASD bundle on  $Gr_2(\mathbf{C}^{n+2})$  with structure group  $SU(n+1)$ . The total Chern class  $c(E_2)$  of  $E_2$  is assumed to be equal to  $\{1 + (-x + y)\} \cdot (1 - x)^{-1}(1 + y)^{-1}$ . More precisely, the Chern classes are expressed as  $c_i(E_2) = -\sum_{k=1}^{i-1} x^{i-k} (-y)^k$ , where  $i = 1, 2, \dots, n+1$ .

**REMARK.** In the case  $n = 1$ ,  $E_0$  is nothing but a 1 instanton on  $S^4$  and  $E_1$  and  $E_2$  are isomorphic to each other, because  $Sp(1) \cong SU(2)$ , and our relations for generators of the cohomology yield that  $c(E_1) = c(E_2)$ . The last two bundles are 1 instantons on  $\mathbf{C}P^2$ .

Let  $\mathcal{M}_0, \mathcal{M}_1$  and  $\mathcal{M}_2$  be the moduli spaces of ASD connections on  $E_0, E_1$  and  $E_2$ , respectively. From the viewpoint of [6],

(0)  $\mathcal{M}_0$  is identified with an open ball in  $\wedge^2 \mathbf{C}^{2n+2}$ , where  $\wedge^2 \mathbf{C}^{2n+2}$  is the corresponding representation of  $Sp(n+1)$ .

(1)  $\mathcal{M}_1$  is identified with an open cone over  $\mathbf{P}(\wedge^2 \mathbf{C}^{n+2})$ , where  $\wedge^2 \mathbf{C}^{n+2}$  is the corresponding representation of  $SU(n+2)$ .

(2)  $\mathcal{M}_2$  is identified with an open cone over  $\mathbf{P}(\mathbf{C}^{n+2})$ , where  $\mathbf{C}^{n+2}$  is the standard representation space of  $SU(n+2)$ .

The completeness of these moduli spaces is proved in [1], [2], [8] and [4]. In particular, the theory of monads is available for proving the completeness of the moduli ([3], [2], [8] and [4]). From the viewpoint of the theory of monads, the boundary point of the moduli spaces represents a singular ASD connection with a singular set, which we denote by  $S$ . These singular sets are described in [1], [7] and [4].

**THEOREM 2.2.** *In each case, the singular set is one of the following quaternion submanifolds of  $\mathbf{HP}^n$  or  $Gr_2(\mathbf{C}^{n+2})$ :*

- (0)  $\mathbf{HP}^i$ , where  $i = 0, 1, \dots, n + 1$ ,
- (1)  $\mathbf{HP}^i$ , where  $i = 0, 1, \dots, [(n + 2)/2]$  and
- (2)  $Gr_2(\mathbf{C}^{n+1})$ ,

where  $[m]$  is the greatest integer not greater than  $m$ .

**THEOREM 2.3 (Main Theorem).** *The Poincaré dual of the homology class represented by the (real)  $4(n - i)$ -dimensional singular set is the 2ith Chern class of the ASD bundle on which the singular ASD connection is defined.*

• Table (Singular sets)

Bundles	Singular sets	Poincaré dual
$E_0$	1 point, $\mathbf{HP}^1, \dots, \mathbf{HP}^{n-1}$	$c_{2n}(E_0), c_{2n-2}(E_0), \dots, c_2(E_0)$
$E_1$	1 point, $\mathbf{HP}^1, \dots, \mathbf{HP}^{[n/2]}$	$c_{2n}(E_1), c_{2n-2}(E_1), \dots,$ $c_n(E_1)(n: \text{even}), c_{n+1}(E_1)(n: \text{odd})$
$E_2$	$Gr_2(\mathbf{C}^{n+1})$	$c_2(E_2)$

### 3. Zero Loci of Sections

The following two sections are devoted to the proof of the Main Theorem. Since our strategy is common to all three cases, we shall now describe it.

First of all, we determine a holomorphic vector bundle  $V$  on the twistor space which has a (holomorphic) section  $s$ . It is shown that the zero locus of the section  $s$  is the inverse image  $\tilde{S}$  by the twistor fibration of the singular set  $S$  and is transverse to the zero section. Hence, Poincaré dual of  $\tilde{S}$  equals the top Chern class of  $V$ .

Next, it is proved that the top Chern class of  $V$  equals the pull-back of the appropriate Chern class of the ASD bundle.

Finally, the Gysin sequence or fibre integration yields the desired results and we omit the details of this procedure.

3.1. THE CASE OF  $E_0$ . We may choose  $V$  to be the direct sum of line bundles of degree 1. To put it more accurately, we denote by  $[z_0; z_1; \dots; z_{2n+1}]$  the homogeneous coordinates on  $\mathbf{CP}^{2n+1}$  and by  $\mathcal{O}(d)$  the line bundle of degree  $d$  on  $\mathbf{CP}^{2n+1}$ . Now  $V$  is the direct sum  $\mathcal{O}(1)^{\oplus 2i}$ , the top Chern class of  $V$  is  $x^{2i}$ , and so it equals  $c_{2i}(E_0)$ . The section  $s$  of  $V$  which we should choose is  $(z_{2(n-i)+1}, z_{2(n-i)+2}, \dots, z_{2n}, z_{2n+1})$ .

3.2. THE CASE OF  $E_1$  AND  $E_2$ . Since  $F^{2n+1}$  is a Fano manifold, the Picard group is  $H^2(F^{2n+1}, \mathbf{Z})$ . We denote by  $\mathcal{O}(p, q)$  the line bundle of which the first Chern class is  $px + qy$ . As mentioned in the previous section,  $F^{2n+1}$  has two holomorphic fibrations:

$$p_1 : F^{2n+1} \rightarrow \mathbf{C}P^{n+1} \quad \text{and} \quad p_2 : F^{2n+1} \rightarrow \mathbf{C}P^{n+1*}.$$

From our definition of the cohomology classes  $x$  and  $y$ , we obtain  $p_1^*\mathcal{O}(1) \cong \mathcal{O}(1, 0)$  and  $p_2^*\mathcal{O}(1) \cong \mathcal{O}(0, 1)$ . For brevity, we fix a unitary basis  $\{e_1, \dots, e_{n+2}\}$  of the standard representation space  $\mathbf{C}^{n+2}$  of  $SU(n+2)$  and the dual basis  $\{e^1, \dots, e^{n+2}\}$  of  $\mathbf{C}^{n+2*}$ . The corresponding homogeneous coordinates on  $\mathbf{C}P^{n+1}$  and  $\mathbf{C}P^{n+1*}$  are denoted by  $[z_1; \dots; z_{n+2}]$  and  $[w_1; \dots; w_{n+2}]$ , respectively. Then  $F^{2n+1}$  can be identified with the divisor on  $\mathbf{C}P^{n+1} \times \mathbf{C}P^{n+1*}$  defined by the equation  $\sum_{i=1}^{n+2} z_i w_i = 0$ .

Let  $t$  be a holomorphic section of  $\mathcal{O}(1, 0)$ . If  $t$  is the pull-back section of  $\mathcal{O}(1)$  on  $\mathbf{C}P^{n+1}$  which corresponds to  $\sum a^i z_i$ , we shall say that  $t$  is the section corresponding to  $\sum a^i z_i$ . We also use a similar terminology for a section of  $\mathcal{O}(0, 1)$ .

LEMMA 3.1. *Let  $s$  be a section of the direct sum  $\mathcal{O}(1, 0) \oplus \mathcal{O}(0, 1)$ . If  $s$  corresponds to  $(z_{n+2}, w_{n+2})$ , then the zero locus of  $s$  is  $F^{2n-1}$  which is the twistor space of the quaternion submanifold  $Gr_2(\mathbf{C}^{n+1})$  of  $Gr_2(\mathbf{C}^{n+2})$ .*

PROOF. Using our notation, we define divisors  $\mathbf{C}P^n = \{z_{n+2} = 0\}$  and  $\mathbf{C}P^{n*} = \{w_{n+2} = 0\}$  on  $\mathbf{C}P^{n+1}$  and  $\mathbf{C}P^{n+1*}$ , respectively. Then we have  $s^{-1}(0) = p_1^{-1}(\mathbf{C}P^n) \cap p_2^{-1}(\mathbf{C}P^{n*})$ .  $\square$

In this case, it is clear that  $s^{-1}(0)$  is transverse to the zero section. The second Chern class of  $\mathcal{O}(1, 0) \oplus \mathcal{O}(0, 1)$  is  $xy$ , and so it is equal to  $c_2(E_2)$ . Thereby, in the case of  $E_2$ , the proof is completed.

Pulling back the Euler sequence on  $\mathbf{C}P^{n+1*}$ , we obtain the exact sequence:

$$0 \rightarrow \mathcal{O}(1, -1) \rightarrow \mathcal{O}(1, 0)^{\oplus n+2} \rightarrow \mathcal{Q} \rightarrow 0.$$

Using weights for example, we also obtain the exact sequence:

$$0 \rightarrow V \rightarrow \mathcal{Q} \rightarrow \mathcal{O}(2, 0) \rightarrow 0.$$

(The vector bundle  $V$  is the “twisted tautological bundle”.) By the Bott-Borel-Weil theorem,  $H^0(F^{2n+1}, V)$  is identified with  $\wedge^2 \mathbf{C}^{n+2*}$  as representations of  $SU(n+2)$ .

For brevity, we replace  $SU(n+2)$  by  $G$  and the isotropy subgroup  $S(U(1) \times U(n) \times U(1))$  by  $K_Z$ . As a  $C^\infty$  vector bundle,  $V$  is identified with the associated vector bundle with  $G$  of which the typical fibre is the representation space  $Ce^1 \otimes C^{n^*}$  of  $K_Z$ , where  $C^{n^*}$  is spanned by  $e^2, \dots, e^{n+1}$ . We denote the orthogonal projection by  $\pi: \wedge^2 C^{n+2^*} \rightarrow Ce^1 \otimes C^{n^*}$ , which is explicitly given by

$$\pi(\phi) = e^1 \wedge \iota(e_1)\phi - h(\phi, e^1 \wedge e^{n+2})e^1 \wedge e^{n+2},$$

where  $\phi \in \wedge^2 C^{n+2^*}$ ,  $\iota$  is the interior product and  $h$  is the induced Hermitian inner product on  $\wedge^2 C^{n+2^*}$ . Using  $\pi$ , the isomorphism between  $\wedge^2 C^{n+2^*}$  and  $H^0(F^{2n+1}, V)$  is expressed as

$$s_\phi[g] = [g, \pi(g^{-1}\phi)],$$

where  $[g]$  is the point of  $F^{2n+1}$  represented by  $g \in G$ , and  $[g, \pi(g^{-1}\phi)]$  is the element of  $V$  represented by  $(g, \pi(g^{-1}\phi))$ .

**PROPOSITION 3.2.** *We assume that  $n$  is even. If  $\phi \in \wedge^2 C^{n+2^*}$  is non-degenerate, then the zero locus of  $s_\phi$  is expressed as*

$$s_\phi^{-1}(0) = \{([v], [\iota(v)\phi]) \in CP^{n+1} \times CP^{n+1^*}\} \cong CP^{n+1}.$$

**PROOF.** With our notation,  $s_\phi^{-1}(0) = \{[g] \in F^{2n+1} \mid \pi(g^{-1}\phi) = 0\}$ . From the definition of  $\pi$ ,  $\pi(g^{-1}\phi) = 0$  if and only if  $\phi(ge_l, ge_l) = 0$  for an arbitrary  $l = 2, \dots, n+1$  and so, there exists a non-zero constant  $\alpha$  such that  $\iota(g^{-1}e_1)\phi = \alpha ge^{n+2}$ , because  $\phi$  is non-degenerate.  $\square$

If  $n$  is odd, then any  $\phi \in \wedge^2 C^{n+2^*}$  is degenerate, but in the generic case, the rank of  $\phi$  is equal to  $n+1$ . Note that if  $\phi$  is of rank  $n+1$ , then there exists a non-zero vector  $v_0 \in C^{n+2}$  such that  $\iota(v_0)\phi = 0$ .

**PROPOSITION 3.3.** *We assume that  $n$  is odd. If  $\phi \in \wedge^2 C^{n+2^*}$  is of rank  $n+1$  and satisfies  $\iota(v_0)\phi = 0$  for non-zero vector  $v_0 \in C^{n+2}$ , then the zero locus of  $s_\phi$  is isomorphic to the blow up of  $CP^{n+1}$  at one point. More explicitly, the zero locus is expressed as*

$$\begin{aligned} s_\phi^{-1}(0) = & \{([v], [\iota(v)\phi]) \in CP^{n+1} \times CP^{n+1^*} \mid [v] \neq [v_0]\} \\ & \cup \{([v_0], [\psi]) \in CP^{n+1} \times CP^{n+1^*} \mid \psi(v_0) = 0\} \end{aligned}$$

**PROOF.** Under the same notation as in the proof of Proposition 3.2, we

obtain the same condition  $\phi(ge_1, ge_l) = 0$  for an arbitrary  $l = 2, \dots, n+1$ . This implies the explicit expression of the zero locus in the Proposition.

To see that this set is identified with the blow up of  $\mathbf{C}P^{n+1}$  at  $[v_0]$ , we may assume that  $v_0 = e_{n+2}$  and  $\phi = e^1 \wedge e^2 + e^3 \wedge e^4 + \dots + e^n \wedge e^{n+1}$ . Under the assumption  $z_{n+2} \neq 0$ , we introduce the inhomogeneous coordinate  $\xi_m = z_m/z_{n+2}$  for  $m = 1, \dots, n+1$ . After a ‘‘compatible’’ transformation on  $\mathbf{C}P^{n+1}$  ( $e_1 \mapsto e_2, e_2 \mapsto -e_1, \dots, e_{n+1} \mapsto -e_n$ , and  $e^{n+2} \mapsto e^{n+2}$ ) related to  $\phi$ , on a neighbourhood of  $([v_0], [e^1]) \in F^{2n+1} \subset \mathbf{C}P^{n+1} \times \mathbf{C}P^{n+1*}$ , the zero locus can be expressed as  $([\xi_1; \dots; \xi_{n+1}; 1], [w_1; \dots; w_{n+1}; 0])$ , where there exists  $\lambda \in \mathbf{C}$  such that  $(\xi_1, \dots, \xi_{n+1}) = \lambda(w_1, \dots, w_{n+1})$ .  $\square$

**LEMMA 3.4.** *If  $\phi \in \wedge^2 \mathbf{C}^{n+2*}$  is of full rank, then the corresponding section  $s_\phi \in H^0(F^{2n+1}, V)$  is transverse to the zero section.*

**PROOF.** It is sufficient to show that there exists  $X \in \mathfrak{su}(n+2)$  such that  $\phi(Xge_1, ge_l) + \phi(ge_1, Xge_l) \neq 0$  for an arbitrary  $l = 2, \dots, n+1$  under the assumption  $\phi(ge_1, ge_l) = 0$  for an arbitrary  $l = 2, \dots, n+1$ . In the case that  $n$  is even, we may choose  $X_l$  such that  $X_l ge_1 = \sqrt{-1}ge_1$  and  $X_l ge_l = ge_{n+2}$ .

If  $n$  is odd and  $ge_1 = v_0$ , then  $X_l$  may be chosen such that  $\phi(X_l ge_1, ge_l) \neq 0$ , because  $\phi$  is non-degenerate on the orthogonal complement of  $\mathbf{C}v_0$  in  $\mathbf{C}^{n+2}$ .  $\square$

**PROPOSITION 3.5.** *We now assume that  $n$  is even. If  $\phi \in \wedge^2 \mathbf{C}^{n+2*}$  defines a compatible quaternion structure with the Hermitian inner product  $h$  on  $\mathbf{C}^{n+2}$ , then the zero locus of the section  $s_\phi$  of  $V$  is the twistor space of a quaternion submanifold  $\mathbf{H}P^{(n+2)/2}$  of  $Gr_2(\mathbf{C}^{n+2})$ .*

**PROOF.** From the assumption, there exists a unitary basis  $\{e_1, \dots, e_{n+2}\}$  of  $\mathbf{C}^{n+2}$  such that  $\phi = e^1 \wedge e^2 + e^3 \wedge e^4 + \dots + e^{n+1} \wedge e^{n+2}$ . The corresponding quaternion structure is denoted by  $j$ .

We denote by  $\sigma_F$  the real structure of  $F^{2n+1}$  [9]. If  $([v], [l(v)\phi])$  is in the zero locus of the section  $s_\phi$ , we obtain  $\sigma_F([v], [l(v)\phi]) = ([-jv], [l(jv)\phi])$ . This means that the real structure  $\sigma_F$  can be restricted to the zero locus and corresponds to the real structure of  $\mathbf{C}P^{n+1}$  as the twistor space of  $\mathbf{H}P^{(n+2)/2}$ .  $\square$

In the case that  $n$  is odd, we define a vector bundle  $V'$  as  $V' = V \oplus \mathcal{O}(1, 0)$ . Using Lemma 3.1 and Proposition 3.3, we can find a section whose zero locus is holomorphically isomorphic to  $\mathbf{C}P^n$ . Then, the following proposition is obtained in a similar way.

**PROPOSITION 3.6.** *In the case that  $n$  is odd, it is assumed that  $\phi \in \wedge^2 \mathbf{C}^{n+2^*}$  defines a compatible quaternion structure with the Hermitian inner product  $h$  on  $\mathbf{C}^{n+1}$ , where  $\mathbf{C}^{n+1}$  is the orthogonal complement of  $\mathbf{C}v_0$  and  $v_0 \in \mathbf{C}^{n+1} \setminus \{0\}$  satisfies  $\iota(v_0)\phi = 0$ . The corresponding section  $(s_\phi, s_{v^0})$  of  $V'$  is defined in the following way. The section  $s_\phi$  is the corresponding section of  $V$ . The linear form  $v^0$  is defined as  $\text{Ker } v^0 = \mathbf{C}^{n+1}$  and  $v^0(v_0) = 1$ . The section  $s_{v^0}$  is the section of  $\mathcal{O}(1, 0)$  corresponding to  $v^0$ .*

*Then the zero locus of the section  $(s_\phi, s_{v^0})$  of  $V'$  is the twistor space of a quaternion submanifold  $\mathbf{HP}^{(n+1)/2}$  of  $Gr_2(\mathbf{C}^{n+2})$ .*

Using Lemma 3.1 and Propositions 3.5 and 3.6, we know that the vector bundles  $V \oplus (\mathcal{O}(1, 0) \oplus \mathcal{O}(0, 1))^{\oplus k}$  has a section of which the zero locus is the twistor space of  $\mathbf{HP}^{(n+2)/2-k}$  in the case that  $n$  is even and the vector bundle  $V' \oplus (\mathcal{O}(1, 0) \oplus \mathcal{O}(0, 1))^{\oplus k}$  has a section whose zero locus is the twistor space of  $\mathbf{HP}^{(n+1)/2-k}$  in the case that  $n$  is odd. Lemma 3.4 yields that each section is transverse to the zero section.

#### 4. The Chern Classes

Under our notation,  $V$  is isomorphic to  $G \times_{K_Z} \mathbf{C}e^1 \otimes \mathbf{C}^{n^*}$ . Theorem 3.4 in [6] assures that the vector bundle  $G \times_{K_Z} \mathbf{C}^{n^*}$  is isomorphic to the pull-back of an instanton bundle  $E$  on  $Gr_2(\mathbf{C}^{n+2})$ . The pull-back of  $E$  can be obtained by the monad [5]:

$$\mathcal{O}(0, -1) \rightarrow \underline{\mathbf{C}^{n+2^*}} \rightarrow \mathcal{O}(1, 0).$$

Hence we have  $c_p(E) = \sum_{i=0}^p (-x)^i y^{p-i}$ .

**LEMMA 4.1.** *The top Chern class  $c_n(V)$  of  $V$  on  $F^{2n+1}$  equals  $\sum_{p=0}^{\lfloor n/2 \rfloor} x^{2p} y^{n-2p}$ .*

**PROOF.** Since  $V \cong E \otimes \mathcal{O}(1, 0)$  and  $c_1(\mathcal{O}(1, 0)) = x$ , it follows that  $c_n(V) = \sum_{p=0}^n c_p(E)x^{n-p} = \sum_{p=0}^n \sum_{i=0}^p (-1)^i x^{n-p+i} y^{p-i}$ . Consequently, we obtain  $c_{n+1}(V^{n+1}) = c_n(V)x + \sum_{i=0}^{n+1} (-x)^i y^{n+1-i}$ , where  $V^{n+1}$  is defined on  $F^{2(n+1)+1}$ . The induction yields our result.  $\square$

**4.1.  $n$ : even.** The relation  $x^{n+1} - x^n y + \dots + (-1)^{n+1} y^{n+1}$ , the definition of  $E_1$  and Lemma 4.1 imply that  $c_n(V) = c_n(E_1)$  and  $x c_n(V) = y c_n(V)$ . On the other hand,

$$c_{n+2k}(V \oplus (\mathcal{O}(1, 0) \oplus \mathcal{O}(0, 1))^{\oplus k}) = x^k y^k c_n(V),$$



$$c_{n+2k}(E_1) = c_n(E_1)y^{2k} + x^{2m} \sum_{q=1}^k x^{2q}y^{2k-2q}.$$

The relation  $x^{n+2} = 0$  yields  $c_{n+2k}(E_1) = c_n(E_1)y^{2k} = c_n(V)y^{2k} = x^k y^k c_n(V)$ .  
 Thereby,  $c_{n+2k}(V \oplus (\mathcal{O}(1, 0) \oplus \mathcal{O}(0, 1))^{\oplus k}) = c_{n+2k}(E_1)$ .

4.2.  **$n$ : odd.** Lemma 4.1 yields  $c_{n+1}(V') = xy \sum_{p=0}^{(n-1)/2} x^{2p}y^{n-1-2p}$ . This equation, the relation  $x^{n+1} - x^n y + \dots + (-1)^{n+1} y^{n+1}$  and the definition of  $E_1$  imply  $c_{n+1}(V') = c_{n+1}(E_1)$ . The relation  $y^{n+2} = 0$  yields  $yc_{n+1}(E_1) = xc_{n+1}(V') = xc_{n+1}(E_1)$  and  $c_{n+1+2k}(E_1) = x^{2k}c_{n+1}(E_1)$ . Hence

$$\begin{aligned} c_{n+1+2k}(V' \oplus (\mathcal{O}(1, 0) \oplus \mathcal{O}(0, 1))^{\oplus k}) &= x^k y^k c_{n+1}(V') = x^{2k} c_{n+1}(E_1) \\ &= c_{n+1+2k}(E_1). \end{aligned}$$

## References

- [1] H. Doi and T. Okai, Moduli space of 1-instantons on a quaternionic projective space  $HP^n$ , Hiroshima. Math. J. **19** (1989), 251–258
- [2] Y. Kametani and Y. Nagatomo, Construction of  $c_2$ -self-dual Bundles on a Quaternionic Projective space, Osaka. J. Math. **32** (1995), 1023–1033
- [3] M. Mamone Capria and S. M. Salamon, Yang-Mills fields on quaternionic spaces, Nonlinearity **1** (1988), 517–530
- [4] Y. Nagatomo, Another type of of instanton bundles on  $Gr_2(\mathbb{C}^{n+2})$ , Tokyo J. Math. **21** (1998), 267–297
- [5] Y. Nagatomo, Examples of vector bundles admitting unique ASD connections on quaternion-Kähler manifolds, Proc. Amer. Math. Soc. **127** (1999), 3043–3048
- [6] Y. Nagatomo, Representation theory and ADHM-construction on quaternion symmetric spaces, Trans. Amer. Math. Soc. **353** (2001), 4333–4355
- [7] Y. Nagatomo and T. Nitta, Moduli of 1-instantons on  $G_2(\mathbb{C}^{n+2})$ , Differential Geom. Appl. **7** (1997), 115–122
- [8] Y. Nagatomo and T. Nitta,  $k$ -instantons on  $G_2(\mathbb{C}^{n+2})$  and stable vector bundles, Math. Z **232** (1999), 721–737
- [9] S. M. Salamon, Quaternionic Kähler Manifolds, Invent. Math. **67** (1982), 143–171

Institute of Mathematics  
 University of Tsukuba  
 Tsukuba-shi, Ibaraki 305, Japan

Mathematical Institute, 24-29 St. Giles'  
 Oxford, OX1 3LB, UK

Current Address: Faculty of Mathematics  
 Kyushu University, Ropponmatsu, Fukuoka, 810-8560, Japan  
*E-mail address:* nagatomo@math.kyushu-u.ac.jp