


DB  
290  
1985  
④ C:415.2



DIMENSION THEORY OF GENERAL SPACES

BY

KÔICHI TSUDA

A thesis submitted in partial fulfillment of the  
requirement for the degree of

DOCTOR OF SCIENCE

(*Mathematics*)

at the

UNIVERSITY OF TSUKUBA

SEPTEMBER, 1985

86318859

# DIMENSION THEORY OF GENERAL SPACES

BY

Kôichi Tsuda

- Chap. 0 Introduction
- Chap. 1 Closed embeddings of complete metric spaces
- Chap. 2 Non-existence of universal spaces for some stratifiable spaces
- Chap. 3 Normal spaces  $X \times Y$  satisfying the inequality  $\dim (X \times Y) > \dim X + \dim Y$

Summary. We shall solve the following three problems, which come from the attempt to develop a harmonious dimension theory:

- (1) Find a non-separable complete metric space which plays a role of the space of real line.
- (2) Find universal spaces for generalized metric spaces within the class of stratifiable spaces.
- (3) Find strongly zero dimensional spaces  $X$  and  $Y$  such that dimension  $\dim (X \times Y)$  of the product is arbitrary high.

## CONVENTIONS.

The Greek letters  $\alpha, \beta$  will indicate ordinal numbers, and each ordinal will be taken to be the set of its predecessors. Thus,  $\alpha \in \beta$  and  $\alpha < \beta$  means the same thing. These same Greek letters will also stand for cardinal numbers.  $\omega$  is the first infinite ordinal, and  $\omega_1$  is the first uncountable ordinal. We also use  $\omega$  to denote the set of positive integers. No difficulty will result from these ambiguity. The cardinality of continuum is sometimes denoted by  $c$ .

A sequence  $\{A_i\}$  of sets is called decreasing if  $A_i \supset A_{i+1}$  for each  $i$ .

All spaces considered are assumed to be Tychonoff, and all maps are continuous. The dimension  $\dim X$  of a normal space  $X$  means the covering dimension of it, which goes back to the idea of Čech [1] and Lebesgue [1]. (When  $X$  is non-normal, we adapt the dimension function due to Katětov [1].) In particular, we say that  $X$  is strongly 0-dimensional when  $\dim X = 0$ .

We shall use several different topologies  $\tau$  on the Cantor-set  $C$ , and we denote it by  $(C, \tau)$ . In particular, by  $(C, \varepsilon_0)$  we denote the usual Euclidean topology on  $C$ . For a topology  $\tau$  on  $C$  and for a subset  $U$  of  $C$ , we say sometimes that  $U$  is  $\tau$ -open if  $U$  is open in  $(C, \tau)$ .

We say that  $U$  is clopen if it is closed and open simultaneously. For a subset  $A$  of  $(C, \tau)$   $\tau|A$  denotes its relative topology. The unit interval with the Euclidean topology is denoted by  $I$  (the letter  $\varepsilon$  stands for its topology).  $\text{Exp}(X)$  denotes the hyperspace consisting of all closed sets of a topological space  $X$  with the Vietoris topology.

For the undefined terminology, we refer to the reader to Engelking [1] and Kunen [1].

## Introduction.

It is difficult to determine the origin of the concept of dimension (see Duda [1]). But, it is no doubt to believe that the discovery by Peano [1] (that is, a continuous map on the unit interval, the image of which is the full square) reminded the people that it is necessary to define the concept of dimension precisely.

Nowadays, we have some satisfactory dimension functions for a space  $X$ ;  $\text{ind } X$ ,  $\text{Ind } X$ , and  $\text{dim } X$ .

One of the most remarkable facts concerning their definitions is that we can assign dimensions to fairly general spaces. For example, we can define the above three dimension functions for every (not necessarily, metrizable) normal spaces.

On the other hand, we believe that our dimension functions are satisfactory, since for separable metric spaces we have a beautiful harmonious dimension theory (see Hurewicz and Wallman [1]).

Hence, one of the main problems in dimension theory is to construct a harmonious dimension theory for more general spaces.

From this point of view we have remarkable developments for metrizable spaces (see Morita [2] and Nagata [7]), and have important results for some stratifiable spaces (Nagami [6,7] and Oka [1]).

There are, however, many theorems, which are valid for separable metric spaces, but are open for more general (even, complete metric) spaces in dimension theory. In this thesis we shall discuss three such theorems, and shall solve them completely.

(1) There is no doubt to believe that the real line  $R$  is one of the most important and useful 1-dimensional spaces. Our first problem is to seek a non-separable complete metric space, which plays a role of the space  $R$ .

To specify our problem, we quickly review the strongly 0-dimensional case. For the strongly 0-dimensional spaces we believe that we have a satisfactory one. It is the Baire's 0-dimensional space. The reason, why it is, is that the Baire's 0-dimensional space  $B(\alpha)$  of weight  $\alpha$  satisfies not only that  $B(\omega)$  (that is, the separable case) is homeomorphic to the familiar space, "the space of irrational numbers", but also that

(1) every strongly 0-dimensional complete metric space of weight  $\alpha$  is homeomorphic to a closed subset of  $B(\alpha)$  (Stone [1]).

Closed embeddings are frequently used in topology (for example, it is indispensable for ANR theory). In particular, the above result (1) is essentially used

in the recent work of Jayne and Rogers [1] to obtain their representation theorem.

On the other hand, it is known that every finite dimensional separable locally compact space can be closed-embedded in the finite iterated product of  $\mathbb{R}$  (Isbell [1]).

(Note that we cannot embed an  $n$ -dimensional space in the same dimensional product  $\mathbb{R}^n$  in general. We can, however, embed it into the  $2n+1$  dimensional product  $\mathbb{R}^{2n+1}$ .)

Therefore, we can specify our problem as follows:

PROBLEM. For a given infinite cardinal number  $\alpha$  find a 1-dimensional complete metric space  $Z_\alpha$  of weight  $\alpha$  such that every 1-dimensional complete metric space of weight  $\alpha$  can be closed-embedded in a finite iterated product of  $Z_\alpha$ .

In chapter 1 we can obtain more general results together with the solution of this problem:

THEOREM 1.2. For a given infinite cardinal number  $\alpha$  and every non-negative integer  $n$  there exists an  $n$ -dimensional complete metric space  $Z_{n,\alpha}$  of weight  $\alpha$  such that every  $n$ -dimensional complete metric space of weight  $\alpha$  can be closed-embedded in the product  $(Z_{n,\alpha})^2$ .

(2) It is one of the most important theorems in dimension theory that there exist universal spaces for several subclasses  $\mathcal{C}$  of separable metric spaces. (A space  $X$  is called a universal space for a class  $\mathcal{C}$  if

- (a)  $X$  is a member of  $\mathcal{C}$  ;
- (b) every member of  $\mathcal{C}$  is homeomorphic to a subset of  $X$ .)

For example, the countable infinite product of the real lines  $\mathbb{R}^\omega$  is a universal space for all separable metric spaces. It is also known that there are universal spaces for several (non-separable) metric spaces (Nagata [7]).

One of the important properties of universal spaces is that it reduces a general problem to a special problem for a subset of a particular space. (That is, when we want to prove a problem for every separable metric space, it suffices to show it for a subset of  $\mathbb{R}^\omega$ . Hence, we can use some special properties of  $\mathbb{R}^\omega$  for this problem.)

We believe that one of the reasons, why we cannot complete a harmonious dimension theory for stratifiable spaces, is that it was not known that whether or not there exist universal spaces.

Hence, we specify our problem as follows:



PROBLEM. For every infinite cardinal number  $\alpha$  find a universal space for strongly 0-dimensional stratifiable spaces of netweight  $\alpha$ .

(Note that the netweight of a space  $X$  is equal to its weight when  $X$  is metrizable, and that it is known that when we classify stratifiable spaces, it is convenient to use its netweight instead of its weight.)

In contrast with the known results for metric spaces we shall give a complete negative solution to this problem in chapter 2 (Theorem 2.1). We shall also show that there are none for almost all of known important subclasses of stratifiable spaces. (When we use their weight instead of their netweight, we obtain similar results in Corollary 2.1.)

Therefore, our results show that there are unexpected deep gaps between the class of metric spaces and that of stratifiable spaces.

(3) The third problem is concerned with the inequality

$$(*) \dim (X \times Y) \leq \dim X + \dim Y,$$

which is called the product theorem in dimension theory.

It is known that the equality in the above formula is not valid in general. Namely, it is Pontrjagin [1] who constructed a 3-dimensional compact metric space  $X \times Y$  with 2-dimensional factor spaces  $X$  and  $Y$  for the first time in the literature.

It is also known that there are higher dimensional examples (Kuz'minov [1]): For a given triple  $(m,n,k)$  of positive integers which satisfies a certain condition (that is, the Bockstein condition) there exist compact metric spaces  $X$  and  $Y$  such that  $\dim X = m$ ,  $\dim Y = n$ , but  $\dim (X \times Y) = m+n - k$ .

There exists an algebraic characterization in terms of the homology groups of  $X$  that the equality is valid for every compact metric space  $Y$  (independently, Boltyanskii (see Kuz'minov [1]) and Kodama [1]).

On the other hand, it seems that almost all theorems, which say that the inequality is valid, have been established in a way parallel to the discovery of the theorems, which say that the product is normal (for example, Morita [1]). Hence, there had been long (for about 20 years) standing the following problem:

PROBLEM. When  $X \times Y$  is normal does the inequality (\*) hold ?

The first counter-example for the inequality (\*), and hence which solves the above problem negatively, was constructed by Wage [1] , under the assumption of continuum hypothesis (CH). After that, Przymusiński [3] eliminates CH, and constructed a counter-example within ZFC.

Following their results, we specify our problem as follows.

HIGHER DIMENSIONAL PROBLEM. For a given pair of non-negative integers  $(m,n)$  and  $k = 1,2,\dots, \infty$ , is there any normal product space  $X \times Y$  such that  $\dim X = m$ ,  $\dim Y = n$ , while  $\dim (X \times Y) = m+n+k$  ?

We shall solve this problem affirmatively. In the case  $k = \infty$  our example also solves a problem raised in Fedorčuk [2] under CH.

Therefore, we now have fairly complete information about the behaviour of the dimension function on the product spaces.

Chapter 1. Closed embeddings of complete metric spaces.

Definition 1.1. ....	9
Remark 1.1. ....	10
Lemma 1.1. ....	11
Theorem 1.1. ....	12
Theorem 1.2 (Main Theorem) ....	15
Remark 1.2. ....	16
Theorem 1.3. ....	18
Theorem 1.4. ....	18
Theorem 1.5. ....	18

## 1 Closed embeddings of complete metric spaces.

DEFINITION 1.1. We say that a metric space  $X$  has the type  $(n, \alpha)$  if  $X$  is of dimension  $n$  and of weight  $\alpha$ . A (complete) metrizable space  $X$  of type  $(n, \alpha)$  is called (complete)  $(n, \alpha)$ -universal if it satisfies that

- (i) every (completely) metrizable space of type  $(n, \alpha)$  can be embedded in  $X$  as a (closed) subset.

When  $X$  is not necessarily of type  $(n, \alpha)$ , we say that a space  $X$  is (closed)  $(n, \alpha)$ -embeddable if  $X$  satisfies (i).

There is no doubt to believe that the real line  $\mathbb{R}$  is one of the most important useful 1-dimensional spaces among separable metric spaces. Our present attempt is to seek a non-separable 1-dimensional complete metric space  $T$  of weight  $\alpha$  satisfies that

- (\*) for every  $n$  there exists an  $m < \infty$  such that  $T^m$  is closed  $(n, \alpha)$ -embeddable;
- (\*\*)  $T$  has some nice properties such as linearly orderable or ANR.

REMARK 1.1. (a) One of the frequently used non-separable 1-dimensional space is the hedgehog (or, star) space  $J(\alpha)$  of weight  $\alpha$  (see for example, Nagata [7, Definition IV. 8]). Since every  $J(\alpha)$  is  $\sigma$ -locally compact, the space of irrationals  $P$  cannot be embedded as a closed subset in any finite iterated products of it (it is only known that the countable product  $J(\alpha)^\omega$  is complete  $(\infty, \alpha)$ -universal). Therefore,  $J(\alpha)$  does not serve our purpose.

(b) On the other hand, we have satisfactory spaces for the 0-dimensional case. We believe that the Baire's 0-dimensional space  $B(\alpha)$  of weight  $\alpha$  is the space we want. It is because  $B(\omega)$  (that is, the separable case) is homeomorphic to  $P$  and every  $B(\alpha)$  is complete  $(0, \alpha)$ -universal (Stone [1]).

(c) It is also known (Nagata [7, Theorem VI. 10]) that there are  $(n, \alpha)$ -universal spaces for every  $(n, \alpha)$ . Finally, we note that Lipscomb [1] defined a complete metric space  $L(\alpha)$  of weight  $\alpha$  such that  $n+1$  fold product  $L(\alpha)^{n+1}$  contains an easy definable  $(n, \alpha)$ -universal spaces. But, these embeddings are not necessarily closed.

We begin with the following lemma which is a slight variation of a lemma due to Nagata. Its proof is achieved in a parallel manner of Nagata [7, VI. 2 A)], so that we omit it.

LEMMA 1.1. Let  $M$  be an  $n$ -dimensional metric space, and let  $\{U_i : i \in \omega\}$  and  $\{F_i : i \in \omega\}$  be a collection of open, and of closed subsets of  $M$ , respectively, with  $U_i \supset F_i$  for each  $i$ . Then, for every pair  $u_i > v_i$  of real numbers there exists a collection

$$\mathcal{V}_i = \{V_i : u_i > t > v_i, t \text{ is rational}\}$$

of open sets in  $M$  for each  $i$  satisfying

- (i)  $F_i \subset V_{i,r} \subset \bar{V}_{i,r} \subset V_{i,s} \subset \bar{V}_{i,s} \subset U_i$  if  $r < s$ ;
- (ii)  $\bar{V}_{i,r} = \cap \{V_{i,s} : s > r\}$  and  $V_{i,r} = \cup \{\bar{V}_{i,s} : s < r\}$ ;
- (iii)  $\text{ord} \{\partial V : V \in \mathcal{V}\} \leq n$ , where  $\mathcal{V} = \cup_{i=1}^{\infty} \{V_i\}$ .

Next, we shall prove the following theorem. We introduce the following spacial subset  $Z_n$  of  $R^\infty$ . Let  $Z_n$  be the set of points in  $R^\infty$  at most  $n$  of whose coordinates are rational. Then,  $Z_n$  is completely metrizable, since it is a  $G_\delta$ -subset of  $R^\infty$ . It is also known that  $Z_n$  is  $n$ -dimensional (cf. Nagata [7, p. 155]).

THEOREM 1.1. Let  $M$  be any completely metrizable space which is  $(n, \alpha)$ -universal. Then, the product space  $M \times Z_n$  is closed  $(n, \alpha)$ -universal.

Proof. Let  $X$  be any completely metrizable space of type  $(n, \alpha)$ . Since  $M$  is  $(n, \alpha)$ -universal, we may think of  $X$  as a subspace of  $M$ . Then, by Lavrentiev's theorem (Kuratowski [1, § 35, II])  $X$  is a  $G_\delta$ -subset of  $M$ . If  $X = M$ , our proof is complete, since  $M \times \{q\}$  is a closed subspace homeomorphic to  $X$ , where  $q$  is a point of  $Z_n$ . So we may assume that  $M \setminus X \neq \emptyset$ . Put

$$M \setminus X = \cup_{i=1}^{\infty} F_i \text{ and } F_i \subset F_{i+1},$$

where each  $F_i$  is a closed subset of  $M$ . First, we shall define a continuous map  $f_i: X \rightarrow \mathbb{R}$  for each  $i$  such that

- (1) for each number  $r > 1$  and every point  $z \in F_i$  there exists a  $\delta > 0$  such that  $f_i(x) > r$  for any  $x \in X \cap B_\delta(z)$ ;
- (2)  $f = \pi_{i=1}^{\infty} f_i: X \rightarrow \mathbb{R}^\omega$  is a  $Z_n$ -valued map (that is,  $f$  is a continuous map into  $Z_n$ ).

Let  $\{r_m\}$  be a strictly decreasing sequence of irrational numbers which converges to 0. Put

$$F_{i,m} = \text{Cl}_M(B_{r_{m+1}}(F_i)), \text{ and } U_{i,m} = B_{r_m}(F_i).$$

Then, for the collections  $\{F_{i,m}: i, m \in \omega\}$ ,  $\{U_{i,m}: i, m \in \omega\}$ , and each pair of irrationals  $u_{i,m} = r_m$  and  $v_{i,m} = r_{m+1}$  we can apply Lemma 1.1 to get a collection



$V_{i,m} = \{V_{i,m,t} : r_m > t > r_{m+1}, t \text{ is rational}\}$   
of open sets in  $M$  which satisfies (i) - (iii) of the  
lemma for each pair of integer  $i$  and  $m$ . For each  $i$  we  
shall define a map  $f_i: X \rightarrow R$  as follows.

$$(3) \quad f_i(x) = 1/r_1 \text{ if } x \notin U_{i,1},$$

$$f_i(x) = 1/r_m \text{ if } x \in U_{i,m} \setminus \cup V_{i,m},$$

and

$$f_i(x) = 1/\inf \{r : x \in V_{i,m,r}\} \text{ if } x \in \cup V_{i,m} \setminus F_{i,m}.$$

Then, it is easy to see (Nagata [1]) that  $f_i$  is  
continuous and that

$$(4) \quad f_i(x) = \text{rational if and only if } x \in \partial V \text{ for some}$$

$$V \in \mathcal{V}_i, \text{ where } \mathcal{V}_i = \cup_{m=1}^{\infty} V_{i,m}.$$

From the definition (3) of  $f_i$ , we have  $f_i(x) \geq 1/r_m$  for  
every  $x \in U_{i,m}$ , so that (1) holds. We can also show  
that (2) holds, since  $\{r_m\}$  is strictly decreasing, (4)  
holds, and  $\text{ord } \partial V \leq n$  by (iii). Let

$$g : X \rightarrow M \times Z_n \text{ be } g(x) = (x, f(x)).$$

Then, it is easy to see that  $g$  is an embedding  
(cf. Kuratowski [1, §21 XIII, Theorem]). Let us show  
that  $g$  is a closed embedding. Take any point

$$(y, z) \notin g(X).$$

At first, we consider the case  $y \in X$ . Then, we have  
 $z \neq f(y)$ , since  $(y, z) \neq g(y) = (y, f(y))$ . Therefore,  
there exists an  $i$  such that  $z_i \neq f_i(y)$ , where  
 $z = (z_i) \in Z_n$ . Take two open sets  $U$  and  $V$  in  $R$  such

that  $z_i \in U$ ,  $f_i(y) \in V$  and  $U \cap V = \emptyset$ . Since  $f_i$  is continuous, there exists a  $\delta > 0$  such that  $f_i(X \cap B_\delta(y)) \subset V$ . Put

$$W = B_\delta(y) \times \pi_i^{-1}(U),$$

where  $\pi_i: Z_n \rightarrow \mathbb{R}$  is a restriction of the natural projection from  $\mathbb{R}^\omega$  to the  $n$ -th factor space. Then,  $W$  is an open neighborhood of  $(y, z)$  and it is easy to see that  $W \cap g(X) = \emptyset$ .

Next, we consider the case  $y \notin X$ . In this case we shall use the property (1). Since  $y \notin X$ , there exists an  $i$  with  $y \in F_i$ . Then, for  $r = |z_i| + 1$  there exists a  $\delta > 0$  such that  $f_i(x) > r$  for any  $x \in X \cap B_\delta(y)$ . Put

$$W = B_\delta(y) \times \pi_i^{-1}(-\infty, r).$$

Then,  $W$  is an open neighborhood of  $(y, z)$  and  $W \cap g(X) = \emptyset$ . Thus, the proof is complete.

THEOREM 1.2. For every non negative integer  $n$  and every infinite cardinal  $\alpha$ , there exists a  $(n, \alpha)$ -universal space  $Z_{n, \alpha}$  such that the  $2n$ -dimensional space  $(Z_{n, \alpha})^2$  is closed  $(n, \alpha)$ -embeddable.

Proof. For the case  $\alpha = \omega$  (that is, for the separable case), put  $M = Z_n$ . Then, since  $Z_n$  is  $(n, \omega)$ -universal (e.g. Nagata [7, Theorem IV. 8.]),  $Z_n$  satisfies Theorem 1.1, so that  $Z_{n, \omega} = Z_n$  satisfies the theorem for the case  $\alpha = \omega$ .

For the case  $\alpha > \omega$ , let  $S(A)$  be the hedgehog space with its weight  $\alpha = |A|$  (see Nagata [7, Definition VI. 6] for the definition of the hedgehog space). Let  $E_1 = [0, 1]_1$  and  $E_2 = [0, 1]_2$  be some distinct segments in  $S(A)$ . Put

$$T(A) = S(A) \setminus \{1_i : i = 1, 2\},$$

where each  $1_i$  is the end point of  $E_i$ ,  $i = 1, 2$ . Put

$$Z(A) = \prod_{m=1}^{\infty} T(A)_m,$$

where each  $T(A)_m$  is a copy of  $T(A)$ . Since

$(E_1 \cup E_2) \cap T(A)$  is homeomorphic to the real line  $\mathbb{R}$ , we may think of  $\mathbb{R}^\omega$  as a closed subset of the completely metrizable space  $Z(A)$ . Let  $Z_n(A)$  be the set of points in  $Z(A)$  at most  $n$  of whose non-zero coordinates are rational. Then,  $Z_n(A)$  is  $(n, \alpha)$ -universal, since it contains a topological copy of the universal space  $K_n(A)$  defined in [7, Theorem VI. 10], and vice versa.

On the other hand,  $Z_n(A)$  is completely metrizable, since it is a  $G_\delta$ -subset of  $Z(A)$ . Since  $R^\omega \cap Z_n(A)$  is homeomorphic to  $Z_n$ ,

(5)  $Z_n(A)$  contains  $Z_n$  as a closed subspace.

Put

$$M = Z_n(A).$$

Then,  $M$  satisfies Theorem 1.1, so that  $Z_{n,\alpha} = Z_n(A)$  also satisfies our theorem from (5). This completes the proof.

REMARK 1.2. (a) Since each Baire's 0-dimensional space  $B(\alpha)$  of weight  $\alpha \geq \omega$  contains the space of irrationals  $Z_n$  as a closed subset and  $B(\alpha)^2$  is homeomorphic to itself, Stone's result in Remark 1 (b) follows from Theorem 1.2.

(b) Recently, above Stone's theorem is used by Jayne and Rogers [1] to obtain their representation theorem.

(c) Let  $S(\omega)$  be the hedgehog space of weight  $\omega$ . Put  $F_i = \{1_k : k \leq i\}$ , where  $1_k$  is the end point of each segment of  $S(\omega)$ . For the space  $X = S(\omega) \setminus \cup_{i=1}^{\infty} F_i$  put

$$f_i(x) = 1/\rho(x, F_i) \text{ for each } x \in X.$$

Then,  $f(0) = (1, 1, \dots)$ , where  $0$  is the origin of  $S(\omega)$  and  $f = \prod_{i=1}^{\infty} f_i : X \rightarrow R^\omega$ . Therefore, the closed embeddings used in Kuratowski [1, §21 XIII, Theorem] does not necessarily satisfy our condition (2).

(d) For spaces which are not necessarily completely metrizable, the corresponding results are not valid, even for separable spaces. In fact, assume that there exists a finite dimensional separable metric space  $Z$  such that every 0-dimensional separable metric space can be embedded as a closed subset in  $Z$ . Since  $Z$  is separable, the power of the set of all its closed subsets is at most the cardinality of continuum  $c$ . But, it is known that there exists a family of power  $2^c$  which is homeomorphic to any other (Kuratowski [1, §35, I, Theorem 5]). This is a contradiction.

(e) The following problem is communicated from R. Brown:

If  $Z_{n,\alpha}$  is  $(n,\alpha)$ -universal, then  $(Z_{n,\alpha})^2$  is closed  $(n,\alpha)$ -universal ?

Since there exist compact  $(n,\omega)$ -universal spaces, it is not the case when  $\alpha = \omega$ .

(f) Professor R. Engelking [3] has kindly communicated the problem whether or not there exists a complete  $(n,\alpha)$ -universal space. He also announced in [3] that R. Pol showed that it is the case for  $\alpha = \omega$ . Quite recently, A. Waśko [1] showed that for every infinite cardinal number  $\alpha$  our space  $Z_{n,\alpha}$  in Theorem 1.2 is complete  $(n,\alpha)$ -universal.

Using her result, we have the following theorem (Tsuda [5, Theorem 2]) which improves a result of J. Nagata.

THEOREM 1.3. The  $n+1$  fold product of 1-dimensional space  $Z_{1,\alpha}$  is closed  $(n,\alpha)$ -embeddable.

Therefore, our space  $Z_{1,\alpha}$  has the property (\*). On the other hand, though our space cannot satisfy the property (\*\*), we have the following theorem, using a result of Kodama [5]:

THEOREM 1.4 (Tsuda [5, Corollary 3]). There exists 2-dimensional complete ANR metric space whose  $n+1$  fold product is closed  $(n,\alpha)$ -embeddable.

We conclude this chapter with the following simple improvement of a theorem of Fréchet (see, Engelking [2, Problem 1.3. G (b)]), which will be used in the next chapter.

THEOREM 1.5. Every metric space consisting of countable points can be embedded in the space of rational numbers  $\mathbb{Q}$  as a closed subset.

Proof. Let  $X$  be arbitrary metric space consisting of countable points. Then, by the theorem of Fréchet we may think  $X$  is a subspace of  $\mathbb{Q}$ . Put

$$Z = \mathbb{Q}^2 \setminus \{0\} \times (\mathbb{Q} \setminus X).$$

Then,  $X$  is a closed subset of  $Z$ . On the other hand, by a theorem of Sierpinski (Engelking [2, Problem 1.3.6 (b)])  $Z$  is homeomorphic to  $\mathbb{Q}$ . This completes the proof.

Chapter 2. Non-existence of universal spaces for  
some stratifiable spaces.

Theorem 2.1 (Main Theorem). . . . .	21
Corollary 2.1. . . . .	22
Theorem 2.2. . . . .	22
Corollary 2.2. . . . .	22
Proposition 2.1. . . . .	23
Remark 2.1. . . . .	25
Problem 2.1. . . . .	28
Proposition 2.2. . . . .	28
Problem 2.2. . . . .	29
Theorem 2.3. . . . .	30
Remark 2.3. . . . .	31
Problem 2.3. . . . .	31



## 2 Non-existence of universal spaces for some stratifiable spaces.

One of remarkable properties of metric spaces is that there are universal spaces with a given weight (for example, the Baire's 0-dimensional space and generalized Hilbert spaces). The purpose of this chapter is to show that there are none for some stratifiable spaces. We also discuss some positive result at the end of this chapter.

THEOREM 2.1. There are no universal spaces for the following subclasses of stratifiable spaces with a given network weight  $\alpha \geq \omega$ ;

- (1)  $\sigma$ -discrete stratifiable spaces,
- (2)  $M_0$ -spaces in the sense of Heath and Junnila [1],
- (3) stratifiable  $\mu$ -spaces (Mizokami [1]),
- (4) stratifiable spaces with encircling nets (Oka [1]),
- (5) (strongly 0-dimensional) stratifiable spaces,
- (6) (strongly 0-dimensional) L-spaces in the sense of Nagami [6],
- (7) (strongly 0-dimensional) free L-spaces in the sense of Nagami [7].

When we use a given weight instead of a given network weight, we have a similar result:

COROLLARY 2.1. For any given infinite cardinal number  $\alpha$  there exists a cardinal number  $\beta > \alpha$  such that there exist no universal spaces for the above classes (1) - (7) with a given weight  $\beta$ .

When we consider more restricted subclasses, contained in Lašnev spaces or Nagata spaces, we also have a similar result:

THEOREM 2.2. There are no universal spaces for either countable Lašnev spaces or first-countable, separable, strongly 0-dimensional stratifiable spaces.

COROLLARY 2.2. There are no universal spaces for separable Lašnev spaces.

We start with the following proposition.

PROPOSITION 2.1. For every infinite cardinal number  $\alpha$  there exists a family  $S$  of cardinality  $2^{2^\alpha}$ , none of whose members are homeomorphic to each other, and which consists of  $\sigma$ -discrete stratifiable spaces  $S$  with  $|S| = nw(S) = \alpha$  and  $w(S) \leq 2^\alpha$ .

Proof. Let  $\{\alpha_i\}$  be countable disjoint copies of  $\alpha$ . Then, for each ultrafilter  $p$  on  $\alpha$ , let  $S_p$  be the space  $\alpha_0 \cup \{p\}$ , where  $\alpha_0 = \cup \{\alpha_i\}$ , with the following topology  $\tau_p$ .

- (i)  $U \in \tau_p$  if and only if  $p \notin U$ , or  $p \in U$  and there exist an  $n \in \omega$  and an  $F \in p$  such that  $U \cap \alpha_i \supset F$  for every  $i \geq n$ .

Then, since each  $\alpha_i$  is a clopen discrete subspace of  $S_p$ , it is readily seen that  $S_p$  is a  $\sigma$ -discrete stratifiable space and that  $|S_p| = nw(S_p) = \alpha$  and  $w(S_p) \leq 2^\alpha$ . Next, we shall show that

- (ii) the family of all  $S_q$ 's which are homeomorphic to a fixed  $S_p$  is of the cardinality at most  $2^\alpha$ .

Let  $S_q$  and  $S_r$  be homeomorphic to a fixed  $S_p$ . Then, take any homeomorphisms  $h_q: S_p \rightarrow S_q$  and  $h_r: S_p \rightarrow S_r$ .

To show (ii) it suffices to show that

- (iii)  $h_q|_{\alpha_0} \neq h_r|_{\alpha_0}$  if  $q \neq r$ .

On the contrary, assume that there exists a permutation  $h: \alpha_0 \rightarrow \alpha_0$  such that  $h = h_q|_{\alpha_0} = h_r|_{\alpha_0}$ . Since  $q \neq r$ , we can assume without loss of generality that there exists an  $F \in q \setminus r$ . Put

$F_0 = \cup_{i=1}^{\infty} F_i$ , where each  $F_i$  is a copy of  $F$  in  $\alpha_i$ , and put

$$U_0 = \{q\} \cup F_0.$$

Then,  $U_0$  is a neighborhood of  $q$  by the definition (i) of  $\tau_q$ . Hence,  $h_r h_q^{-1}(U_0)$  is a neighborhood of  $r$ , and by the definition (i) there exist an  $n \in \omega$  and a  $G \in r$  such that  $G \subset \alpha_i \cap h_r h_q^{-1}(U_0)$  for  $i \geq n$ . Hence,  $F_n \supset G$ , since  $\alpha_n \cap h_r h_q^{-1}(U_0) = \alpha_n \cap h h^{-1}(U_0) = F_n$ .

Since  $F_n$  is a copy of  $F$  and  $r$  is an ultrafilter, we have  $F \in r$ . This contradiction shows that (iii) holds. Since (ii) holds and there exist  $2^{2^\alpha}$  many ultrafilters on  $\alpha$  (Engelking [1, Theroem 3.6.11]), there exists a family  $S = \{S_p : p \in \Lambda\}$ , with  $|\Lambda| = 2^{2^\alpha}$ , none of whose members are homeomorphic to each other. Thus, our family  $S$  is the required one.

REMARK 2.1. For the case  $\alpha = \omega$  we can show the above proposition much easier, using subspaces  $\omega \cup \{p\}$  of the Stone-Čech compactification  $\beta\omega$  (cf. Nagami [6] and Shelah and Rudin [1]).

Proofs of Theorem 2.1 and Corollary 2.1. The proofs for all the cases of Theorem 2.1 and Corollary 2.1 are derived simultaneously from the following observation, since every class (i+1) contains the preceding class (i) except  $i = 5$ , the class (3) contains the class (7), and the collection  $\mathcal{S}$  in Proposition 2.1 consists of  $\sigma$ -discrete L-spaces (cf. references cited in Theorem 2.1).

Let us denote by  $[A]^\kappa$  the family consisting of all subsets of a set  $A$  having cardinality  $\kappa$ . If  $\kappa$  and  $\lambda$  are infinite and  $\kappa \leq \lambda$ , then  $|[\lambda]^\kappa| = \lambda^\kappa$ . It follows that if  $X$  is a stratifiable space (more generally, a paracompact  $\sigma$ -space) with  $nw(X) \leq 2^\kappa$ , where  $\kappa$  is infinite, then  $|X| \leq |[2^\kappa]^\omega| = |(2^\kappa)^\omega| = 2^\kappa$ ; consequently,  $|[X]^\kappa| \leq 2^\kappa$ , and we see that  $X$  cannot contain copies of all the spaces in Proposition 2.1, with  $\alpha = \kappa$ .

Proofs of Theorem 2.2 and Corollary 2.2. It suffices to show that there exists a family  $\mathcal{T}$  (respectively,  $\mathcal{U}$ ) of cardinality  $2^c$ , where  $c = 2^\omega$ , none of whose members are homeomorphic to each other, and which consists of countable Lašnev spaces (respectively,  $\sigma$ -compact, first countable, separable, strongly 0-dimensional, stratifiable spaces).

Let  $C$  be the Cantor set in the real line with usual Euclidean topology, and let  $0 \in C$ . Take a countable dense subset  $D$  of  $C$ , and let  $\{x_n\} \subset C$  be a sequence converging to 0. At first, we show the existence of  $\mathcal{T}$ . Put

$F = \{0\} \times C$ , and  $A = \{(x_n, d_k) : k \leq n, \text{ and } n \in \omega\}$ , where  $D = \{d_k : k \in \omega\}$ . Then,  $A$  is a countable discrete subset in  $C^2$  with  $\text{Cl}(A) \supset F$ . For every subset  $T \subset F$  let

$$Y_T = T \cup A.$$

Take a point  $p_T \notin A$ , and let

$$X_T = \{p_T\} \cup A.$$

Define a function  $\phi_T: Y_T \rightarrow X_T$  as

$$\phi_T(T) = p_T \text{ and } \phi_T(a) = a \text{ for every } a \in A.$$

We topologize the set  $X_T$  as

$$U \text{ is open in } X_T \text{ if } \phi_T^{-1}(U) \text{ is open in } Y_T.$$

By the definition of  $\phi_T$ ,  $\phi_T$  is a closed map between  $Y_T$  and  $X_T$ . Thus, each  $X_T$  is a countable Lašnev space.

By the proof of Proposition 2.1, one can show that there exists a family  $\mathcal{T} = \{X_T: T \in F\}$ , with  $|F| = 2^c$ , none of whose members are homeomorphic to each other.

Next, we show the existence of  $u$ . We shall modify the examples constructed in van Douwen and Przymusiński [1]. Let  $F$  be a free ultrafilter on  $\omega$ . Then, enumerate the family of all elements of  $F$  as  $\{F_s: s \in C\}$ , and for each  $s \in C$  and  $m \in \omega$  choose a  $q_s(m) \in D$  such that  $0 < |s - q_s(m)| < 1/m$ , and put

$$D_s = \{q_s(m): m \in \omega\}, \text{ and } E_s = \{X_n: n \in F_s\}.$$

We topologize the set

$$\Delta_F = (C \times \{0\}) \cup (D \times \{X_n: n \in \omega\})$$

as follows. Points of  $D \times \{X_n: n \in \omega\}$  are isolated, and basic neighborhoods of point  $(s, 0) \in C \times \{0\}$  have the form

$$B_m(s) = \{(x, y) \in \Delta_F: |s-x| < 1/m\} \setminus (D_s \times E_s \cup \{s\} \times \{X_n: n \in \omega\}),$$

for  $m \in \omega$ . Then, it is known (van Douwen and Przymusiński [1]) that each  $\Delta_F$  is first countable,  $\sigma$ -compact, strongly 0-dimensional, cosmic. We can show without difficulty that each  $\Delta_F$  is stratifiable. Here, we show moreover that it admits a free L-structure.

Let  $\mathcal{V} = \{V_i: i \in \omega\}$  be a countable clopen base of  $C$ .

Put

$$L_0 = C \times \{0\}, \quad L_{2i} = V_i \times C, \quad \text{and } L_{2i+1} = \{u_i\},$$

where  $\Delta_F \setminus L_0 = \{u_i: i \in \omega\}$ .

Put

$$u_{L_0} = \{u\} : u \notin L_0, \text{ and } u_{L_i} = \{ \Delta_F \setminus L_i \}$$

for each  $L_i \neq L_0$ . Then, one can easily check that

$$(L = \{L_i : i \in \omega\}, u_L : L \in L)$$

is a free  $L$ -structure of  $\Delta_F$ . Again, by the proof of Proposition 2.1, we can show that there exists a family  $u = \{\Delta_F : F \in \Lambda\}$ , with  $|\Lambda| = 2^c$ , none of whose members are homeomorphic to each other. This completes the proof of Theorem 2.2.

The proof for Corollary goes on a way parallel in Corollary 2.1.

Professor Junnila [2] has kindly communicated to the author the following problem.

PROBLEM 2.1. Does there exist a universal space for closed images of countable metric spaces?

Though the above problem is still open, the following proposition shows that it is enough to consider the case of "closed images of the space of natural numbers  $\mathbb{Q}$ ".

PROPOSITION 2.2. Every closed map  $f: X \rightarrow Y$ , where  $X$  is a countable metric space, is a restriction of a closed map from  $\mathbb{Q}$  to a countable space containing  $Y$ .



Proof. By Theorem 1.5 we can assume that  $X$  is a closed subset of  $Q$ . Then, consider the following decomposition of  $Q$ ;

$$\{\{x\} : x \notin X\} \cup \{f^{-1}(y) : y \in Y\}.$$

Then, the decomposition space  $S$  and the natural quotient map  $q: Q \rightarrow S$  satisfies that

$$q|_X = f.$$

It is easy to see that  $q$  is a closed map. This completes the proof.

The following problem is communicated by T. Nogura:

PROBLEM 2.2. Can every closed images of countable metric space be embedded in the  $\mathbb{E}$ -product  $\mathbb{E}_0$  of countably many  $Q$ 's ?

It is known that (San-ou [1])  $\mathbb{E}_0$  is a stratifiable space consisting of countable points.

Finally, we discuss some positive result. We believe that the class of simplicial complexes with weak topology  $|K|_W$  is one of the most important classes among those of generalized metric spaces. In particular, it is known that (Ceder [1]) every  $|K|_W$  is stratifiable, and that (Cauty [1]) every  $|K|_W$  is ANR(stratifiable).

For a given cardinal  $\alpha$  let  $|K(\alpha)|_{\mathcal{W}}$  be the full complex with its vertices  $A$ , where the cardinality  $|A|$  is equal to  $\alpha$ . Let  $Z_\alpha$  be the countable Cartesian product of  $|K(\alpha)|_{\mathcal{W}}$ . Then, we consider the class  $\mathcal{C}$  defined as follows:

$X \in \mathcal{C}$  if and only if  $X$  is a subset of  $Z_\alpha$  for some cardinal number  $\alpha$ .

It is easy to see that

- (1) if  $A \subset X \in \mathcal{C}$ , then  $A \in \mathcal{C}$  (that is, subset hereditary);
- (2) if  $X_i \in \mathcal{C}$ , then  $\prod X_i \in \mathcal{C}$  (that is, countably productive);
- (3) every (not necessary full) complex belongs to  $\mathcal{C}$ .

Here, we show that every metric space belongs to  $\mathcal{C}$ :

**THEOREM 2.3.** Every metric space of weight  $\alpha$  can be embedded in  $Z_\alpha$ .

Proof. It suffices to see that the hedgehog space  $S(A)$  of weight  $\alpha = |A|$  can be embedded in  $Z_\alpha$ . Put  $S(A) = \cup_{\lambda \in A} I_\lambda$ , where each  $I_\lambda$  is a copy of unit interval  $I$ , and  $I_\lambda \cap I_\mu = \{0\}$  for distinct  $\lambda$  and  $\mu$ .

On the other hand, for each vertex  $\lambda \in A$  and an integer  $i \in \omega$ , where  $\lambda$  is distinct from a fixed vertex  $0$ , put  $\lambda_i = (0, \dots, 0, \lambda, 0, \dots) \in Z_\alpha$ .

Since each factor  $|K|_{\mathcal{W}}$  of  $Z_\alpha$  has the ordinary convex structure, we can define

$t\lambda_i + (1-t)\lambda_{i+1}$  for each  $\lambda_i, \lambda_{i+1}$  and  $0 \leq t \leq 1$ .

Then, put also

$$J_{\lambda, i} = \{t\lambda_i + (1-t)\lambda_{i+1} : 0 \leq t \leq 1\}.$$

Note that  $J_\lambda = \cup_{i=1}^{\infty} J_{\lambda, i} \cup \{(0, 0, \dots)\}$  is homeomorphic to the unit interval. It is not difficult to see that  $\cup_{\lambda \in A} J_\lambda$  is homeomorphic to  $S(A)$ . This completes the proof.

REMARK 2.3. (a) By the definition of  $\mathcal{C}$  we see that the space  $Z_\alpha$  is a universal space for the members in  $\mathcal{C}$  of weight  $\alpha$ .

(b) The class  $\mathcal{C}$  is not contained in either the class of L-spaces or free L-spaces, since the full complex is not a free L-space by Nagami and Tsuda [1].

On the other hand, every (strongly 0-dimensional) member of it is an  $EM_3(M_0)$ -space respectively, since every member of  $\mathcal{C}$  is a  $\mu$ -space.

PROBLEM 2.3. Find an inner characterization of the members of the class  $\mathcal{C}$ .

Chapter 3. Normal spaces  $X \times Y$  satisfying the inequality  
 $\dim (X \times Y) > \dim X + \dim Y$ .

3.	A 1-dimensional Wage's metric space. ....	37
4.	$n$ -dimensional Wage's metric spaces. ....	49
5.	Some consequences of Kunen construction and those of van Douwen construction. ....	56
6.	Wage's factorization technique. ....	64
7.	Przymusiński's method. ....	72
8.	An application to inverse sequences. ....	77
9.	Perfect normality of products. ....	82
10.	Local compactness and collectionwise normality of products. ....	88
11.	With a Lindelöf factors and normality of products. .....	99
12.	Non-normal Wage-type examples. ....	105
	Appendix. Several product theorems. ....	116

In Chapter 3 we shall discuss the counter-examples for the inequality

$$(*) \dim (X \times Y) \leq \dim X + \dim Y.$$

Here, we mainly discuss the problem under the condition that  $X \times Y$  is normal, since it seems that there is still room to discuss what the suitable dimension function for non-normal spaces is (see Nagata [7], and compare the two treatments in Engelking [1 and 2] of the dimension function due to Katětov [1].)

We believe, however, that it is an interesting area of dimension theory to treat the cases when the product spaces are non-normal. Hence, we shall mention it in the final section.

The problem whether or not the inequality (\*) is valid if  $X \times Y$  is normal has been standing (at least 20 years) (for example, see Gillman and Jerison [1]).

It is Wage [1], who constructed for the first time the following counter-example for the inequality (\*), assuming Continuum Hypothesis(CH):

EXAMPLE 1 (CH). There exist two spaces  $X$  and  $Y$  such that  $\dim (X \times Y) > \dim X + \dim Y = 0$  and that  $X \times Y$  is perfectly normal and locally compact.

Eliminating the assumption CH, Przymusiński [1,3] constructed the following counter-example, without any set-theoretic assumptions beyond ZFC:

EXAMPLE 2. There exists a separable, first-countable, Lindelöf space  $X$  such that  $\dim X = 0$  and  $X^2$  is normal, while  $\dim X^2 > 0$ .

Since Wage [1] discovered a counter-example for the inequality (\*) for the first time, we say that a triple  $(X, Y, X \times Y)$  is a Wage-type example when it does not satisfy the inequality (\*).

The construction of the Wage-type examples in Examples 1 and 2 can be outlined as follows.

At first, we show that there exists a "special" separable complete metric space  $\tilde{T}$ , which we want to call the 1-dimensional Wage's metric space. Then, we extend it to a metric space  $T$ . Next, we approximate  $T$  by a product topology of  $X \times Y$ . Finally, using some properties of our approximation, we calculate the dimensions of factors  $X$  and  $Y$ , and of the product  $X \times Y$ .

To solve our problem, we follow the above procedure. That is:

§§3,4. We shall find at first an  $n$ -dimensional Wage's metric space and next find its extension.

§§§ 5,6,7. We shall perform a delicate and complicated approximation of it.

§§§ 9,10,11. We can show that not only  $\dim(X \times Y)$  is positive, but also it is exactly equal to  $n$ . Hence, we must work very hard, using the properties derived from our approximations.

One of the main points of our construction is to find the metric spaces in §4. Such spaces will be produced in Theorems 4.1 and 4.2. Their completeness is heavily related to the elimination of CH. We shall explain this point in some detail.

A space which satisfies all the conditions in Theorem 4.1 except completeness would be easily obtained by a simple modification of Rubin, Scori, and Walsh [1] construction. But, if we use such a space, we must assume CH throughout our constructions.

Moreover, the fact that our space satisfies the condition (1) in Theorem 4.1 is deeply related to make  $\dim(X \times Y) \geq n$ . So, we must try to find a space which has the property (1) and completeness, simultaneously.

Since the space we found is satisfactory for our purpose, we can use both of approximation techniques


due to Wage [1,2] (which is also called the factorization technique in there), and due to Przymusiński [3].

(There is, however, some room to improve our space for producing other Wage-type examples (see the appendix and Remark 4.2(b)).)

Though we can proceed our approximation in a way parallel to their ones, rough constructions sometimes cause some errors to make our product space being normal (see the footnote 7 in Przymusiński [6]). Therefore, we must do the delicate and complicated inductive constructions very carefully. The techniques used here go back to the idea of Kunen (as cited in Przymusiński [5]) and that of van Douwen [1], respectively. In Chapter 3 we will construct both of higher dimensional version of Examples 1 and 2.



### 3. A 1-dimensional Wage's metric space.



### 3. A 1-dimensional Wage's metric space.

The first Wage's metric space was constructed in Wage [1] as a 1-dimensional subspace of 2-dimensional cube  $I^2$ . In [2] Wage gives a more elementary one as follows:

THEOREM 3.1. There exists a 1-dimensional separable metric topology  $\tilde{\rho}$  on  $C$  such that

- (1)  $\tilde{\rho}$  is the topology generated by the subbase  $\varepsilon_0 \cup \{h^{-1}(U) : U \in \varepsilon\}$ , where  $h:C \rightarrow I$  is a lower semi-continuous function;
- (2) there exists a pair of disjoint closed sets  $A_1$  and  $B_1$  in  $(C, \tilde{\rho})$  such that every separation  $S$  between  $A_1$  and  $B_1$  has the cardinality of continuum.

Since we shall show the above result as the special case of Theorem 4.1, there is no need to prove it here. But we believe that Wage's function  $h$  in [2] is defined more explicitly than ours, and is interesting in itself (see also explanation of computer graphics I & II).

Therefore, we give here the definition of the function  $h$  in [2] together with the proof of (2). We begin with the definition of the function  $h$ . Let  $Q$  be a countable dense subset in the open interval  $(0,1)$ . Let

$$f : C \rightarrow \prod_{q \in Q} \{0,1\}_q$$

be a natural homeomorphism between the Cantor set and the countable product of 2-point set  $\{0,1\}$ . Then, define  $h:C \rightarrow I$  as follows.

$$h(x) = \text{Sup} \{q: \pi_q f(x) = 1\} ,$$

where  $\pi_q: \prod_{q \in \mathbb{Q}} \{0,1\}_q \rightarrow \{0,1\}_q$  is the natural projection. Then, it is easy to see that  $h$  is lower semi-continuous.

Proof of (2) (due to Wage [1]). Put

$$A_1 = h^{-1}(0) \text{ and } B_1 = h^{-1}(1).$$

Let  $S$  be any separation between  $A_1$  and  $B_1$ . Then, there exist two open sets  $U$  and  $V$  in  $X$  such that

$$U \cup V = X \setminus S, \quad U \cap V = \emptyset, \quad U \supset A, \quad \text{and } V \supset B .$$

Fix  $x \in U$  and  $\alpha$  positive integer  $n$ . Consider the following procedure for generating  $x'$ ,  $x''$ ,  $B'$ , and  $B''$ :

Let  $B$  be an  $\varepsilon_0$ -clopen neighborhood of  $x$  with its diameter  $< 1/n$ . Define

$$\lambda = \text{Sup} \{ h(y) : y \in B \cap U \} .$$

It follows from the definition of  $h$  that  $\lambda > h(x)$ .

Fix two distinct points  $x'$  and  $x''$  of  $B \cap U$  such that  $|h(x') - \lambda|$  and  $|h(x'') - \lambda|$  are less than  $|h(x) - \lambda|/2$ .

(Note that  $h(x'), h(x'') > h(x)$ .) Since  $h$  is lower semi-continuous, choose disjoint  $\varepsilon_0$ -clopen sets  $B'$  and  $B''$  such that diameters of  $B'$  and  $B''$  are less than  $1/(n+1)$ ,  $x' \in B'$ ,  $x'' \in B''$ ,  $B' \cup B'' \subset B$ , and  $h(t) > h(x)$  for every  $t \in B' \cup B''$ .

Fix  $x_0 \in U$  and  $n_0 = 1$ . We can inductively generate sequences  $\{x_n\}$  and  $\{B_n', B_n''\}$  from  $x_0$  and  $n_0$  using the above procedure: Having chosen  $x_m$  for each  $m < k$ , fix  $n_k = k$  and  $x_k$  as either  $x_{k-1}'$  or  $x_{k-1}''$ . Since we have two choices at each step, there are  $c$  possible such sequences. The construction guarantees that each sequence has a different (unique) limit point in the Euclidean topology  $\epsilon_0$ . Let  $x = \lim x_n$ . By construction,  $\{h(x_n)\}$  is monotone increasing. Hence,  $h(x) \leq \lim h(x_n)$ , since  $h$  is lower semi-continuous. On the other hand, by the definition of  $h$ ,  $h(x) \geq \lim h(x_n)$ .

Therefore,  $h(x) = \lim h(x_n)$ , and hence  $x \in \text{Cl}_{\tilde{\rho}}(U)$ , since  $x_n \in U$ . We complete the proof of (2) by showing that  $x \notin U$ . Suppose that  $x \in U$ . Then, there are real numbers  $a, b$ , and an  $\epsilon_0$ -clopen set  $B_n$  such that

$$x \in B_n \cap h^{-1}(a, b) \subset U.$$

This implies that  $\lambda \geq b$  for each  $\lambda$  defined in the procedure above in the construction of  $\{x_m\}_{m \geq n}$ . But, this contradicts at some step, the condition that  $|h(x') - \lambda|$  and  $|h(x'') - \lambda|$  are less than  $|h(x) - \lambda|/2$ .

REMARK 3.1 (a) (due to Wage [1]). It is probably easiest to think of  $\tilde{\rho}$  as the restriction of the Euclidian topology on  $I^2$  to the subset

$$\tilde{X} = \{(x, h(x)) : x \in C\}$$

( $h(x)$  stands for the "height" of  $x$  in  $I^2$ ).

(b) Comparing two Wage's metric spaces in [1, 2], we would say that the latter is better, since topologies on Cantor set  $C$  are more convenient to determine the dimension of itself, the factors, and products (see §§§ 9, 10, 11).

## Explanation of computer graphics I.

We shall explain here the following computer graphics given in the next page, which is performed by Mrs. K. Nogura. We believe that the graphs show some aspects of Wage's metric space given in the preceding section.

Let  $\{q_i: 1 \leq i \leq N\}$  be a collection of  $N$  rational numbers. For each  $N$ -tuple  $x = (x_1, \dots, x_N)$  consisting of 0, 1 sequence we assign  $\tilde{\phi}(x)$  by the following formula:

$$\tilde{\phi}(x) = \text{Max} \{q_i: 1 \leq i \leq N \text{ and } x_i = 1\}.$$

We also put

$$x^* = \sum_{i=1}^N x_i / 2^i.$$

Let

$$C^* = \{x^*: x \in \{0,1\}^N\}.$$

Then, the graphs consist of

$$\tilde{G}^* = \{(x^*, \tilde{\phi}(x)) : x^* \in C^*\},$$



where  $N = 8$  in the bottom half, and  $N = 10$  in the upper half, respectively. (We put

$$(q_1, \dots, q_{10}) = (1/2, 1/3, 1/4, 2/3, 1/5, 1/6, 2/5, 3/4, 3/5, 1/7).)$$

We may think that the graph  $\tilde{G}^*$  is the restriction of the Wage's metric space in Theorem 3.1 to the following subset

$$\{(x, h(x)) : x \in \{0,1\}^N \times \{0\} \times \{0\} \times \dots\},$$

where the first  $N$ -terms of  $\{q_i\}$  in Theorem 3.1 are equal to the above ones.



For each  $i$  let  $B_i$  be a finite disjoint clopen cover of  $(C, \epsilon_0)$  with its mesh  $1/i$ , and  $\bigwedge_{i \in \mathbb{N}^+}$  each  $B_{i+1}$  refines  $B_i$ . Then, since  $h$  is lower semi-continuous, there exists a monotone increasing continuous functions  $\{h_i: C \rightarrow I\}$  such that  $h_i(x) \uparrow h(x)$  and  $h_i|_B$  is constant for each  $B \in B_i$ . Now, we define a function  $H: C^2 \rightarrow I$  as follows.

$$H(x,y) = \begin{cases} h(x) & \text{if } x = y, \\ 0 & \text{if } (x,y) \notin \bigcup \{B^2: B \in B_1\}, \\ h_i(x) = h_i(y) & \text{if } (x,y) \in \bigcup \{B^2: B \in B_i\} \setminus \bigcup \{B^2: B \in B_{i+1}\}. \end{cases}$$

Let  $\rho$  be the topology on  $C^2$  generated by the subbase  $\epsilon_0 \cup \{H^{-1}(U) : U \in \mathcal{E}\}$ . Then,  $(C^2, \rho)$  is homeomorphic to the set  $\{(x,y), H(x,y) : (x,y) \in C^2\}$  with the restriction of the Euclidean topology on  $C^2 \times I$ . Hence,

**THEOREM 3.2.**  $(C^2, \rho)$  is a complete separable metric space satisfying

- (1)  $\tilde{\rho} = \rho|_{\Delta}$ , where  $\Delta = \{(x,x) : x \in C\}$ ;
- (2)  $\rho|_{C^2 \setminus \Delta}$  is homeomorphic to usual Euclidean topology on it;
- (3) both topologies  $\rho|_{(\{t\} \times C)}$  and  $\rho|_{(C \times \{t\})}$  are homeomorphic to  $\epsilon_0$ ;
- (4)  $\rho$  is coarser than both topologies  $\tilde{\rho} \times \epsilon_0$  and  $\epsilon_0 \times \tilde{\rho}$ .



We say that the above extension  $(C^2, \rho)$  is a natural extension of Wage's metric space  $(C, \tilde{\rho})$ . In Wage [1,2] different kinds of extensions are used, and the property (4) is not mentioned there. We give here its proof, since it is deeply concerned in constructing our examples with a pseudocompact (or, metric) factor.

Proof of (4). We shall show that  $\rho$  is coarser than  $\tilde{\rho} \times \varepsilon_0$ , (the proof for  $\varepsilon_0 \times \tilde{\rho}$  goes on in a parallel way). It suffices to show that for any point  $(x, x)$  and a given  $\delta > 0$  there exists a  $\rho$ -open neighborhood  $U$  of  $x$  and an  $\varepsilon_0$ -open neighborhood  $B$  of  $x$  such that

(i)  $|H(y, z) - H(x, x)| < \delta$  for every  $(y, z) \in U \times B$ .

Since  $h_i(x) \uparrow h(x)$ , there exists an  $n$  such that  $0 \leq h(x) - h_n(x) < \delta/2$ . Let  $B \in \mathcal{B}_j, j \geq n$ , be an  $\varepsilon_0$ -open neighborhood of  $x$  such that  $|h_n(y) - h_n(x)| < \delta/2$  for every  $y \in B$ . Let  $U \subset B$  be a  $\tilde{\rho}$ -neighborhood of  $x$  such that

$|h(x) - h(z)| < \delta/2$  for every  $z \in U$ . Then, note that  $H(y, z) \leq h(y)$  and  $|h_n(x) - h(y)| < \delta$  for  $(y, z) \in U \times B$ .

Thus, for  $(y, z) \in (Y \times B) \cap \Delta$ , (i) holds, since  $H(y, z) = h(y)$ . Let  $(y, z) \in U \times B \setminus \Delta$ , and put

$H(y, z) = h_k(y) = h_k(z)$  for  $k \geq j$ . Then, if  $h(y) > h(x)$ , (i) holds, since  $h_n(y) \leq H(y, z) \leq h(y) \leq h(x) + \delta/2$ .

If  $h(y) \leq h(x)$ , then (i) holds, since

$$h_n(y) \leq H(y, z) \leq h(y) \leq h(x).$$

REMARK 3.2. The above natural extension, using an increasing sequence of continuous functions, is a slight modification of the construction given by Przymusiński [3]. The extension suggested in Wage [1,2] are too complicated to verify the property (4) in Theorem 3.2.

## Explanation of computer graphics II.

We shall explain here the following computer graphics in the next page, together with the ones given in the front pages of every section. We believe that they show some outlines of the natural extension given in the preceding section as well as the extension given in Wage [2].

We shall use same notations given in Explanation I.

For each  $x, y \in \{0,1\}^N$ , we define

$$\phi(x,y) = \text{Min} \{ \phi(z) : \text{for all } z \text{ such that } x^* \leq z^* \leq y^* \}.$$

Then, the graph consists of

$$G^* = \{ (x^*, y^*, \phi(x,y)) : x, y \in \{0,1\}^N \}.$$



The one given in the upper half of the next page is performed for the case  $N = 4$ . The bottom half is the corresponding Wage's metric space given in explanation I.

The graphics in every section are the ones performed for the case  $N = 10$ . In this case, however, they only show the graphs around the diagonal (they consist of 9 pieces), since the whole graphs need the capacity beyond our computer.

We may regard that the graph  $G^*$  is the restriction of the natural extension of Wage's metric space to the subset


$$\{ (x,y,H(x,y)) : x,y \in \{0,1\}^N \times \{0\} \times \{0\} \times \dots \}.$$

12:



4. n-dimensional Wage's metric spaces.

32: S882



#### 4. n-dimensional Wage's metric spaces.

We begin with the following two lemmas due to Mazurkiewicz [1].

LEMMA 4.1. Let  $h: C \rightarrow I^n$  be a Baire function of class 1. Then, the graph  $G = \{(x, h(x)) : x \in C\}$  is a  $G_\delta$ -set of  $I^{n+1}$  and hence, is completely metrizable.

LEMMA 4.2. There exists a function  $\eta: \text{Exp}(I^n) \rightarrow I^n$  such that

- (a)  $\eta(M) \in M$  for each  $M \in \text{Exp}(I^n)$ ,
- (b) if  $M_k \supset M_{k+1}$  and  $M_k \in \text{Exp}(I^n)$  for each  $k \in \omega$ , then  
$$\lim_{k \rightarrow \infty} \eta(M_k) = \eta\left(\bigcap_{k=1}^{\infty} M_k\right) \text{ in } I^n.$$

Proof of Lemma 4.2. Let  $\psi: C \rightarrow I$  be an order preserving continuous surjective map (for example, see Engelking [2, Problem 1.3.D]). Put

$$\phi = \psi^n: C^n \rightarrow I^n.$$

Let us regard  $C^n$  as a subset of the real line. Put for each  $M \in \text{Exp}(I^n)$

$$(*) \quad \eta(M) = \phi(\min(\phi^{-1}(M))),$$

where minimum is taken with respect to the usual order of the real line. Note that our definition (\*) is well-defined, since  $\min(\phi^{-1}(M)) \in C^n$  for every above  $M$ . Then, one can show that  $\eta$  satisfies (a), (b).

THEOREM 4.1. (An n-dimensional Wage's metric space).  
 For every  $n = 1, 2, \dots, \infty$ , there exists a separable completely metrizable, n-dimensional space  $(C, \tilde{\rho}(n))$  such that

- (0)  $\tilde{\rho}(n)$  is the topology generated by the subbase  $\varepsilon_0 \cup \{h^{-1}(U) : U \in \varepsilon_0\}$ , where  $h: C \rightarrow I^n$  is a Baire function of class 1, and  $h$  is lower semi-continuous when  $n = 1$ ;
- (1) there exists a collection  $\{(A_i, B_i) : i = 1, \dots, n\}$  of n-pairs of disjoint closed sets which satisfies that, if for each  $i$ , a closed set  $S_i$  separates  $A_i$  and  $B_i$ , then the set  $\bigcap_{i=1}^n S_i$  has the cardinality of continuum.

Proof. At first, we consider the case  $n < \infty$ . Put

$$\tilde{A}_{n+1} = \pi_1^{-1}(0) \text{ and } \tilde{B}_{n+1} = \pi_1^{-1}(1),$$

where  $\pi_1: I^{n+1} \rightarrow I_1$  is the natural projection to the first factor. Let  $\tau$  be the collection of all continua meeting both  $\tilde{A}_{n+1}$  and  $\tilde{B}_{n+1}$  in  $I^{n+1}$ . Then,  $\tau$  is closed in  $\text{Exp}(I^{n+1})$ , and is compact metrizable, and hence there exists a continuous map  $g$  from  $(C, \varepsilon_0)$  onto  $\tau$  such that

- (2)  $|g^{-1}(T)| = c$  for any  $T \in \tau$ .

By the definition of  $\tau$  we note that

- (3) for each  $p \in C$  we have  $T \cap \pi_1^{-1}(p) \neq \emptyset$  for any  $T \in \tau$ .

Let  $\pi^n : I^{n+1} \rightarrow I^n$  be the projection to the last  $n$  factors. Then, using Lemma 4.2 and property (3) we can obtain a function  $h: C \rightarrow I^n$  such that

$$h(p) = \eta(\Pi^n(g(p) \cap \pi_1^{-1}(p)))$$

for each  $p \in C$ . Put

$$\tilde{X} = \{(p, h(p)) : p \in C\}.$$

Then, from (2) and (3) we have

$$(4) \quad |T \cap \tilde{X}| = c \text{ for each } T \in \mathcal{T}.$$

To see that  $h$  satisfies (0), let  $\{B_i\}$  be a sequence of clopen disjoint covers of  $(C, \varepsilon_0)$  with its mesh  $1/i$  and we assume that  $B_{i+1}$  refines  $B_i$ . Define a function  $h_k : C \rightarrow I^n$  by the following formula:

$$h_k(t) = \eta(\Pi^n(\cup \{g(p) \cap \pi_1^{-1}(p) : p \in B\}))$$

for each  $t \in B \in B_k$ . Since  $g$  is continuous and  $B$  is clopen, the set  $\cup \{g(p) \cap \pi_1^{-1}(p) : p \in B\}$  is closed in  $I^{n+1}$ . Hence  $h_k : (C, \varepsilon_0) \rightarrow I^n$  is well-defined and is continuous. By Lemma 4.2(b) we can see that  $h$  is the limit function of  $\{h_k\}$ , so that  $h$  is a Baire function of class 1, and hence  $X$  is a  $G_\delta$ -set in  $I^{n+1}$  by Lemma 4.1.

When  $n = 1$ , one can show without difficulty that  $h$  is lower semi-continuous, since  $\{h_k\}$  is monotone increasing in this case by the definition (\*) in Lemma 4.2.



To verify the property (1), let

$A_k = \tilde{X} \cap \pi_{k+1}^{-1} [0, 1/7]$  and  $B_k = \tilde{X} \cap \pi_{k+1}^{-1} [6/7, 1]$  for each  $k = 1, \dots, n$ . Then, each  $A_k$  and  $B_k$  are non-empty, since  $\pi_{k+1}^{-1}(0)$  and  $\pi_{k+1}^{-1}(1)$  are continua meeting both  $\tilde{A}_{n+1}$  and  $\tilde{B}_{n+1}$ . Let  $S_k$  be a separation of  $A_k$  and  $B_k$  for each  $k$  and let  $\tilde{S}_k$  be a separation of

$$\tilde{A}_k = \pi_{k+1}^{-1}(0) \text{ and } \tilde{B}_k = \pi_{k+1}^{-1}(1)$$

which extends  $S_k$  (that is,  $\tilde{S}_k$  separates  $\tilde{A}_k$  and  $\tilde{B}_k$  in  $I^{n+1}$  and  $\tilde{S}_k \cap \tilde{X} \subset S_k$ ). Then, since  $\cap \tilde{S}_k$  contains a continuum meeting both  $\tilde{A}_{n+1}$  and  $\tilde{B}_{n+1}$  by Rubin-Schori-Walsh [1], there exists a  $T \in \mathcal{T}$  such that

$$T \subset \cap \tilde{S}_k \text{ and } T \cap \tilde{X} \subset \cap \tilde{S}_k \cap \tilde{X} \subset \cap S_k.$$

From (4)  $|\tilde{X} \cap T| = c$  so that  $|\cap S_k| = c$ , and hence for the case  $n < \infty$  the theorem has been established.

For the case  $n = \infty$  we replace  $I^{n+1}$  by the Hilbert cube  $I^\infty$ . Then, the construction goes on analogously.

REMARK 4.1. (a) The above theorem is a slight improvement of Tsuda [1, Theorem 2.1]. Since  $(C, \tilde{\rho}(1))$  in our Theorem 4.1 satisfies Theorem 3.1, we believe that our Theorem 4.1 is an  $n$ -dimensional version of Theorem 3.1.

(b) Using Lemma 4.1, Mazurkiewicz [1] has constructed  $n$ -dimensional totally disconnected, separable, complete metric space. The lemma is also used elsewhere efficiently (see Krasinkiewicz [1]).

(c) Comparing our spaces with other spaces used in Engelking-Pol [1] or Charalambous [2], ours have several advantages. In the former paper they used a separable complete metric space due to R. Pol [1] which is not  $A$ -weak infinite dimensional. In their setting their space satisfies our theorem (see Section 5), and we believe that the property (1) is the key point in their case, too. We also note that we cannot apply their method for the finite dimensional case (see Section 5 why), because of lacking the property (1).

On the other hand, though we can obtain his result much easier (see Section 8), we cannot use the space in Charalambous [2] to produce Wage-type examples, because his space is not completely metrizable, and hence we cannot obtain the following natural extension for his space.


We shall embed  $(C, \tilde{\rho}(n))$  in the natural extension  $(C^2, \rho(n))$  as follows: Using the functions  $\{h_k\}$  in the proof of Theorem 4.1, we get a function  $H : C^2 \rightarrow I^n$  and the topology  $\rho(n)$  in the same manner given in § 3. Therefore, we have:

- THEOREM 4.2. For every  $n = 1, 2, \dots, \infty$ ,  $(C^2, \rho(n))$  is a complete separable metric space satisfying
- (1)  $\rho(n)|_{\Delta} = \tilde{\rho}(n)$ , where  $\Delta = \{(x, x) : x \in C\}$ ;
  - (2)  $\rho(n)|_{C^2 \setminus \Delta}$  is homeomorphic to the usual Euclidean topology on it;
  - (3) both topologies  $\rho(n)|_{(\{t\} \times C)}$  and  $\rho(n)|_{(C \times \{t\})}$  are homeomorphic to  $\varepsilon_0$ ;
  - (4) when  $n = 1$ ,  $\rho(n)$  is coarser than both  $\rho(n) \times \varepsilon_0$  and  $\varepsilon_0 \times \rho(n)$ .

REMARK 4.2. (a) The extension space given in Engelking & Pol[1] is some modification of the one used in Wage [1]. Therefore, when  $n = 1$ , our method has an advantage (see §3).

(b) It is an open problem whether or not the  $n$ -fold product  $(C, \rho)^n$  of Wage's metric space  $(C, \rho)$  in Theorem 3.1 satisfies the condition (2) in Theorem 4.1. If it were the case, there could be a space satisfying Theorem 4.2 together with the condition (4) for every  $n$ .

- Some consequences of Kunen construction and those of van Douwen construction.



5. Some consequences of Kunen construction and those of van Douwen construction.

In the following 1) we assume the continuum hypothesis, CH. While, in 2) we do not assume any set-theoretic assumptions beyond ZFC.

1) Some consequences of Kunen construction.

In this place we review the outline of the machine given in Juhász, Kunen & Rudin [1] for refining given topologies. While it is the technique for some general topologies, we specify it for our Wage's metric topologies  $\tilde{p}(n)$  on  $C$ . Let  $\{x_\alpha : \alpha < \omega_1\}$  be a well-ordering of  $C$ , and put  $X_\alpha = \{x_\beta : \beta < \alpha\}$ . Then, we have:

LEMMA 5.1 (CH). There exists an enumeration  $\{S_\alpha : \alpha < \omega_1\}$  of all countable subsets of  $C$  (respectively,  $C^2$ ) such that  $S_\alpha \subset X_\alpha$  (respectively,  $S_\alpha \subset (X_\alpha)^2$ ).

Proof. Let  $\tau = \{T_\alpha : \alpha < \omega_1\}$  be a well-ordering of the collection of all countable subsets of  $C$ . Then, for each  $\alpha$  we define an ordinal  $f(\alpha) < \omega_1$  by the following formula:

$$f(\alpha) = \min \{ \beta : \beta \notin \{f(\gamma) : \gamma < \alpha\} \text{ and } T_\alpha \subset X_\beta \}.$$

Then, since each  $T_\alpha$  is countable,  $f(\alpha)$  is well-defined. It is easy to see that  $f: \omega_1 \rightarrow \omega_1$  is bijective. Put

$$S_\alpha = T_{f^{-1}(\alpha)}.$$

Then, it is easy to see that  $\{S_\alpha: \alpha < \omega_1\}$  is a required well-ordering. The assertion for  $C^2$  follows in a way parallel.

It is shown in Juhász, Kunen & Rudin[1] that (see also § 6) there exists a topology  $\tilde{\nu}$  on  $C$  such that

- (i)  $\tilde{\nu}$  is finer than  $\tilde{\rho}(n)$ ,
- (ii) for each  $\alpha < \beta$ , if  $x_\beta \in \text{Cl}_{\tilde{\rho}(n)}(S_\alpha)$ , then  $x_\beta \in \text{Cl}_{\tilde{\nu}}(S_\alpha)$ .

From (ii) we see (Juhász, Kunen & Rudin[1]) that

- (#) if  $A$  is  $\tilde{\nu}$ -closed, then the set  $\text{Cl}_{\tilde{\rho}(n)}(A) \setminus A$  is countable.

We believe that the above property (#) is one of the most important properties of Kunen construction, so that we conclude this section to see the following lemma, which uses (#) extensively:

LEMMA 5.2.  $\dim(C, \tilde{\nu}) = \text{Ind}(C, \tilde{\nu}) = n$ . Moreover, for the case  $n = \infty$   $(C, \tilde{\nu})$  is strongly infinite dimensional.

Proof. Let  $\{(A_i, B_i) : 1 \leq i \leq n\}$  be the  $n$ -pairs of disjoint  $\tilde{\rho}(n)$ -closed sets in Theorem 4.1 (1). Then, take  $n$ -pairs of  $\tilde{\rho}(n)$ -open sets  $\{(U_i, V_i)\}$  such that

(2)  $A_i \subset U_i$ ,  $B_i \subset V_i$ , and  $M_i \cap N_i = \emptyset$ , where

$$M_i = \text{Cl}_{\tilde{\rho}(n)}(U_i) \text{ and } N_i = \text{Cl}_{\tilde{\rho}(n)}(V_i).$$

Then,  $M_i$  and  $N_i$  are  $\tilde{\nu}$ -closed, since (i) holds. Let  $T_i$  be a closed separation between  $M_i$  and  $N_i$  in  $(C, \tilde{\nu})$ . Then, put  $M_i \subset G_i$ ,  $N_i \subset H_i$ ,  $C \setminus T_i = G_i \cup H_i$ , for some open disjoint sets  $G_i$  and  $H_i$ . Let

$$S_i = \text{Cl}_{\tilde{\rho}(n)}(T_i) \cup (\text{Cl}_{\tilde{\rho}(n)}(G_i) \cap \text{Cl}_{\tilde{\rho}(n)}(H_i)).$$

Then,  $S_i$  is a  $\rho(n)$ -closed separation between  $A_i$  and  $B_i$  from (2). It follows from (#) that

(3) the set  $S_i \setminus T_i$  is countable.

Hence, the set  $\cap S_i$  has the cardinality of continuum from Theorem 4.1 (1). Therefore, the set  $\cap T_i$  also has the cardinality of continuum from (3). Hence,  $\dim(C, \tilde{\nu}) \geq n$ . On the other hand, it follows from Fedorčuk [1, Lemmas 7,9] that  $\dim(C, \tilde{\nu}) \leq \text{Ind}(C, \tilde{\nu}) \leq n$ . This completes the proof.

REMARK 5.1. (a) We follow Engelking & Pol [1, p. 16] to say that the construction given here "Kunen construction". It is also called "Juhász-Kunen-Rudin construction" in E. Pol [1].

(b) As mentioned in Remark 4.1 (c) our condition (1) in Theorem 4.1 is crucial at least for finite dimensional cases. Indeed, let  $(C, \rho)$  be the  $n$ -dimensional space in Charalambous [2] or that of Engelking & Pol [1], which does not necessarily satisfy (1). Then, we cannot

determine exactly the dimension of Kunen construction  $(C, \nu)$  for it. As is shown in Fedorčuk [1] (see also Gruenhage & Pol [1]), we can only say that

$$n-1 \leq \dim (C, \nu) \leq n \text{ and } n-1 \leq \text{Ind} (C, \nu) \leq n.$$

(c) Let  $(\Delta, \rho)$  be the strongly infinite dimensional complete separable metric space given in Engelking & Pol [1]. Then, it is known that its Kunen construction is strongly infinite dimensional (see Gruenhage & Pol [1]). Moreover, we have:

PROPOSITION 5.1. Every strongly infinite dimensional complete metric space satisfies the property (1) in Theorem 4.1.

Proof. Let  $\{(A_i, B_i) : i \in \omega\}$  be a countable collection of pairs of disjoint closed sets satisfying (4) if a closed set  $S_i$  separates  $A_i$  and  $B_i$  for each  $i$ , then  $\bigcap_{i \geq 1} S_i \neq \emptyset$ .

We show that the collection  $\{(A_{i+1}, B_{i+1}) : i \in \omega\}$  satisfies (1) in Theorem 4.1. Indeed, let  $S_i$  be a closed separation between  $A_i$  and  $B_i$ . We must show that the set  $\bigcap_{i \geq 1} S_i$  has the cardinality of continuum.

On the contrary, assume that  $|\bigcap_{i \geq 1} S_i| < c$ . Then,  $\bigcap_{i \geq 1} S_i$  is countable, since every uncountable complete separable metric space has the cardinality of continuum. Hence,



$\dim \left( \bigcup_{i \geq 1}^{\omega} S_i \right) \leq 0$ , and therefore, there exists a separation  $S_0$  between  $A_0$  and  $B_0$  which does not meet the set  $\bigcup_{i \geq 1}^{\omega} S_i$ . Then, the collection  $\{S_i : i \in \omega\}$  does not satisfy (4). This contradiction completes the proof.

2) Some consequences from van Douwen construction.

In this place we review the techniques of van Douwen [1], which does not require the continuum hypothesis. As in the preceding section, we specify the construction for our Wage's metric topologies  $\tilde{\rho}(n)$ , though it can be applied for some general topologies (see also § 8).

One of the key points of van Douwen construction is

- (i) it is a topology  $\tilde{\tau}$  on  $C$  finer than  $\tilde{\rho}(n)$ ,
- (ii) if  $A$  and  $B$  are disjoint  $\tau$ -closed sets, then the set  $Cl_{\tilde{\rho}(n)}(A) \cap Cl_{\tilde{\rho}(n)}(B)$  is countable.

Though the property (ii) is weaker than (ii) in Kunen's, we can deduce from it that  $(C, \tilde{\tau})$  is countably paracompact, collectionwise normal (see §10). Here, we shall show the following lemma, using (ii):

LEMMA 5.3 (Tsuda [1, Lemma 5.6]). When  $n = \infty$   $(C, \tilde{\tau})$  is not strongly countable dimensional.

Proof. If we assume that  $(C, \tilde{\tau})$  is strongly countable-dimensional, there exists a countable collection  $\{H_i\}$  of closed subspaces of  $(C, \tilde{\tau})$  such that  $C = \cup H_i$ , and  $\dim H_i \leq m_i$  for some integer  $m_i$  for each  $i$ . By Theorem 4.1 there exists a countable collection  $\{(A_i, B_i)\}$  of pairs of disjoint  $\tilde{\rho}(n)$ -closed sets such that

(1) if for each  $i$ ,  $\tilde{\rho}(n)$ -closed set  $S_i$  separates  $A_i$  and  $B_i$ , then  $| \cap S_i | = c$ .

Take disjoint  $\tilde{\rho}(n)$ -open sets  $U_i$  and  $V_i$  satisfying Lemma 5.2 (2). Then, since  $\dim H_i \leq m_i$ , for each  $m_i+1$  pairs

$$\{(M_k, N_k) : i + \sum_{j=1}^{i-1} m_j \leq k \leq i + \sum_{j=1}^i m_j\},$$

there exists a zero set separation  $T_k$  of  $H_i \cap M_k$  and  $H_i \cap N_k$  in  $H_i$  such that

$$(2) \quad \cap \{T_k : i + \sum_{j=1}^{k-1} m_j \leq k \leq i + \sum_{j=1}^i m_j\} = \emptyset.$$

By (i) we have a  $\tilde{\rho}(n)$ -closed separation  $\tilde{S}_k$  of  $A_k \cap H_i$  and  $B_k \cap H_i$  such that

$$(3) \quad |\tilde{S}_i \setminus T_i| \leq \omega, \text{ and } \tilde{S}_i \subset (C \setminus (M_i \cup N_i)).$$

By the hereditarily normality of  $\tilde{\rho}(n)$ , there exists a  $\tilde{\rho}(n)$ -closed separation  $S_k$  of  $A_k$  and  $B_k$  such that

$$\tilde{S}_k = S_k \cap H_i. \text{ Then, from (1) } | \cap S_k | = c, \text{ and from (3)}$$

$| \cap \tilde{S}_k | = c$ . Since  $C = \cup H_i$ , there exists an  $H_i$  such that  $|H_i \cap (\cap S_k)| = c$ . On the other hand, for each

$k$  such that  $i + \sum_{j=1}^{i-1} m_j \leq k \leq i + \sum_{j=1}^i m_j$  we have

$$|\tilde{S}_k \cap H_i \setminus T_k| \leq \omega \text{ from (2), and hence from (3) we have}$$

$|H_i \cap (\cap S_k)| \leq \omega$ . This contradiction shows that  $(C, \tilde{\nu})$  is not strongly countable-dimensional.


REMARK 5.1. Recently, we learned in Engelking & Pol [1] that every normal strongly countable-dimensional space is A-weakly infinite-dimensional. Moreover, it is known that (van Douwen [2, Theorem 3.4 (a)]) a normal space which is the union of countably many closed A-weak infinite dimensional subspaces is A-weak infinite dimensional. Therefore, our lemma also follows from their results.

By a way parallel in the above lemma we have:

LEMMA 5.4.  $\dim (C, \tilde{\tau}) = \text{Ind} (C, \tilde{\tau}) = n$  when  $n < \infty$ .

6. Wage's factorization technique.

34: S884



## 6. Wage's factorization technique.

In this section we approximate  $\rho(n)$  in §4 by the method of Wage suggested in Wage [1,2]. (Note that we follow the improved method due to Przymusiński [5]. See also Remark 6.1 (a).) Our aim is to show that there exist two topologies  $\nu(i)$  on  $C$  such that

- (i)  $\nu = \nu(1) \times \nu(2)$  is finer than  $\rho(n)$ , and
- (ii) if  $A$  and  $B$  are disjoint  $\nu$ -closed sets, then

$$|Cl_{\rho(n)}(A) \cap Cl_{\rho(n)}(B)|_2 \leq \omega, \text{ where } |S|_2$$

denotes the 2-cardinality of the set  $S \subset C^2$   
defined in Przymusiński [2].

Since we want to make our factors same (that is,  $X = Y$ ), we also make our topologies satisfy

- (iii) if  $A$  and  $B$  are disjoint  $\nu(i)^2$ -closed sets, then

$$|Cl_{\varepsilon_0^2}(A) \cap Cl_{\varepsilon_0^2}(B)|_2 \leq \omega.$$

To achieve  $\nu$  satisfying (i)-(iii), we need the following lemma, which is also used for other examples in §§10,11.

LEMMA 6.1. Let  $G_m$  be the  $(1/m)$ -neighborhood of  $z_0 = (x_0, x_0)$  with respect to the metric  $\rho(n)$ , and let  $D$  be a countable  $\rho(n)$ -dense set in  $C$ . Let  $(r_m)$  and  $(z_m = (s_m, t_m))$  be two sequences convergent to the point  $z_0$  in  $\rho(n)$  and in  $\varepsilon_0^2$ , respectively. Then, there exist two sequences  $(x_m), (y_m)$  in  $C$  and three sequences of decreasing  $\varepsilon_0$ -clopen sets

$\{H_m\}, \{U_m\}, \{V_m\}$  such that

- (1)  $(x_0, y_m) \in H_m \times V_m \subset G_m, (x_m, x_0) \in U_m \times H_m \subset G_m;$
- (2)  $D^2 \cap (X_m \times Y_m)$  and  $R_m \cap (X_m \times Y_m)$  are non-empty;
- (3)  $Z_m \cap X_m^2, Z_m \cap Y_m^2$  are also non-empty,

where  $P_m$  denotes the set  $\{p_k : k \geq m\}$ , for a sequence  $(p_k)$ .

Proof. We show the assertion by induction on  $k$ , and suppose that we have chosen  $x_i, y_i, H_i, U_i, V_i$ , for  $i < 6k$ . We shall choose them for all  $m$  satisfying  $6k \leq m < 6(k+1)$ . By Thoerm 4.2(3) we can always choose  $\varepsilon_0$ -clopen set  $H_m' \subset H_{m-1}$  such that

$$H_m' \times \{x_0\} \cup \{x_0\} \times H_m' \subset G_m.$$

(a) For  $m = 6k$  (respectively,  $m = 6k+1$ ) we define  $x_m$  and  $y_m$  (respectively,  $x_{m+1}$  and  $y_{m+1}$ ) by taking any point  $r_k = (x_m, y_m) \in R_m \cap G_m \cap (H_m')^2$  (respectively, any point  $(x_m, y_m) \in G_m \cap D^2 \cap (H_m')^2$ ), since  $(r_k)$  is  $\rho(n)$ -converging (respctively,  $D^2$  is  $\rho(n)$ -dense). Then, take any  $H_{m+i}, U_{m+i}, V_{m+i}$  satisfying (1),  $i = 0, 1$ .

(b) For  $m = 6k+2$ , we define  $x_m, x_{m+1}$  by taking any point  $z_k = (x_m, x_{m+1}) \in Z_m \cap (H_m')^2$ . Take  $\varepsilon_0$ -clopen sets  $H_{m+i}, U_{m+i}$  such that  $H_m' \supset H_m \supset H_{m+1}$ ,  $(x_{m+i}, x_0) \in U_{m+i} \times H_{m+i} \subset G_{m+i}$ . Take any point  $y_{m+i} = t_k \in H_{m+i}$ , since  $(t_k)$  is  $\varepsilon_0$ -converging to  $x_0$ . Let  $V_{m+i}$  be any  $\varepsilon_0$ -clopen set satisfying (1),  $i = 0, 1$ .

(c) For  $m = 6k+4$ , define  $(y_m, y_{m+1})$  by taking any point  $z_k = (y_m, y_{m+1}) \in Z_m \cap (H_m')^2$ . Take  $\varepsilon_0$ -clopen sets  $H_{m+i}, V_{m+i}$  such that  $H_m' \supset H_m \supset H_{m+1}$ ,  $(x_0, y_{m+i}) \in H_{m+i} \times V_{m+i} \subset G_{m+i}$ . Then, take any point  $x_{m+i} = s_k \in H_{m+i}$ , since  $(s_k)$  is  $\varepsilon_0$ -converging to  $x_0$ . Let  $U_{m+i}$  be any  $\varepsilon_0$ -clopen set satisfying (1),  $i = 0, 1$ .

This completes the proof of our lemma.

In the sequel, we assume CH and show that there is a topology satisfying (i)-(iii).

Since we assume CH, let  $\{x_\alpha : \alpha < \omega_1\}$  be a well-ordering of  $C$  and let

$$X_\alpha = \{x_\beta : \beta < \alpha\}.$$

It follows from Lemma 5.1 that the family of all countable subsets  $\{A_\alpha : \alpha < \omega_1\}$  of  $C^2$  can be well-ordered so that

$$A_\alpha \subset (X_\alpha)^2.$$

For each  $x_\alpha$  we define two neighborhood bases  $\{N_m(x_\alpha)_i\}$

$i = 1, 2$ , so that the following inductive assumptions

(0)-(3) are satisfied:

(0)  $N_m(x_\alpha)_i \cap D \neq \emptyset$  for each  $m, i$ , where  $D$  is a fixed countable  $\tilde{\rho}(n)$ -dense set.

(1) The  $\rho(n)$ -diameter of  $B_m = N_m(x_\alpha)_1 \times N_m(x_\alpha)_2$  is less than  $1/m$ .

- (2) The  $\varepsilon_0$ -diameter of  $N_m(x_\alpha)_i$  is less than  $1/m$ ,  
 $N_m(x_\alpha)_i$  is a compact set consisting of countable points, and for any  $y \in N_m(x_\alpha)_i$  there exists an integer  $k$  such that  $N_k(y)_i \subset N_m(x_\alpha)_i$ .
- (3) Let  $\nu(i, \alpha)$  be the topology on  $X_\alpha$  whose local base is  $\{N_m(x_\beta)_i : \beta < \alpha \text{ and } m \in \omega\}$ . Then, for every  $i = 1, 2$ , and  $\beta, \gamma < \alpha$
- (a) if  $(x_\alpha, x_\alpha) \in Cl_{\rho(n)}(A_\gamma)$ , then  
 $A_\gamma \cap B_m \neq \emptyset$  for any  $m \in \omega$ ;
- (b) if  $(x_\alpha, x_\alpha) \in Cl_{\varepsilon_0^2}(A_\gamma)$ , then  
 $A_\gamma \cap (N_m(x_\alpha)_i)^2 \neq \emptyset$  for any  $m \in \omega$ ;
- (c) if  $(x_\alpha, x_\beta) \in Cl_{\varepsilon_0 \times \nu(i, \alpha)}(A_\gamma)$ , then  
 $A_\gamma \cap B_m \neq \emptyset$  for any  $m \in \omega$ ;
- (d) if  $(x_\beta, x_\alpha) \in Cl_{\nu(i, \alpha) \times \varepsilon_0}(A_\gamma)$ , then  
 $A_\gamma \cap B_m \neq \emptyset$  for any  $m \in \omega$ .

Suppose that two such neighborhood bases have been constructed for each  $x_\beta$  such that  $\beta < \alpha$ .

Let  $R_\alpha^1$  be the collection of  $D$  and all  $A_\gamma$  such that  $\gamma < \alpha$  and  $(x_\alpha, x_\alpha) \in Cl_{\rho(n)}(A_\gamma)$ .

Let  $R_\alpha^2$  be the collection of all  $A_\gamma$  such that  $\gamma < \alpha$ ,  $A_\gamma \notin R_\alpha^1$ , and  $(x_\alpha, x_\alpha) \in Cl_{\varepsilon_0^2}(A_\gamma)$ .

Let  $s_\alpha^i$  ( $i = 1, 2$ ) be the collection of all  $A_\gamma$  such that  $\gamma < \alpha$ , and  $(x_\alpha, x_\beta) \in Cl_{\varepsilon_0 \times \nu(i, \alpha)}(A_\gamma)$  for some  $\beta < \alpha$ .

Let  $\tau_\alpha^i$  ( $i = 1, 2$ ) be the collection of all  $A_\gamma$  such that  $\gamma < \alpha$ , and  $(x_\beta, x_\alpha) \in Cl_{\nu(i, \alpha) \times \varepsilon_0}(A_\gamma)$  for some  $\beta < \alpha$ .



Put

$$K_\alpha = \bigcup_{i=1}^2 R_\alpha^i \cup S_\alpha^i \cup T_\alpha^i.$$

Then, take a convergent set  $\lambda(A)$  for each  $A \in K_\alpha$ , which converges to one of the points  $(x_\alpha, x_\alpha)$ ,  $(x_\alpha, x_\beta)$ , and  $(x_\beta, x_\alpha)$  according to the collection it belongs to. Put

$$\zeta(A) = \pi_1(\lambda(A)) \text{ and } \xi(A) = \pi_2(\lambda(A)),$$

where  $\pi_i: C^2 \rightarrow C$  is the projection to the  $i$ -th factor.

Then, by Lemma 6.1 we can choose a sequence  $\langle (x_m, x_m) \rangle$  converging to  $(x_\alpha, x_\alpha)$ , which satisfies the following (4):

(4) Let  $G_m$  be the  $(1/m)$ -neighborhood of  $(x_\alpha, x_\alpha)$  with respect to the metric  $\rho(n)$ . Then, there exist

$\varepsilon_m$ -clopen sets  $H_m$ ,  $U_m$ , and  $V_m$  such that

(a)  $(x_m, y_m) \in G_m$ ,  $(x_\alpha, y_m) \in H_m \times V_m \subset G_m$ , and

$(x_m, x_\alpha) \in U_m \times H_m \subset G_m$ ;

(b)  $H_m \supset H_{m+1}$  and  $U_m \supset U_{m+1}$ ,  $V_m \supset V_{m+1}$ ;

(c)  $x_m \neq x_k$  and  $y_m \neq y_k$  for  $m \neq k$ ;

(d) let  $\lambda = \{(x_m, y_m)\}$ ,  $\zeta = \{x_m\}$ ,  $\xi = \{y_m\}$ , then

(1)  $\lambda \cap \lambda(A) = \emptyset$  for each  $A \in R_\alpha^1$ ,

(2)  $\zeta^2 \cap \lambda(A) \neq \emptyset$  and  $\xi^2 \cap \lambda(A) \neq \emptyset$  for each  $A \in R_\alpha^2$ ,

(3)  $\zeta \cap \zeta(A) \neq \emptyset$  for each  $A \in S_\alpha^i$ ,

(4)  $\xi \cap \xi(A) \neq \emptyset$  for each  $A \in T_\alpha^i$ ;

(e) when  $S_\alpha^i = \emptyset$  or  $T_\alpha^i = \emptyset$ , we put  $U_m = V_m = \emptyset$ , and we do not define  $x_m$ ,  $y_m$  for the corresponding  $m$ .

Since  $x_m, y_m < x_\alpha$ , there exist two integers  $i(m)$  and  $j(m)$  by our inductive assumptions such that

$$N_{i(m)}(x_m)_1 \subset U_m, N_{j(m)}(y_m)_2 \subset V_m, \text{ and} \\ N_{i(m)}(x_m)_1 \times N_{j(m)}(y_m)_2 \subset G_m.$$

Put

$$N_m(x_\alpha)_1 = \{x_\alpha\} \cup (\cup \{N_{i(k)}(x_k)_1 : k \geq m\}),$$

and

$$N_m(x_\alpha)_2 = \{x_\alpha\} \cup (\cup \{N_{j(k)}(y_k)_2 : k \geq m\}).$$

Then, we shall show that (1) holds. By our construction

$$L_1 = \bigcup_{k \geq m} N_{i(k)}(x_k)_1 \times N_{j(k)}(y_k)_2 \subset \bigcup_{k \geq m} G_k,$$

$$L_2 = N_{i(m)}(x_m)_1 \times (\bigcup_{k > m} N_{j(k)}(y_k)_2) \subset N_{i(m)}(x_m)_1 \times H_m,$$

$$L_3 = \bigcup_{k > m} (N_{i(k)}(x_k)_1 \times (\bigcup_{m \leq s < k} N_{i(s)}(y_s)_2)) \subset \bigcup_{m \leq s < k} H_s \times V_s,$$

and

$$L_4 = \bigcup_{k > m} (N_{i(k)}(x_k)_1 \times (\bigcup_{s > k} N_{j(k)}(y_k)_2)) \subset \bigcup_{k > m} U_k \times H_k.$$

Therefore, (1) holds, since  $B_m = \{(x_\alpha, x_\alpha)\} \cup (\bigcup_{i=1}^4 L_i)$ .

It is easy to see that  $\{N_m(x_\beta)_i : \beta \leq \alpha, m \in \omega\}$ ,  $i = 1, 2$ ,

satisfy all the remaining conditions. Thus, our

inductive construction is completed. Let  $\nu(i)$  be the

topology whose local base is  $\{N_m(x)_i : x \in C\}$ . Then,

they satisfy that

$$(5) \quad \nu = \nu(1) \times \nu(2) \text{ is finer than } \rho(n);$$

and

$$(6) \quad |Cl_{\rho(n)}(A) \setminus A|_2 \leq \omega \text{ for any } \nu\text{-closed set } A.$$

From (3)(b)

$$(7) \quad |Cl_{\varepsilon_0^2}(A) \setminus A|_2 \leq \omega \text{ for any } \delta_i^2\text{-closed set } A,$$

where  $\delta_i = \nu(i)$ ,  $i = 1, 2$ .


REMARK 6.1. (a) In the above construction we use the method due to Przymusiński [5], since the procedure suggested in Wage [1] is not sufficient (see, Remark 9.1).

(b) There are some applications of Wage's factorization technique outside of dimension theory (Rudin [2], Wage [1,2]).

(c) Be careful that there are other famous "factorization" theorems in dimension theory (for example, that of Mardešić [1] and that of Pasynkov [3]). The meaning "factorization" in there is different from ours.

7. Przymusiński's method.

S885



## 7. Przymusiński's method.

In this section we shall define topologies  $\tau_i$  on  $C$  so that  $\tau = \tau_1 \times \tau_2$  approximates  $\rho(n)$ . Our method is due to Przymusiński [3], which eliminated the continuum hypothesis from the argument in the preceding section, and is based on a technique of van Douwen instead of Kunen.

For our inductive construction we need the following subset  $P$  of  $C^2$ . Let  $F = \{(A, B)_\lambda : \lambda \in \Lambda\}$  be the collection of all pairs of countable subsets of  $C^2$  such that

$$(0) \quad |Cl_{\rho(n)}(A) \cap Cl_{\rho(n)}(B)|_2 > \omega.$$

Note that for any  $(A, B)_\lambda \in F$  we have by Przymusiński [2]

$$|Cl_{\rho(n)}(A) \cap Cl_{\rho(n)}(B)|_2 = c.$$

In the sequel let  $<$  be a well-ordering of  $C$  and let fix a  $\rho(n)$ -dense countable set  $D$ . Then, define for  $x \in C$

$$I(x) = \{y \in C : y < x\} \cup D.$$

Then, for each  $\lambda \in \Lambda$ , since  $|A \cup B| = \omega$ , there exists an  $x_\lambda \in C$  such that  $A \cup B \subset I(x_\lambda)^2$ , where  $(A, B) = (A, B)_\lambda$ .

Hence, we can construct a transfinite sequence  $\langle p_\lambda \rangle$  of points in  $C^2$  satisfying

$$p_\lambda \in Cl_{\rho(n)}(A) \cap Cl_{\rho(n)}(B) \setminus I(x_\lambda)^2, \text{ where } (A, B) = (A, B)_\lambda,$$

and

if  $\lambda \neq \mu$ , then  $\hat{p}_\lambda \cap \hat{p}_\mu = \emptyset$ , where  $\hat{p}$  denotes the set  $\{p_1, p_2\}$  of  $C$  for  $p = (p_1, p_2)$ . Put

$$P = \{p_\lambda : \lambda \in \Lambda\}.$$

Since  $D$  is a countable  $\rho(n)$ -dense set of  $C$ , in a way parallel in the preceding section we can choose a sequence  $\langle (x_m, y_m) \rangle$  for each  $p \in P$ , satisfying the following condition (1).

- (1) Let  $G_m$  be the  $(1/m)$ -neighborhood of  $p = p_\lambda = (x_0, y_0)$  with respect to  $\rho(n)$ . Then,
  - (a)  $(x_m, y_m) \in G_m$ ,  $x_m \neq x_k$  and  $y_m \neq y_k$  for  $m \neq k$ , and  $\{(x_{3m}, y_{3m})\} \subset A$ ,  $\{(x_{3m+1}, y_{3m+1})\} \subset B$ , where  $(A, B) = (A, B)_\lambda$  and  $\{(x_{3m+2}, y_{3m+2})\} \subset D$ ;
  - (b) there exist  $\epsilon_0$ -clopen sets  $H_m, K_m, U_m$ , and  $V_m$  such that  $p \in H_m \times K_m$ ,  $(x_0, y_0) \in H_m \times V_m \subset G_m$ , and  $(x_m, y_0) \in U_m \times K_m \subset G_m$ ;
  - (c)  $H_m \supset H_{m+1}$ ,  $K_m \supset K_{m+1}$ ,  $H_m \supset U_{m+1}$ , and  $K_m \supset V_{m+1}$ .

Now, by transfinite induction on  $\alpha$  we assign to each point of  $C$  two neighborhood bases  $\{B_m(x)_i\}$ ,  $i = 1, 2$  so that the following inductive assumptions (2) and (3) are satisfied.

- (2) Each  $B_m(x)_i$  is a compact set consisting of countable points, and for any  $y \in B_m(x)_i$  there exists a  $k$  such that  $B_m(x)_i \supset B_k(y)_i$ .
- (3) The diameter of  $B_m = B_m(x)_1 \times B_m(x)_2$  is less than  $1/m$  with respect to the metric  $\rho(n)$ .

Let each point of  $D$  be isolated, and suppose that  $x \in C$  and such bases of neighborhoods have been constructed for each  $y < x$ . Then, we consider the following cases.

- (a)  $x = p_1 = p_2$  for some (unique)  $p \in P$ .
- (b)  $x = p_i$  for some (unique)  $p \in P$  and  $p_1 \neq p_2$ .
- (c)  $x \neq p_i$  for any  $p \in P$  and  $i$ .

In the cases (a) and (b) we have  $x_m, y_m < p_i$  for  $i = 1, 2$  and  $m \in \omega$ , because  $p_\lambda \notin I(x_\lambda)^2$  and  $x_m, y_m \in I(x_\lambda)$ . Thus inductive assumptions can be applied so that there exist integers  $k(m)$  for each  $m$  such that

$$B_{k(m)}(x_m)_1 \subset U_m, \quad B_{k(m)}(y_m)_2 \subset V_m, \quad \text{and}$$

$$B_{k(m)}(x_m)_1 \times B_{k(m)}(y_m)_2 \subset G_m.$$

The last inclusion follows from the following observation.

If  $x_m = y_m$ , then the inclusion follows from (3). If  $x_m \neq y_m$ , then  $(x_m, y_m) \in \text{Int}_{\varepsilon_0^2}(G_m)$ . Therefore, the inclusion follows from the fact that both  $x_m$  and  $y_m$  have neighborhood bases finer than  $\varepsilon_0$ .

Case (a). Put for each  $i = 1, 2$  and  $m \in \omega$

$$B_m(x)_i = \{x\} \cup \left( \bigcup_{t \geq m} B_{k(t)}(x_t)_i \right).$$

Case (b). If  $x = p_1$ , put for each  $m \in \omega$

$$B_m(x)_1 = \{x\} \cup \left( \bigcup_{t \geq m} B_{k(t)}(x_t) \right), \quad \text{and} \quad B_m(x)_2 = \{x\}.$$

If  $x = p_2$ , put for each  $m \in \omega$

$$B_m(x)_1 = \{x\}, \quad \text{and} \quad B_m(x)_2 = \{x\} \cup \left( \bigcup_{t \geq m} B_{k(t)}(y_t)_2 \right).$$


Case (c). Put  $B_m(x)_i = \{x\}$  for each  $i = 1, 2$  and  $m \in \omega$ .

Because there does not exist any decreasing chains in a well-ordered set and (1) holds, we can show that  $\{B_m(x)_i\}$  satisfies (2) and (3) for each  $i$  in a way parallel in the preceding section. Thus, inductive construction has completed.

From (2) and (3)  $\{B_m(x)_i : m \in \omega, x \in C\}$  constitutes a local base on  $C$  and it produces a Hausdorff topology  $\tau_i$  for each  $i = 1, 2$ . This completes the constructions of  $\tau_1$  and  $\tau_2$ .



8. An application to inverse sequences.



## 8. An application to inverse sequences.

The purpose of this section is to give some application of the Wage's metric spaces in §4 and of the technique of van Douwen in §5. We begin with the following theorem due to Charalambous [1]:

THEOREM 8.1. There exists an inverse sequence  $\{X_i, f_{ij}\}$  with the limit space  $X$  such that each  $X_i$  is 0-dimensional, first countable, separable, Lindelöf, while  $X$  is normal with  $\dim X = 1$ .

REMARK 8.1. (a) We do not give here the proof of the theorem, since we shall show an  $n$ -dimensional version of it below which contains the above result as its corollary.

(b) Before Charalambous it is already known in Engelking [1, Problem 6.3.25] that

(#) the limit of an inverse sequence of strongly

0-dimensional spaces need not be strongly 0-dimensional.

Comparing these two constructions, Charalambous construction has the advantage of nice factor spaces  $X_i$ .

On the other hand, by Engelking's construction we have a sequence with a nice limit space: there exists an inverse sequence, which satisfies (#), and whose limit is metrizable.

Indeed, let  $X$  be the Roy's 1-dimensional space. Since  $X$  has a clopen base of cardinality of continuum  $c$ , we can assume that  $X$  is a subspace of the generalized Cantor-set  $C^C$ . Take a suitable sequence  $\{x_i\}$  of points in  $C^C$  such that the sequence  $\{\Sigma(x_i)\}$  of  $\Sigma$ -products with base point  $x_i$  is mutually disjoint. Put

$$X_i = X \cup \{\Sigma(x_k) : k \geq i\}.$$

Then, each  $X_i$  is strongly 0-dimensional, since  $X_i$  is  $C^*$ -embedded in  $C^C$  (Engelking [1, Problem 3.11.23(c)]).

Let  $f_{ij} : X_j \rightarrow X_i$  be the inclusion map. Then, it is readily seen that  $X$  is the limit of the inverse sequence  $\{X_i, f_{ij}\}$ .

**THEOREM 8.2.** For every  $k = 1, 2, \dots, \infty$  there exists an inverse sequence  $\{X_i, f_{ij}\}$  with the limit space  $X$  such that each  $X_i$  is 0-dimensional, first countable, Lindelöf, while  $X$  is <sup>(separable)</sup> countably paracompact, collection-wise normal with  $\dim X = \text{Ind } X = k$ .

Proof. Let  $(C, \tilde{\rho}(k))$  be the  $k$ -dimensional Wage's metric space in Theorem 4.1. Then, take a sequence  $\{E_i : i \in \omega\}$  of pairwise disjoint Bernstein sets satisfying that

- (1) the set  $A \cap E_i$  has the cardinality of continuum for any  $\tilde{\rho}(k)$ -closed uncountable set  $A$ .

Let  $\tau_i$  be the approximation of the topology  $\mu_i = \tilde{p}(k)|E_i$  on  $E_i$  due to van Douwen (see §5, 2). Thus,

- (2)  $\tau_i$  is first countable, separable, having a clopen base, collectionwise normal, countably paracompact, and the set  $Cl_{\mu_i}(A) \cap Cl_{\mu_i}(B)$  is countable whenever  $A$  and  $B$  are disjoint  $\tau_i$ -closed sets.

Now, we shall define topologies  $\nu_i$  on  $C$ , which is generated by the subbase  $\varepsilon_0 \cup \{\tau_j: j \leq i\}$ . Let  $\nu_\infty$  be the topology on  $C$  which is generated by  $\{\tau_j: j \in \omega\}$ . Put for each  $i$

$$X_i = (C, \nu_i), \text{ and let}$$

$$f_{ij}: X_j \rightarrow X_i \text{ be the identity function.}$$

Then, by (1) and (2) each  $X_i$  is first countable, 0-dimensional, separable, and Lindelöf. By the definition,  $f_{ij}$  is continuous. Let  $X_\infty$  be the limit of the inverse sequence  $\{X_i, f_{ij}\}$ . Then, it can be seen from the construction that

$$X_\infty \cong (C, \nu_\infty) \cong \overset{\text{separable}}{\bigoplus_{i=1}^{\infty}} (E_i, \nu_i).$$

Hence, by (2)  $X_\infty$  is countably paracompact, collectionwise normal, and  $\dim X_\infty = \text{Ind } X_\infty = k$  by the proof of Lemma 5.4.


REMARK 8.2. (a) The above construction is a slight improvement of the one given in Tsuda [1] (separability is not achieved there). In Charalambous [2] similar examples are given, independently. Comparing ours and his construction, we believe that our construction is much simpler. For example, we can show that the limit space is collectionwise normal and countably paracompact much easier than he does in Charalambous [2].

(b) When we consider more special case when the limit space is  $X^\omega$  and its factors are  $X^i$ , we have no Wage-type example from the following theorem due to Nagami.

THEOREM 8.1 (Nagami [5]). Let each  $X_i$  be a normal space with  $\dim X_i \leq n$ . Let its limit  $X_\infty$  be countably paracompact and each bonding map  $f_{ij}$  be open. Then,  $X_\infty$  is a normal space with  $\dim X_\infty \leq n$ .

9. Perfect normality of products.

7



## 9. Perfect normality of products.

We shall show the following theorem under continuum hypothesis, CH:

THEOREM 9.1 (CH). For every  $n = 1, 2, \dots, \infty$ , there exists a space  $X(n)$  such that

$$\dim X(n)^2 = \text{Ind } X(n)^2 = n > \dim X(n) = 0,$$

and  $X(n)^2$  is perfectly normal, separable, first-countable, and locally compact. Moreover,  $X(\infty)^2$  is not countable dimensional.

Proof. Put

$$X(n) = (C, \nu(1)) \oplus (C, \nu(2)),$$

where  $\nu(i)$  is defined in §6. Hence,  $X(n)^2$  is separable, first-countable, locally compact, regular and 0-dimensional by the definition. Let us show that at first  $X(n)^2$  is perfectly normal. We begin with the following lemma essentially due to Kunen.

LEMMA 9.1. For every  $i$  and every subset  $A$  of  $C$  there exists a  $\gamma < \omega_1$  such that

$$\text{if } \alpha \geq \gamma \text{ and } x_\alpha \in \text{Cl}_{\varepsilon_0}(A), \text{ then } x_\alpha \in \text{Cl}_{\nu(i)}(A).$$

LEMMA 9.2. For every  $A \subset \mathbb{C}^2$  there exists a  $\lambda < \omega_1$  such that if  $\alpha \neq \beta$ ,  $\alpha, \beta \geq \lambda$ , and  $(x_\alpha, x_\beta) \in Cl_{\varepsilon_0^2}(A)$ , then  $(x_\alpha, x_\beta) \in (Cl_{\nu(k) \times \varepsilon}(A)) \cap (Cl_{\varepsilon \times \nu(k)}(A))$ ,  $k = 1, 2$ .

LEMMA 9.3. For every  $A \subset \mathbb{C}^2$  there exists a  $\gamma$  such that if  $(x_\alpha, x_\beta) \in Cl_{\rho(n)}(A)$  and  $\alpha, \beta \geq \gamma$ , then  $(x_\alpha, x_\beta) \in Cl_\nu(A)$ .

LEMMA 9.4. For every  $i$  and  $A \subset \mathbb{C}^2$  there exists a  $\gamma$  such that if  $(x_\alpha, x_\beta) \in Cl_{\varepsilon_0^2}(A)$  and  $\alpha, \beta \geq \gamma$ , then  $(x_\alpha, x_\beta) \in Cl_{\nu(i)^2}(A)$ .

Proof of Lemma 9.1. Let  $L$  be a countable Euclidean dense subset of  $A$ . Then, by (3) (a) in §6, if  $x_\alpha \in Cl_{\varepsilon_0}(A)$  with  $\alpha > \gamma$ , then  $(x_\alpha, x_\alpha) \in Cl_{\rho(n)}(A^2) = Cl_{\rho(n)}(L^2)$ . Since  $L^2 = A_\gamma$  for some  $\gamma$ , each neighborhood of  $x_\alpha$  in  $\nu(i)$  contains a subset of  $L$  by the definition of  $\nu(i)$  and (3) (a) in §6.

The proofs for the remaining lemmas follow from (5), (6) in §6 and by ways parallel to Przymusiński [5, Lemmas 1,2,3].

Now, we shall show that  $X(n)^2$  is perfectly normal. At first, we show that  $X(n)$  is perfectly normal by a method due to Juhász, Kunen and Rudin [1].



It suffices to see that each  $\nu(i)$  is perfectly normal. Let  $A$  be a  $\nu(i)$ -closed set. Then, the set  $\text{Cl}_{\varepsilon_0}(A) \setminus A$  is countable by Lemma 9.1. Since  $\text{Cl}_{\varepsilon_0}(A)$  is  $\varepsilon_0$ - $G_\delta$  and hence  $\nu(i)$ - $G_\delta$ , so  $A$  is a  $\nu(i)$ - $G_\delta$ . Let  $H, K$  be  $\nu(i)$ -closed disjoint. To see that they can be separated, it is sufficient to produce a countable cover of  $(C, \nu(i))$  by  $\nu(i)$ -open sets  $U$  such that  $\text{Cl}_{\nu(i)}(U)$  intersects at most one of  $H$  and  $K$ ; call such  $U$  "nice". By Lemma 9.1 the set  $S = \text{Cl}_{\varepsilon_0}(H) \cap \text{Cl}_{\varepsilon_0}(K)$  is countable, and around each its points we may put a nice  $U$ . Since  $C \setminus S$  is  $\varepsilon_0$ -Lindelöf, we may cover it with a countable collection of nice  $U$  (which are in fact  $\varepsilon_0$ -open and whose  $\varepsilon_0$ -closure intersect at most one of  $H$  and  $K$ ). Thus, these two collections together produce the desired cover. Therefore,  $X(n)$  is perfectly normal.

We see that  $X(n)^2$  is perfectly normal by a way parallel to the above observation, since  $\nu$  is finer than  $\rho(n)$ ,  $|\text{Cl}_{\rho(n)}(A) \setminus A|_2 \leq \omega$  for any  $\nu$ -closed set  $A$ , and  $|\text{Cl}_{\varepsilon_0^2}(A) \setminus A|_2 \leq \omega$  for any  $\nu(i)^2$ -closed set  $A$  (see §6 (i) - (iii)).

By Lemma 5.2 we have  $\dim X(n)^2 \geq n$ . (It is easy to see that  $\dim X(n) = 0$ .) By Fedorčuk [1, Lemmas 7, 9]  $\dim X(n)^2 \leq \text{Ind } X(n)^2 \leq n$ . By Lemma 5.2  $X(\infty)^2$  is not countable dimensional. This completes the proof.

REMARK 9.1. (1) Theorem 9.1 solves a problem raised in Fedorčuk [2, Problem 6] affirmatively. Similar example is outlined in Engelking and Pol [1, Example 4.1],

(2) Our present construction of  $X(n)$  is a slight improvement of the construction given in Tsuda [2]. The main differences are

(i) our space  $X(n)$  is separable,

(ii) while two different factor spaces  $X$  and  $Y$  are constructed in Tsuda [2], we can construct here one satisfying that  $X = Y$ .

(3) Since the product of  $X(n)^2$  and  $k$ -dimensional cube  $I^k$  is perfectly normal (Engelking [1, Problem 4.5.16(b)]),

we have: For every  $n = 1, 2, 3, \dots, \infty$ , and

for every pair of non-negative integers  $(\ell, m)$

there exist spaces  $X$  and  $Y$  such that

(a)  $\dim X = \ell$  and  $\dim Y = m$ , while  $\dim (X \times Y) = m+n+\ell$ ;

(b)  $X \times Y$  is locally compact, separable, and perfectly normal.


(4) As mentioned in Wage[2] the first Wage-type example in the literature is the one satisfying Theorem 9.1 for the case  $n = 1$ . The construction suggested in Wage [1] remains correct if we do not force  $X = Y$  (in other word, in our terminology his proof that  $\nu(1) \times \nu(2)$  is normal is correct, though the method to make  $X^2$  being normal was incorrect (cited in Przymusiński [6, Footenote 7]).

Since he said nothing about Lemma 9.4, we cannot conclude that  $v(i)^2$  is normal. By the lemma we can show that it is normal.

(5) It is announced in Wage [1] that Theorem 9.1 holds for general dimension.

10. Local compactness and collectionwise normality of products.

88



10. Local compactness and collectionwise normality of products.

In this section we shall prove the following theorem without any set-theoretic assumptions beyond ZFC:

THEOREM 10.1. For every  $n = 1, 2, \dots, \infty$ , there exist two first-countable, separable, locally compact, locally countable spaces  $X$  and  $Y$  such that

- (a)  $X \times Y$  is countably paracompact, collectionwise normal;
- (b)  $\dim X = \dim Y = 0$ , while  $\dim (X \times Y)$  and  $\text{Ind} (X \times Y)$  are exactly equal to  $n$ . Moreover, our space  $X \times Y$  in the case  $n = \infty$  is not strongly countable dimensional.

Proof. We put

$$X = (C, \tau_1) \text{ and } Y = (C, \tau_2),$$

where  $\tau_i$  is defined in §7. By the construction it follows that  $X \times Y$  is first-countable, separable, locally compact, and locally countable. We shall show at first that it is countably paracompact and collectionwise normal. We need the following lemma.

LEMMA 10.1. (a) If  $A$  and  $B$  are closed disjoint sets in  $X$  or  $Y$ , then the set  $\text{Cl}_{\varepsilon_0}(A) \cap \text{Cl}_{\varepsilon_0}(B)$  is countable.

(b) Let  $A$  and  $B$  be closed disjoint sets in  $X \times Y$ . Then,

$$|\text{Cl}_{\rho(n)}(A) \cap \text{Cl}_{\rho(n)}(B)|_2 \leq \omega.$$

Proof. (a) The assertion follows from the fact that if  $|Cl_{\varepsilon_0}(A) \cap Cl_{\varepsilon_0}(B)| = c$ , then  $|Cl_{\rho(n)}(A^2) \cap Cl_{\rho(n)}(B^2)|_2 > \omega$ .

(b) It follows from the definition of  $\tau$  (see §7).

By the above lemma we can prove that each of  $X$  and  $Y$  is countably paracompact, collectionwise normal, and  $\omega_1$ -compact (that is, every closed discrete sets in it is countable). Hence, we can show that  $X \times Y$  is also countably paracompact and collectionwise normal by the technique of Przymusiński [1,3].

Next, we determine the dimension of our products and factors. We need the following lemma.

LEMMA 10.2. (a) Let  $U$  be a cozero-set of  $X$  or  $Y$ . Then, the set  $U \cap Cl_{\varepsilon_0}(C \setminus U)$  is countable.

(b) Let  $Z$  be a zero-set of  $(\Delta, \tau|_{\Delta})$ . Then, the set  $Cl_{\tilde{\rho}(n)}(Z) \setminus Z$  is countable.

(c) Let  $A$  be closed in  $X \times Y$  and  $U \subset A$  be a cozero-set in  $A$ . Then,  $|U \cap Cl_{\rho(n)}(A \setminus U)|_2 \leq \omega$ .

The proof goes on similarly as Lemma 10.1, so we omit it.

Now, we show that  $\dim X = \dim Y = 0$ . Let  $U$  be a cozero-set in  $X$  (or,  $Y$ ). Then, by Lemma 10.2(a) the set  $C \setminus Cl_{\varepsilon_0}(C \setminus U)$  is countable. Then, there exists an  $\varepsilon_0$ -clopen collection  $\{G_i: i \in \omega\}$  such that

$$\bigcup_{i \in \omega} G_i = C \setminus Cl_{\varepsilon_0}(C \setminus U) \text{ and } |U \setminus \bigcup G_i| \leq \omega.$$

Therefore,

$$U = \bigcup \{G_i: i \in \omega\} \cup \{B(y): y \in U \setminus \bigcup G_i\},$$

where  $B(y)$  is a clopen neighborhood of  $y$  in  $X$  (or  $Y$ ) satisfying  $B(y) \subset U$ . Hence,  $\dim X = \dim Y = 0$  by Terasawa [1, Theorem 1].

Next, we shall show that

LEMMA 10.3.  $\dim (X \times Y) \geq n$ .

Proof. Since  $\Delta$  is closed in  $X \times Y$ , it suffices to show that  $\dim (\Delta, \tau|\Delta) \geq n$ . Let  $Z = (\Delta, \tau|\Delta)$ . Put

$$A_k' = \Delta \cap \pi_{k+2}^{-1}[0, 1/3] \text{ and } B_k' = \Delta \cap \pi_{k+2}^{-1}[2/3, 1]$$

for each  $1 \leq k \leq n$ . (Remember that  $\rho(n) = \check{\rho}(n)|\Delta$ .) Then, it is obvious that each  $A_k'$  and  $B_k'$  are disjoint zero-set in  $Z$ . We shall show that if for each  $k$ , a zero-set  $S_k'$  separates  $A_k'$  and  $B_k'$  in  $Z$ , then  $\bigcap S_k' \neq \emptyset$ .

At first, for such a separation  $S_k'$  we shall show that there exists a closed separation  $S_k$  of  $A_k$  and  $B_k$  in  $(\Delta, \check{\rho}(n))$  such that

(1)  $S_k \setminus S_k'$  is a countable set for each  $k$ .

Because the zero-set  $S_k'$  separates  $A_k'$  and  $B_k'$ , there exist two cozero-sets  $U_k'$  and  $V_k'$  for each  $k$  such that  $U_k' \cap V_k' = \emptyset$ ,  $U_k' \supset A_k'$ ,  $V_k' \supset B_k'$ , and  $\Delta \setminus S_k' = U_k' \cup V_k'$ . Since  $U_k' \cup S_k'$  is a zero-set, it follows from Lemma 10.2(b) that

- (2) the set  $\text{Cl}_{\tilde{\rho}(n)}(U_k') \setminus (U_k' \cup S_k')$  is countable, and  $B_k \cap (\text{Cl}_{\tilde{\rho}(n)}(U_k')) = \emptyset$  for each  $k = 1, \dots, n$ .

(Remember that

$$B_k = \Delta \cap \pi_{k+2}^{-1}[6/7, 1] \text{ and } \text{Cl}_{\tilde{\rho}(n)}(U_k') \subset \Delta \cap \pi_{k+2}^{-1}[0, 2/3].)$$

Put

$$V_k = \Delta \setminus \text{Cl}_{\tilde{\rho}(n)}(U_k').$$

Then,  $V_k$  is open in  $(\Delta, \tilde{\rho}(n))$ ,  $V_k \supset B_k$  and

$$A_k \cap \text{Cl}_{\tilde{\rho}(n)}(V_k) = \emptyset. \text{ Finally, put}$$

$$S_k = \text{Cl}_{\tilde{\rho}(n)}(V_k) \setminus V_k.$$

Then, the set  $S_k \setminus S_k'$  is countable, since the inclusion  $\Delta \cap \pi_{k+2}^{-1}[1/3, 1] \supset V_k' \cup S_k' \supset V_k$  holds, (2) and Lemma 10.2 (b) holds. Thus,  $S_k$  is a closed separation of  $A_k$  and  $B_k$  satisfying (1). On the other hand, by Theorem 4.1 (1) we can see that  $|\cap S_k| = c$ . From (1) and the fact that  $(\cap S_k) \setminus S_k' \subset S_k \setminus S_k'$ , we have  $|(\cap S_k) \cap (\cap S_k')| = c$ , so that  $\cap S_k' \neq \emptyset$ . This completes the proof of Lemma 10.3.

Finally, we shall show that  $\text{Ind}(X \times Y) \leq n$ . For this purpose we need the following lemma.



LEMMA 10.4. Let  $A$  be a closed set in  $X \times Y$  such that  $A = T \cup S$  for some  $\rho(n)$ -closed set  $T$  of  $|T|_2 \leq \omega$ , and a set  $S$  satisfying  $\dim(S, \rho(n)|S) \leq m$  for some non-negative integer  $m$ . Then,  $\dim(A, \tau|A) \leq \text{Ind}(A, \tau|A) \leq m$ .

Proof. Since  $A$  is normal, it suffices to show that  $\text{Ind}(A, \tau|A) \leq m$ . We shall prove it by induction on  $m$ . Because  $|T|_2 \leq \omega$ ,  $\dim(S, \rho(n)|S) \leq m$ , and the countable sum theorem holds, we first note that

$$(1) \quad \dim(A, \rho(n)|A) \leq m.$$

(Remember that  $\tilde{\rho}(n)|\{t\} \times C$  and  $\tilde{\rho}(n)|C \times \{t\}$  are homeomorphic to  $(C, \varepsilon_0)$  for each  $t \in C$  by Theorem 4.2.)

For the case  $m = 0$  we take arbitrary cozero-set  $U$  in  $(A, \tau|A)$ . Then, the 2-cardinality of the set

$$F = U \cap \text{Cl}_{\rho(n)}(A \setminus U)$$

is countable by Lemma 10.2 (c). For the  $\rho(n)|A$ -open set

$$U \setminus F = A \setminus \text{Cl}_{\rho(n)}(A \setminus U)$$

we have a collection  $\mathcal{u}$  of countably many  $\rho(n)|A$ -clopen sets such that

$$\cup \mathcal{u} = U \setminus F$$

from (1). For the set  $F$  we take a countable set  $W \subset C$  such that

$$W \times C \cup C \times W \supset F.$$

Then, for each  $y \in W$  we shall show that there exist two collections  $\mathcal{u}(y)$  and  $\mathcal{v}(y)$  consisting of countably

many  $\tau|A$ -open sets such that

$$(2) \quad U \supset \cup u(y) \supset U \cap (\{y\} \times C) \text{ and } U \supset \cup v(y) \supset U \cap (C \times \{y\}).$$

Let  $G$  be a  $\tau$ -open set such that  $U = G \cap A$ . Since  $U$  is a  $F_\sigma$ -set in  $X \times Y$ , put  $U = \cup K_i$ , where each  $K_i$  is  $\tau$ -closed. Because  $\dim Y = 0$ , we have a clopen set  $Z_i$  in  $Y$  for each  $K_i$  and  $G$  such that

$$(3) \quad K_i \cap (\{y\} \times C) \subset \{y\} \times Z_i \subset G \cap (\{y\} \times C).$$

Let  $\{G_s : s \in \omega\}$  be a countable clopen base of  $y$  in  $Y$ .

Put

$$U_s = \cup \{V : V \text{ is open in } Y \text{ and } G_s \times V \subset G\}.$$

Then, from (3)  $\{U_s : s \in \omega\}$  is a countable open cover of  $Z_i$ . Because  $\dim Z_i = 0$  and  $Z_i$  is countably paracompact normal, there exists a disjoint clopen cover  $\omega = \{W_s : s \in \omega\}$  of  $Z_i$  such that  $W_s \subset U_s$ . Put

$$u_i = \{A \cap (G_s \times W_s) : s \in \omega\}.$$

Then,  $u_i$  is a countable  $\tau|A$ -clopen cover of the set  $K_i \cap (\{y\} \times C)$  satisfying  $\cup u_i \subset U$ . Similarly, for each  $i$ , we have a countable  $\tau|A$ -clopen collection  $v_i$  satisfying

$$K_i \cap (C \times \{y\}) \subset \cup v_i \subset U.$$

Finally, put

$$u(y) = \cup u_i \text{ and } v(y) = \cup v_i.$$

Thus, (2) holds. Then,  $\dim (A, \tau|A) \leq 0$  by Terasawa [1, Theorem 1], since

$$U = \cup u \cup \{u(y), v(y) : y \in W\}.$$

This completes the proof for the case  $m = 0$ . Suppose that our lemma is true for  $m-1$ . For the case  $m$  let  $H$  and  $F$  be disjoint closed set in  $(A, \tau|_A)$ , and let  $U$  and  $V$  be disjoint cozero-sets in it such that

$$(4) \quad U \supset H, \quad V \supset F, \quad \text{and} \quad \text{Cl}_\tau(U) \cap \text{Cl}_\tau(V) = \emptyset.$$

Put

$$T' = \text{Cl}_{\rho(n)}(U) \cap \text{Cl}_{\rho(n)}(V).$$

Then,  $|T'|_2 \leq \omega$  by Lemma 10.1(b). Because  $\dim (A, \rho(n)|_A) \leq m$  from (1), we have an at most  $(m-1)$ -dimensional closed separation  $S'$  in the space  $(A \setminus T', \rho(n)|_{A \setminus T'})$  between the sets  $\text{Cl}_{\rho(n)}(U) \setminus T'$  and  $\text{Cl}_{\rho(n)}(V) \setminus T'$ . (Note that  $S' \cap (U \cup V) = \emptyset$ .) Put

$$A' = S' \cup (T' \setminus U \cup V).$$

Then,  $A'$  is a closed separation of  $H$  and  $F$  in  $(A, \tau|_A)$  from (4) and the construction of  $S'$ . Hence, inductive assumption can be applied to the set  $A'$  so that  $\text{Ind}(A', \tau|_{A'}) \leq m-1$ , and therefore,  $\text{Ind}(A, \tau|_A) \leq m$ . This completes the proof of our lemma.

To show that  $\text{Ind}(X \times Y) \leq n$  take two disjoint closed sets  $H$  and  $F$  in  $X \times Y$ . Then, by taking cozero-sets  $U$  and  $V$  satisfying (4), we have a closed separation  $A$  of  $H$  and  $F$  which satisfies all the conditions in Lemma 10.4 for the case  $m = n-1$ . Hence,  $H$  and  $F$  are separated by a closed set  $A$  of  $\text{Ind} A \leq n-1$ , therefore  $\text{Ind}(X \times Y) \leq n$ .

The assertion for the case  $n = \infty$  follows from Lemma 5.3, since the topology  $\tilde{\tau} = \tau|_{\Delta}$  satisfies the conditions i) and ii) in §5, 2).

REMARK 10.1. (a) For the case  $n = 1$  the existence of our example was suggested in Przymusiński [3].  
(b) Our present examples are slight improvements of the ones given in Tsuda [1] (our examples are separable).  
(c) Since  $X \times Y$  is countably paracompact normal, we have: for every  $n = 1, 2, \dots, \infty$ , and for every pair of non-negative integers  $(\ell, m)$  there exist two spaces  $X$  and  $Y$  such that  
(i)  $\dim X = \ell$  and  $\dim Y = m$ , while  $\dim (X \times Y) = \ell + m + n$ ;  
(ii)  $X \times Y$  is locally compact, separable, countably paracompact and collectionwise normal.

One of the remaining problems in this section is:

PROBLEM 10.1. Whether or not we can construct one with  $X = Y$ .

In Przymusiński [3] or Engelking and Pol [1] it is announced that it is the case.



11. With Lindelöf factors and normality of products.

In this section we prove the following theorem without any set-theoretic assumptions beyond ZFC:

THEOREM 11.1. For every  $n = 1, \dots, \infty$  and integer  $m \geq 1$  there exists a separable and first-countable space  $X$  such that

- (i)  $X^m$  is Lindelöf and  $\dim X^m = 0$ ;
- (ii)  $X^{m+1}$  is normal but  $\dim X^{m+1} = n$ . Moreover,  $X^{m+1}$  is not strongly countable dimensional when  $n = \infty$ .

COROLLARY 11.1. For every  $n = 1, 2, \dots, \infty$ , there exists a separable and first-countable Lindelöf space  $X$  such that

- (i)  $\dim X = 0$ ;
- (ii)  $X^2$  is normal but  $\dim X^2 = n$ . Moreover,  $X^2$  is not strongly countable dimensional when  $n = \infty$ .

At first, we shall show the following lemma, which is some version of the natural extension of  $\tilde{p}(n)$  (cf. Theorem 4.2).

LEMMA 11.1. There exists a complete separable metric space  $(C^{m+1}, \bar{\rho}(n))$  satisfying

- (1)  $\bar{\rho}(n)|_{\Delta} = \tilde{\rho}(n)$ , where  $\Delta = \{(x, x, \dots, x) : x \in C\}$ ;
- (2)  $\bar{\rho}(n)|_{C^{m+1} \setminus \Delta}$  is homeomorphic to the usual Euclidean topology on it;
- (3) for each  $t \in C$   $(m+1)$  topologies  $\bar{\rho}(n)|_{\pi_i^{-1}(t)}$  are homeomorphic to  $\varepsilon_0$ , where  $\pi_i: C^{m+1} \rightarrow C$  is the projection to the  $i$ -th factor.

Proof. Let  $h: C \rightarrow I^n$  and  $B_i$  be the function and the development given in Theorem 4.1, respectively. We shall extend  $h$  to the function  $\bar{H}: C^{m+1} \rightarrow I^n$  as follows.

$$\bar{H}((x_i)) = \begin{cases} h(x_i) & \text{if } (x_i) \in \Delta, \\ 0 & \text{if } (x_i) \in \cup \{B^{m+1}; B \in B_1\}, \\ h_i(x_i) & \text{if } (x_i) \in \cup \{B^{m+1}; B \in B_i\} \setminus \cup \{B^{m+1}; B \in B_{i+1}\}. \end{cases}$$

Since  $h|B$  is constant for each  $B \in B_i$ , the above definition of  $\bar{H}$  is well-defined. Let  $\bar{\rho}(n)$  be the topology on  $C^{m+1}$  generated by the subbase

$$\varepsilon_0 \cup \{\bar{H}^{-1}(U) : U \in \varepsilon_0\}.$$

Then,  $(C^{m+1}, \bar{\rho}(n))$  is homeomorphic to the set

$$\{(x_i, \bar{H}((x_i))) : (x_i) \in C^{m+1}\}$$

with the restriction of the Euclidean topology on  $C^{m+1} \times I^n$ .

By an argument parallel in Theorem 4.2 we see that (1)

and (2) hold and that  $\bar{p}(n)$  is separable complete metric. Let us show that (3) holds. By the definition of  $H$ , we see that there exists a sequence of  $\varepsilon_0$ -open sets  $\{U_i\}$  in  $\pi_i^{-1}(t)$  such that  $\cup U_i = \pi_i^{-1}(t) \setminus \{(t, \dots, t)\}$ ,  $\bar{H}|_{U_i}$  is constant, and  $\varepsilon_0^m$ -distance between the point  $(t, \dots, t)$  and  $U_i$  is less than  $1/i$ .

Since  $\lim_{i \rightarrow \infty} \bar{H}(U_i) = \bar{H}(t, \dots, t) = h(t)$ ,  $\bar{H}|_{\pi_i^{-1}(t)}$  is  $\varepsilon_0$ -continuous. Hence, (3) holds. This completes the proof.

Next, we need the following lemma.

LEMMA 11.2. There exist  $m+2$  disjoint subsets  $E_0, \dots, E_{m+1}$  in  $C$  such that the  $m+1$ -cardinality of the set  $A \cap (E_i)^{m+1}$  is the cardinality of continuum for any  $\bar{p}(n)$ -closed set  $A$  in  $C^{m+1}$  when  $A$  has uncountable  $m+1$ -cardinality.

Proof. Since  $\bar{p}(n)$  is complete separable metric, the assertion follows from Przymusiński [2, Theorem 2].

On the other hand, we choose the following collection  $\bar{P}$  of points from  $(E_0)^{m+1}$ , which plays the role of  $P$  given in section 7.



Let  $F = \{(A, B)_\lambda : \lambda \in \Lambda\}$  be the collection of all pairs of countable subsets of  $C^{m+1}$  such that

$$|Cl_{\bar{\rho}(n)}(A) \cap Cl_{\bar{\rho}(n)}(B)|_{m+1} > \omega.$$

Let  $<$  be a well-ordering of  $C$  and for each  $x \in C$  we define  $I(x) = \{y \in C : y \in x\}$ . Let  $D$  be some fixed countable  $\bar{\rho}(n)$ -dense set in  $C$ . Then, by an argument parallel in Lemma 7.3 we can choose a transfinite sequence  $\langle p_\lambda : \lambda \in \Lambda \rangle$  of points in  $(E_0)^{m+1}$  satisfying the following (4) and (5).

(4)  $p_\lambda \in (E_0)^{m+1} \cap Cl_{\bar{\rho}(n)}(A) \cap Cl_{\bar{\rho}(n)}(B) \setminus ((I(x_\lambda) \cup D)^{m+1})$ , where  $(A, B) = (A, B)_\lambda$ , and  $x_\lambda$  is some element of  $C$  such that  $A \cup B \subset I(x_\lambda)^{m+1}$ .

(5) If  $\lambda \neq \mu$ , then  $\bar{p}_\lambda \cap \bar{p}_\mu = \emptyset$ , where  $\bar{p} = \{p_0, \dots, p_m\}$  for  $p = (p_0, \dots, p_m)$ .

Put  $\bar{P} = \{p_\lambda : \lambda \in \Lambda\}$ .

Finally, we can show by a way parallel to §§10, 11 that

(6) Let  $G_k$  be the  $(1/k)$ -neighborhood of  $p_\lambda = (t_i)$  with respect to the metric  $\bar{\rho}(n)$ . Then, there exists a sequence  $\{z_k = (z_{k,i})\}$  such that

(a)  $z_k \in G_k$ ,  $z_{k,i} \neq z_{s,i}$  for  $s \neq k$ ,  $\{(z_{3k,i})\} \subset A$ ,  $\{(z_{3k+1,i})\} \subset B$ ,  $z_{3k+2} \in D^{m+1}$ , where  $(A, B) = (A, B)_\lambda$ ;

(b) there exist  $2(m+1)$   $\varepsilon_0$ -clopen sets  $H_{k,i}$  and  $U_{k,i}$  such that

$$p_\lambda \in \prod_{i=1}^{m+1} H_{k,i}, \quad (t_1, \dots, z_{k,i}, \dots, t_{m+1}) \in \prod_{j \neq i} H_{k,j} \times U_{k,i} \subset G_k$$

for every  $1 \leq i \leq m+1$ .

Proof of Theorem 11.1. 1) Construction of  $X$ .

At first, we construct  $m+1$  local bases

$$\{H_k(x)_i : k \in \omega, x \in C\}, \quad 1 \leq i \leq m+1,$$

satisfying the following conditions as inductive assumptions.

- (7) Each  $H_k(x)_i$  is  $\varepsilon_0$ -closed, and for any  $y \in H_k(x)_i$  there exists an  $s \in \omega$  such that  $H_s(y)_i \subset H_k(x)_i$ .
- (8) The diameter of  $H_k(x)_1 \times \dots \times H_k(x)_{m+1}$  is less than  $1/k$  with respect to the metric  $\bar{\rho}(n)$ .

Suppose that  $x \in C$  and such bases of neighborhoods have been constructed for each  $y < x$ .

We consider the following four cases:

- (c)  $(x, \dots, x) = p_\lambda \in \Delta$  for some (unique)  $\lambda \in \Lambda$ . Put for each  $k \in \omega$

$$H_k(x)_i = \{x\} \cup \bigcup_{t \geq k} H_{s(t)}(z_{t,i})_i, \quad 1 \leq i \leq m+1,$$

where  $k(t)$  is defined by a way parallel to the one in §11.

- (d)  $x = p_{\lambda,i}$ , and  $p_\lambda \in \Delta$ . Put for each  $k \in \omega$

$$H_k(x)_i = \{x\} \text{ if } x \neq p_{\lambda,j}, \text{ and}$$

$$H_k(x)_i = \{x\} \cup \bigcup_{t \geq k} H_{s(t)}(z_{t,i})_i \text{ if } x = p_{\lambda,i}.$$

- (e)  $x \in E_i$ ,  $1 \leq i \leq m+1$ . Put  $H_k(x)_j = \{x\}$  for  $j \neq i$ , and  $H_k(x)_i$  is an  $\varepsilon_0$ -clopen neighborhood of  $x$  whose diameter with respect to  $\varepsilon_0$  is less than  $1/s$ .

- (f)  $x$  does not belong to the above three cases. Let  $H_k(x)_i = \{x\}$  for  $1 \leq i \leq m+1$ , and  $k \in \omega$ .

It is easy to see that  $\{H_k(x)_i : k \in \omega, x \in C\}$  satisfies (7). Thus, inductive construction has completed. From (7) and (8) they constitute local bases on  $C$ , and they produce Hausdoeff topologies  $\sigma_i$ . This completes the construction of  $\sigma_i$ .

2) Desired properties of  $X$ . We put

$$X = \bigoplus_{i=1}^{m+1} (C, \sigma_i).$$

Then, it follows from Przymusiński [1] that  $X^m$  is first-countable, separable, Lindelöf. We have  $\dim X^m = 0$ , and  $\dim X^{m+1} = \text{Ind } X^{m+1} = n$  by a way parallel to Theorem 10.1 (b). The assertion for  $n = \infty$  follows from an argument parallel to Theorem 10.1 (b), also. This completes the proof.

REMARK 11.1. (1) For the case  $(m,1)$  our result was stated in Przymusiński [3] without proof. For the case  $(2,\infty)$  our result was outlined in Engelking and Pol [1, Example 4.2]. Corollary 11.1 is a slight improvement of Tsuda [1, Theorem 1.2]. Here, our space is separable. It is also announced in Charalambous [2, Footnote at p. 648] that Engelking and Pol have independently showed this corollary.

(2). We have no Wage-type examples for the case  $m = \infty$  (see also section 8):


THEOREM 11.2 (Nagami [5]). Let  $X^\omega$  be normal. If each  $X^i$  is paracompact (respectively, Lindelöf), then  $X^\omega$  is paracompact (respectively, Lindelöf).

One of remaining problems in this section is:

PROBLEM 11.1. For every triple of non-negative integers  $(m,k,n)$  find a space  $X$  such that

- (i)  $\dim X^i = k$ , and  $X^i$  is Lindelöf for each  $i \leq m$ ;
- (ii)  $X^{m+1}$  is normal but  $\dim X^{m+1} = k+m$ .

12. Non-normal Wage-type examples.



## 12. Non-normal Wage-type examples.

1) The purpose of this section is to show the following theorem due to Wage [2]. We also discuss some related problems.

**THEOREM 12.1.** There exists a separable metric space  $X$  and a first-countable separable Lindelöf space  $Y$  such that

$$\dim X = \dim Y = 0, \text{ and } \dim (X \times Y) = 1.$$

Proof. 1) Constructions of  $X$  and  $Y$ . Let  $\tilde{\rho}$  and  $\rho$  be the Wage's metric topologies given in Theorem 3.1, and Theorem 3.2, respectively. Let  $E_1$  and  $E_2$  be two Bernstein sets given in Lemma 11.2 which also satisfy  
(0)  $|A \cap (E_1 \times E_2)|_2 = c$  for any  $\rho$ -closed set  $A \subset C \setminus \Delta$ .  
Finally, let  $\tau_0$  be the approximation by the method of van Douwen (see § 5) of the topology  $\tilde{\rho}_0$  of the restriction of  $\tilde{\rho}$  to the subset  $E_1$ . That is,  $\tau_0$  is first-countable, locally countable, locally compact, separable topology on  $E_1$  finer than  $\tilde{\rho}_0$  such that  
(1) if  $A$  and  $B$  are disjoint  $\tau_0$ -closed sets, then the set  $Cl_{\tilde{\rho}_0}(A) \cap Cl_{\tilde{\rho}_0}(B)$  is countable.

Put

$$X = (E_1, \varepsilon'), \text{ and } Y = (C, \upsilon),$$

where  $\varepsilon' = \varepsilon_0|E_1$  and  $\upsilon$  is the topology generated by the subbase  $\varepsilon_0 \cup \tau_0$ .

2) Dimension of factors and the product  $X \times Y$ . Since  $\varepsilon_0$  is 0-dimensional separable metric topology,  $X$  is also 0-dimensional separable metrizable. On the other hand, since  $E_1$  is a Bernstein set,  $Y$  is Lindelöf (see § 11). By the definition of  $Y$ , we see that  $Y$  is separable, first-countable, and has a clopen base. We shall show that  $\dim(X \times Y) = 1$ .

At first, we note that from Theorem 3.2 (4) the topology  $\varepsilon' \times \upsilon$  is finer than the topology  $\tilde{\rho}|E_1 \times C$ . Hence, since  $H|X \times Y$  is continuous, it can be seen in a parallel way as in § 3 that  $\dim(X \times Y) > 0$ . Therefore, it suffices to see that  $\dim(X \times Y) \leq 1$ . Let  $u = \{U_i\}$  be a finite cozero-sets cover of  $X \times Y$ , and let  $F = \{F_i\}$  be a finite zero-set shrinking of  $u$  such that  $U_i \supset F_i$ . We shall show at first that  $\Delta' = \{(x, x) : x \in E_1\}$  is covered by finite cozero-sets of order at most 2. Put

$$T = \cup (\text{cl}_{\tilde{\rho}_0}(\Delta' \setminus F_i) \cap \text{Cl}_{\tilde{\rho}_0}(\Delta' \setminus U_i)).$$

Then, by (1)  $T$  is a countable set in  $\Delta'$ , and is  $\tau_0$ -closed, so that when we consider the finite  $\tilde{\rho}_0$ -open cover  $\{\Delta' \setminus \text{Cl}_{\tilde{\rho}_0}(\Delta' \setminus U_i)\}$  of the space  $\Delta' \setminus T$ , we have a  $\tilde{\rho}_0$ -open cover  $\nu' = \{V_i'\}$  of order at most 2 such that  
 (1)  $V_i' \subset \Delta' \setminus \text{Cl}_{\tilde{\rho}_0}(\Delta' \setminus U_i)$  and  $\cup \nu' = \Delta' \setminus T$ .

Let  $\rho_0$  be the restriction of the topology  $\rho$  to the subset  $E_1 \times C$ . Then, by the following theorem we have a finite family of  $\rho_0$ -open sets  $\mathcal{G}$  satisfying the following (2).

THEOREM (K. Kuratowski [1, p. 226]). Given a family  $\mathcal{V}'$  of  $\tilde{\rho}_0$ -open sets, there exists a family  $\mathcal{G}$  of  $\rho_0$ -open sets such that

(2)  $\Delta' \cap G_i = V_i'$  and the condition  $V_{i_1}' \cap \dots \cap V_{i_k}' = \emptyset$  implies  $G_{i_1} \cap \dots \cap G_{i_k} = \emptyset$ , for every (finite) system of indices  $i_1, \dots, i_k$ .

Then,  $\mathcal{G}$  is a cozero-set collection of  $X \times Y$  such that

(3)  $(\cup \mathcal{G}) \cap \Delta' = \cup \mathcal{V}' = \Delta' \setminus T$ .

For each  $t \in T$  we can take a clopen set

$$N(t) = G(t) \times B(t),$$

where  $G(t)$  is an  $\varepsilon'$ -clopen neighborhood of  $t$  and  $B(t)$  is a  $\tau_0$ -clopen neighborhood of  $t$  consisting countable points such that  $N(t) \subset U_i$  for some  $U_i \in \mathcal{U}$ . Note that  $N(t)$  is Lindelöf. Put

$$S = \cup \{N(t) : t \in T\}.$$

Then,  $S$  is a 0-dimensional Lindelöf space, because  $N(t)$  is Lindelöf and  $T$  is countable. Since  $S$  is a cozero-set containing the zero-set  $\Delta'$  in  $X \times Y$ , there exists a finite clopen collection  $\mathcal{S} = \{S_i\}$  in  $X \times Y$  such that

$$S_i \subset U_i \text{ and } \cup \mathcal{S} \supset T.$$



Hence, the set  $W = \cup G \cup \cup S$  is a cozero-set containing the set  $\Delta'$ . Hence, there exists a zero-set  $Z$  and a cozero-set  $G$  in  $X \times Y$  such that

$$\Delta' \subset G \subset Z \setminus W.$$

Because space  $X \times Y \setminus \Delta'$  is 0-dimensional Lindelöf by (0), we can choose a clopen set  $K$  in  $X \times Y \setminus \Delta'$  such that

$$K \cap Z = \emptyset \text{ and } K \subset X \times Y \setminus W.$$

Since  $\text{Cl}_{X \times Y}(K) \subset X \times Y \setminus G$ ,  $K$  is also clopen in  $X \times Y$ .

Put

$$V_i'' = (G_i \cup S_i \setminus K \cup \cup_{j < i} S_j).$$

Then,  $V_i''$  is a cozero-set of  $X \times Y$  such that

$$\cup V_i'' = X \times Y \setminus K \supset \Delta',$$

and its order is at most 2 by (2), (3). Because  $K$  is strongly 0-dimensional, there exists a finite disjoint collection  $\mathcal{K} = \{K_i\}$  of clopen sets such that

$$K_i \subset U_i \text{ and } \cup \mathcal{K} = K.$$

At last, put

$$V_i = V_i'' \cup K_i$$

for each  $i$ . Then, it can be seen that  $\mathcal{V} = \{V_i\}$  is a cozero-sets refinement of  $\mathcal{U}$  and its order is at most 2. This completes the proof of our theorem.

REMARK 12.1. (a) Our present construction of  $X$  and  $Y$  is a slight improvement of the construction given in Wage [2]. In there some approximation on  $E_1$  is used, which is defined more easier than ours. But, it seems that its definition is too simple to see that it satisfies our condition (1). Therefore, he can only announce that  $\dim(X \times Y)$  is positive. In our case, however, we can determine the dimension of our product space exactly, using the property (1) and Lemma 10.2 (b).

(b) One of the main characteristic properties of Wage-type examples with a metric factor is that its product cannot be normal (see Kodama [4]). In this point of view we can consider our space  $Y$  as some variation of the well-known Michael line  $C_{E_1}$ : Its topology is defined by declaring the set of the form  $U \cup K$ , where  $U$  is  $\varepsilon_0$ -open and  $K \subset C \setminus E_1$ , as its open set. (It is some times called the Hannerization in  $C$  with respect to  $E_1$ ). Since it is known (Terasawa [1]) that  $\dim(X \times C_{E_1}) = 0$  for every strongly 0-dimensional metric space  $X$ , Michael line does not serve our purpose in the view of Wage-type examples. (From our constructions we see that the discrete topology on  $E_1$  in  $C_{E_1}$  is too strong to construct Wage-type example with a metric factor.)

(c) There is another well-known Michael line  $C_Q$ , which is the Hannerization in  $C$  with respect to the countable

dense subset  $Q$  in  $C$ . Its advantage is that its product with the space of irrational numbers  $P$  is not normal. Hence, we can raise a following problem. (Note that our space  $X$  in Theorem 12.1 cannot be complete metric.)

PROBELM 12.1. Are there any Wage-type examples with a complete metric factor ?

We also note that we cannot replace complete metric by compact metric in the above problem, since  $\dim (X \times I) = \dim X + 1$  for every  $X$  (see, Chiba and Chiba [1, Lemma 1] or Morita [4] ).

We also note that there exists a strongly 0-dimensional Dwoker space (Dow and van Mill [1]).

We can specify the above problem as follows.

PROBELM 12.2. Are there any Wage-type examples with an irrational factor ?

As mentioned at the end of Remark 12.1 (b), when we attempt to prove this problem by modifying the Michael line, I believe that one of the key points is (\*) whether or not there exists a topology on  $P$  finer than  $\tilde{\rho}$ , which is 1-dimensional paracompact and has a clopen base.

We can raise one more problem for higher dimensional examples:

PROBLEM 12.3. Does there exist for every non-negative integer  $n$  a separable metric space  $X$  and a first-countable separable Lindelöf space  $Y$  such that

$$\dim X = \dim Y = 0, \text{ and } \dim (X \times Y) = n ?$$

2) With a pseudocompact factor.

One of the most remarkable facts among the previously known Wage-type examples is that all of them satisfy the first-countability axiom (hence are k-spaces), and some of their products are also locally compact. The purpose of this section is to show that there exists another Wage-type example which cannot be a k-space.

By a theorem due to H. Tamano [1], we can specify our problem as follows.

PROBLEM 12.4. Are there any counter-examples for the inequality (\*) with a pseudocompact factor ?

The following theorem solves this problem affirmatively:

THEOREM 14.2 (Tsuda [3]). Without any set-theoretic assumptions beyond ZFC there exist a hereditarily separable, hereditarily Lindelöf space  $X$  and a first-countable, locally compact, separable, pseudocompact space  $Y$  such that

$$\dim X = \dim Y = 0, \text{ while } \dim (X \times Y) > 0.$$

To prove our theorem we need the following another factorization technique:

THEOREM 12.2 (Tsuda [3, Key Lemma]). For each point  $(x, x) \in C$  and each countable set  $\lambda$  which converges to  $x$  in the usual Euclidean topology on  $C$ , and for a given  $\rho$ -open neighborhood  $G$  of  $(x, x)$ , there exists a  $\tilde{\rho}$ -clopen set  $B$  and a finite set  $F$  such that

$$(x, x) \in \underbrace{(\lambda \setminus F)}_{\cup \{x\}} \times B \subset G.$$

Now, we define our space  $X$  as the set  $C$  with the topology  $\mu$  determined by assigning to each point  $x \in C$  a base of neighborhoods consisting of those subset  $U$  of  $C$  which satisfy the condition :

(1)  $U$  contains some  $\tilde{\rho}$ -clopen set  $B$  with  $x \in B$ .

On the other hand, the space  $Y$  is a space which is an  $N \cup R$  for some maximal almost disjoint collection (m.a.d. collection for short)  $R$ . By  $N \cup R$  we mean a space which is defined in the following way on the set-theoretic union of a countable infinite set  $N$  and an almost disjoint collection  $R$  of infinite subsets of  $N$ : each point of  $N$  is isolated and  $\lambda \in R$  has a neighborhood base

$$\{ \{\lambda\} \cup (\lambda \setminus F) : F \text{ is a finite subset of } N \}.$$

REMARK 12.2. (a) Spaces  $N \cup R$  have been introduced by Mrówka [1], and have been found to have interesting properties Mrówka [2,3], Terasawa [2,3]. The use of spaces  $N \cup R$  for dimension theory is not new. One of the remarkable results among them is the one of Terasawa [3].

(b) Our present example was given in Tsuda [3]. It gives a partial answer to the problem given in Przymusiński [6, Problem 18], that whether or not there exists a Wage-type example with a countably compact factor.


(c) We remark that our factor space  $X$  is perfectly normal, since it is a  $\sigma$ -space. We cannot, however, improve  $X$  to be also pseudocompact, since  $Y$  is first-countable, locally compact. We also note that  $Y$  is neither normal nor countably compact, since  $Y$  has uncountable closed discrete subset. Hence, we can raise the following problem:

PROBLEM 12.5. Are there any Wage-type example with a pseudocompact factor whose product are also normal ?

We can raise one more problem (see also, Remark 8.1):

PROBLEM 12.6. For a given positive integer  $n$  are there any Wage-type examples with a pseudocompact factor which satisfy that  $\dim X = \dim Y = 0$ , while  $\dim (X \times Y) = n$  ?

Appendix, Several product theorems.





Appendix. Several product theorems.

Here, we review some positive results for the inequality (\*), because we can conclude from them some topological properties of our examples in Chapter 3 (for example, non-paracompactness of products, with non-metric factor, or with non-pseudocompact factor). It gives also some motivations of the attempt to look for other Wage-type examples in Chapter 3. In the sequel, we do not assume the normality of products unless otherwise specified. Hence, by the dimension of a space we mean the covering dimension due to Katětov [1]. Therefore, it coincides with the ordinal covering dimension when the space is normal.

We begin with the relatively strong theorem due to Pasynkov. For this purpose we need the following interesting and important concept of the "(piecewise) rectangularity" of a product (Pasynkov [1, - 5]).

Definition A.1. A subset of the product space  $X \times Y$  is said to be (piecewise) cozero rectangular if it is (a clopen subset of the set) of the form  $U \times V$ , where  $U$  and  $V$  are cozero sets of  $X$  and  $Y$ , respectively.

The product space is said to be (piecewise) rectangular if any finite cozero cover of it has a

$\sigma$ -locally finite refinement consisting of (piecewise) cozero rectangular subsets.

THEOREM A.1 (Pasyнков [1, 4]). Every (piecewise) rectangular product satisfies the inequality (\*).

In each of the following cases (a) through (c), and (d) the product is rectangular and piecewise rectangular, respectively.

- (a)  $X$  is a metric space and  $X \times Y$  is countably paracompact and normal (Kodama [4]). Moreover,  $X$  is a paracompact  $p$ -space and  $X \times Y$  is countably paracompact and normal (Filippov [1] and Pasyнков [1,3]).
- (b)  $X$  is locally compact and paracompact (Morita [4]).
- (c) The projection  $p_X: X \times Y \rightarrow X$  is a  $z$ -closed map (that is,  $p_X(Z)$  is closed in  $X$  for every zero set  $Z$  of  $X \times Y$ ) (Filippov [1], Nagami [8], Pasyнков [2]).
- (d)  $X \times Y$  is completely paracompact. (That is, for any open cover  $\mathcal{u}$  there exists a sequence  $\mathcal{v}_i$  of star-finite open cover such that  $\cup \mathcal{v}_i$  contains a refinement of  $\mathcal{u}$ . In particular, every Lindelöf space is completely paracompact.) Zolotarev [1].

REMARK A.1. (a) By a result of Rudin and Starbird [1] we can remove the assumption countably paracompactness in the case (a) when it is a metric factor. But, we cannot remove the normality of the product (see §12).

(b) When  $X$  is compact, the result is due to Terasawa (see Chiba and Chiba [1, Lemma 1]).

(c) The concept of the  $z$ -closed map was introduced by Ishiwata. The proof of Theorem 1.1 in Ishiwata [1] contains a proof that the inequality (\*) is valid in the case (c), implicitly. In particular,  $p_X$  is  $z$ -closed, when  $X$  is a pseudocompact and  $Y$  is a  $k$ -space (implicitly, H. Tamano [1]).

(d) More general results can be found in Filippov [2], and Pasynkov [5].

The Pasynkov's theorem is relatively strong, but it is known that

(1) there exist non-rectangular strongly 0-dimensional products (Hoshina and Morita [1], Ohta [1],

Przymusiński [7], K. Tamano [1], and Terasawa [1]). Moreover, Ohta [1] showed a machine to produce normal non-rectangular products  $X \times Y$  which satisfy the inequality (\*) for every normal non-paracompact (not necessary strongly 0-dimensional) space  $X$ .

One of the remarkable consequences of the introduction of the piecewise rectangularity is that it is a necessary and sufficient condition for the validity of the inequality (\*) when the factor spaces are strongly 0-dimensional. Hence, all the examples in (1) are piecewise rectangular. In Tsuda [7] we showed that for every positive integer  $n$  there exists an  $n$ -dimensional, collectionwise normal, non-piecewise rectangular product which satisfies the inequality (\*).

The idea we used there goes back to Chiba and Chiba [2], and Chiba [1].

On the other hand, there is a theorem which cannot be deduced from Theorem A.1.

THEOREM A.2 (Morita [6]). Let  $X$  be  $\sigma$ -locally compact paracompact. Then, the inequality (\*) is valid.

The following problem is communicated by T. Goto.

PROBLEM A.1. Is the inequality (\*) valid for the normal product with a Lašnev factor ?

It is known (Hoshina [1]) that the normality of product and countably paracompactness of it is equivalent in this case.

Finally, we mention infinite products. The following definition is suggested by Pasynkov [5] and Yajima [3].

DEFINITION A.2. A Cartesian product  $X = \prod_{\lambda \in \Lambda} X_\lambda$  is said to be (piecewise) cylindrical if each finite cozero cover of  $X$  has a  $\sigma$ -locally finite refinement by sets of (a clopen subset of) the form  $\pi_\xi^{-1}(U)$ , where  $\xi$  is a finite subset of  $\Lambda$  and  $U$  is a cozero set in  $X_\xi = \prod_{\lambda \in \xi} X_\lambda$ .

Then, the following theorem is obtained in a way parallel to Pasynkov [5] and Yajima [3].

THEOREM A.3. If  $X$  is piecewise cylindrical, then  
 (\*\*)  $\dim X = \text{Sup} \{ \dim X_\xi : \xi \text{ is a finite subset of } \Lambda \}$ .

Note that our definition is slight different from the piecewise rectangularity of infinite product due to Pasynkov [5]. The following result shows that it is sometimes easy to see piecewise cylindricality instead of piecewise rectangularity.

Theorem A.4 (Filippov [2]). Every Cartesian product of paracompact  $p$ -spaces is piecewise cylindrical.

REMARK A.2. Filippov [2] announced that it is indeed piecewise rectangular in the sense of Pasynkov. But, the proof given there is the one for our Theorem A.3.

COROLLARY A.1 (E. Pol [2]). Every Cartesian product of metric spaces satisfies (\*\*).

REMARK A.3. It is known in Anderson and Kleisler [1] that for every  $n$  there exists an  $n$ -dimensional separable metric space  $X$  such that every finite product  $X^z$  is (and, hence countable product  $X^\omega$  is)  $n$ -dimensional.

We conclude this section with the following problem:

PROBLEM A.2. Does every Cartesian product of paracompact  $\Sigma$ -spaces satisfy (\*\*)?

When we assume countable tightness, it is the case (Yajima [2]). It is also known that every finite product of paracompact  $\Sigma$ -spaces is rectangular (Pasynkov [2], and see also, Nagami [2]).

## Acknowledgements.

My first experience of really understanding and doing mathematics came in the seminar given in Tokyo by Professor Y. Kodama. It is Professor K. Nagami, who gave me the courage to study Wage-type examples. It is my pleasure to acknowledge my indebtedness to Professors Kodama and Nagami. I would also like to sincerely thank the following people for their cheerful instruction and information, their enthusiasm, and their unstinting encouragement: R. Cauty, K. Chiba, E. K. van Douwen, R. Engelking, T. Goto, Y. Hattori, J. E. Jayne, H. J. K. Junnila, Y. Kodama, S. Mardešić, V. Mardešić, K. Nagami, J. Nagata, K. Nogura, T. Nogura, H. Ohta, E. Pol, M. E. Rudin, S. Spieź, A. H. Stone, K. Tamano, J. Terasawa, Y. Yajima, and A. Waśko. Special thanks are due to my father Professor M. Tsuda, who has read the manuscript through, and has suggested many improvements for this thesis.

BIBLIOGRAPHY



R. D. Anderson & J. E. Keisler

- 1 An example in dimension theory, Proc. Amer. Math. Soc. 18(1967)709-713.

C. R. Borges

- 1 On stratifiable spaces, Pacific J. Math. 17(1966)1-16.

K. Borsuk

- 1 Some remarks on the Cartesian product of two 1-dimensional spaces, Bull. Acad. Polon. Sci. 23 (1975)971-973.

K. Borsuk, S. Nowak & S. Spieź

- 1 Remarks on the  $n$ -dimensional geometric measure of compacta, Fund. Math. 121(1983)59-71.

R. Cauty

- 1 Convexité topologique et prolongement des fonctions continues, Compositio Math. 27(1973)233-271.
- 2 A letter to the author on June 20, 1984.

E. Čech

- 1 Contribution to dimension theory, Časopis Pěst. Mat. Fys. 62(1933)277-291.

J. G. Ceder

- 1 Some generalizations of metric spaces, Pacific J. Math. 11(1961)105-126.

M. B. Charalambous

- 1 An example concerning inverse limit spaces of normal spaces, Proc. Amer. Math. Soc. 78(1980)605-607.
- 2 The dimension of inverse limit and N-compact spaces, ibid. 85(1982)648-652.

K. Chiba

- 1 Two remarks on the normality of product spaces, Rep. Fac. Sci. Shizuoka Univ. 11(1976)17-22.
- 2 A letter to the author on January 9, 1985.

T. Chiba & K. Chiba

- 1 Q-paracompactness and closed mappings, Sci. Rep. Tokyo Kyoiku Daigaku, Ser. A11(1972)230-234.
- 2 A note on normality of product spaces, ibid. 12(1974) 165-173.

E. K. van Douwen

- 1 A technique for constructing honest, locally compact,  $\beta$ -submetrizable examples, unpublished manuscript.
- 2 Mild infinite dimensionality of  $\beta X$  and  $\beta X - X$  for metrizable  $X$ , manuscript.

E. K. van Douwen & T. C. Przymusiński

- 1 First countable, and countable spaces all compactifications which contain  $\beta N$ , Fund. Math. 102(1979)229-234.

A. Dow & J. van Mill

- 1 An extremally disconnected Dowker space, Proc. Amer. Math. Soc. 86(1982)669-672.

R. Duda

- 1 The origins of the concept of dimension, Colloq. Math. 42(1979)95-110.

R. Engelking

- 1 General Topology (PWN, Warsaw, 1977).
- 2 Dimension Theory (North-Holland, Amsterdam, 1978).
- 3 A letter to the author on October 27, 1983.

R. Engelking & E. Pol

- 1 Countable-dimensional spaces: a survey, Dissertationes Math. 216(1983)1-45.

V. V. Fedorčuk

- 1 On the dimension of hereditarily normal spaces, Proc. London Math. Soc. 36(1978)153-175.
- 2 Infinite dimensional compact Hausdorff spaces, Izv. Akad. Nauk 42(1978)1162-1178 = Math. USSR-Izv. 13 (1979)445-460.

V. V. Filippov

- 1 On the dimension of products of topological spaces, Fund. Math. 105(1980)181-212.
- 2 On normally situated subspaces, in: P. S. Alexandrov (ed.), Topology (Moscow 1979) Trudy Mat. Inst. Steklov AN SSSR 154(1983)239-251.

L. Gillman & H. Jerison

- 1 Rings of continuous functions (van Nostrand, New York, 1960).

G. Gruenhage & E. Pol

- 1 Perfectly normal spaces and its applications to dimension theory, Fund. Math. 118(1983)213-222.

R. W. Heath & H. J. K. Junnila

- 1 Stratifiable spaces as subspaces and continuous images of  $M_1$ -spaces, Proc. Amer. Math. Soc. 83(1981) 146-148.

T. Hoshina

- 1 Products of normal spaces with Lašnev space, Fund. Math. 124(1984)143-153.

T. Hoshina & K. Morita

- 1 On the rectangular products of topological spaces, Top. and Appl. 11(1980)47-57.

W. Hurewicz & H. Wallman

1 Dimension Theory (Princeton Univ. Press, Princeton, 1941).

J. R. Isbell

1 Uniform spaces, Amer. Math. Soc. Math. Surveys 12  
(Providence, R. I., 1964).

T. Ishiwata

1 Z-mappings and  $C^*$ -embeddings, Proc. Japan Acad. 45  
(1969)889-893.

M. Ito

1  $M_0$ -spaces are  $\mu$ -spaces, Tsukuba J. Math. 8(1984)77-80.

J.E. Jayne & C. A. Rogers

1 Upper semi-continuous set valued function, Acta Math.  
149(1982)87-125.

I. Juhász, K. Kunen & M. E. Rudin

1 Two more hereditarily separable non-Lindelöf spaces,  
Canad. J. Math. 28(1976)998-1005.

H. J. K. Junnila

1 A characterization of  $M_0$ -spaces, Proc. Amer. Math.  
Soc. 91(1984)481-484.

2 A letter to the author on August 29, 1984.

M. Katětov

1 A theorem on the Lebesgue dimension, Časopis Pěst.  
Mat. Fys. 75(1950)79-87.

Y. Kodama

- 1 On a problem of Alexandroff concerning the dimension of product spaces. I&II, J. Math. Soc. Japan 10(1958) 380-404 & ibid. 11(1959)94-111.
- 2 Test spaces for homological dimension, Duke Math. J. 29(1962)41-50.
- 3 Some characterizations of homological dimension, Michigan Math. J. 9(1962)167-172.
- 4 On subset theorems and the dimension of products, Amer. J. Math. 91(1969)486-497.
- 5 On embeddings of spaces into ANR and shape, J. Math. Soc. Japan 27(1975)533-544.
- 6 Several problems in General Topology (in Japanese), 30th Topology Symposium Abstracts (1982. 7. 20-22, Yamanashi Univ.)116-125.

K. Kuratowski

- 1 Topology I (Academic Press, New York, 1966).

J. Krasinkiewicz

- 1 Essential mappings onto products of manifolds, preprint.

K. Kunen

- 1 Set Theory (North-Holland, Amsterdam, 1980).

V. I. Kuz'minov

- 1 Homological dimension theory, Uspekhi Mat. Nauk 23 (1968)3-49 = Russian Math. Surveys 23(1968)1-45.

H. Lebesgue

- 1 Sur la non applicabilité de deux domaines appartenant respectivement à des espaces, de  $n$  et  $n+p$  dimension, Math. Ann. 70(1911)166-168.

S. Lipscomb

- 1 On imbedding finite-dimensional metric spaces, Trans. Amer. Math. Soc. 211(1975)143-160.

S. Mardešić

- 1 On covering dimension and inverse limits of compact spaces, Illinois J. Math. 4(1960)278-291.

S. Mazurkiewicz

- 1 Sur les problèmes  $\kappa$  et  $\lambda$  de Urysohn, Fund. Math. 10 (1927)311-319.

E. Michael

- 1 The product of a normal space and a metric space need not be normal, Bull. Amer. Math. Soc. 69(1963) 375-376.
- 2 Paracompactness and the Lindelöf property in finite and countable Cartesian products, Compositio Math. 23(1971)119-214.

T. Mizokami

- 1 On  $M$ -structures, Top. and Appl. 17(1984)63-89.

K. Morita

- 1 On the dimension of product spaces, Amer. J. Math. 75(1953)205-223.
- 2 Normal families and dimension for metric spaces, Math. Ann. 128(1954)350-362.
- 3 A condition for the metrizability of topological spaces and for  $n$ -dimensionality, Sci. Rep. Tokyo Kyoiku Daigaku Sec. A 5(1955)33-36.
- 4 On the dimension of the product of Tychonoff spaces, Gen. Top. and Appl. 3(1973)125-133.
- 5 Čech cohomology and covering dimension for topological spaces, Fund. Math. 87(1975)31-52.
- 6 On the dimension of the product of topological spaces, Tsukuba J. Math. 1(1977)1-6.

S. Mrówka

- 1 On completely regular spaces, Fund. Math. 41(1954) 105-106.
- 2 Some comments on the author's example of non- $R$ -compact spaces, Bull. Acad. Polon. Sci. 18(1970)443-448.
- 3 Some set-theoretic constructions in topology, Fund. Math. 94(1977)83-92.
- 4 Realcompactness,  $N$ -compactness, and dimension theory, Abstracts Amer. Math. Soc. 3(1982)p. 95 No. 792-54-48.



K. Nagami

- 1 Mappings of finite order and dimension theory, Jap. J. of Math. 30(1960)25-54.
- 2 A note on the large inductive dimension of totally normal spaces, J. Math. Soc. Japan 21(1969)282-290.
- 3 A note on the normality of inverse limits and products, Proc. Intern. Symp. Topol. Appl. Herceg-Novri 1968, Beograd (1969)261-264 (Academic Press, New York).
- 4 Dimension Theory (Academic Press, New York, 1970).
- 5 Countable paracompactness of inverse limits and products, Fund. Math. 73(1972)261-270.
- 6 The equality of dimension, *ibid.* 106(1980) 239-246.
- 7 Dimension of free L-spaces, *ibid.* 108(1980) 211-224.
- 8 Dimension of non-normal spaces, *ibid.* 109(1980) 113-121.

K. Nagami & K. Tsuda

- 1 Complexes and L-structures, J. Math. Soc. Japan 33 (1981)639-648.

J. Nagata

- 1 Note on dimension theory for metric spaces, *Fund. Math.* 45(1958)143-181.
- 2 Product theorems in dimension theory. I, *Bull. Acad. Polon. Sci.* 15(1967)439-448.
- 3 A survey of dimension theory, in: J. Novak (ed.) *General Topology and its Relations to Modern Analysis and Algebra II*, Proc. Pragh Topol. Symp. (1966) 259-270 (Academic Press, New York).
- 4 A survey of dimension theory II, *Gen. Top. and Appl.* 1(1971)439-448.
- 5 Recent developments in general topology I & II, *Eleutheria* 1(1978)59-71 & *ibid.* 2(1979)447-458.
- 6 A survey of dimension theory III, in: P. S. Alexandrov (ed.), *Topology (Moscow, 1979)* *Trudy Mat. Inst. Steklov AN SSSR* 154(1983)186-199.
- 7 *Modern dimension theory (revised and extended edition)*, (Herderman Verlage, Berlin, 1983).

N. Noble

- 1 A note on  $z$ -closed projections, *Proc. Amer. Math. Soc.* 23(1969)73-76.

H. Ohta

- 1 On normal, non-rectangular products, *Quart. J. Math. Oxford (2)*, 32(1981)229-344.
- 2 Rectangular products with a metric factor, preprint.

S. Oka

- 1 Dimension of stratifiable spaces, Trans. Amer. Math. Soc. 275(1983)231-243.

B. A. Pasynkov

- 1 On the dimension of products of normal spaces, Dokl. Akad. Nauk 209(1973)792-794 = Soviet Math. Dokl. 14 (1973)530-533.
- 2 On the dimension of rectangular products, Dokl. Akad. Nauk 221(1975)291-294 = Soviet Math. Dokl. 16(1975)344-347.
- 3 Factorization theorems in dimension theory, Uspekhi Mat. Nauk 36(1981)147-175 = Russian Math. Surveys 36(1981) 175-209.
- 4 On the monotonicity of dimension, Dokl. Akad. Nauk 267 = Soviet Math. Dokl. 26(1982)654-658.
- 5 On dimension theory, in: I. M. James & E. H. Kronheimer (ed.) Aspects of Topology (In memory of Hugh Dowker 1912-1982) London Math. Soc. Lecture Note Ser. 93 (1985)227-250.

G. Peano

- 1 Sur une courbe qui remplit toute une aire plane, Math. Ann. 36(1890)157-160.

A. R. Pears

- 1 Dimension theory of general spaces (Cambridge Univ. Press, Cambridge 1975).

E. Pol

- 1 A remark about the Juhász-Kunen-Rudin construction of a hereditarily separable non-Lindelöf space, Bull. Acad. Polon. Sci. 24(1976)749-751.
- 2 On the dimension of the product of metrizable spaces, *ibid.* 24(1978)525-534.

R. Pol

- 1 A weakly infinite-dimensional compactum which is not countable-dimensional, Proc. Amer. Math. Soc. 82 (1981)634-636.

L. S. Pontrjagin

- 1 Sur une hypothèse fondamentale de la théorie de la dimension, C. R. Acad. Paris 190(1930)1105-1107.

T. C. Przymusiński

- 1 Topological properties of product spaces and the notion of  $n$ -cardinality, Topol. Proceedings 2(1977) 233-241.
- 2 On the notion of  $n$ -cardinality, Proc. Amer. Math. Soc. 69(1978)333-338.
- 3 On the dimension of product spaces and an example of M. Wage, *ibid.* 76(1979)315-321.
- 4 Normality and paracompactness in finite and countable Cartesian products, Fund. Math. 105(1980)87-104.

- 5 Product of perfectly normal spaces, *ibid.* 108(1980) 129-136.
- 6 Product spaces, in : G.M. Reed (ed.), *Surveys in Gen. Topol.* (Academic Press, New York, 1980)399-429.
- 7 A solution to a problem of E. Michael, *Pacific J. Math.* 114(1984)235-242.

L. Rubin, R. Scori & J. Walsh

- 1 New dimension-theory techniques for constructing infinite dimensional examples, *Gen. Top. and Appl.* 10(1979)93-102.

M.E. Rudin

- 1 A normal space  $X$  for which  $X \times I$  is not normal, *Fund. Math.* 73(1971)179-186.
- 2 A letter to the author on August 7, 1984.

M. E. Rudin & M. Starbird

- 1 Products with a metric factor, *Gen. Top. and Appl.* 5(1975)235-248.

S. Shelah & M. E. Rudin

- 1 Unordered types of ultrafilters, *Topol. Proceedings* 3(1978)199-204.

M. Starbird

- 1 Extending maps from products, in: N. M. Stavrakas & K. R. Allen (ed.), *Studies in Topology* (Academic Press, New York, 1975) 559-565.

A. H. Stone

- 1 Non-separable Borel sets, *Rozprawy Mat.* (= *Dissertationes Math.*) 28(1962)1-40.
- 2 A letter to the author on April 22, 1985.

H. Tamano

- 1 A note on the pseudocompactness of product of two spaces, *Mem. Coll. Sci. Univ. Kyoto Ser. A Math.* 33 (1960/61) 225-230.

K. Tamano

- 1 A note on E. Michael's example and rectangular products, *J. Math. Soc. Japan* 34(1982)187-190.

J. Terasawa

- 1 On the zero-dimensionality of some non-normal product spaces, *Sci. Rep. Tokyo Kyoiku Daigaku Ser. A* 11(1972) 95-102.
- 2 Spaces  $N \cup R$  need not be strongly 0-dimensional, *Bull. Acad. Polon. Sci.* 25(1977)279-281.
- 3 Spaces  $N \cup R$  and their dimension, *Top. and Appl.* 11 (1980)93-102.

K. Tsuda

- 1 Some examples concerning the dimension of product spaces, *Math. Japonica* 27(1982)177-195.
- 2 An  $n$ -dimensional version of Wage's example, *Colloq. Math.* 49(1984)15-19.
- 3 A Wage-type example with a pseudocompact factor, to appear in *Top. and Appl.*
- 4 A note on closed embeddings of metric spaces, to appear in *Bull. London Math. Soc.*
- 5 A note on closed embeddings of metric spaces II, submitted.
- 6 Non-existence of universal spaces for some stratifiable spaces, *Topol. Proceedings* 9(1984)165-171.
- 7 Rectangularity versus piecewise rectangularity of product spaces, preprint.

M. Wage

- 1 The dimension of product spaces, unpublished manuscript, 1976, Department of Mathematics, Yale University.
- 2 The dimension of product spaces, *Proc. Natl. Acad. Sci. U. S. A.* 75(1978)4671-4672.
- 3 Products of Radon spaces, *Uspekhi Mat.* 35(1980)151-153  
= *Russian Math. Surveys* 35(1980)185-187.

J. J. Walsh

- 1 Infinite dimensional compacta containing no  $n$ -dimensional ( $n \geq 1$ ) subsets, *Topology* 18(1979) 91-95.

A. Waśko

- 1 Spaces universal under embeddings for finite dimensional complete metric spaces, preprint.

Y. Yajima

- 1 Topological games and products II, *Fund. Math.* 117 (1983)47-60.
- 2 On  $\Sigma$ -products of  $\Sigma$ -spaces, *Fund. Math.* 123(1984) 29-37.
- 3 On the dimension of limits of inverse systems, *Proc. Amer. Math. Soc.* 91(1984)461-466.

V. P. Zolotarev

- 1 On the dimension of subspaces, *Vestnik Moskovskogo Universiteta. Matematika* 30(1975)10-12 = *Moscow Math. Bull.* 30(1975)7-8.