

**Significant Correlation between Endothelial nitric oxide synthase (eNOS) Expression and Alveolar Repair in Elastase-induced Rat Pulmonary Emphysema**

**ST-2011-1151-EO.R2**

**Running head: Increased eNOS and Alveolar Repair**

Hiromichi Ito <sup>1</sup>, Shonosuke Matsushita <sup>1</sup>, Shigemi Ishikawa<sup>2</sup>, Yukinobu Goto<sup>1</sup>, Mitsuaki Sakai<sup>1</sup>, Masataka Onizuka<sup>1</sup>, Yukio Sato<sup>1</sup>, and Yuzuru Sakakibara<sup>1</sup>

<sup>1</sup>Graduate School of Comprehensive Human Science, Department of Thoracic Surgery, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan

<sup>2</sup>Department of Chest Surgery, International University of Health and Welfare Hospital, Nasu-Shiobara, Tochigi 329-2763, Japan

Reprint requests to: S. Matsushita

Telephone: +81-29-853-7627

Fax: +81-29-853-3097

e-mail address: [shomatsu@md.tsukuba.ac.jp](mailto:shomatsu@md.tsukuba.ac.jp)

First author: [hiromi396@aol.com](mailto:hiromi396@aol.com)

**Key words**

neovascularization, emphysema, eNOS, angiogenic factor, G-CSF

## **Abstract**

### **Background and Purpose:**

Angiogenic factors, such as endothelial nitric oxide synthase (eNOS), are thought to play an important role in the repair of pulmonary emphysema (PE); yet, the correlation of the factors involved has not been investigated. We conducted this study to clarify the positive correlation between eNOS expression and alveolar repair in PE recovery.

### **Methods:**

We used elastase to induce PE in rats, which were divided into Groups A (Control), B (G-CSF), C (PE) and D (PE + G-CSF). G-CSF was injected for 12 days, 4 weeks after which the alveolar walls, arterioles, and angiogenic factors including eNOS were examined histopathologically and by western blotting.

### **Results:**

In comparing Groups A, B, C, and D, the alveolar density was  $2.4\pm 0.2$ ,  $2.4\pm 0.1$ ,  $1.8\pm 0.1$ ,  $2.5\pm 0.1$  per  $100\ \mu\text{m}^2$ , respectively (C vs. others;  $p < 0.00001$ ) and the number of arterioles was  $4.5\pm 1.0$ ,  $5.6\pm 0.6$ ,  $3.2\pm 0.5$ ,  $5.5\pm 0.7$  / $\text{mm}^2$ , respectively (C vs. others;  $p < 0.05$ ). Immunohistochemical staining (IHC) revealed different eNOS expression in Group D vs. Group C ( $p < 0.0001$ ) and western blotting revealed different eNOS, VEGF, and FLT-1 expression in Group D vs. Group C ( $p < 0.01$ ,  $p < 0.05$ ,  $p < 0.001$ ), reflecting the contribution of angiogenesis to PE repair. eNOS showed a significantly positive correlation to alveolar density and arteriole repair.

### **Conclusion:**

Alveolar repair was correlated positively to eNOS expression by vascular regeneration in elastase-induced rat pulmonary emphysema.

## **Introduction**

Pulmonary emphysema (PE) is one of the most common causes of respiratory failure and a frequent cause of death. Severe PE can predispose to pneumonias and the perioperative management of respiratory conditions complicating severe PE is difficult in thoracic surgery. Postoperatively, persistent PE leads to poor pulmonary function and dyspnea on effort, restricting the activities of daily life. Patients who undergo lung resection with PE suffer even further compromised quality of life (QOL)<sup>1</sup>. PE involves decreased elasticity of the airway and alveolar wall destruction, including chronic inflammation, with remodeling of the alveolar area<sup>2</sup>. The precise mechanisms behind these pathological changes and lung repair remain unclear<sup>3</sup>. A better understanding of the mechanism of regeneration from PE will make lung resection safer and improve QOL after surgery.

Recent studies have found that bone marrow-derived cells and angiogenic cytokines are related to lung repair in PE. Cytokines such as G-CSF, vascular endothelial growth factor (VEGF), angiopoietin-1, and erythropoietin are increased by the inflammation and ischemia in acute lung disease, triggering the mobilization of EPCs from the bone marrow to the injured lung tissue<sup>4</sup>. The importance of vascular regeneration in lung repair in PE has also been shown<sup>5-7</sup>. Endothelial nitric oxide synthase (eNOS), one of the most important angiogenic factors, is located downstream of VEGF. Reduced eNOS expression and fewer endothelial cells in the pulmonary vasculature of PE have been reported<sup>8</sup>. Conversely, enhanced eNOS expression in the residual lung after pneumonectomy with alveolar wall regeneration has been reported<sup>9</sup>. However, although the importance of eNOS expression in lung repair is recognized, the

correlation between direct alveolar repair and eNOS expression through neovascularization remains unclear. We hypothesized that alveolar repair in PE is positively correlated to eNOS expression by vascular regeneration. In this study, G-CSF was used as the angiogenic factor inducing mobilizing endothelial progenitor cells (EPCs)<sup>10</sup> and enhancing eNOS.

## **Methods**

### ***Animals***

The Animal Research Committee at the University of Tsukuba approved the experimental protocols. Rats were cared for in accordance with the Guiding Principles for the Care and Use of Animals based on the Helsinki Declaration of 1964. The 6-week-old male Wistar rats (BW: 200g, n=38, Charles River Japan, Yokohama, Japan) were divided into four groups according to the induction of PE and the use of G-CSF (GRAN<sup>R</sup>, Kirin Pharma, Japan) as follows: Group A (Control; non-PE treated with normal saline (NS), n=9), Group B (non-PE treated with G-CSF, n=11), Group C (PE treated with NS, n=9), and Group D (PE treated with G-CSF, n=9).

### ***Surgical Procedure***

Anesthesia was induced by diethyl ether (Wako Chemicals, Japan) inhalation, followed by an intraperitoneal injection of pentobarbital (Somnopentyl<sup>R</sup>: 50 mg/kg, Kyoritsu Seiyaku Corporation, Japan). A skin incision was made to expose the trachea, the front of which was punctured by a 27-gauge needle to inject porcine-derived elastase (200 U/kg, ELASTIN PRODUCTS COMPANY. INC, MO, U.S.A.). Rats were

temporarily placed upright at 70 degrees. The porcine-derived elastase was dissolved in 0.1 ml normal saline and injected over 60 s in Groups C and D. The same volume of normal saline was injected in the control non-PE groups (Groups A and B).

### ***Experimental Protocol***

Three weeks after the injection of elastase, G-CSF (100 µg/kg/day) was injected subcutaneously for 12 days. The rats were killed 9 weeks after the injection of elastase and the lungs were removed and fixed with an intratracheal injection of 10% formalin solution under constant pressure of 20 cmH<sub>2</sub>O. We examined the specimens histopathologically (HE stain, eNOS immunohistochemistry) and also stored frozen specimens for western blotting.

### ***Examination of the Alveolar Wall and Arterioles***

PE was evaluated by the alveolar density per unit (septum frequency / 100 µm<sup>2</sup>) and the mean linear intercept (Lm) using modified point counting methods<sup>11</sup>, as follows: *Alveolar density per unit* (septum frequency / 100 µm<sup>2</sup>): Alveolar density was based on the modified point counting method of Leuwerke<sup>9, 12</sup>. Using HE staining, the points at which the alveolar walls crossed the borders of 10 × 10 lattices were counted continuously. The number of crossing points was averaged by the number of lattices in efficient alveolar areas and this process was performed using light microscopy in 100 magnifications. The lattices were 100 µm in size. Lattices in non-alveolar areas containing collapsed or inflamed alveolar areas and bronchi, cartilage, and vessels were eliminated (Fig. 1a). We calculated 15 randomly selected fields.

*Mean Linear Intercept (Lm)*: The extent of emphysematous lesions was assessed

by measuring the mean linear intercept (Lm) using the method of Thulbeck with modification<sup>13, 14</sup>. Briefly, Lm was defined as the linear sum of all lines in all frames counted and divided by the number of intercepts defined as an alveolar septa intersecting with a counting line. A minimum of 10 fields and 200 intercepts was measured for each rat<sup>15</sup>.

Arterioles were defined as arteries more than 50  $\mu\text{m}$  and less than 400  $\mu\text{m}$  in diameter. Arterioles were evaluated by the number within 1  $\text{mm}^2$  in each lattice (Fig.1a).

### ***Immunohistochemistry***

Paraffin-embedded sections of the lung were deparaffinized with xylene and rehydrated with ethanol. Antigens were activated by thermal microwave radiation. Sections were blocked by 3%  $\text{H}_2\text{O}_2$  for 15 min and incubated overnight with anti-eNOS antibody (Lab Vision Corporation, Fremont, CA, USA); then for another 30 min with the universal immuno-peroxidase polymer for rat tissue sections (Nichirei, Tokyo, Japan). The antibody-polymer-peroxidase complexes were visualized using DAB (Simple Stain DAB solution, Nichirei, Tokyo, Japan) and nuclei were counterstained with hematoxylin. Images were captured using a high-resolution digital camera mounted on a microscope (Olympus BX-40F, Tokyo, Japan). Analysis was performed on approximately eight capillaries per section of the left middle lobe of each rat lung at 1,000 times magnification. Each capillary image was converted into black and white using the graphic software Photoshop Element (Microsoft, Seattle, WA. USA). The 256-grayscale value of capillary endothelial cell bodies, excluding the nuclei, was evaluated using the graphic software ImageJ (NIH, USA). The difference in the

grayscale value of each group was regarded as the difference in eNOS expression for semi-quantitative analysis. High grayscale values indicate a high density of eNOS.

### ***Western Blotting***

The homogenized lung tissue was lysed on ice with RIPA buffer. Samples were run on SDS-PAGE, using 7.5-10% polyacrylamide gel (Wako Pure Chemical Industries, Ltd, Osaka, Japan), and electrotransferred to a polyvinylidene difluoride (PVDF, Paul Corporation, Pensacola, FL, U.S.A.) filter membrane. The PVDF membrane was blocked for 1 h at room temperature with 1% bovine serum albumin (Wako Pure Chemical Industries, Ltd, Osaka, Japan) in PBS containing 0.1% Tween 20 (TPBS). Thereafter, the PVDF membrane was incubated overnight at 4°C with primary antibodies in TPBS. After washing with TPBS, the PVDF membrane was incubated with horseradish peroxidase-conjugated anti-rabbit antibody (GE Healthcare Japan, Tokyo, Japan), and diluted to 1:6000 in TPBS at room temperature for 60 min. The blots were visualized with an enhanced chemiluminescence detection system (Amersham Biosciences, UK), exposed to X-ray film, and analyzed by the graphic software, ImageJ, using the 256-grayscale value. The following antibodies were used: Anti-eNOS antibody (AnaSpec, Inc, San Jose, CA, USA), anti-VEGF antibody (Santa Cruz Biotechnology, Santa Cruz, CA, USA), anti-Flt-1 antibody (Thermo Fisher Scientific, Fremont, CA, USA), and anti- $\beta$ -actin antibody (AnaSpec, Inc. San Jose, CA, USA).

### ***Statistical Analysis***

All variables are expressed as means  $\pm$  SD. All groups were analyzed statistically by the Student's unpaired t-test with Bonferroni correction. Specific groups were compared using an unpaired Student's test. Simple linear regression analysis was

performed for each principal factor. A difference of  $p < 0.05$  was considered significant. We used the statistical software SPSS (SPSS Inc., Chicago, Ill. USA).

## **Results**

### ***Evaluation of Alveolar Change***

#### *Alveolar Density (Septum Frequency)*

The number of alveolar density crossing points on lattices in Groups A, B, C, and D was  $2.4 \pm 0.2$ ,  $2.4 \pm 0.1$ ,  $1.8 \pm 0.1$ , and  $2.5 \pm 0.1$  (unit: points/100  $\mu\text{m}^2$ ), respectively. There were no significant differences between Groups A and B (NS), but the number of lattices was significantly higher in Group D than in Group C ( $P < 0.00001$ ) and significantly lower in Group C than in the other three groups ( $p < 0.00001$ ; Fig. 1b).

#### *Mean linear intercept (Lm)*

The Lm in Groups A, B, C, and D were  $92.4 \pm 7.0$ ,  $95.2 \pm 4.7$ ,  $125.1 \pm 8.8$ , and  $91.7 \pm 4.4$  ( $\mu\text{m}$ ), respectively. There were no significant differences between Groups A and B (NS), but the Lm was significantly lower in Group D than in Group C ( $P < 0.00001$ ) and significantly higher in Group C than in the other three groups ( $p < 0.00001$ ; Fig. 1c).

### ***Number of Arterioles***

The average number of arterioles in Groups A, B, C, and D was  $4.5 \pm 1.0 /\text{mm}^2$ ,  $5.6 \pm 0.6 /\text{mm}^2$ ,  $3.2 \pm 0.5 /\text{mm}^2$ , and  $5.5 \pm 0.7 /\text{mm}^2$ , respectively. Although there were significantly fewer arterioles in Group C than in Groups A or B, there were significantly more arterioles in Group D than in Group C (Group C vs. Group A, group B, and Group

D;  $p < 0.05$ ,  $p < 0.0001$ , and  $p < 0.0001$ , respectively; Fig. 1d).

### ***eNOS Expression According to Immunohistochemistry and Western Blotting of the Angiogenic Factors***

The levels of eNOS expression in the capillary endothelial cells in Groups A, B, C, and D were  $59.1 \pm 8.9$ ,  $62.9 \pm 13.6$ ,  $35.1 \pm 6.4$ , and  $61.7 \pm 7.3$ , respectively (Group C vs. Groups A and D;  $p < 0.0001$ ,  $p < 0.0005$ , respectively; Fig. 2). Enhanced eNOS expression in the alveolar capillaries was accompanied by a higher number of arterioles in Group D than in Group C (Figs 1e, 2, 3). The eNOS expression in Groups C and D according to western blotting was  $0.13 \pm 0.06$  and  $0.6 \pm 0.2$ , respectively ( $p < 0.01$ ; Fig. 3a). The VEGF expression in Groups C and D according to western blotting was  $0.13 \pm 0.06$  and  $0.70 \pm 0.20$ , respectively ( $p < 0.05$ ; Fig. 3b). FLT-1 expression was similar, being  $0.25 \pm 0.1$  in Group C vs.  $1.56 \pm 0.2$  in Group D ( $p < 0.001$ ; Fig. 3c).

### ***Evaluation by Simple Linear Regression***

Simple linear regression analyses showed significant correlations between eNOS expression and alveolar wall counts, being alveolar density ( $r = 0.648$ ;  $p < 0.005$ ; Fig. 4a); arteriole counts and alveolar wall counts, being alveolar density ( $r = 0.546$ ;  $p < 0.005$ ; Fig. 4b); and eNOS expression and arteriole counts ( $r = 0.670$ ;  $p < 0.005$ ; Fig. 4c). Moreover, e-NOS expression, the number of arterioles, and the number of alveolar walls were all positively correlated with each other.

## **Discussion**

Elastase-induced emphysema is a simple experimental model, designed to allow us

to study emphysema<sup>16</sup>. It can be established within approximately 2 weeks and is particularly useful for evaluating regeneration to assess the effectiveness of drugs administered after the development of emphysema<sup>3</sup>.

In this experiment, injecting elastase into the trachea induced PE by disrupting the alveolar wall and reducing the number of arterioles with lowered eNOS expression and VEGF axis. We administered G-CSF as an angiogenic promoter, resulting in the repair of arteriole structure and alveolar wall density. Alveolar wall density and vascular development (arteriole number) was found to be significantly correlated with the eNOS expression of the alveolar capillary endothelium. It was recently recognized that apoptosis of endothelium, leading to the loss of capillaries, may be a central mechanism in emphysema<sup>5</sup>. Other reports have described impaired endothelial function in the small pulmonary arteries from patients with COPD<sup>17, 18</sup>. Accordingly, fewer arterioles were detected in our group C than in the other groups.

G-CSF as a potent angiogenic promoter acts by mobilizing endothelial progenitor cells (EPCs)<sup>19</sup>, which are essential for vasculogenesis<sup>20</sup>. This activation of angiogenesis, might have contributed to the recovery from PE. Nitric oxide (NO) is an important cellular messenger molecule with pleiotropic effects including vasodilatation, anti-oxidation, anti-inflammation, mobilization of EPC, and neovascularization<sup>21</sup>. The activation of eNOS and the production of NO, which is triggered by a serine/threonine protein kinase (Akt) signaling axis, is one of the most potent endothelial functions<sup>22</sup>. In endothelial cells, the Akt signaling axis is activated by a variety of stimuli and growth factors such as VEGF or angiopoietin-1 that regulate angiogenesis, endothelial cell survival, and vessel integrity by controlling NO synthesis<sup>22</sup>.

Arteriogenesis is reported to be enhanced by increased eNOS expression<sup>23</sup>,

whereas G-CSF is reported to induce eNOS in endothelial cells<sup>10</sup>. In this experiment, eNOS contributed to vascular regeneration in addition to the VEGF axis. In relation to PE, it has been speculated that eNOS is capable of regenerating the alveolar wall<sup>9</sup>. Consequently, we focused on the role of eNOS in lung repair from PE from the following perspectives: the pleiotropic effects of NO; eNOS and NO as essential factors for angiogenesis; and the regenerative effects of eNOS on cells, including cells in the alveolar wall.

Although eNOS expression in PE repair has been emphasized<sup>3,8</sup>, its correlation has not been proven (Fig. 4). Thus, eNOS expression in the capillaries was positively correlated with the number of alveolar walls and arterioles. Fig. 4 shows that the angiogenic effect is one of the most important aspects of eNOS in lung regeneration from PE. Other studies have found that statins have a protective effect on alveolar and capillary apoptosis in PE<sup>8</sup>. This effect is mediated by Akt and eNOS activation<sup>3,7</sup>. The importance of the angiogenic function of eNOS on PE repair was also apparent in our study.

The findings of this study provide solid evidence that eNOS plays an important role in vascular regeneration in lung repair. Drugs enhancing eNOS may become a strategy for the treatment of PE, focusing on regeneration of the pulmonary vasculature and may help surgical patients with PE.

## **Conclusion**

Alveolar repair is correlated positively with eNOS expression through vascular regeneration in elastase-induced rat pulmonary emphysema.

## **Acknowledgment**

We thank Mr. Avi Landau for his language consultation. This study was supported in part by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (grant C-20591471).

### **Conflict of Interest Statement**

We have no conflicts of interest.

## References

1. Fernandes EO, Teixeira C, Silva LC. Thoracic surgery: risk factors for postoperative complications of lung resection. *Rev Assoc Med Bras* 2011;57:292-8.
2. Burrows B, Fletcher CM, Heard BE, Jones NL, Wootliff JS. The emphysematous and bronchial type of chronic airways obstruction. A clinicopathological study of patient in London and Chicago. *Lancet* 1996;1:1355-60.
3. Takahashi S, Nakamura H, Seki M, Shiraishi Y, Yamamoto M, Furuuchi M. Reversal of elastase-induced pulmonary emphysema and promotion of alveolar epithelial cell proliferation by simvastatin in mice. *Am J Physiol Lung Cell Mol Physiol* 2008;294:882-90.
4. Aicher A, Heeschen C, Mildner-Rihm C, Urbich C, Ihling C, Technau-Ihling K, et al. Essential role of endothelial nitric oxide synthase for mobilization of stem and progenitor cells. *Nat Med* 2003;9:1370-6.
5. Siafakas NM, Antoniou KM, Tzortzaki EG. Role of angiogenesis and vascular remodeling in chronic obstructive pulmonary disease. *Int J Chron Obstruct Dis* 2007;2:453-62.
6. Magee F, Wright JL, Wiggs BR, Paré PD, Hogg JC. Pulmonary vascular structure and function in chronic obstructive pulmonary disease. *Thorax* 1988;43:183-89.
7. Barberà JA, Peinado VI, Santos S, Ramirez J, Roca J, Rodriguez-Roisin R. Reduced

expression of endothelial nitric oxide synthase in pulmonary arteries of smokers. *Am J Respir Crit Care Med* 2001;164:709-13.

8. Lee JH, Lee DS, Kim EK, Choe KH, Oh YM, Shim TS, et al. Simvastatin inhibits cigarette smoking-induced emphysema and pulmonary hypertension in rat lungs. *Am J Respir Crit Care Med* 2005;172:987-93.

9. Leuwerke SM, Kaza AK, Tribble CG, Kron IL, Laubach VE. Inhibition of compensatory lung growth in endothelial nitric oxide synthase-deficient mice. *Am J Physiol Lung Cell Mol Physiol* 2002;282:L1272-8.

10. Park KW, Kwon YW, Cho HJ, Shin JI, Kim YJ, Lee SE, et al. G-CSF exerts dual effects on endothelial cells - opposing actions of direct eNOS induction versus indirect CRP elevation. *J Mol Cell Cardiol* 2008;45:670-8.

11. Dunnill, M.S. Evaluation of a simple method of sampling the lung for quantitative histological analysis. *Thorax* 1964;19:443-8.

12. Kaza AK, Kron IL, Kern JA, Long SM, Fiser SM, Nguyen RP, et al. Retinoic acid enhances lung growth after pneumonectomy. *Ann Thorac Surg* 2001;71:1645-50.

13. Thurlbeck WM. Internal surface area and other measurements in emphysema. *Thorax* 1967;22:483-96.

14. Heemskerk-Gerritsen BA, Dijkman JH, Ten Have-Opbroek AA. Stereological methods: a new approach in the assessment of pulmonary emphysema. *Microvasc Res Tech* 1996;34:556-62.
15. Yamada M, Kubo H, Kobayashi S, Ishizawa K, Numasaki M, Ueda S, et al. Bone marrow-derived progenitor cells are important for lung repair after lipopolysaccharide-induced lung injury. *J Immunol* 2004;172:1266-72.
16. Massaro GD, Massaro D. Retinoic acid treatment abrogates elastase-induced pulmonary emphysema in rats. *Nat Med* 1997;3:675-7.
17. Peinado VI, Barbera JA, Ramirez J, Gomez FP, Roca J, Jover L, et al. Endothelial dysfunction in pulmonary arteries of patients with mild COPD. *Am J Physiol* 1998;274:L908-13.
18. Kasahara Y, Tuder RM, Cool CD, Lynch DA, Flores SC, Voelkel NF. Endothelial cell death and decreased expression of vascular endothelial growth factor and vascular endothelial growth factor receptor 2 in emphysema. *Am J Respir Crit Care Med* 2001;163:737-44.
19. Ishizawa K, Kubo H, Yamada M, Kobayashi S, Numasaki M, Ueda S, et al. Bone marrow-derived cells contribute to lung regeneration after elastase-induced pulmonary emphysema. *FEBS Lett* 2004;556:249-52.

20. Werner N, Kosiol S, Schiegl T, Ahlers P, Walenta K, Link A, et al. Circulating endothelial progenitor cells and cardiovascular outcomes. *N Engl J Med* 2005;353:999-1007.
21. Duda DG, Fukumura D, Jain RK. Role of eNOS in neovascularization: NO for endothelial progenitor cells. *Trends Mol Med* 2004;10:143-5.
22. Dimmeler S, Aicher A, Vasa M, Mildner-Rihm C, Adler K, Tiemann M, et al. HMG-CoA reductase inhibitors (statins) increase endothelial progenitor cells via the PI 3-kinase/Akt pathway. *J Clin Invest* 2001;108:391-7.
23. Cai WJ, Kocsis E, Luo X, Schaper W, Schaper J. Expression of endothelial nitric oxide synthase in the vascular wall during arteriogenesis. *Mol Cell Biochem* 2004;264:193-200.

## Figure legends

**Fig.1** Alveolar walls and arterioles. **a** (upper section). Typical alveolar images of Groups A to D (HE stain, X100). The number of alveolar walls (alveolar wall density, septum frequency) were counted by point counting methods. **b, c.** G-CSF induced a recovery of the alveolar walls (Group C vs. Group D). **d.** The number of arterioles recovered significantly well following the administration of G-CSF to the rats with induced pulmonary emphysema (PE) (Group C vs. Group D,  $p < 0.05$ ).


**Fig.2** Immunohistochemical findings of eNOS expression. eNOS was dyed brown inside the vascular endothelial cells (arrow). The average 256 grayscale value of eNOS expression was significantly higher in the G-CSF treated rats than in the non-treated rats with pulmonary emphysema (PE; Group C vs. Group D,  $p < 0.00005$ ).

**Fig.3** Quantification of angiogenic factors. Pulmonary emphysema (PE) with G-CSF treatment significantly increased the angiogenic factors, VEGF and eNOS, vs. PE without G-CSF (Group C vs. D; Fig3a and 3b). Flt-1 expression showed the same tendency (Fig3c).


**Fig.4** Evaluation by simple linear regression. Significant correlations were found between the alveolar walls and eNOS expression (a), the alveolar walls and arterioles (b), and eNOS expression and the arterioles (c). From this series of correlations, we concluded that higher eNOS expression enhances alveolar repair through recovery of the arterioles.

**Figure 1**


**Fig.1a** Counting of Alveolar walls and arterioles HE X100




**Fig.1c** mean linear intercept (Lm)  
(Bonferroni correction)



**Fig.1b** alveolar wall density (septum frequency)  
(Bonferroni correction)



**Fig.1d** arterioles  
(Bonferroni correction)



**Figure 2**





Figure 3




Figure 4

