

LEVI-PARALLEL HYPERSURFACES IN A COMPLEX SPACE FORM

By

Jong Taek CHO

Abstract. In this paper, we classify a Hopf hypersurface in a non-flat complex space form whose Levi-form is parallel with respect to the generalized Tanaka-Webster connection.

1. Introduction

Let $\tilde{M} = (\tilde{M}^n, J, \tilde{g})$ be a complex n -dimensional Kählerian manifold with complex structure J and Kählerian metric \tilde{g} . Let M be an oriented real hypersurface in \tilde{M} , g be the induced metric and η be the 1-form defined by $\eta(X) = g(X, \xi)$ where $\xi = -JN$ and N is a unit normal vector field on M . Then M has an (integrable) CR-structure associated with the complex structure of the ambient space. Let TM be the tangent bundle of M and D be the subbundle of TM (or the $(2n - 2)$ -dimensional distribution) which is defined by $\eta = 0$. We denote by $CD = D \otimes \mathbb{C}$ its complexification. Then we see that D is *holomorphic* (or maximally invariant by J) and

$$\mathcal{H} = \{X - iJX : X \in D\}$$

defines an *CR-structure* on M . That is, \mathcal{H} satisfies the following properties:

- (i) each fiber \mathcal{H}_x ($x \in M$) is of complex dimension $n - 1$,
- (ii) $\mathcal{H} \cap \overline{\mathcal{H}} = \{0\}$,
- (iii) $[\mathcal{H}, \mathcal{H}] \subset \mathcal{H}$ (integrability).

Furthermore, we have $CD = \mathcal{H} \oplus \overline{\mathcal{H}}$. We call $\{D, J\}$ the real representation of \mathcal{H} . Then for $\{D, J\}$ we define the Levi form by

$$L : D \times D \rightarrow \mathcal{F}(M), \quad L(X, Y) = d\eta(X, JY)$$

Mathematics Subject Classification (2000): 53C15, 53C40, 53D15.

This study was financially supported by research fund of Chonnam National University in 2003.

Key words and phrases. Tanaka-Webster connection, Levi parallel hypersurfaces, complex space forms.

Received February 2, 2005.

where $\mathcal{F}(M)$ denotes the algebra of differentiable functions on M . If the Levi form is hermitian, then the CR-structure is called *pseudo-hermitian*, in addition, in the case that the Levi form is non-degenerate (positive or negative definite, resp.), then the CR structure is called a *non-degenerate (strongly pseudo-convex, resp.) pseudo-hermitian CR structure*. Recently, Y. T. Siu [14] proved the nonexistence of compact smooth Levi-flat hypersurfaces in complex projective spaces of dimension ≥ 3 . Here, it is remarkable that the assumption of compactness in Siu's theorem has a crucial role. Actually, there are non-complete examples which are Levi-flat in a complex projective space (see section 2). Anyway, the examples of Levi-flat hypersurfaces which are known so far are not Hopf. In this situation, we prove that there does not exist a Levi-flat Hopf hypersurface (Theorem 3).

On the other hand, the Tanaka-Webster connection ([19], [20]) is defined as a canonical affine connection on a pseudo-hermitian, non-degenerate, integrable CR manifold. For contact metric manifolds, their associated almost CR structures are pseudo-hermitian and strongly pseudo-convex, but they are not in general integrable. For a non-zero real number k , the author [7] defined the generalized Tanaka-Webster connection (in short, the *g.-Tanaka-Webster connection*) $\hat{\nabla}$ for real hypersurfaces in Kählerian manifolds. The *g.-Tanaka-Webster connection* $\hat{\nabla}$ coincides with the Tanaka-Webster connection if real hypersurfaces satisfy $\phi A + A\phi = 2k\phi$ (Proposition 2). The covariant differentiation of the Levi form L with respect to the *g.-Tanaka-Webster connection* $\hat{\nabla}$ is well-defined:

$$(\hat{\nabla}_X L)(Y, Z) = XL(Y, Z) - L(\hat{\nabla}_X Y, Z) - L(Y, \hat{\nabla}_X Z)$$

for any $X, Y, Z \in D$. Then we say that M is *Levi-parallel with respect to the g.-Tanaka-Webster connection* or shortly *Levi-parallel* if M satisfies

$$g((\hat{\nabla}_X L)(Y, Z)) = 0$$

for any vector fields $X, Y, Z \in D$. We note that a Levi-flat hypersurface is Levi-parallel (see (2) in Remark 1).

A complex n -dimensional complete and simply connected Kählerian manifold of constant holomorphic sectional curvature c is called a complex space form, which is denoted by $\tilde{M}_n(c)$. A complex space form consists of a complex projective space $P_n\mathbf{C}$, a complex Euclidean space $E_n\mathbf{C}$ or a complex hyperbolic space $H_n\mathbf{C}$, according as $c > 0$, $c = 0$ or $c < 0$. R. Takagi [16, 17] classified the homogeneous real hypersurfaces of $P_n\mathbf{C}$ into six types. T. E. Cecil and P. J. Ryan [6] extensively studied a real hypersurface whose structure vector ξ is a principal curvature vector, which is realized as tubes over certain submanifolds in $P_n\mathbf{C}$, by using its focal map. A real hypersurface of a complex space form is said to

be a *Hopf hypersurface* if its structure vector is a principal curvature vector. By making use of those results and the mentioned work of R. Takagi, M. Kimura [9] proved the local classification theorem for Hopf hypersurfaces of $P_n\mathbf{C}$ whose all principal curvatures are constant. For the case $H_n\mathbf{C}$, J. Berndt [3] proved the classification theorem for Hopf hypersurfaces whose all principal curvatures are constant.

The main purpose of the present paper is to classify real hypersurfaces of $\tilde{M}_n(c)$, $c \neq 0$ whose Levi form is parallel with respect to the generalized Tanaka-Webster connection. More specifically, in section 4, we prove

MAIN THEOREM. *Let M be a Hopf hypersurface of a complex space form $\tilde{M}_n(c)$, $c \neq 0$. Suppose that M is Levi-parallel with respect to the g -Tanaka-Webster connection. Then we have the following.*

- (I) *If $\tilde{M}_n(c) = P_n\mathbf{C}$, then M is locally congruent to one of:*
 - (A₁) *a geodesic hypersphere of radius r , where $0 < r < \frac{\pi}{2}$,*
 - (A₂) *a tube of radius r over a totally geodesic $P_k\mathbf{C}$ ($1 \leq k \leq n - 2$), where $0 < r < \frac{\pi}{2}$,*
 - (B) *a tube of radius r over a complex quadric Q_{n-1} , where $0 < r < \frac{\pi}{4}$.*
- (II) *If $\tilde{M}_n(c) = H_n\mathbf{C}$, then M is locally congruent to one of:*
 - (A₀) *a horosphere,*
 - (A₁) *a geodesic hypersphere or a tube over a complex hyperbolic hyperplane $H_{n-1}\mathbf{C}$,*
 - (A₂) *a tube over a totally geodesic $H_k\mathbf{C}$ ($1 \leq k \leq n - 2$),*
 - (B) *a tube over a totally real hyperbolic space $H_n\mathbf{R}$.*

2. The Generalized Tanaka-Webster Connection for Real Hypersurfaces

In this paper, all manifolds are assumed to be connected and of class C^∞ and the real hypersurfaces are supposed to be oriented. First, we give a brief review of several fundamental concepts and formulas on almost contact structure. An odd-dimensional smooth manifold M^{2n+1} has an *almost contact structure* if it admits a vector ξ , a 1-form η and a (1,1)-tensor field φ satisfying

$$\eta(\xi) = 1 \quad \text{and} \quad \varphi^2 X = -X + \eta(X)\xi.$$

Then there exists a compatible Riemannian metric g :

$$g(\varphi X, \varphi Y) = g(X, Y) - \eta(X)\eta(Y)$$

for all vector fields X and Y on M . We call (η, ξ, φ, g) an almost contact metric structure of M and $M = (M; \eta, \xi, \varphi, g)$ an almost contact metric manifold. For

an almost contact metric manifold M we define its fundamental 2-form Φ by $\Phi(X, Y) = g(\varphi X, Y)$. If

$$(1.1) \quad \Phi = d\eta,$$

M is called a contact metric manifold. We refer to [4] on contact metric geometry for more detail.

For an almost contact metric manifold M , the tangent space $T_p M$ of M at each point $p \in M$ is decomposed as $T_p M = D_p \oplus \{\xi\}_p$ (direct sum), where we denote $D_p = \{v \in T_p M \mid \eta(v) = 0\}$. Then $D : p \rightarrow D_p$ defines a distribution orthogonal to ξ . The restriction $\bar{\varphi} = \varphi|_D$ of φ to D defines an almost complex structure to D . If the associated Levi form L , defined by

$$L(X, Y) = d\eta(X, \bar{\varphi}Y),$$

$X, Y \in D$, is hermitian, then $(\eta, \bar{\varphi})$ is called a pseudo-hermitian CR structure and in addition, if its Levi form is non-degenerate (positive or negative definite, resp.), then $(\eta, \bar{\varphi})$ is called a non-degenerate (strongly pseudo-convex, resp.) pseudo-hermitian CR structure. Moreover, if the following conditions are satisfied:

$$(1.2) \quad [\bar{\varphi}X, \bar{\varphi}Y] - [X, Y] \in D$$

and

$$(1.3) \quad [\bar{\varphi}, \bar{\varphi}](X, Y) = 0$$

for all $X, Y \in D$, where $[\bar{\varphi}, \bar{\varphi}]$ is the Nijenhuis torsion of $\bar{\varphi}$, then the pair $(\eta, \bar{\varphi})$ is called a pseudo-hermitian, non-degenerate, (strongly pseudo-convex, resp.) integrable CR structure associated with the almost contact metric structure (η, ξ, φ, g) . In particular, for a contact metric manifold its associated CR structure is pseudo-hermitian, strongly pseudo-convex but is not in general integrable. For further details about CR structures, we refer for example to [2], [5], [18].

Let M be a real hypersurface of a Kählerian manifold $\tilde{M} = (\tilde{M}; J, \tilde{g})$ and N a global unit normal vector on M . By $\tilde{\nabla}$, A we denote the Levi-Civita connection in \tilde{M} and the shape operator with respect to N , respectively. Then the Gauss and Weingarten formulas are given respectively by

$$\tilde{\nabla}_X Y = \nabla_X Y + g(AX, Y)N, \quad \tilde{\nabla}_X N = -AX$$

for any vector fields X and Y tangent to M , where g denotes the Riemannian metric of M induced from \tilde{g} . An eigenvector (resp. eigenvalue) of the shape operator A is called a principal curvature vector (resp. principal curvature). For any vector field X tangent to M , we put

$$(2.1) \quad JX = \varphi X + \eta(X)N, \quad JN = -\xi.$$

We easily see that the structure (η, ξ, φ, g) is an almost contact metric structure on M . From the condition $\tilde{\nabla}J = 0$, the relations (2.1) and by making use of the Gauss and Weingarten formulas, we have

$$(2.2) \quad (\nabla_X \varphi)Y = \eta(Y)AX - g(AX, Y)\xi,$$

$$(2.3) \quad \nabla_X \xi = \varphi AX.$$

By using (2.2) and (2.3), we see that a real hypersurface in a Kählerian manifold always satisfies (1.2) and (1.3), the integrability condition of the associated almost CR structure. From (1.1) and (2.3) we have

PROPOSITION 1. *Let $M = (M; \eta, \xi, \varphi, g)$ be a real hypersurface of a Kählerian manifold. The almost contact metric structure of M is contact metric if and only if $\varphi A + A\varphi = \pm 2\varphi$, where \pm is determined by the orientation.*

The Tanaka-Webster connection ([19], [20]) is the canonical affine connection defined in a non-degenerate integrable CR manifold. Tanno ([18]) defined the generalized Tanaka-Webster connection for contact metric manifolds by the canonical connection which coincides with the Tanaka-Webster connection if the associated almost CR structure is integrable. We define the generalized Tanaka-Webster connection (in short, the g.-Tanaka-Webster connection) for real hypersurfaces of Kählerian manifolds by the naturally extended one of Tanno's generalized Tanaka-Webster connection for contact metric manifolds.

We recall Tanno's generalized Tanaka-Webster connection $\hat{\nabla}$ for contact metric manifolds:

$$\hat{\nabla}_X Y = \nabla_X Y + (\nabla_X \eta)(Y)\xi - \eta(Y)\nabla_X \xi - \eta(X)\varphi Y$$

for all vector fields X and Y .

Taking account of (2.3), the g.-Tanaka-Webster connection for real hypersurfaces of Kählerian manifolds, which is denoted by the same symbol $\hat{\nabla}$ as the one for contact metric manifolds, is naturally defined by (cf. [7])

$$(2.4) \quad \hat{\nabla}_X Y = \nabla_X Y + g(\varphi AX, Y)\xi - \eta(Y)\varphi AX - k\eta(X)\varphi Y,$$

where k is a non-zero real number. We put $F_X Y = g(\varphi AX, Y)\xi - \eta(Y)\varphi AX - k\eta(X)\varphi Y$. Then the torsion tensor \hat{T} is given by $\hat{T}(X, Y) = F_X Y - F_Y X$. Also, by using (1.2), (1.3), (2.2), (2.3) and (2.4) we can see that

$$(2.5) \quad \hat{\nabla}\eta = 0, \quad \hat{\nabla}\xi = 0, \quad \hat{\nabla}g = 0, \quad \hat{\nabla}\varphi = 0.$$

and

$$\hat{T}(X, Y) = 2 d\eta(X, Y)\xi, \quad X, Y \in D.$$

We note that the associated Levi form is

$$L(X, Y) = \frac{1}{2}g((\bar{\varphi}A + A\bar{\varphi})X, \bar{\varphi}Y),$$

where we denote by \bar{A} the restriction A to D . If M satisfies $\varphi A + A\varphi = 2k\varphi$, then we see that the associated CR structure is pseudo-hermitian, strongly pseudo-convex and further satisfies $\hat{T}(\xi, \varphi Y) = -\varphi\hat{T}(\xi, Y)$, and hence the generalized Tanaka-Webster connection $\hat{\nabla}$ coincides with the Tanaka-Webster connection. Namely, we have (cf. [7])

PROPOSITION 2. *Let $M = (M; \eta, \xi, \varphi, g)$ be a real hypersurface of a Kählerian manifold. If M satisfies $\varphi A + A\varphi = 2k\varphi$, then the associated CR-structure is pseudo-hermitian, strongly pseudo-convex, integrable, and further the generalized Tanaka-Webster connection $\hat{\nabla}$ coincides with the Tanaka-Webster connection.*

Since the structure vector field ξ is $\hat{\nabla}$ -parallel, we see that $\hat{\nabla}_X Y$ for $X, Y \in D$ still belongs to D . We define the covariant differentiation of the Levi form L with respect to the g.-Tanaka-Webster connection $\hat{\nabla}$ as follows:

$$(2.6) \quad (\hat{\nabla}_X L)(Y, Z) = XL(Y, Z) - L(\hat{\nabla}_X Y, Z) - L(Y, \hat{\nabla}_X Z)$$

for any $X, Y, Z \in D$.

3. Real Hypersurfaces of a Complex Space Form

Let $\tilde{M} = \tilde{M}_n(c)$ be a non-flat complex space form of constant holomorphic sectional curvature $c(\neq 0)$ and let M a real hypersurface of \tilde{M} . Then we have the following Gauss and Codazzi equations:

$$(3.1) \quad R(X, Y)Z = \frac{c}{4}\{g(Y, Z)X - g(X, Z)Y \\ + g(\varphi Y, Z)\varphi X - g(\varphi X, Z)\varphi Y - 2g(\varphi X, Y)\varphi Z\} \\ + g(AY, Z)AX - g(AX, Z)AY,$$

$$(3.2) \quad (\nabla_X A)Y - (\nabla_Y A)X = \frac{c}{4}\{\eta(X)\varphi Y - \eta(Y)\varphi X - 2g(\varphi X, Y)\xi\}$$

for any tangent vector fields X, Y, Z on M . We now suppose that M is a Hopf hypersurface, that is, $A\xi = \alpha\xi$. Then we already know that α is constant (see [8]). Differentiating this covariantly along M , and then by using (2.3) we have

$$(\nabla_X A)\xi = \alpha\varphi AX - A\varphi AX,$$

and further by using (3.2) we obtain

$$(\nabla_\xi A)X = \frac{c}{4}\varphi X + \alpha\varphi AX - A\varphi AX$$

for any vector field X on M . From this, we have

$$2A\varphi AX - \frac{c}{2}\varphi X = \alpha(\varphi A + A\varphi)X.$$

Here, we assume that $AX = \lambda X$ for a unit vector field X orthogonal to ξ , then

$$(3.3) \quad (2\lambda - \alpha)A\varphi X = \left(\alpha\lambda + \frac{c}{2}\right)\varphi X.$$

Now, we prove

THEOREM 3. *There does not exist a Levi-flat Hopf hypersurface in a non-flat complex space form.*

PROOF. Suppose that M is Hopf and Levi-flat. Then $A\xi = \alpha\xi$ and we get

$$\varphi AX + A\varphi X = 0$$

for any $X \in D$. We assume $AX = \lambda X$. Since ξ is a principal curvature vector by using (3.3) we have $2\lambda^2 + \frac{c}{2} = 0$, which shows $c < 0$. Then we see that M has at most three constant principal curvatures λ, μ and α , and further we see that $\mu = -\lambda$. But, Corollary 1 in [3] states that $\lambda\mu + c/4 = 0$. Thus, we have a contradiction. \square

We remark here that there are examples of Levi-flat hypersurfaces which are not Hopf. We say that M is a *ruled real hypersurface* of $\tilde{M}_n(c)$, $c \neq 0$ if there is a foliation of M by complex hyperplanes $\tilde{M}_{n-1}(c)$. In other words, M is ruled if and only if D is integrable and its integral manifold is a totally geodesic sub-

manifold $\tilde{M}_{n-1}(c)$. Then we easily see that a ruled real hypersurface is Levi-flat. In fact, its shape operator may be written down as following:

$$A\xi = \alpha\xi + \mu U \quad (\mu \neq 0),$$

$$AU = \mu\xi,$$

$$AZ = 0$$

for any $Z \in D, \perp U$, where U is unit vector orthogonal to ξ , α and μ are functions on M . M. Kimura [10] constructed ruled real hypersurfaces in complex projective space. Let \bar{M} be a hypersurface in S^{2n+1} defined by

$$\left\{ \left(re^{it} \cos \theta, re^{it} \sin \theta, \sqrt{1-r^2}z_2, \dots, \sqrt{1-r^2}z_n \right) \in \mathbf{C}^{n+1} \mid \sum_{j=2}^n |z_j|^2 = 1, 0 < r < 1, 0 \leq t, \theta < 2\pi, \right\}.$$

Then the Hopf image M of \bar{M} is a minimal ruled hypersurface in $\mathbf{C}P^n$. We note that the above example of a ruled real hypersurface is not complete. In a similar way, in [1] the authors gave a minimal ruled real hypersurfaces in complex hyperbolic space. For more details about ruled real hypersurfaces we may refer to [13].

From Proposition 2, together with the results in [12] (in case of $P_n\mathbf{C}$) and [15] (in case of $H_n\mathbf{C}$) we get easily

THEOREM 4. *Let M be a real hypersurface of $\tilde{M}_n(c)$, $c \neq 0$. Suppose that M satisfies $\varphi A + A\varphi = 2k\varphi$ for some non-zero constant k . Then the CR-structure is pseudo-hermitian and strongly pseudo-convex. Furthermore we have the following:*

(I) *in the case $\tilde{M}_n(c) = P_n\mathbf{C}$ with the Fubini-Study metric of $c = 4$, then M is locally congruent to one of the following:*

(A₁) *a geodesic hypersphere of radius r , where $0 < r < \frac{\pi}{2}$,*

(B) *a tube of radius r over a complex quadric Q_{n-1} , where $0 < r < \frac{\pi}{4}$.*

(II) *in the case $\tilde{M}_n(c) = H_n\mathbf{C}$ with the Bergman metric of $c = -4$, then M is locally congruent to one of the following:*

(A₀) *a horosphere,*

(A₁) *a geodesic hypersphere or a tube over a complex hyperbolic hyperplane $H_{n-1}\mathbf{C}$,*

(B) *a tube over a totally real hyperbolic space $H_n\mathbf{R}$.*

REMARK 1. (1) Together with Proposition 1, we see that the almost contact metric structure on M which appears in the above theorem is a contact metric structure only for the very special case determined by $k = \pm 1$, where \pm depends on the orientation. More precisely, with the help of the tables in [3] and [16], we see that the almost contact metric structures are contact metric only for a geodesic hypersphere of radius $\frac{\pi}{4}$ in $P_n\mathbf{C}$, for a horosphere in $H_n\mathbf{C}$. Hence for real hypersurfaces appearing in Theorem 4, except those just mentioned, they do not admit contact structure but their associated CR structures are pseudo-hermitian, strongly pseudo-convex and further the g.-Tanaka-Webster connection $\hat{\nabla}$ defined on them coincides with the Tanaka-Webster connection.

(2) From (2.6), it follows that Levi-flat hypersurface is Levi-parallel. Leaving the Levi-flat case aside, we find that real hypersurfaces stated in Theorem 4 are also Levi-parallel.

We prepare some more results which are needed to prove our Main Theorem.

THEOREM 5 ([9]). *Let M be a Hopf hypersurface of $P_n\mathbf{C}$. Then M has constant principal curvatures if and only if M is locally congruent to one of the following:*

- (A₁) *a geodesic hypersphere of radius r , where $0 < r < \frac{\pi}{2}$,*
- (A₂) *a tube of radius r over a totally geodesic $P_k\mathbf{C}$ ($1 \leq k \leq n-2$), where $0 < r < \frac{\pi}{2}$,*
- (B) *a tube of radius r over a complex quadric Q_{n-1} , where $0 < r < \frac{\pi}{4}$,*
- (C) *a tube of radius r over $P_1\mathbf{C} \times P_{(n-1)/2}\mathbf{C}$, where $0 < r < \frac{\pi}{4}$ and $n(\geq 5)$ is odd,*
- (D) *a tube of radius r over a complex Grassmann $G_{2,5}\mathbf{C}$, where $0 < r < \frac{\pi}{4}$ and $n = 9$,*
- (E) *a tube of radius r over a Hermitian symmetric space $SO(10)/U(5)$, where $0 < r < \frac{\pi}{4}$ and $n = 15$.*

THEOREM 6 ([3]). *Let M be a Hopf hypersurface of $H_n\mathbf{C}$. Then M has constant principal curvatures if and only if M is locally congruent to one of the following:*

- (A₀) *a horosphere,*
- (A₁) *a geodesic hypersphere or a tube over a complex hyperbolic hyperplane $H_{n-1}\mathbf{C}$,*
- (A₂) *a tube over a totally geodesic $H_k\mathbf{C}$ ($1 \leq k \leq n-2$),*
- (B) *a tube over a totally real hyperbolic space $H_n\mathbf{R}$.*

THEOREM 7 ([11], [15]). *Let M be a Hopf hypersurface of a non-flat complex space form $\tilde{M}_n(c)$, $c \neq 0$. Suppose that the shape operator A is η -parallel (i.e., $g((\nabla_X A)Y, Z) = 0$) for any tangent vectors X, Y and Z which are orthogonal to ξ . Then we have the following.*

(I) *In case that $\tilde{M}_n(c) = P_n\mathbf{C}$, then M is locally congruent to one of real hypersurfaces of type (A_1) , (A_2) and (B) ;*

(II) *In case that $\tilde{M}_n(c) = H_n\mathbf{C}$, then M is locally congruent to one of real hypersurfaces of type (A_0) , (A_1) , (A_2) and (B) .*

4. Levi-parallel Hopf Hypersurfaces in a Complex Space Form

In this section we shall prove our Main Theorem. Suppose that M is a Levi-parallel Hopf hypersurface of a complex space form $\tilde{M}_n(c)$ with respect to g-Tanaka-Webster connection. Then by using (2.5) and (2.6) we have

$$g((\varphi(\hat{\nabla}_Z A) + (\hat{\nabla}_Z A)\varphi)X, \varphi Y) = 0$$

for any vector fields X, Y, Z orthogonal to ξ on M . It follows easily that

$$g((\hat{\nabla}_Z A)X, Y) - \eta((\hat{\nabla}_Z A)X)\eta(Y) + g((\hat{\nabla}_Z A)\varphi X, \varphi Y) = 0$$

for any $X, Y, Z \in D$.

Together with (2.4), we have

$$(4.1) \quad g((\nabla_Z A)X, Y) - \eta(AX)g(\varphi AZ, Y) - g(\varphi AZ, X)\eta(AY) \\ + g((\nabla_Z A)\varphi X, \varphi Y) - \eta(A\varphi X)g(\varphi AZ, \varphi Y) - g(\varphi AZ, \varphi X)\eta(A\varphi Y) = 0$$

for any $X, Y, Z \in D$. We now suppose that $A\xi = \alpha\xi$. Then (4.1) reduces to

$$(4.2) \quad g((\nabla_Z A)X, Y) - g(\varphi(\nabla_Z A)\varphi X, Y) = 0$$

where $X, Y, Z \in D$. Assume $X \in V_\lambda$, that is, $AX = \lambda X$, where we denote by V_λ the eigenspace of A associated with a principal curvature λ . Taking account of (3.3), we divide our arguments into two cases: (i) $2\lambda \neq \alpha$ and $2\lambda = \alpha$. First, we consider the case (i). Then for any $Z \in D$, we get

$$(\nabla_Z A)X = \nabla_Z(AX) - A(\nabla_Z X) \\ = (Z\lambda)X + (\lambda I - A)(\nabla_Z X).$$

So we have

$$(4.3) \quad g((\nabla_Z A)X, X) = Z\lambda + g((\lambda I - A)\nabla_Z X, X) \\ = Z\lambda + g(\nabla_Z X, (\lambda I - A)X) = Z\lambda.$$

Similarly, by using (3.3), we have

$$(4.4) \quad g((\nabla_Z A)\varphi X, \varphi X) = -(Z\lambda) \frac{\alpha^2 + c}{(2\lambda - \alpha)^2}.$$

From (4.2), (4.3) and (4.4) we obtain

$$(Z\lambda) \left(\lambda^2 - \alpha\lambda - \frac{c}{4} \right) = 0.$$

Since α is constant, this shows that

$$(4.5) \quad Z\lambda = 0 \quad \text{for any } Z \in D.$$

Also, it follows from the equation of Codazzi (3.2) that

$$(\nabla_Z A)\xi - (\nabla_\xi A)Z = -\frac{c}{4}\varphi Z \quad \text{for any } Z \in D.$$

On the other hand, from (2.3) and (3.3) we find

$$\begin{aligned} (\nabla_Z A)\xi - (\nabla_\xi A)Z &= \nabla_Z(A\xi) - A\nabla_Z\xi - \nabla_\xi(AZ) + A(\nabla_\xi Z) \\ &= (\alpha I - A)\varphi AZ - (\xi\lambda)Z - (\lambda I - A)\nabla_\xi Z \\ &= \lambda \left(\alpha - \frac{\alpha\lambda + \frac{c}{2}}{2\lambda - \alpha} \right) \varphi Z - (\xi\lambda)Z - (\lambda I - A)\nabla_\xi Z \end{aligned}$$

for any unit vector $Z \in V_\lambda$. From the above two equations, we obtain

$$(4.6) \quad \xi\lambda = 0$$

where we have used $g(\varphi Z, Z) = 0$ and $g((\lambda I - A)\nabla_\xi Z, Z) = 0$. Hence from (4.5) and (4.6) we see that λ is constant. Next, in the case (ii) $2\lambda = \alpha$, since α_1 is constant, λ must be constant.

Thus, by virtue of Theorems 5 and 6 we can see that M is locally congruent to one of six types (A_1) , (A_2) , (B) , (C) , (D) and (E) in $P_n\mathbf{C}$ or (A_0) , (A_1) , (A_2) and (B) in $H_n\mathbf{C}$. Conversely, by using Theorem 7, we check that real hypersurfaces of types (A_1) , (A_2) , (B) in $P_n\mathbf{C}$ or (A_0) , (A_1) , (A_2) and (B) in $H_n\mathbf{C}$ are Levi-parallel (with respect to the g.-Tanaka-Webster connection).

Now, we shall prove M of types (C) , (D) and (E) in $P_n\mathbf{C}$ is not Levi parallel. For M of type (C) , (D) or (E) in $P_n\mathbf{C}$, M has five distinct constant principal curvatures, say $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ and α so that $TM = V_{\lambda_1} \oplus V_{\lambda_2} \oplus V_{\lambda_3} \oplus V_{\lambda_4} \oplus \{\xi\}_{\mathbf{R}}$. We put $x = \cot(\theta - \frac{\pi}{4})$ ($\frac{\pi}{4} < \theta < \frac{\pi}{2}$). Then we may express (cf. [11])

$$(4.7) \quad \lambda_1 = x, \quad \lambda_2 = -\frac{1}{x}, \quad \lambda_3 = \frac{x+1}{1-x}, \quad \lambda_4 = \frac{x-1}{x+1}, \quad \alpha = \frac{-4x}{x^2-1}.$$

We note that

$$(4.8) \quad 0 < x < 1 \quad \text{and} \quad \phi V_{\lambda_1} = V_{\lambda_2}, \quad \phi V_{\lambda_2} = V_{-\lambda_1}, \quad \phi V_{\lambda_a} = V_{\lambda_a}, \quad a = 3, 4.$$

We first prove the following

LEMMA 1. *Let M be a real hypersurface M of types (C), (D) and (E) in $P_n\mathbf{C}$. If M is Levi-parallel, then*

$$(4.9) \quad \begin{aligned} (1) & \text{ for } X \in V_{\lambda_i} \ (i = 1, 2); \nabla_Z X = (\nabla_Z X)_{\lambda_i} - g(X, \phi AZ)\xi, \\ (2) & \text{ for } X \in V_{\lambda_a} \ (a = 3, 4); \nabla_Z X = (\nabla_Z X)_{\lambda_a} - g(X, \phi AZ)\xi. \end{aligned}$$

for any $Z \in D$, where X_λ denotes the V_λ -component of the vector X .

PROOF. For $X \in V_\lambda$ and $Y \in V_\mu$, we get

$$g((\nabla_Z A)X, Y) = (\lambda - \mu)g(\nabla_Z X, Y).$$

If we put $\bar{\lambda} = \frac{\alpha\lambda+2}{2\bar{\lambda}-\alpha}$, then $\phi X \in V_{\bar{\lambda}}$ and $\phi Y \in V_{\bar{\mu}}$. Together with (2.2) we get

$$\begin{aligned} g((\nabla_Z A)\phi X, \phi Y) &= (\bar{\lambda} - \bar{\mu})g(\nabla_Z(\phi X), \phi Y) \\ &= (\bar{\lambda} - \bar{\mu})g(\phi(\nabla_Z X), \phi Y) \\ &= (\bar{\lambda} - \bar{\mu})g(\nabla_Z X, Y) \end{aligned}$$

Suppose that M is Levi-parallel. Then from (4.2) we obtain

$$(4.10) \quad [(\lambda - \mu) + (\bar{\lambda} - \bar{\mu})]g(\nabla_Z X, Y) = 0.$$

From (4.7) and (4.10) we calculate the following:

$$(4.11) \quad \text{for } X \in V_{\lambda_i} \ (i = 1, 2), Y \in V_{\lambda_3}; \frac{(x+1)(x^2+1)}{x(x-1)}g(\nabla_Z X, Y) = 0,$$

$$\text{for } X \in V_{\lambda_i}, Y \in V_{\lambda_4}; \frac{(x-1)(x^2+1)}{x(x+1)}g(\nabla_Z X, Y) = 0,$$

$$\text{for } X \in V_{\lambda_1}, Y \in V_{\lambda_2}; 2xg(\nabla_Z X, Y) = 0,$$

$$\text{for } X \in V_{\lambda_2}, Y \in V_{\lambda_1}; -2xg(\nabla_Z X, Y) = 0,$$

$$\text{for } X \in V_{\lambda_3}, Y \in V_{\lambda_4}; \frac{(x+1)(x^2+1)}{x(1-x)}g(\nabla_Z X, Y) = 0,$$

$$\text{for } X \in V_{\lambda_4}, Y \in V_{\lambda_3}; \frac{(1-x)(x^2+1)}{x(x+1)}g(\nabla_Z X, Y) = 0.$$

Since $g(\nabla_Z X, \xi) = -g(X, \varphi AZ)$, from (4.8) and (4.11), we may express $\nabla_Z X$ as (4.9). \square

Secondly, we also prove

LEMMA 2. *Let M be a real hypersurface M of type (C), (D) and (E) in $P_n\mathbb{C}$. Then we have*

$$(4.12) \quad \nabla_\xi Z \in V_{\lambda_i} \oplus \{\varphi Z\}_{\mathbb{R}} \quad \text{for } Z \in V_{\lambda_i} \quad (i = 1, 2).$$

PROOF. For any unit vector $Z \in V_\lambda$, from (2.3) and Proposition 3 it follows that

$$\begin{aligned} (\nabla_Z A)\xi - (\nabla_\xi A)Z &= \nabla_Z(A\xi) - A\nabla_Z\xi - \nabla_\xi(AZ) + A(\nabla_\xi Z) \\ &= (\alpha I - A)\varphi AZ - (\xi\lambda)Z - (\lambda I - A)\nabla_\xi Z \\ &= \lambda \left(\alpha - \frac{\alpha\lambda + 2}{2\lambda - \alpha} \right) \varphi Z - (\lambda I - A)\nabla_\xi Z. \end{aligned}$$

On the other hand, from (3.2) we get

$$(\nabla_Z A)\xi - (\nabla_\xi A)Z = -\varphi Z.$$

Hence we obtain

$$(4.13) \quad (\lambda I - A)\nabla_\xi Z = \left[\lambda \left(\alpha - \frac{\alpha\lambda + 2}{2\lambda - \alpha} \right) \right] \varphi Z \quad \text{for } Z \in V_\lambda.$$

Since $\varphi V_{\lambda_1} = V_{\lambda_2}$, from (4.13) we can find (4.12). \square

Thus, it follows from Proposition 3 and (4.13) that for $i = 1, 2$,

$$\left\{ \lambda_i - \frac{\alpha\lambda_i + 2}{2\lambda_i - \alpha} \right\} g(\nabla_\xi Z, \varphi Z) = \left[\lambda_i \left(\alpha - \frac{\alpha\lambda_i + 2}{2\lambda_i - \alpha} \right) \right] g(\varphi Z, \varphi Z)$$

or

$$(4.14) \quad 2(\lambda_i^2 - \alpha\lambda_i - 1)g(\nabla_\xi Z, \varphi Z) = \alpha(\lambda_i^2 - \alpha\lambda_i - 1)g(\varphi Z, \varphi Z).$$

But, for a real hypersurface M which is locally congruent to one of types (C), (D) and (E) we know that $\lambda^2 - \alpha\lambda - 1 \neq 0$. (We note that the equation $\lambda^2 - \alpha\lambda - 1 = 0$ holds if and only if M is locally congruent to a real hypersurface of type (A_1) or (A_2) .) Therefore from (4.14) we get

$$(4.15) \quad g(\nabla_{\xi}Z, \varphi Z) = \frac{\alpha}{2}g(\varphi Z, \varphi Z) \quad \text{for } Z \in V_{\lambda_i}, i = 1, 2.$$

For $X \in V_{\lambda_1}$ and $Z \in V_{\lambda_3}$, by using (1) and (2) in (4.9), we have

$$(4.16) \quad \begin{aligned} R(Z, \varphi Z)X &= \nabla_Z(\nabla_{\varphi Z}X) - \nabla_{\varphi Z}(\nabla_ZX) - \nabla_{[Z, \varphi Z]}X \\ &= \nabla_Z\{(\nabla_{\varphi Z}X)_{\lambda_1} - \lambda_3g(X, \varphi^2Z)\xi\} \\ &\quad - \nabla_{\varphi Z}\{(\nabla_ZX)_{\lambda_1} - \lambda_3g(X, \varphi Z)\xi\} \\ &\quad - \nabla_{\{(\nabla_Z\varphi Z)_{\lambda_3} - \lambda_3\xi\}}X + \nabla_{\{(\nabla_{\varphi Z}Z)_{\lambda_3} + \lambda_3\xi\}}X \\ &= (\nabla_Z(\nabla_{\varphi Z}X)_{\lambda_1})_{\lambda_1} - \lambda_3g((\nabla_{\varphi Z}X)_{\lambda_1}, \varphi Z)\xi \\ &\quad - (\nabla_{\varphi Z}(\nabla_ZX)_{\lambda_1})_{\lambda_1} + \lambda_3g((\nabla_ZX)_{\lambda_1}, \varphi^2Z)\xi \\ &\quad - (\nabla_{(\nabla_Z\varphi Z)_{\lambda_3}}X)_{\lambda_1} + \lambda_3g(X, \varphi(\nabla_Z\varphi Z)_{\lambda_3})\xi + \lambda_3\nabla_{\xi}X \\ &\quad + (\nabla_{(\nabla_{\varphi Z}Z)_{\lambda_3}}X)_{\lambda_1} - \lambda_3g(X, \varphi(\nabla_{\varphi Z}Z)_{\lambda_3})\xi + \lambda_3\nabla_{\xi}X. \end{aligned}$$

The equations (4.15) and (4.16) show that

$$g(R(Z, \varphi Z)X, \varphi X) = 2\lambda_3g(\nabla_{\xi}X, \varphi X) = \alpha\lambda_3g(\varphi X, \varphi X).$$

On the other hand, since $\varphi X \in V_{\lambda_2}$ and $\varphi Z \in V_{\lambda_3}$, the equation of Gauss (3.1) gives

$$(4.17) \quad g(R(Z, \varphi Z)X, \varphi X) = -2g(\varphi Z, \varphi Z)g(\varphi X, \varphi X).$$

From this, together with (4.7), we have $\frac{-4x}{x^2-1} \cdot \frac{1+x}{1-x} = -2$, that is, $x^2 + 1 = 0$. This is a contradiction.

Thus, we have our Main Theorem. \square

References

- [1] S.-S. Ahn, S.-B. Lee and Y. J. Suh, CO_n ruled real hypersurfaces in a complex space form, Tsukuba J. Math. **17**(2) (1993), 311–322.
- [2] A. Bejancu, Geometry of CR-submanifolds, Mathematics and Its Application, D. Reidel Publ. Comp. (1986).
- [3] J. Berndt, Real hypersurfaces with constant principal curvatures in complex hyperbolic space, J. Reine Angew. Math. **395** (1989), 132–141.
- [4] D. E. Blair, Riemannian geometry of contact and symplectic manifolds, Progress in Mathematics, Birkäuser **203** (2001).
- [5] A. Bogges, CR manifolds and the tangential Cauchy-Riemann complex, Press, Boca Raton (1991).
- [6] T. E. Cecil and P. J. Ryan, Focal sets and real hypersurfaces in complex projective space, Trans. Amer. Math. Soc. **269** (1982), 481–499.

- [7] J. T. Cho, CR structures on real hypersurfaces of a complex space form, *Publ. Math. Debrecen* **54(3-4)** (1999), 473–487.
- [8] U.-H. Ki and Y. J. Suh, On real hypersurfaces of a complex space form, *Math. J. Okayama* **32** (1990), 207–221.
- [9] M. Kimura, Real hypersurfaces and complex submanifolds in complex projective space, *Trans. Amer. Math. Soc.* **296** (1986), 137–149.
- [10] M. Kimura, Sectional curvatures of holomorphic planes on a real hypersurface in $P^n(\mathbb{C})$, *Math. Ann.* **276** (1987), 487–497.
- [11] M. Kimura and S. Maeda, On real hypersurfaces of a complex projective space, *Math. Z.* **202** (1989), 299–311.
- [12] M. Kon, Pseudo-Einstein real hypersurfaces of complex space forms, *J. Differential Geometry* **14** (1979), 339–354.
- [13] M. Lohnherr and H. Reckziegel, On ruled hypersurfaces in complex space forms, *Kyungpook Math. J.* **35(3)** (1995).
- [14] Y. T. Siu, Nonexistence of smooth Levi-flat hypersurfaces in complex projective spaces of dimensions ≥ 3 , *Ann. of Math. (2)* **151** (2000), 1217–1243.
- [15] Y. J. Suh, On real hypersurfaces of a complex space form with η -parallel Ricci tensor, *Tsukuba J. Math.* **14** (1990), 27–37.
- [16] R. Takagi, On homogeneous real hypersurfaces in a complex projective space, *Osaka J. Math.* **19** (1973), 495–506.
- [17] R. Takagi, Real hypersurfaces in a complex projective space with constant principal curvatures I, II, *J. Math. Soc. Japan* **27** (1975); 43–53, 507–516.
- [18] S. Tanno, Variational problems on contact Riemannian manifolds, *Trans. Amer. Math. Soc.* **314** (1989), 349–379.
- [19] N. Tanaka, On non-degenerate real hypersurfaces, graded Lie algebras and Cartan connections, *Japan J. Math.* **2** (1976), 131–190.
- [20] S. M. Webster, Pseudohermitian structures on a real hypersurface, *J. Diff. Geometry* **13** (1978), 25–41.

Department of Mathematics
Chonnam National University
CNU The Institute of Basic Sciences
Kwangju 500-757, Korea
(Visiting Scholar in:)
University of Washington Mathematics Department
Box 354350 Seattle
WA 98195-4350, USA.