

1 Hybrid Sex Pheromones of the Hibiscus Flower-bud borer, *Rehimena surusalis*

2
3
4 HIROSHI HONDA^{1,*} RYOKUHEI YAMASAKI², YOKO SUMIUCHI², TAKUYA
5 UEHARA²,
6 SHIGERU MATSUYAMA¹, TETSU ANDO³, HIDESHI NAKA⁴
7

8 ¹*Faculty of Life and Environmental Sciences, University of Tsukuba; Tsukuba Ibaraki 305-8572,*
9 *Japan*

10 ²*Graduate School of Life and Environmental Sciences, University of Tsukuba; Tsukuba, Ibaraki*
11 *305-8572, Japan*

12 ³*Graduate School of Bio-Applications and Systems Engineering; Tokyo University of*
13 *Agriculture and Technology, Koganei, Tokyo 184-8588, Japan*

14 ⁴*Faculty of Agriculture, Tottori University; Koyama Minami, Tottori 680-8553, Japan*
15

16 **To whom correspondence should be addressed at E-mail: honda.hiroshi.ew@alumni.tsukuba.ac.jp*
17
18

19 **Abstract**—The sex pheromone of the hibiscus flower borer *Rehimena surusalis* (Walker)
20 (Lepidoptera: Crambidae) was analyzed by gas chromatography with electroantennographic
21 detection (GC-EAD) and GC-mass spectrometry (GC-MS). Three EAD-active components
22 were found in crude pheromone gland extracts of calling females. GC-MS and GC analyses
23 using synthetic chemicals and derivatization of the extracts identified three components as
24 (10*E*,12*Z*)-hexadeca-10,12-dienal (E10,Z12-16:Ald.), (10*E*,12*E*)-hexadeca-10,12-dienyl
25 acetate (E10,Z12-16:OAc) and (3*Z*,6*Z*,9*Z*)-tricoso-3,6,9-triene (Z3,Z6,Z9-23:HC). In field
26 tests, male moths were remarkably attracted to a ternary blend of E10,Z12-16:Ald, E10,Z12-
27 16:OAc and Z3,Z6,Z9-23:HC at a ratio of 1:5:14, but single and binary blend of either
28 compound showed only weak or no attraction activity.
29

30 **Key Words**—Hibiscus flower-bud borer, *Rehimena surusalis*, (10*E*,12*Z*)-10,12-
31 hexadecadienal, (10*E*,12*Z*)-10,12-hexadecadienyl acetate, (Z3,Z6,Z9)-3,6,9-tricosatrinene.
32

33 INTRODUCTION

34 Hibiscus flower-bud borer *Rehimena surusalis* (Walker) (Lepidoptera: Crambidae) is
35 widely distributed in Africa, Australia, China, India, Indonesia, Taiwan, Korea and Japan

36 (Shibuya, 1928, 1929; Inoue et al., 1982; Liu, 1990; Shin, 2001; Ades and Kendrick, 2004;
37 Herbison-Evans and Crossley, 2013) and is a continual pest of Malvaceae garden and street
38 trees including *Hibiscus syriacus* (rose of Sharon), *H. mutabilis* (cotton rose), *H. rosa-sinensis*
39 (Chinese hibiscus) *H. tiliaceus* and *H. glaber* (Sea Hibiscus) (Anonymous 1994, 2006). In
40 Japan and Korea, *H. syriacus* is particularly damaged by *R. surusalis*. *H. syriacus*
41 (mugunghwa in Korean) is authorized as the national flower of Korea, and *R. surusalis* has
42 been reported to eat the seed of this plant (Lee et al., 2005; Kim et al., 2013; Bea 2012). The
43 larvae bore into the developed flowers and flower buds. Because of the larval feeding habit as
44 a typical borer, it is difficult to control this pest with cover sprays of insecticides. To control
45 insects with a perforative lifestyle in the larval stage, pheromones are advantageous to
46 monitor the flying adults, and disrupt their mating, resulting in a reduction in oviposition
47 (Witzgall et al. 2010).

48 In this study, we identified components of the female sex pheromone of *R. surusalis* and
49 demonstrated sex pheromone activity of synthetics in the field. We also discussed a
50 commonality of the hybrid-type of sex pheromone in Pyraloidea.

51
52

MATERIALS AND METHODS

53 *Insects.* Colonies of *R. surusalis* were maintained as laboratory cultures. Mated females were
54 allowed to lay eggs in small plastic cylinders that were lined with felt cloth impregnated with
55 methanol extracts of *H. syriacus* flower buds. Because of heavy cannibalism, larvae of *R.*
56 *surusalis* were individually reared on an artificial diet composed of Insecta[®] F-II (Nosan
57 Corporation, Japan) and dried leaf powder of *H. syriacus* at a ratio of 8:2. Adults were sexed
58 at the pupal stage and kept separately in cages at $25 \pm 2^\circ\text{C}$, 60–70% relative humidity (RH)
59 and a 15L9D photoperiod, and provided with a 10% sugar solution from cotton pads. A red
60 lamp was used for observations during scotophase.

61

62 *Extracts and chemicals.* For identification of pheromone components, pheromone extracts
63 were obtained from 2 to 7 day old calling females, whose abdominal tips were cut with
64 ophthalmology scissors after half of scotophase by extraction with redistilled *n*-hexane for 20
65 min. Pooled extracts (60 female equivalents, FE) were stored at -20°C until use for chemical
66 analyses and bioassays. Aliquot of the extracts were subjected into GC analysis for
67 quantitative determination of pheromone candidates in 5 replications. Each four geometric
68 isomers of synthetic 10,12-hexadecadienals (Z10,E12-16:Ald, E10,Z12-16:Ald, Z10,Z12-
69 16:Ald and E10,E12-16:Ald) and 10,12-hexadecadienyl acetates (Z10,E12-16:OAc, E10,Z12-
70 16:OAc, Z10,Z12-16:OAc and E10,E12-16:OAc), and (3Z,6Z,9Z)-tricoso-3,6,9-triene
71 (3Z,6Z,9Z-23:CH) were supplied by coauthors T. A. or S. M. The isomeric purity of all

72 compounds was confirmed by GC to be $\geq 97\%$.

73

74 *Chemical analysis.* Pheromone extracts were subjected to GC-EAD analyses using a HP-5890
75 series II GS (Agilent Technologies, California, USA) equipped with an HP-5MS capillary
76 column (30 m \times 0.32 mm ID, film thickness 0.25 μm ; Agilent Technologies, USA) and
77 helium as a carrier gas (37 cm/s). Oven temperature was programmed at 130°C for 2 min,
78 then increased at a rate of 5°C /min to 250°C and held at the final temperature for 10 min. The
79 temperature of the detector and injector was 250°C, and that of the outlet for the EAD was
80 maintained at 300°C. Extracts were injected in splitless mode and chromatographed using
81 helium as a carrier gas (37 cm/s). GC effluent from the column was slit in a 1:1 ratio between
82 the flame ionization detector (FID) and the EAD. The effluent was delivered in humidified
83 air (23°C) to the antennal preparation connected to an EAG probe (Type PRG-2, Syntech, The
84 Netherlands) via Ag-AgCl electrodes with 0.1.M KCl. EAD responses of male antenna were
85 recorded in PC with GC-EAD 2010 software (Ver. 4.60, Syntech) via GC-EAD signal
86 acquisition controller (IDAC-2, Syntech).

87 Analyses of EAD active components in the extracts by GC-MS employed a MS-600H
88 mass spectrometer (JEOL Ltd., Japan) coupled with HP-6890N GC (Agilent), which was
89 equipped with a DB-5MS (25 m \times 0.25 mm ID, film thickness 0.25 μm , Agilent) capillary
90 column, and operated in electron impact ionization mode (70 eV). GC oven temperature was
91 programmed at 100°C for 1 min, then increased at a rate of 10°C /min to 320°C and held at
92 the final temperature for 17 min.

93 GC analyses were conducted with GC-17A (Shimadzu Co., Ltd., Japan) and GC-6890N
94 (Agilent) fitted with a nonpolar HP-5MS column and a polar DB-23 column (30 m \times 0.25 mm
95 ID, film thickness 0.15 μm ; Agilent), respectively. GC oven temperature of the nonpolar
96 column was programmed at 130°C for 2 min, then increased at a rate of 5°C /min to 250°C
97 and held at the final temperature for 10 min. GC oven temperature of the polar column was
98 programmed at 80°C for 2 min, then increased at the rate of 3°C /min to 250°C and held at the
99 final temperature for 5 min.

100 To determine the position of conjugated double bonds, pheromone candidates in the
101 extracts were reacted with 4-methyl-1,2,4-triazoline-3,5-dione (MTAD), followed by GC-MS
102 analysis of the resulting derivatives. Kováts retention indices (KRI) (Kováts, 1958; Dool and
103 Kratz, 1963) of EAD-active components and authentic chemicals were determined with
104 retention times of standard hydrocarbons. The GC peak area of each component on the HP-
105 5MS column was used to determine the ratio of EAD-active components in the pheromone
106 extracts.

107

108 *Laboratory and field tests.* Pheromone activity of candidate components, E10,Z12-16:Ald,
109 E10,Z12-16:OAc and Z3,Z6,Z9-23:HC and their blends were examined by laboratory and
110 field assays. Laboratory cage tests were conducted in a mesh cage (30 cm×25 cm×30 cm)
111 with 10 males at the second half of scotophase that the most of calling by males were
112 observed. Pheromone extracts or synthetics were applied on a filter paper (1cm x 3cm) in 1 µl
113 hexane as solvent. Filter paper was suspended 10 cm from the ceiling with a wire clip.
114 Amounts of synthetics were adjusted to 1 female equivalent (FE)/µl. Crude extracts were
115 concentrated to 1 FE/µl under a gentle N₂ stream. Numbers of males showing orientation
116 flight (OF) by hovering to pheromone source and source contact (SC) were counted for 3 min
117 with 5 ~ 7 replications and the cumulative numbers compared in single, binary and ternary
118 blends of the candidate compounds.

119 Field experiments were conducted in fields with *H. syriacus* plantations on the campus of
120 University of Tsukuba (36.1°N, 140.1°E) during June and August in 2013. Similar sets of
121 synthetic blends with those used in the laboratory assays were loaded on gray rubber septa
122 (West Corp., Singapore) at 500 µg / trap. In addition to the regular blend, blends with two and
123 five times excessive Z3,Z6,Z9-23:HC (750 µg and 1750 µg/ trap) were also tested. Each
124 rubber septum was placed on a sticky board trap with a triangle roof (SE-trap, 30 cm in length
125 x 27 cm in width x 10 cm in height; Sankei Chemical Co., Ltd., Kagoshima, Japan). Traps
126 were hung ca. 1.5 m above the ground on tree branches with at least 10 m intervals, and were
127 set in a completely randomized design, and the lure were renewed once a week. Positions of
128 traps were rotated one position every three days to avoid positional effects. As a control,
129 empty traps were also tested. Numbers of captured males in each trap were counted every 3
130 days.

131

132 *Statistical analyses.* Results of laboratory and field assays were analyzed using one-way
133 analysis of variance (ANOVA), followed by a Tukey-Kramer's honestly significant difference
134 (HSD) test. Numbers of captured males (x) in field tests were transformed $\sqrt{x+0.5}$ prior to
135 ANOVA. Software package R 3.0.1 (R Core Team 2013), was used for the statistical analyses.

136

137

RESULTS

138 *Chemical analysis.* GC-EAD analyses of crude pheromone gland extracts of female *Rehimena*
139 *surusalis* showed three active components **A** (Rt 11.28 min) **B** (Rt 14.66 min) and **C** (Rt
140 18.52 min) on FID chromatogram (Fig. 1). In GC-MS analyses, spectra of the active
141 component **A** showed putative parental ion at m/z 236 (M^+ , 36 %), and fragment ions at m/z

142 67 ($[\text{C}_5\text{H}_7]^+$, base peak), m/z 95 ($[\text{C}_7\text{H}_{11}]^+$, 41 %), m/z 96 ($[\text{C}_7\text{H}_{12}]^+$, 42 %) and m/z 109
143 ($[\text{C}_8\text{H}_{13}]^+$, 28 %). The ion peaks spaced by m/z 14 and peaks at m/z 96 and 109 suggested the
144 double bonds at the 10- and 12- ($\omega 4$, $\omega 6$) positions in a straight carbon chain (Ando et al.
145 1998). From these spectral data, the
146 structure of compound **A** was consistent to 10, 12-hexadecadienal ($\text{C}_{16}\text{H}_{28}\text{O}$). Relatively high
147 intensity of molecular ion peak (m/z 236) also supported this identification for component **A**.

148 GC-MS analysis of component **B** showed ion peaks at m/z 280 (M^+ , 38 %), m/z 61
149 ($[\text{CH}_3\text{COOH}+2\text{H}$, 5%], m/z 67 ($[\text{C}_5\text{H}_7]^+$, base peak), m/z 95 ($[\text{C}_7\text{H}_{11}]^+$, 48 %), m/z 96
150 ($[\text{C}_7\text{H}_{12}]^+$, 58 %), m/z 109 ($[\text{C}_8\text{H}_{13}]^+$, 29 %), and m/z 220 ($[\text{M}-\text{CH}_3\text{COOH}]^+$, 16%). Mass
151 spectra with ion peaks spaced by m/z 14 and two prominent peaks at m/z 96 and 109
152 suggested a straight carbon chain and double bond positions at 10, 12- ($\omega 4$, $\omega 6$) positions in
153 $\text{C}_{16}\text{H}_{32}\text{O}_2$. Two diagnostic ion peaks at m/z 61 and m/z 220 predicted structure of compound
154 **B** to be 10, 12-hexadecadienyl acetate. Relatively high intensity of molecular ion peak at m/z
155 280 also indicated conjugated double bonds in compound **B**.

156 In GC-MS analysis, component **C** showed ion peaks at m/z 318 (M^+ , 6 %), m/z 79
157 ($[\text{C}_6\text{H}_7]^+$, 79%), m/z 93 ($[\text{C}_7\text{H}_9]^+$, 33%), m/z 107 ($[\text{C}_8\text{H}_{11}]^+$, 15%), m/z 108 ($[\text{C}_8\text{H}_{12}]^+$, base
158 peak), m/z 121 ($[\text{C}_9\text{H}_{13}]^+$, 18%) and m/z 262 ($[\text{M}-\text{C}_4\text{H}_8]^+$, 19%). The fragmentation pattern
159 indicated an unsaturated straight-chain compound, with possible molecular formula of
160 $\text{C}_{23}\text{H}_{42}$, consistent with a tricosatriene (3,6,9-23:HC). In addition, three conspicuous
161 diagnostic ion peaks at m/z 79, m/z 108 and m/z 262 indicated three double bonds at 3, 6 and
162 9-position of compound **C** (Ando et al. 2004).

163 The position of double bonds in **A** and **B** were further confirmed by derivatization with
164 MTAD, which reacts specifically with conjugated dienyl structures. The mass spectra of
165 MTAD reaction products exhibited ions at m/z 349 (M^+ , $[\text{C}_{19}\text{H}_{31}\text{O}_3\text{N}_3]^+$, 17 %), m/z 208
166 ($[\text{C}_{10}\text{H}_{12}\text{O}_2\text{N}_3]^+$, base peak) and m/z 306 ($[\text{C}_{16}\text{H}_{24}\text{O}_3\text{N}_3]^+$, 57%) for compound **A**, and at m/z
167 393 (M^+ , $[\text{C}_{21}\text{H}_{35}\text{O}_4\text{N}_3]^+$, 17 %), m/z 208 ($[\text{C}_{10}\text{H}_{12}\text{O}_2\text{N}_3]^+$, base peak) and m/z 350
168 ($[\text{C}_{18}\text{H}_{28}\text{O}_4\text{N}_3]^+$ for compound **B** supporting two conjugated double bonds at either 3- and 5-
169 positions or 10- and 12-positions in hexadecadienal and hexadecadienyl acetate, respectively.

170 Components **A** and **B** had KRIs similar to those of each four isomers of 10, 12-16: Ald
171 and 10,12-16: OAc on both nonpolar and polar GC columns. The 3, 5-dienes would have
172 been expected to elute much more earlier than 10, 12-dienes on GC (Ando et al., 2004). As
173 shown in Table 1, KRIs of components **A** and **B** corresponded well to those of (10*E*,12*Z*)-
174 hexadeca-10,12-dien-1-al (E10,Z12-16:Ald,) and (10*E*,12*Z*)-hexadeca-10,12-dien-1-yl acetate
175 (E10,Z12-16:OAc), respectively, on both HP-5MS and DB-23 columns. KRI of component **C**
176 was compared with only that of Z3,Z6,Z9-23:HC, because 3,6,9-tricosatrienes as insects

177 pheromones are considered to be biosynthesized from (9Z,12Z,15Z)-octadeca-9,12,15-trienoic
178 acid with elongation of the carbon chain (Ando et al. 2008). The geometric configuration of
179 component **C** was confirmed to be 3Z,6Z,9Z-isomer from agreement with the RI.

180 The amounts of these three components (**A**, **B** and **C**) in the extracts were determined to
181 be 0.77 ± 0.08 ng, 3.60 ± 0.56 ng and 11.1 ± 0.96 ng per female, respectively, at ratio of
182 1:5:14.

183

184 *Laboratory and field tests.* In the laboratory test, pheromone activities of the crude pheromone
185 extract and all of possible combinations of synthetic E10,Z12-16:Ald, E10,Z12-16:OAc and
186 Z3,Z6,Z9-23:HC are summarized in Fig. 2. Three one-component baits and binary blends of
187 E10,Z12-16:Ald and E10,Z12-16:OAc, and Z3,Z6,Z9-23:HC with E10,Z12-16:Ald or
188 E10,Z12-16:OAc showed no pheromone activity in both activity criteria, orientation flight
189 and source contact by male moths, whereas significantly higher activity in orientation flight
190 was observed with binary combination of E10,Z12-16:Ald and E10,Z12-16:OAc though it
191 was still lower than that of the extract. Highest activity in orientation flight was observed with
192 the ternary blend of the above synthetics in natural amounts, and it corresponded well to
193 activity of the extract. In source contact by male moths, only the ternary blend showed
194 significantly different activity from that of the crude extract.

195 In the field tests, the ternary blend of E10,Z12-16:Ald, E10,Z12-16:OAc and Z3,Z6,Z9-
196 23:HC attracted the highest number of male moths in all treatments tested, whereas single and
197 binary blends attracted fewer or no male moths (Fig. 3). Similar to the results of the
198 laboratory tests, the binary blend of E10,Z12-16:Ald and E10,Z12-16:OAc showed also
199 relatively high activity in male attraction. When the amount of Z3,Z6,Z9-23:HC was
200 increased, trap catches somewhat decreased at 700 μ g, and significantly decreased at 1750 μ g
201 (Fig. 3).

202

203

DISCUSSION

204 Three GC-EAD active components were identified as E10,Z12-16:Ald, E10,Z12-16:OAc
205 and Z3,Z6,Z9-23:HC by GC and GC-MS analyses. The ternary blend of these compounds in
206 a ratio of 1:5:14 showed pheromone activity to male moths of *R. surusalis* in laboratory and
207 field bioassays. These results show that the sex pheromone of *R. surusalis* consists of three
208 components in this ratio. 10,12-Hexadecadienals are widely known as major or minor
209 components of sex pheromones of several moth families including Noctuidae (Cork et al.,
210 1988), Sphingidae (Starratt et al., 1979; Bestmann et al., 1992; Uehara et al., 2012, 2015),
211 Pyralidae or Crambidae (Klun et al., 1986; Raina et al 1986; Honda et al.,1994), Saturniidae

212 (Dai et al. 1988; McElfresh and Millar, 1999a,b) and also Bombycidae (Daimon et al., 2012).
213 E10,Z12-16:Ac was also identified as a sex pheromone in Bombycidae (Daimon et al., 2012)
214 and Saturniidae (Dai et al., 1987; McElfresh and Millar 1999a,b,c; 2001).

215 Sex pheromone components can be categorized into Type I and Type II groups depending
216 on whether they have or don't have terminal functional groups in the molecules, and
217 compounds such as E10,Z12-16:Ald, and E10,Z12-16:OAc belong to the Type I group but
218 polyenyl hydrocarbons such as Z3,Z6,Z9-23:HC belong to Type II group (Ando et al. 2004).
219 Recently so-called hybrid type of pheromone systems consisting of Type I and Type II
220 compounds such as that of *R. surusalis*, are reported mainly in Crambid and Pyralid species
221 (Cabrera et al 2001; Millar et al. 2005; Leal et al. 2005; Gibb et al. 2007; Miller et al. 2010;
222 Löfstedt et al. 2012; El-Sayed et al. 2013; Yan et al. 2014).

223 *Rehimena surusalis* male moths showed low but significant orientation flight responses to
224 a binary blend of E10,Z12-16:Ald and E10,Z12-16:OAc, although neither component was
225 active as a single component, in the laboratory cage test and field tests (Fig. 2 and 3),
226 indicating a crucial synergistic function of E10,Z12-16:Ald and E10,Z12-16:OAc in male
227 attraction from a long distance. Z3,Z6,Z9-23:HC significantly increased male catches in the
228 field traps, suggesting synergistic effect to E10,Z12-16:Ald and E10,Z12-16:OAc. However,
229 trap catches decreased when Z3,Z6,Z9-23:HC was mixed with these dienyl components at
230 1:5:70 (25. 125, 1750 µg), showing an optimal ratio of the trienyl hydrocarbon component for
231 the pheromone system in this species.

232 In the laboratory tests, the numbers of source contacts by male moths significantly
233 increased when Z3,Z6,Z9-23:HC was added to the binary blend. In some lepidopteran
234 species, hydrocarbons of body waxes have critical effects, such as a releaser for copulation
235 (Grant et al. 1987) or stimulator for contact to pheromone source (Schlamp et al. 2005; Xiao
236 et al 2010; 2011; 2012), over short range behaviors such as synergistic effects with other high
237 volatile pheromone components. Xiao (2011) showed the possibility that although their
238 actual functions are unknown, homologous polyene hydrocarbons including Z3,Z6,Z9-23:HC
239 also widely exist in body wax of moths other than Crambidae, because similar synergistic
240 activity was observed when body wax extracts of some Noctuidae and Sphingidae species
241 were mixed with the two aldehydes as sex pheromone components.

242 The four families, Noctuidae, Arctiidae, Lymantriidae and Geometridae use Type II
243 compounds as female sex pheromones (Ando, 2014; El-Sayed, 2014). However, Zahiri et al
244 (2010) reconstructed Noctuidae *sensu lato* by molecular phylogeny, and showed traditional
245 Arctiidae and Lymantriidae *sensu* Miller (1991) were included in Erebidae with various Type
246 II-pheromone-using noctuids. This indicated that only Geometroidea and Noctuoidea, which

247 show sister linkages in recent molecular phylogenetic trees (Regier et al 2009) use Type II sex
248 pheromones and also that the origin of Type II pheromones may be from a common ancestor
249 of the two taxa. However, recently hybrid type pheromone system has been reported in
250 several Pyraloidea species (Cabrera et al 2001; Millar et al. 2005; Leal et al. 2005; Gibb et al.
251 2007; Miller et al. 2010; Lofstedt et al. 2012; El-Sayed et al. 2013; Yan et al. 2014). In
252 Pyraustinae *sensu lato*, *R. surusalis* is the 4th species that has a hybrid type pheromone
253 system as shown in two *Conogethes* species (Xiao et al 2010, 2011b, 2012; El-Sayed et al
254 2013) and *Omphisa anastomosalis* (Yan et al 2014). These results suggest that the hybrid
255 type pheromone system is at least common in Pyraloidea, and the origin of Type II
256 pheromones may be a common ancestor of Pyraloidea and Geometroidea +
257 Noctuoidea. However, the Pyraloidea + (Geometroidea + Noctuoidea) clade include some
258 taxa, e.g., Bombycoidea, Lasiocampoidea or Drepanoidea, that have no reports of Type II
259 pheromones (Regier et al 2009). To reveal the origin of Type II pheromones, we must
260 carefully reinvestigate some species which use only Type I compounds for their female sex
261 pheromones, included into the Pyraloidea + (Gemoetridea + Noctuoidea clade), by
262 physiological or molecular biological methods.

263 Three Crambidae species, *Haritalodes derogate*, *H. basipunctalis* and *R. surusalis* use
264 E10,Z12-16:Ald as a sex pheromone component, and occur sympatrically in hibiscus
265 plantations. This sympatric reproductive biology may be allowed by their species-specific
266 pheromone systems, which consist of binary mixtures of E10,Z12-16:Ald and E10,E12-
267 16:Ald at different ratios in the two *Haritalodes* (*Notracha*) species (Honda et al., 1994), and
268 addition of E10,Z12-16:OAc and Z3,Z6,Z9-23:HC in *R. surusalis*.

269

270

Acknowledgments

271 This work was supported in part by a grant-in-aid for scientific research from the Ministry of
272 Education, Culture, Sports, Science, and Technology (MEXT) of Japan [KAKENHI, Grant-
273 in-Aid for Scientific Research (C) No. 25450071]. We thank Dr. DeMar Taylor for improving
274 the English of the manuscript.

275

276

References

277 Ades GWJ and Kendrick RC (Eds) (2004) Hong Kong Fauna- A Checklist of Selected Taxa
278 Fauna Conservation Department Kadoorie Farm & Botanic Garden Corporation, 87pp.

279

280 Ando T (2014) Internet database: <http://www.tuat.ac.jp/~antetsu/LepiPheroList.htm>

281

282 Ando T, Ogura Y, Uchiyama M (1988) Mass spectra of lepidopterous sex pheromones with a
283 conjugated diene system. *Agric Biol Chem* 52:1415–1423.
284

285 Ando T, Inomata S, Yamamoto M (2004) Lepidopteran sex pheromone. *Top Curr Chem* 239:
286 51–96.
287

288 Ando T, Kawai T, Matsuoka K (2008) Epoxyalkenyl sex pheromones produced by female
289 moths in highly evolved groups: biosynthesis and its endocrine regulation. *J Pestic Sci* 33:
290 17–20.
291

292 Anonymous (1994) Check List of Insects from Korea. Kon-kuk University Press. Seoul,
293 744pp.
294

295 Anonymous (2006) Major Insect and Other Pests of Economic Plants in Japan (revised
296 edition). The Japanese Society of Applied Entomology and Zoology, Tokyo, 381pp.
297

298 Bea Y. (2012) The final report of researches for publication of Korean Red List: Insects. *Nat.*
299 *Inst. Biol. Res, Biol. Res. Co. Div.* 164pp. (in Korean)
300

301 Cabrera A, Eiras AE, Gries G, Gries R, Urdaneta N, Mirás B, Badji C, Jaffe K (2001) Sex
302 pheromone of tomato fruit borer, *Neoleucinodes elegantalis*. *J Chem Ecol* 27: 2097–2107.
303

304 Cork A, Chamberlain DJ, Beevor PS, Hall DR, Nesbitt BF, Campion DG, Attique MR (1988)
305 Components of female sex pheromone of spotted bollworm, *Earias vittella* F. (Lepidoptera:
306 Noctuidae): identification and field evaluation in Pakistan. *J Chem Ecol* 14: 929–945.
307

308 Dai XJ, Xu SF, Wang MZ, Zhu YX, Tang XH, Zhu JW, Du JW, Dong TX, Du MZ (1988)
309 10E,12Z-hexadecadienyl acetate - sex pheromone of the mulberry white caterpillar *Rondotia*
310 *menciana* Moore (Lepidoptera: Bombycidae). *Kexue Tongbao* 33: 1575-1576.
311

312 Daimon T, Fujii T, Yago M, Hsu YF, Nakajima Y, Fujii T, Katsuma S, Ishikawa Y, Shimada T
313 (2012) Female sex pheromone and male behavioral responses of the bombycid moth *Trilocho*
314 *varians*: comparison with those of the domesticated silkmoth *Bombyx mori*.
315 *Naturwissenschaften* 99: 201-215. Doi: 10.1007/s00114-012-0887-3
316

317 Dool HD, Kratz PD (1963) A generalization of the retention index system including linear
318 temperature programmed gas—liquid partition chromatography. *J Chromatogr* 11: 463–471
319

320 EI-Sayed AM, Gibb AR, Mitchell VJ, Manning L-AM, Revell J, Thistleton B (2013)
321 Identification of the sex pheromone of *Conogethes pluto*: a pest of *Alpinia*. *Chemoecology*
322 23:93-101.
323

324 Gibb AR, Pinese B, Tenakanai D, Kawi AP, Bunn B, Ramankutty P, Suckling DM (2007) (Z)-
325 11-Hexadecenal and (3Z,6Z,9Z)-tricosatriene: sex pheromone components of the red banded
326 mango caterpillar *Deanolis sublimbalis*. *J Chem Ecol* 33: 579-589.
327

328 Grant GG, Frech D, MacDonald L, Slessor KN, King GGS (1987) Copulation releaser
329 pheromone in body scales of female whitemarked tussock moth, *Orgyia leucostigma*
330 (Lepidoptera: Lymantriidae): Identification and behavioral role. *J Chem Ecol* 13: 345–356.
331

332 Herbison-Evans D and Crossley SA (2013) Australian Spilomelinae - Lepidoptera Larvae of
333 Australia, <http://lepidoptera.butterflyhouse.com.au/spil/spilomelinae.html>
334

335 Honda H, Himeno K, Yoshiyasu Y (1994) Chemotaxonomy of the cotton leaf-roller
336 (Lepidoptera: Pyralidae) in Japan with special reference to differences in sex pheromones.
337 *Appl Entamol Zool* 29: 323–330.
338

339 Howard RW, Blomquist GJ (2004) Ecological, behavioral, and biochemical aspects of insect
340 hydrocarbons. *Annu Rev Entomol* 50: 371–393.
341

342 Inoue H, Sugie S, Kuroko H, Moriuti S, Kawabe A (1982) *Moths of Japan I*. Kodansha,
343 Toky, Japan.
344

345 Kim Y, Cho Y, Kang Y-K, Choi M, Nam S-H (2013) A study of the major insect pest
346 communities associated with *Hibiscus syriacus* (Columniferae, Malvaceae). *J. Ecol. Environ.*
347 36: 125-129.
348

349 Kováts E (1958) Gas-chromatographische charakterisierung organischer verbindungen. teil 1:
350 Retentionsindices aliphatischer halogenide, alkohole, aldehyde und ketone. *Helv Chim Acta*
351 41: 1915-1932.

352

353 Leal WS, Parra-Pedrazzoli AL, Kaissling KE, Morgan TI, Zalom FG, Pesak DJ, Dundulis EA,
354 Burks CS, Higbee BS (2005) Unusual pheromone chemistry in the navel orangeworm: novel
355 sex attractants and a behavioral antagonist. *Naturwissenschaften* 92: 139–146.

356

357 Lee, SY, Nam SH and Seok O (2005) Insect community complex of *Hibiscus syriacus*. 2005
358 ESA Annual meeting, <http://abstracts.co.allenpress.com/pweb/esa2005/document/49852>

359

360 Liu, Y. (1990) The insect fauna at Jianfengling in Hainan Island – Pyralidae. *Forest Research*
361 3: 574-579.

362

363 Löfstedt C, Svensson GP, Jirle EV, Rosenberg O, Roques A, Millar JG (2012)
364 3Z,6Z,9Z,12Z,15Z)-pentacosapentaene and (9Z,11E)-tetradecadienyl acetate: sex pheromone
365 of the spruce coneworm *Dioryctria abietella* (Lepidoptera: Pyralidae). *J Appl Entomol* 136:
366 70–78.

367

368 McElfresh JS, Millar JG (1999a) Sex pheromone of the common sheep moth, *Hemileuca*
369 *elganterina*, from the San Gabriel Mountains of California. *J Chem Ecol* 25: 687-709.

370

371 McElfresh JS, Millar JG (1999b) Sex attractant pheromone saturniida moth, *Coloradia velda*.
372 *J Chem Ecol* 25: 1067-1078.

373

374 McElfresh JS, Millar JG (1999c) Geographic variation in sex pheromone blend of *Hemileuca*
375 *electra* from southern California. *J Chem Ecol* 27: 1409-1422.

376

377 McElfresh JS, Hammond AM, Millar JG (2001) Sex pheromone components of the buck
378 moth *Hemileuca maja*. *J Chem Ecol* 25: 2505-2525.

379

380 Millar JG, Grant GG, McElfresh JS, Strong W, Rudolph C, Stein JD, Moreira JA (2005)
381 Pheromone Component of the Fir Coneworm Moth, *Dioryctria abietivorella*. *J Chem Ecol*
382 31: 1229-1234.

383

384 Miller JS (1991) Cladistics and classification of the Notodontidae (Lepidoptera, Noctuoidea)
385 based on larval and adult morphology. *Bulletin of the American Museum of Natural History*
386 204: 1-230.

387

388 Miller DR, Millar JG, Mangini A, Crowe CM, Grant GG (2010) Pentacosapentaene and (Z)-
389 11-Hexadecenyl Acetate: Sex Attractant Blend for *Dioryctria amatella* (Lepidoptera:
390 Pyralidae). J Econ Entomol 103: 1216-1221.

391

392 Schlamp KK, Gries R, Khaskin G, Brown K, Khaskin E, Judd GTR, Gries G (2005)
393 Pheromone components from body scales of female *Anarsia lineatella* induce contacts by
394 conspecific males. J Chem Ecol 31: 2897–2911.

395

396 Shibuya J (1928) The Systematic Study on the Formosan Pyralidae. J Fac Agric Hokkaido
397 Imp Univ 21(1): 1-300.

398

399 Shibuya J (1929) On the known and unrecorded species of the Japanese Pyraustinae (Lepid.).
400 J Fac Agric Hokkaido Imp Univ 25(3): 151-242.

401

402 Shimizu K, Tamaki Y (1980) Releasers of male copulatory attempt in the smaller tea tortrix
403 moth (Lepidoptera: Tortricidae). Appl Entomol Zool 15: 140–150.

404

405 Shin YH (2001) Coloured Illustrations The Moths of Korea. Academy Book Publishing Co.,
406 Seoul. (In Korean.)

407

408 R core team (2013) R: A language and environment for statistical computing. R Foundation
409 for Statistical Computing, Vienna, URL [http:// www.R-project.org/](http://www.R-project.org/)

410

411 Regier JC, Zwick A, Cummings MP, Kawahara AY, Cho S, Weller S, Roe A, Baixeras J,
412 Brown JW, Parr C, Davis DR, Epstein M, Hallwachs W, Hausmann A, Janzen DH, Kitching
413 IJ, Solis MA, Yen SH, Bazinet AL, Mitter C (2009) Toward reconstructing the evolution of
414 advanced moths and butterflies (Lepidoptera: Ditrysia): an initial molecular study. BMC
415 Evol. Biol. 9: 280.

416

417 Uehara T, Naka H, Matsuyama S, Ando T, Honda H (2012) Identification and field evaluation
418 of sex pheromones in two hawk moths *Deilephila elpenor lewisii* and *Theretra oldenlandiae*
419 *oldenlandiae* (Lepidoptera: Sphingidae). Appl Entomol Zool 47: 227–232.

420

421 Uehara T, Naka H, Matsuyama S, Ando T, Honda H (2015) Sex Pheromone of the diurnal hawk
422 moth, *Hemaris affinis*. J Chem Ecol 41: 9–14.

423

424 Witzgall P, Kirsch P, Cork A (2010) Sex pheromones and their impact on pest management. J
425 Chem Ecol 36: 80–100.

426

427 Xiao W, Honda H (2010) Non-polar body waxes enhance sex pheromone activity in the
428 yellow peach moth, *Conogethes punctiferalis* (Guene'e) (Lepidoptera: Crambidae). Appl
429 Entomol Zool 45: 449–456.

430

431 Xiao W (2011) Identification and Function of Body Wax Hydrocarbons as a Sex Pheromone
432 of the Yellow Peach Moth. PhD dissertation of the University of Tsukuba, 97pp.

433

434 Xiao W, Honda H, Matsuyama S (2011) Monoenyl hydrocarbons in female body wax of the
435 yellow peach moth as synergists of aldehyde pheromone components. J Chem Ecol 46: 239-
436 246.

437

438 Xiao W, Matsuyama S, Ando T, Millar JG, Honda H (2012) Unsaturated cuticular
439 hydrocarbons synergize responses to sex attractant pheromone in the yellow peach moth,
440 *Conogethes punctiferalis*. J Chem Ecol 38: 1143-50.

441

442 Yan Q, Vang LV, Khanh CNQ, Naka H, Ando T (2014) Reexamination of the female sex
443 pheromone of the sweet potato vine borer moth: Identification of tricosatriene and its field
444 evaluation. J Chem Ecol 40: 590-598.

445

446 Zahiri R, Kitching IJ, Lafontaine JD, Mutanen M, Kaila L, Holloway JD, Wahlberg N (2010)
447 A new molecular phylogeny offers hope for a stable family level classification of the
448 Noctuoidea (Lepidoptera). Zool. Scr. 40(2): 158-173.

449

450

451

452

453 Figure legends

454

455 **Fig.1** GC/EAD analysis of a crude pheromone extract from *R. surusalis* on HP-5MS GC
456 column (upper trace EAD, lower trace GC)

457

458 **Fig.2** Cumulative number of male exhibiting orientation flight (OF) to pheromone source and
459 source contact (SC) in laboratory assays. The amount of the synthetic components in the
460 respective baits are shown under the bars. Bars with the same letters are not significantly
461 different at $P<0.05$ by Tukey–Kramer’s HSD test after ANOVA (OF: $N=5$, $F=56.75$, $P<0.01$;
462 SC: 21.31 , $P<0.01$). The number of trapped males was transformed to $\sqrt{(x+0.5)}$ prior to the
463 test.

464

465 **Fig.3** Field catches of male *R. surusalis* in traps baited with synthetic E10,Z12-16:Ald(Ald),
466 E10,Z12-16:OAc(OAC) and Z3,Z6,Z9-23:HC(HC) and their mixtures. Bars with the same
467 letters are not significantly different at $P<0.05$ by Tukey–Kramer’s HSD test after ANOVA
468 ($N=9$, $F=5.838$, $P<0.01$). The number of trapped males was transformed to $\sqrt{(x+0.5)}$ prior to
469 the test.

470

471 **Fig.4** Type of female sex pheromone and molecular phylogenetics in the crade Ditrysia
472 (Lepidoptera). Type II pheromone was identified from 3 taxonomic groups (Geometroidea,
473 Geometridae and Noctuoidea: Erebidae and Pyraloidea). Papilionoidea etc. indicates a crade
474 (((((Nymphalidae + Pieridae) + (Hesperioidea + Hedyloidea)) + Thyridoidea) + (Papilionidae
475 + Calliduloidea)) + (Copromorphoidea + Hyblaeoidea). Alucitoidea, Urodoidea and
476 Choreutoidea were omitted from the phylogenetic tree that was modified from Regier et al
477 (2009).

478

Fig.1 Honda et al.




Table 1

Retention indices of EAD-active components and synthetic compounds on GC columns with different polarities

Compounds	Kovats Retention Index (KRI)	
	HP-5MS	DB-23
Compound A	1862	2252
Compound B	2051	2369
Compound C	2276	2391
Z10,E12-16:Ald	1853	2243
E10,Z12-16:Ald	1862	2252
Z10,Z12-16:Ald	1874	2254
E10,E12-16:Ald	1880	2257
Z10,E12-16:OAc	2039	2360
E10,Z12-16:OAc	2051	2369
Z10,Z12-16:OAc	2063	2372
E10,E12-16:OAc	2069	2374
Z3,Z6,Z9-23:HC	2276	2390

Fig.2 Honda et al.




Fig.3 Honda et al.




Fig.4 Honda et al.

