

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Differential Induction of Type I Interferons in Macaques by Wild-type Measles Virus or Wild-type Measles Virus with the Hemagglutinin Protein of the Edmonston Vaccine strain

Nguyen Van Nguyen¹, Sei-ich Kato¹, Kyosuke Nagata¹, and Kaoru Takeuchi²

¹Department of Infection Biology and ²Laboratory of Environmental Microbiology, Faculty of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan

Running title: Interferon induction by measles virus

Corresponding author: Kaoru Takeuchi

Division of Environmental Microbiology, Faculty of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan.

E-mail: ktakeuch@md.tsukuba.ac.jp

Tel: +81-29-853-3472, fax: +81-29-853-3472

Subject Section : Virology

Specified Fields : Vaccines and antiviral agents

23

24 List of Abbreviations :

25 BAL, bronchoalveolar lavage ; DI, defective interfering ; EGFP, enhanced green
26 fluorescent protein ; ELISA, enzyme-linked immunosorbent assay; H protein,
27 hemagglutinin protein ; IFN, interferon ; ingLNs, inguinal lymph nodes ; MDA5,
28 melanoma differentiation-associated gene 5; MV, measles virus ; PBMCs, peripheral
29 blood mononuclear cells ; pDC, plasmacytoid DCs; PVRL4, poliovirus receptor-like
30 protein 4 ; RIG-I, retinoic acid-inducible gene-I; RT-PCR, reverse transcriptase PCR ;
31 SLAM, signaling lymphocyte activation molecule

32 **ABSTRACT**

33 Measles vaccines are highly effective and safe, but the mechanism underlying
34 their attenuation has not been well understood. In this study, type I interferons (IFNs)
35 (IFN- α and IFN- β) induction in macaques infected with measles virus (MV) strains
36 was examined. Type I IFNs were not induced in macaques infected with wild-type MV.
37 However, IFN- α was sharply induced in most of macaques infected with the
38 recombinant wild-type MV bearing the hemagglutinin (H) protein of the Edmonston
39 vaccine strain. These results indicate that the H protein of MV vaccine strains may
40 have a role in MV attenuation.

41

42 Key words

43 interferon, hemagglutinin protein, macaque, measles virus

44

45 Measles is highly infectious and remains a major cause of childhood morbidity
46 and mortality worldwide despite the availability of effective vaccines (1). Measles
47 vaccines were generated by successive passages of field isolates of measles virus (MV)
48 in cells from different origins (1). Measles vaccines are highly effective and safe, but
49 the mechanism underlying their attenuation has not been well understood. Major
50 difference between wild-type and vaccine strains of MV is receptor specificity of the
51 hemagglutinin (H) protein in vitro. The H proteins of wild-type MV strains recognize
52 the signaling lymphocyte activation molecule (SLAM), also known as CD150, which
53 is expressed in certain immune system cells (2), and nectin-4, also known as poliovirus
54 receptor-like protein 4 (PVRL4), which is expressed in epithelial cells in trachea, skin,
55 lung, prostate and stomach, as cellular receptors (3, 4). On the other hand, the H
56 proteins of vaccine MV strains recognize CD46, which is ubiquitously expressed on all
57 nucleated human and monkey cells (5, 6), in addition to SLAM and nectin-4 as cellular
58 receptors.

59 To examine the contribution of the H protein to the MV attenuation, an enhanced
60 green fluorescent protein (EGFP)-expressing recombinant wild-type MV bearing the H
61 protein of the Edmonston vaccine strain (EdH-EGFP₂) was generated using a reverse
62 genetics system based on the pathogenic wild-type IC-B strain (7), and cynomolgus
63 monkeys between 4 to 5 years old (three animals per each strain) were intranasally
64 infected with wild-type MV (IC323-EGFP₂) or EdH-EGFP₂ (8). IC323-EGFP₂ and
65 EdH-EGFP₂ have the EGFP gene between the N and P genes. All animal experiments
66 were performed in compliance with the guidelines of National Institute of Infectious

67 Disease (permission number 510008). Interestingly, the replication of EdH-EGFP₂ in
68 tissues and lymphocytes of infected macaques were significantly lower than those of
69 the wild-type MV. From these results we speculated that type I interferons (IFNs) may
70 affect the growth of EdH-EGFP₂ in macaques, as type I IFNs are induced by many
71 viruses and play central roles in the host defense against viral infection (9).

72 In this study, we examined type I IFNs induction in those macaques to investigate
73 the mechanism for the growth attenuation of EdH-EGFP₂ in macaques. For this
74 purpose, we first examined the presence of defective interfering (DI) RNA in virus
75 stocks, because it is known that DI RNAs, especially the 5' copy-back DI RNAs, in
76 virus stocks of MV is able to induce type I IFN through interaction with the RNA
77 helicases retinoic acid-inducible gene-I (RIG-I)/ melanoma differentiation-associated
78 gene 5 (MDA5) (10, 11, 12, 13). Viral RNA was extracted from virus stocks using
79 QIAamp Viral RNA Mini kit (QIAGEN), and DI RNA was detected using reverse
80 transcriptase PCR (RT-PCR) as previously described (10). DI RNAs were detected in
81 the Edmonston (laboratory strain) and IC-V (wild-type strain isolated in Vero cells)
82 stocks as reported (10), while no DI RNA was detected in IC323-EGFP₂ and
83 EdH-EGFP₂ virus stocks used in this experiment (Fig. 1).

84 To examine the interferon responses elicited by IC323-EGFP₂ and EdH-EGFP₂ in
85 macaques, we compared the transcription of IFN- α and IFN- β genes in peripheral
86 blood mononuclear cells (PBMCs) of infected monkeys as previously reported (14).
87 PBMCs were collected at 0, 3, 7, and 10 days post infection (dpi), and stored in
88 RNAProtect Animal Blood Tubes (QIAGEN, Hilden, Germany) at -30°C. Tissues of

89 inguinal lymph nodes (IngLN) were collected at 7 days prior infection and at 10 dpi,
90 and stored in RNAlater solution (QIAGEN) at -30°C. Tissues of lung were collected at
91 10 dpi and stored in RNAlater solution at -30°C. Plasma was collected at 7 days prior
92 infection, and at 0, 3, 7, and 10 dpi, and stored at -80°C. Bronchoalveolar lavage
93 (BAL) were collected at 10 dpi and stored at -80°C. Total RNA was isolated from
94 PBMCs and tissues by using RNeasy mini kit and RNase-free DNase (QIAGEN)
95 according to the manufacturer's protocol, reverse transcribed using oligo (dT) primer
96 and PCR amplified with a Thermal Cycler Dice TP800 (Takara, Tokyo, Japan) by
97 using FastStart SYBR Green Master (Roche, Mannheim, Germany). In macaques
98 infected with IC323-EGFP₂ (no. 5058, 5062, and 5069), IFN- α and IFN- β
99 transcription was transiently down-regulated at day 3 (Fig. 2A and B). Then, the levels
100 of IFN- α and IFN- β transcription were returned to the baseline at day 7. In macaques
101 infected with EdH-EGFP₂ (no. 5056, 5057, and 5068), IFN- α and IFN- β transcription
102 was gradually induced from day 0 to day 7. At day 10, the levels of IFN- α and IFN- β
103 transcription were decreased in all monkeys infected with both strains.

104 Type I IFN responses elicited by IC323-EGFP₂ and EdH-EGFP₂ were also examined
105 in several tissues of infected monkeys. However, the levels of IFN- α and IFN- β
106 transcription in inguinal lymph nodes were not significantly changed between at 7 days
107 before infection and at day 10 for all monkeys (Fig. 2C and D). Day 10 may be too late
108 to detect the change of the levels of IFN- α and IFN- β transcription. Similar levels of
109 IFN- α and IFN- β transcription were detected in lung of monkeys infected with both
110 strains (Fig. 2C and D).

111 Next, we examined plasma levels of IFN- α using an enzyme-linked
112 immunosorbent assay (ELISA) kit. IFN- α levels in the plasma and BAL were
113 measured by VeriKineTM cynomolgus/rhesus IFN- α serum ELISA kit (PBL,
114 Piscataway, USA) according to the manufacturer's protocol. Plasma levels of IFN- α
115 were not significantly changed in wild-type IC323-EGFP₂-infected macaques (no.
116 5058, 5062, and 5069), although slight induction was observed in one macaque (no.
117 5062) at day 7 (Fig. 3A). On the other hand, plasma levels of IFN- α were sharply
118 elevated by 4- to 5-fold in two (no. 5056 and 5057) out of three macaques infected
119 with EdH-EGFP₂ at day 7, and then declined by day 10. To confirm the IFN- α
120 induction in EdH-EGFP₂-infected macaques, we examined plasma collected in former
121 experiments in which we infected macaques with recombinant MV strains. In the first
122 group, two old (10 years old) macaques (no. 4568 and 4569) were used. One macaque
123 (no. 4568) was infected with wild-type MV (IC323-EGFP) having the EGFP gene
124 between the leader sequence and the N gene (8), and the other (no. 4569) was infected
125 with IC323-EGFP₂. In the second group, seven juvenile (1 year old) macaques (no.
126 4848, 4849, 4850, 4858, 4859, 4860, and 4865) were used. Three of them (no. 4850,
127 4860, and 4865) were infected with IC323-EGFP₂, and the other four (no. 4848, 4849,
128 4858, and 4859) were infected with EdH-EGFP₂. Again, plasma levels of IFN- α were
129 sharply elevated in two EdH-EGFP₂-infected macaques (no. 4848 and 4849) at day 7
130 but not in IC323-EGFP- and IC323-EGFP₂-infected macaques (no. 4568, 4569, 4850,
131 4860 and 4865) (Fig. 3B). Macaques (no. 4860, 4865, 4858 and 4859) and macaques
132 (no. 4850, 4848 and 4849) were sacrificed at day 3 and 7, respectively. Therefore, their

133 samples were not available hereafter. Induction of IFN- α was not observed in BAL of
134 all infected monkeys at 10 days (Fig. 3A).

135 Although many studies indicated the IFNs production in vitro by MV, little is
136 known about the IFNs production in measles patients. Previous clinical study using a
137 sensitive radioimmunoassay indicated that IFN- α was not induced in plasma of
138 measles patients (15). In another clinical study, Yu et al. found that IFN- α expression
139 was suppressed in PBMCs of measles patients (16). In vivo study using macaques,
140 Devaux et al. reported that expression of type I IFN genes were well regulated (14). In
141 addition, Shivakoti et al. recently found that type I IFNs were not induced in macaques
142 infected by wild-type MV (17). We found that IFN- α was not induced in macaques
143 infected with wild-type MV (Fig. 3A and B). Our result is consistent with previous
144 clinical studies (15, 16) and in vivo studies using monkeys (14, 17). These results
145 suggest that MV has way to circumvent the host IFNs production possibly by the C
146 and V proteins (18). Likewise, little is known about the IFNs production in measles
147 vaccinee. In a previous clinical study, IFN was induced after measles vaccination (19).
148 We found that IFN- α was sharply induced in plasma of macaques infected with
149 EdH-EGFP₂ (Fig. 3). Interestingly, it was shown that large amounts of IFN- α were
150 rapidly produced from plasmacytoid DCs (pDCs) after infection of the Edmonston
151 strain mostly independent of the viral infection cycles (20). As pDCs express CD46 but
152 not SLAM (20), pDCs in macaques would be infected with EdH-EGFP₂ by
153 CD46-mediated pathway and may produce large amounts of IFN- α in plasma. In
154 summary, we found that IFN- α was induced in macaques infected with wild-type MV

155 bearing the H protein of the Edmonston vaccine strain but not with wild-type MV. Our
156 results suggest that the H protein of vaccine strains of MV may have a role in the MV
157 attenuation.

158

159 **ACKNOWLEDGEMENTS**

160 We thank T. Ohkura, N. Nagata, and Y. Ami for continuous supports. We also thank
161 M. Ayata and M. Okuwaki for critical readings and valuable comments. This work was
162 supported in part by a grant-in-aid (No. 21022006 and 23659227) from the Ministry of
163 Education, Culture, Sports, Science and Technology of Japan.

164

165 **DISCLOSURE**

166 The authors have no conflicts of interest associated with this study.

167

168 **REFERENCES**

169 1. Griffin D.E. (2013) Measles. In: Knipe D.M, Howley P.M., eds. Fields virology, 6th
170 edn. vol 1. Philadelphia, PA: Lippincott, Williams & Wilkins, pp. 1042-69.

171

172 2. Tatsuo H., Ono N., Tanaka K., Yanagi Y. (2000) SLAM (CDw150) is a cellular
173 receptor for measles virus. *Nature* **406**: 893-7.

174

175 3. Muhlebach M.D., Mateo M., Sinn P.L., Prufer S., Uhlig K.M., Leonard V.H.,
176 Navaratnarajah C.K., Frenzke M., Wong X.X., Sawatsky B., Ramachandran S., McCray

177 P.B. Jr., Cichutek K., von Messling V., Lopez M., Cattaneo R. (2011) Adherens
178 junction protein nectin-4 is the epithelial receptor for measles virus. *Nature* **480**: 530-3.
179

180 4. Noyce R.S., Bondre D.G., Ha M.N., Lin L.T., Sisson G., Tsao M.S., Richardson C.D.
181 (2011) Tumor cell marker PVRL4 (Nectin 4) is an epithelial cell receptor for measles
182 virus. *PLoS Pathog* **7**: e1002240.
183

184 5. Nanche D., Varior-Krishnan G., Cervoni F., Wild T.F., Rossi B., Rabourdin-Combe
185 C., Gerlier D. (1993) Human membrane cofactor protein (CD46) acts as a cellular
186 receptor for measles virus. *J Virol* **67**:6025-32.
187

188 6. Dorig R.E., Marcil A., Chopra A., Richardson C.D. (1993) The human CD46
189 molecule is a receptor for measles virus (Edmonston strain). *Cell* **75**: 295-305.
190

191 7. Takeda M, Takeuchi K, Miyajima N, Kobune F, Ami Y, Nagata N, Suzaki Y, Nagai Y,
192 Tashiro M. (2000) Recovery of pathogenic measles virus from cloned cDNA. *J Virol.*
193 **74**:6643-7.
194

195 8. Takeuchi K., Nagata N., Kato S.I., Ami Y., Suzaki Y., Suzuki T., Sato Y.,
196 Tsunetsugu-Yokota Y., Mori K., Nguyen N.V., Kimura H., Nagata K. (2012) Wild-type
197 measles virus with the hemagglutinin protein of the Edmonston vaccine strain retains
198 wild-type tropism in macaques. *J Virol* **86**: 3027-37.

199

200 9. Borden E.C., Sen G.C, Uze G, Silverman R.H., Ransohoff R.M., Foster G.R., Stark
201 G.R. (2007) Interferons at age 50: past, current and future impact on biomedicine. *Nat*
202 *Rev Drug Discov* **6**:975-90.

203

204 10. Shingai M., Ebihara T., Begum N.A., Kato A., Homma T., Matsumoto K., Saito H.,
205 Ogura H., Matsumoto M., Seya T. (2007) Differential type I IFN-inducing abilities of
206 wild-type versus vaccine strains of measles virus. *J Immunol* **179**:6123-33.

207

208 11. Kessler J.R., Kremer J.R., Muller C.P. (2011) Interplay of measles virus with early
209 induced cytokines reveals different wild type phenotypes. *Virus Res* **155**:195-202.

210

211 12. Shivakoti R., Siwek M., Hauer D., Schultz K.L., Griffin D.E. (2013) Induction of
212 dendritic cell production of type I and type III interferons by wild-type and vaccine
213 strains of measles virus: role of defective interfering RNAs. *J Virol* **87**: 7816-27.

214

215 13. Ho T.H., Kew C., Lui P.Y., Chan C.P., Satoh T., Akira S., Jin D.Y., Kok K.H.
216 (2015) PACT- and RIG-I-dependent activation of type I interferon production by a
217 defective interfering RNA derived from measles virus vaccine. *J Virol* **90**:1557-68.

218

219 14. Devaux P., Hodge G., McChesney M.B., Cattaneo R. (2008) Attenuation of V- or
220 C-defective measles viruses: infection control by the inflammatory and interferon

221 responses of rhesus monkeys. *J Virol* **82**: 5359-67.

222

223 15. Shiozawa S., Yoshikawa N., Iijima K., Negishi K. (1988) A sensitive
224 radioimmunoassay for circulating α -interferon in the plasma of healthy children and
225 patients with measles virus infection. *Clin Exp Immunol* **73**: 366-9.

226

227 16. Yu X.L., Cheng Y.M., Shi B.S., Qian F.X., Wang F.B., Liu X.N., Yang H.Y., Xu
228 Q.N., Qi T.K., Zha L.J., Yuan Z.H., Ghildyal R. (2008) Measles virus infection in
229 adults induces production of IL-10 and is associated with increased CD4+ CD25+
230 regulatory T cells. *J Immunol* **181**: 7356-66.

231

232 17. Shivakoti R., Hauer D., Adams R.J., Lin W.H., Duprex W.P., de Swart R.L., Griffin
233 D.E. (2015) Limited in vivo production of type I or type III interferon after infection of
234 macaques with vaccine or wild-type strains of measles virus. *J Interferon Cytokine Res*
235 **35**: 292-301.

236

237 18. Goodbourn S., Randall R.E. (2009) The regulation of type I interferon production
238 by paramyxoviruses. *J Interferon Cytokine Res* **29**:539-47.

239

240 19. Petralli J.K., Merigan T.C., Wilbur J.R. (1965) Circulating interferon after measles
241 vaccination. *N Engl J Med* **273**: 198-201.

242

243 20. Duhon T., Herschke F., Azocar O., Drulle J., Plumet S., Delprat C., Schicklin S.,

244 Wild T.F., Roubourdin-Combe C., Gerlier D., Valentin H. (2010) Cellular receptors,
245 differentiation and endocytosis requirements are key factors for type I IFN response by
246 human epithelial, conventional and plasmacytoid dendritic infected cells by measles
247 virus. *Virus Res* **152**: 115-125.

248

249 **FIGURE LEGENDS**

250 **Fig. 1. Absence of 5' copy back DI RNA in MV stocks.** Stocks of IC323-EGFP₂
251 and EdH-EGFP₂ used for infection were tested for the absence of 5' copy back DI
252 RNA. 5' copy back DI genomes were detected with primers (JM396;
253 5'-TATAAGCTTACCAGACAAAGCTGGGAATAGAACTTCG-3' and JM403;
254 5'-CGAAGATATTCTGGTGTAAGTCTAGTA-3'). MV standard genomes were
255 detected using primers (JM396 and JM402;
256 5'-TTTATCCAGAATCTCAARTCCGG-3'). The Edmonston and IC-V strains, which
257 are known to contain 5' copy back DI RNA, were used for positive control.

258

259 **Fig. 2. IFN- α / β mRNA expression in PBMCs, inguinal lymph nodes and lungs.**

260 IFN- α (A) and IFN- β (B) mRNA expression in PBMCs from monkeys infected with
261 IC323-EGFP₂ or EdH-EGFP₂ were measured by RT-qPCR. PBMCs were collected at
262 0, 3, 7, and 10 days post infection (dpi). IFN- α (C) and IFN- β (D) mRNA expression
263 in inguinal lymph nodes (IngLN) and lungs from monkeys infected with IC323-EGFP₂
264 or EdH-EGFP₂ were measured by RT-qPCR. IngLN were excised at 7 days prior to
265 infection and at 10 dpi, and lungs were excised at 10 dpi from monkeys. Three

266 monkeys (no. 5058, 5062, and 5069) were infected with IC323-EGFP₂, and three
267 monkeys (no. 5056, 5057, and 5068) were infected with EdH-EGFP₂. For
268 amplification of the IFN- α mRNA, IFN- α F primer
269 5'-GCCTGAAGGACAGACATGACTTT-3' and IFN- α R primer
270 5'-GGATGGTTTGAGCCTTTTGG-3' were used. For amplification of the IFN- β
271 mRNA, IFN- β F primer 5'-TGCCTCAAGGACAGGATGAA-3' and IFN- β R primer
272 5'-ATGGTCCAGGCACAGTGACT-3' were used. For amplification of the 18S rRNA
273 segment, the 18S sense primer 5'-TCAAGAACGAAAGTCGGAGG-3' and 18S
274 antisense primer 5'-GGACATCTAAGGGCATCACA-3' were used. For determining
275 the relative amounts of IFN- α/β mRNA, the amounts of IFN- α/β mRNA in
276 cynomolgus monkey PBMCs infected with Sendai virus (Cantell strain), which is
277 commonly used to induce IFN- α/β , in vitro were set to 10¹.

278

279 **Fig. 3. Plasma and BAL levels of IFN- α**

280 (A) Plasma and BAL levels of IFN- α in monkeys infected with IC323-EGFP₂ or
281 EdH-EGFP₂ were measured by ELISA. Plasma was collected at 7 days prior
282 infection, and at 0, 3, 7, and 10 dpi. BAL was collected at 10 dpi. Three monkeys
283 (no. 5058, 5062, and 5069) were infected with IC323-EGFP₂, and three monkeys (no.
284 5056, 5057, and 5068) were infected with EdH-EGFP₂. (B) Plasma levels of IFN- α in
285 monkeys infected with IC323-EGFP, IC323-EGFP₂ or EdH-EGFP₂ were measured by
286 ELISA. Plasma was collected at 0, 3, 7, and 10 dpi. One monkey (no. 4568) were
287 infected with IC323-EGFP. Four monkeys (no. 4569, 4850, 4860, and 4865) were

288 infected with IC323-EGFP₂, and four monkeys (no. 4848, 4849, 4858, and 4859) were

289 infected with EdH-EGFP₂. Sensitivity of this assay is 0.30 pg/ml. nd, not done.

290

291

Fig. 1




Fig. 2




Fig. 3

