

1 **Title:**

2 *SEXUAL STERILITY* is essential for both male and female gametogenesis in tomato

3

4 **Running Title:**

5 A gene essential for gametogenesis in tomato

6

7 **Corresponding Author:**

8 Hiroshi Ezura, Graduate School of Life and Environmental Sciences, University of Tsukuba

9 Tennodai 1-1-1, Tsukuba, Ibaraki 305-8577, Japan

10 029-853-4710 (Extansion 4710), E-mail address: ezura.hiroshi.fa@u.tsukuba.ac.jp

11

12 **Subject Areas:**

13 [Growth and development], [Regulation of gene expression]

14

15 **Number of figures and tables:**

16 Black and white figures: 2 Color figures: 6 Tables: 2

17 Supplementary figures: 6 Supplementary tables: 5

18

1 *SEXUAL STERILITY* is essential for both male and female gametogenesis in tomato

2

3 A gene essential for gametogenesis in tomato

4

5 Shuhei HAO, Tohru ARIIZUMI, Hiroshi EZURA

6

7 Graduate School of Life and Environmental Science, University of Tsukuba, Tsukuba, Ibaraki

8 305-8577, Japan

9

10 **Abbreviations:**

11 DBA, day before anthesis; MMC, megaspore mother cell; PMC, pollen mother cell; SISES,

12 *Solanum lycopersicum* *SEXUAL STERILITY*; SPL/NZZ, SPOROCTELESS/NOZZLE; WT,

13 wild type

14

15

16

17

1 **Abstract:** Gametogenesis is a key step in the production of ovules or pollen in higher plants.
2 The molecular aspects of gametogenesis are well characterized in the model plant Arabidopsis;
3 however, little information is known in tomato, which is a model plant for fleshy fruit
4 development. In this study, we characterized a tomato (*Solanum lycopersicum* L.) γ -ray mutant,
5 *sexual sterility* (*Slses*), that exhibited both male and female sterility. Morphological analysis
6 revealed that the *Slses* mutant forms incomplete ovules and wilted anthers devoid of pollen
7 grains at the anthesis stage. Genetic and next-generation sequencing analyses revealed that the
8 *Slses* mutant carried a 13-bp deletion within the first exon of a homologue of
9 SPOROCTELESS/NOZZLE (*SPL/NZZ*), which plays an important role in gametogenesis in
10 Arabidopsis. Complementation analysis in which the complete *SISES* genomic region was
11 introduced into the *Slses* mutant fully restored normal phenotypes, demonstrating that
12 *Solyc07g063670* is responsible for the *Slses* mutation. *SISES* probably act as transcriptional
13 repressor because of EAR motif at the C-terminal region. Gene expression levels of *WUSCHEL*
14 (*SIWUS*) and *INNER NO OUTER* (*SIINO*), both of which are required for ovule development,
15 were dramatically reduced in the early stages of pistil development in the *Slses* mutant,
16 suggesting a positive regulatory role of *SISES* in the transcription of gametogenesis genes and
17 differences in the regulation of *INO* (*SIINO*) and integument development by *SPL/NZZ* (*SLSES*)
18 between Arabidopsis and tomato. Taken together, our results indicate that *SISES* is a novel
19 tomato gametogenesis gene essential for both male and female gametogenesis.

1

2 **Keywords:** Anther, Gametogenesis, Mutant, Ovule, Sexual sterility

3 **Introduction:**

4 Tomato is one of the most important crops produced and consumed worldwide and has
5 also been used as a model plant for studying plants in the Solanaceae family and plants bearing
6 fleshy fruits. To date, diverse tomato studies have been conducted, especially tomato genomic
7 studies, given the availability of the complete genome sequence (The Tomato Genome
8 Consortium, 2012), various public databases, and mutant resources that were generated using
9 breeding lines and model tomato lines, including a dwarf cultivar, ‘Micro-Tom’ (Scott and
10 Harbaugh, 1989; Minoia et al. 2010; Saito et al. 2011).

11 Angiosperms form reproductive organs, such as anthers and pistils, that bear pollen
12 grains and ovules, respectively. In this reproductive system, gametogenesis plays an important
13 role in producing offspring. To date, several genes involved in gametogenesis have been
14 identified in the model plant Arabidopsis, and their stage-specific roles in the regulation of
15 reproductive development have been elucidated (Schneitz et al. 1997; Wilson and Zhang 2009;
16 Bencivenga et al. 2011; Plackett et al. 2011).

17 According to Sanders et al. (1999), the development of anthers and pollen in
18 Arabidopsis is divided into 15 stages. During stages 1 to 4, when the anther primordium
19 emerges and establishes a bilateral structure, *EXCESS MICROSPOROCTES1/EXTRA*

1 *SPOROGENOUS CELLS (EMS1/EXS)*, a putative leucine-rich repeat receptor-like kinase, and
2 *TAPETAL DETERMINANT1 (TPD1)*, a ligand for *EMS1/EXS*, both function to determine
3 archesporial cell number. Both the *Arabidopsis ems1/exs* mutant and the *tpd1* mutant are male
4 sterile and form extra meiocytes caused by additional L2 layer cells, and the tapetal and middle
5 cell layers are absent in these mutants (Canales et al. 2002; Zhao et al. 2002; Yang et al. 2003).
6 Differentiation of the endothecium, middle layer, tapetum and microspore mother cells initiate
7 at stage 5. Then, microspore mother cells undergo meiosis between stages 5 and 7 (Sanders et
8 al. 1999), producing microspores or pollen grains within the tapetal cell layer.
9 *DYSFUNCTIONAL TAPETUM1 (DYTI)* encodes a putative basic helix-loop-helix (bHLH)
10 transcription factor that is predicted to act downstream *EMS1/EXS*, and this action is required
11 for the normal expression of *ABORTED MICROSPORES (AMS)* and *MALE STERILITY1*
12 (*MS1*), which are essential for normal tapetal development and viable pollen production during
13 these stages (Wilson et al. 2001; Sorensen et al. 2002; Zhang et al. 2006).

14 On the other hand, many genes related to ovule development have also been identified
15 in *Arabidopsis* (Schneitz et al. 1997; Skinner et al. 2004). Based on cytological aspects, ovule
16 primordia are divided into 3 elements: the funiculus, chalaza and nucellus. Genes specifically
17 expressed within each part have been identified (Schneitz et al. 1997; Sieber et al. 2004a; Kelly
18 and Gasser 2009). The site-specific expression of the homeobox gene *WUSCHEL (WUS)* within
19 the nucellus is critical for integument development and the formation of normal mature ovules

1 in *Arabidopsis* and tomato (Gross-Hardt et al. 2002; Sicard et al. 2008a). Accordingly, *WUS*
2 expression is generally confined to the nucellus during ovule development based on the
3 regulation by specific genes or hormones in *Arabidopsis*. For example, the *bell1* (*bell*) mutant
4 and the *corona* (*cna*)/*phabulosa* (*phb*)/*phavoluta* (*phv*) triple mutant, which showed abnormal
5 ovule development, exhibit an aberrant *WUS* expression pattern that prevents normal
6 integument formation (Brambilla et al. 2007; Bencivenga et al. 2012; Yamada et al. 2016).
7 Furthermore, *WUS* expression expands into the chalaza with exogenous treatment with the plant
8 hormone cytokinin N⁶-benzylaminopurine (BAP), and the treatment results in the formation of
9 new primordia (Bencivenga et al. 2012). In addition, precise mRNA localization of
10 *AINTEGUMENTA* (*ANT*), which encodes an AP2 family transcription factor, and *INNER NO*
11 *OUTER* (*INO*), which encodes a plant-specific YABBY family transcription factor, at the
12 chalaza is required for normal ovule integument development (Elliott et al. 1996; Baker et al.
13 1997; Villanueva et al, 1999).

14 There are several reports that describe anther and ovule development in tomato. For
15 example, tomato *male sterile 10³⁵* (*MS10³⁵*) has been isolated as a homologous gene of *AtDYT1*,
16 and the *ms10³⁵* mutant exhibits dysfunctional meiosis and aberrant tapetum formation (Jeong
17 et al. 2014). The *parthenocarpic fruit* (*pat*) mutant forms aberrant ovules caused by a failure of
18 integument development (Mazzucato et al. 1998). In addition, the mini zinc finger gene
19 *INHIBITOR OF MERISTEM ACTIVITY* (*IMA*) regulates ovule development by activating D-

1 class gene of ACBD model of flower development expression in tomato and Arabidopsis
2 (Sicard et al. 2008ab). However, gametogenesis genes and their associated mechanisms remain
3 largely unknown in tomato.

4 *SPOROCTELESS/NOZZLE (SPL/NZZ)* plays a key role in gametogenesis, both in
5 anther and ovule development, in Arabidopsis (Schiefthaler et al. 1999; Yang et al. 1999), and
6 loss of function of *SPL/NZZ* results in abnormal germ cell production, causing strong sterility
7 in both male and female reproductive organs. *SPL/NZZ* transcripts are observed in the stamen
8 primordia and developing stamens as well as in ovule primordia, which is induced by
9 *AGAMOUS (AG)* or brassinosteroids (Ito et al. 2004; Ye et al. 2010). *AG* is a member of
10 *MADS*-box gene and it is necessary for carpel and stamen development (Pinyopich et al. 2003).
11 Additionally, *SPL/NZZ* physically binds with *CINCINNATA (CIN)*-like *TEOSINTE*
12 *BRANCHED1/CYCLOIDEA/PCF (TCP)* transcription factors or *YABBY* family transcription
13 factors through the N-terminal domain and the *TOPLESS/TOPLESS-RELATED (TPL/TPR)*
14 co-repressor at the EAR motif (Sieber et al. 2004b; Chen et al. 2014; Wei et al. 2015). Then,
15 *SPL/NZZ* functions as an adaptor-like transcriptional repressor through interaction with
16 *TPL/TPR* (Wei et al. 2015). However, very little is known about the function of *SPL/NZZ* in
17 tomato. In this study, we report the identification and characterization of *SISES*, a homologue
18 of *SPL/NZZ*, as an essential factor of gametogenesis development. This study demonstrates that

1 pollen and ovule development were arrested in the very early developmental stages of each
2 reproductive organ in the *Sl ses* mutant.

3

4 **Results:**

5 *The identification of the sterile Sl ses mutant*

6 To uncover the regulatory mechanism underlying reproductive organ development, we
7 screened for mutants that were defective in normal reproductive development. From the tomato
8 mutant populations that were generated by gamma ray-irradiated lines (Saito et al. 2011), one
9 mutant producing seedless fruits was isolated. To characterize the cytological aspects of the
10 mutant, we first counted the number of each type of flower organ. In both the WT and mutant,
11 the inflorescence was typically composed of 5 sepals, petals and anthers and a single pistil
12 (Table 1). Although the WT formed an inflorescence with swollen anthers containing mature
13 viable pollen grains at the anthesis stage, the mutant formed clearly defective anthers; they
14 appeared wilted and small, and no pollen grains were visible at this time (Fig. 1A-D). To
15 confirm the fertility of the ovules and pollen of the mutant, we crossed WT pollen to the mutant
16 stigma or mutant pollen to the WT stigma and examined whether viable seeds were produced.
17 No pollen was visually observed in the mutant. As the result, no viable seeds were produced
18 from either type of cross-pollination; however, the mutant occasionally produced seedless fruit
19 (data not shown). These results indicated that the mutant showed both male and female sterility.

1 Based on this phenotype, the mutant was designated as *sexual sterility* (*Slses*).

2 Then, to clarify genetic nature of the *Slses* mutation, we conducted crossing analysis
3 with wild type Micro-Tom. Given that the *Slses* homozygous mutant exhibited both complete
4 male and female sterility, heterozygous *Slses* plants were crossed with WT, and the resulting F₁
5 seeds and the following F₂ offspring were obtained (Supplementary Fig. S1). The
6 morphological phenotype and fertility of 17 F₁ plants were almost equivalent to those of WT,
7 indicating that the *Slses* mutation was recessive (Table 2). The phenotypes of the F₂ offspring
8 segregated to 118 (phenotype; WT) : 40 (phenotype; *Slses*), which approximately fit the
9 theoretical ratio of the WT and *Slses* phenotypes 3 : 1, suggesting that the *Slses* mutation was
10 monogenic (Table 2).

11 Next, time course analysis of cross-sections during anther development was conducted.
12 In this analysis, flower buds at different developmental stages, which were defined by the size
13 of flower buds, were obtained, and cross-sections of the anther structure were examined under
14 light microscopy. There was no obvious difference in timing of flowering and flower
15 development between WT and the *Slses* mutant. At the stage when the flower buds were 1 to 2
16 mm and 2 to 3 mm, normal formation of the vascular bundle (VB), epidermis (Ep), endothecium
17 (En), middle layer (ML), tapetum (T), and pollen mother cells (PMC, 1-2 mm bud) or meiotic
18 cells (MC, 2-3 mm bud) were observed in WT (Fig. 2E, F). In contrast, the formation of the En,
19 ML, T, PMC and MC were not clearly evident, and we only observed the vascular bundle and

1 epidermis at the stage of 1 to 2 mm and 2 to 3 mm buds in the *Slses* mutant (Fig. 2I, J).
2 Microspores (Msp) were observed at the bud stage of 4 to 5 mm in WT but not in the *Slses*
3 mutant (Fig. 2G, K). At anthesis stage, when the bud size is 6 to 7 mm, many mature pollen
4 grains were clearly observed in the anther locules in the WT, whereas the *Slses* mutant never
5 underwent pollen grain formation inside the anther (Fig. 2A-D, H, L). The anther structure
6 appeared wilted and less colored in the mutant compared with the WT (Fig. 1A, B, Fig. 2A-D).

7 Next, we observed the pistil in the WT and the *Slses* mutant (Fig. 1E-I). The size of
8 the ovary at anthesis in the *Slses* mutant was clearly smaller than that in the WT (Fig. 1I). At
9 the 1 to 2 mm bud size stage, we observed ovule primordia around the placenta in both the WT
10 and the *Slses* mutant (Fig. 3E, I). At the 2 to 3 mm and 4 to 5 mm bud stages, ovule integuments
11 and megaspore mother cells (MMCs) were observed, and the ovule primordia enlarged as the
12 stages proceeded in WT. We could not observe the ovule integuments and MMC, which are
13 essential for complete ovule development, and the enlargement level of ovule primordia was
14 quite low in the *Slses* mutant (Fig. 3 F, G, J, K, M-O). From this stage, ovule development was
15 arrested in the *Slses* mutant. We observed cross-sections of mature ovules at the anthesis stage
16 in WT. The ovules were smaller, and their structure appeared disorganized in the *Slses* mutant
17 (Fig. 3A-D, H, L).

18

19 *The Slses mutant produces seedless fruit*

1 Despite male and female sterility, the *Slses* mutant occasionally bore red mature fruits
2 and the size and weight of these fruits were significantly reduced compared with WT fruits (Fig.
3 4A-D). Most of mature fruits produced in the WT contained 3 locules (average 3.2 locules per
4 fruit), whereas those in the *Slses* mutant contained 2 locules (average 2.2 locules per fruit),
5 indicating that the number of locules was significantly reduced in the *Slses* mutant (Fig. 4E).
6 The mature red fruits produced in the *Slses* mutant were completely seedless, indicating that
7 the *Slses* mutant induces parthenocarpy. However, molecular mechanism of how the *Slses*
8 mutation induces parthenocarpy is still unclear.

9

10 *SISES* encodes *SPL/NZZ*

11 To map the *Slses* mutation, positional cloning was performed using SolCap Infinium
12 carrying a large number of SNPs that could discriminate tomato genotypes (Sim et al. 2012),
13 with 2098 markers available for distinguishing between ‘Micro-Tom’ and ‘Ailsa Craig’. The
14 SolCap array coupled with 28 F₂ mapping populations that were derived from crosses between
15 *Slses* heterozygous plants and Ailsa Craig allowed for narrowing down the candidate area to a
16 2.3 Mbp region corresponding to the physical position between 62,465,682 base pairs (bps) to
17 64,759,097 bps on chromosome 7 based on the protein annotation extracted from version
18 SL2.40 from the SGN database (Fig. 5A). The 2.3 Mbp region spans 334 open reading frames
19 (ORFs), and none of these ORFs are involved in tomato gametogenesis

1 (http://www.kazusa.or.jp/tomato_sbm/). Then, we obtained the whole genome sequence of the
2 *Slses* mutant using Illumina HiSeq 2000 next-generation sequencing, and its genome
3 information was compared with the tomato reference genome sequence according to Ariizumi
4 et al. (2014). Given that the *Slses* mutant was a monogenic recessive mutation, we exclusively
5 focused on homozygous mutations specifically found in the *Slses* mutant. The genome sequence
6 analysis identified 94 homozygous mutations consisting of 9 point mutations and 85 insertion
7 and deletion (indel) mutations within the 2.3 Mbp region. The 9 point mutations included 1
8 nonsynonymous, 1 synonymous, and 7 intergenic mutations. The 85 indels included 2
9 mutations within exon causing modification of amino acid sequence of putative ORFs and 19
10 mutations within intron or UTR regions. In addition, 64 mutations were present within the
11 intergenic region (Supplementary Table S1). Therefore, 1 missense and 1 deletion mutation,
12 which were found in *Solyc07g063670*, and 1 insertion mutation, which was found in
13 *Solyc07g065700*, were considered candidates for the *SISES* gene. Among them, one deletion
14 mutation, which caused a 13-bp deletion from the first exon of *Solyc07g063670*, was primarily
15 selected because the *Slses* mutant was produced by gamma-ray irradiation and caused massive
16 frame shift rather than the insertion mutation. Gamma-ray irradiation often causes genome
17 insertion or deletion rather than SNPs, and we often obtain frame shift caused mutant. The
18 deletion caused a frame shift at the 123rd nucleotide position from the start codon, which was
19 considered the 1st position. As the results of frame shift the putative amino acid was shortened

1 from 353 to 57 in the *Sl ses* mutant (Fig. 5B, C).

2 Genetic linkage analysis using a DNA marker (*SlSES* marker-1; Supplementary Table
3 S2) distinguishes this deletion and the F₂ populations derived from a cross between the WT
4 Micro-Tom and the *Sl ses* mutant, revealing a perfect linkage between the mutation and the
5 phenotypes (Supplementary Fig. S2). The candidate gene of the *Sl ses* mutant, *Solyc07g063670*,
6 encodes a domain of plant transcription factor NOZZLE (29-215 a.a.; NCBI Conserved Domain
7 Search; Fig. 5C). This protein shares 26.4% identity (39.2% similarity) with the SPL/NZZ
8 protein and 21.1% identity (31.8% similarity) with the TCP interactor containing EAR motif 3
9 (TIE3) protein of Arabidopsis (Fig. 5D; Supplementary Table S3). The candidate protein also
10 had an EAR motif (LxLxL; Kagale and Rozwadowski 2011) at the C-terminal region (345-351
11 a.a.) similar to that found in the SPL/NZZ and TIE proteins of Arabidopsis (Fig. 5E). Further
12 sequence analysis indicated that no paralogues of the candidate gene existed in the tomato
13 genome (Fig. 5D), suggesting that *SlSES* exists as a single gene and plays a non-redundant role
14 in the tomato genome.

15

16 *Complementation experiments of the Sl ses mutant*

17 To confirm whether the candidate gene *Solyc07g063670* was responsible for the *Sl ses*
18 mutation, the genomic region of the candidate gene, including the 2.5-kb promoter region,
19 1,310-bp coding sequence, and 700-bp terminator region, were introduced into the offspring of

1 heterozygous *Slses* plants (Supplementary Fig. S3) such that the transgene randomly integrated
2 into plants that were homozygous, heterozygous or azygous for the 13-bp nucleotide deletion
3 within *Solyc07g063670*. We developed a PCR genotyping method that allowed for
4 discrimination of 13-bp nucleotide deletion such that we selected the transgenic plants that
5 carried both the transgene and the endogenous homozygous 13-bp nucleotide deletion (*SISES*
6 marker-2; Supplementary Table S2; Supplementary Fig. S4). Three such transgenic plants were
7 obtained. The plants exhibited swollen anthers, producing fertile pollen grains and fertile
8 ovules; thus, self-pollination produced viable offspring seeds that germinated thereafter
9 (Supplementary Fig. S5), indicating that the phenotypes in the complementation lines were
10 fully restored. These results indicated that the 13-bp deletion found in *Solyc07g063670* was
11 responsible for the *Slses* mutation and that the Arabidopsis *SPL/NZZ* homologue in tomato
12 (*SISES*) functions to regulate normal reproductive organ development.

13

14 *SISES is expressed at early stages of reproductive development and regulates SIWUS and SIINO*
15 *expression*

16 qRT-PCR analysis was conducted to examine the expression pattern of the *SISES* gene
17 (Fig. 6). mRNA was extracted from flower buds 1 mm in size; pistils from developmental stages
18 when the flower bud is 3, 4.5 and 6 mm in size; pistils 1 day before anthesis (DBA); and pistils
19 at the anthesis stage. In this analysis, several genes related to anther or ovule development in

1 tomato or Arabidopsis were also examined regarding the effect of the *Sl ses* mutation on the
2 transcriptional regulation of gametogenesis genes. *SISES* mRNA was highly expressed in 1 mm
3 flower buds and 3 mm pistils. However, mRNA levels decreased along with pistil development
4 and were barely detectable in 6 mm pistils and 1 DBA buds in WT (Fig. 6). mRNA expression
5 of *Male sterile 10³⁵* (*MS10³⁵*), which is involved in both meiosis and programmed cell death of
6 the tapetum during microsporogenesis in tomato (Jeong et al. 2014), was evident in 1 mm WT
7 buds, whereas its mRNA was minimally detectable in the *Sl ses* mutant (Fig. 7A). mRNA
8 expression of *BARELY ANY MERISTEM1* and *2-like* (*BAM1/2-like*) and *EMS1/EXS-like*, which
9 are homologues of Arabidopsis *BAM1/2* and *EMS1/EXS*, respectively, and are essential for early
10 anther development (Zhao et al. 2002; Hord et al. 2006), were significantly downregulated in
11 the *Sl ses* mutant (Fig. 7B, C). mRNA expression of *TOMATO AGAMOUS1* (*TAG1*), a
12 homologue of Arabidopsis *AG1*, which is involved in the transcriptional regulation of *SPL/NZZ*,
13 was decreased in the *Sl ses* mutant compared with the WT (Fig. 7D). In addition, mRNA
14 expression of *WUSCHEL* (*WUS*) and *INNER NO OUTER* (*INO*), which are the important genes
15 for ovule formation, and *GOBLET* (*GOB*), a homologue of Arabidopsis *CUP-SHAPED*
16 *COTYLEDONS1* (*CUC1*) and *CUC2*, which are essential for the formation of carpel margin
17 meristems (CMMs) (Hendelman et al. 2013; Kamiuchi et al. 2014), was decreased in the *Sl ses*
18 mutant compared with WT (Fig. 7E, F, M). The expression of *AINTEGUMENTA-like* (*ANT-*
19 *like*), *PIN-FORMED1-like* (*PINI-like*), *BELL1-like* (*BEL1-like*), *REVOLUTA* (*SIREV*),

1 *CORONA-like (CNA-like)* and *PHABULOSA/PHAVOLUTA-like (PHB/PHV-like)*, which are
2 homologues of Arabidopsis and are required for ovule formation, were downregulated in the
3 pistil from 4.5 mm and buds in the *Sl ses* mutant (Fig. 7G-L).

4

5 **Discussion:**

6 We identified a 13-bp deletion in the *SISES* CDS in the *Sl ses* mutant, and the deletion
7 caused a frame shift, resulting in the loss of the last 296 amino acids. The plant transcription
8 factor NOZZLE motif and EAR motif are completely lost in the *Sl ses* mutant (Fig. 5B, C). In
9 contrast, complementation experiments demonstrated that introduction of the genomic region
10 of *Solyc07g063670* with its native promoter and terminator into the *Sl ses* mutant fully restored
11 these phenotypes (Supplementary Fig. S5). These results indicate that *SISES (Solyc07g063670)*
12 is the gene responsible for the *Sl ses* mutant, and the *SISES* loss-of-function mutation had a
13 drastic effect on the development of reproductive organs.

14 The *SISES* protein sequence exhibits significant homology with the *SPL/NZZ*
15 (Schiefthaler et al. 1999; Yang *et al.* 1999) and *TIE* proteins (Tao et al. 2013) of Arabidopsis
16 (Fig. 5D). These proteins carry an EAR motif and act as transcriptional repressors due to
17 interaction with *TOPLESS* through the EAR motif. *TOPLESS* can repress transcription in vivo
18 (Long et al. 2006; Pan et al. 2010; Causier et al. 2012; Tao et al. 2013; Wei et al. 2015). The
19 putative amino acid sequence of *SISES* exhibited a highly conserved EAR motif at the C-

1 terminus (Fig. 5E), suggesting that the tomato *SISES* protein may interact with TOPLESS and
2 act as transcriptional repressor.

3 The Arabidopsis homologous gene *SPL/NZZ* is one of the key genes involved in anther
4 development, especially in the regulation of sporocyte formation in Arabidopsis (Yang et al.
5 1999; Plackett et al. 2011). En, ML, T, PCM and MC development were observed at the 1 to 3
6 mm bud stages in WT but not in the *Slses* mutant (Fig. 2E, F, I, J). These tissues differentiate
7 from the L2 layer of initial stages of anther development, indicating that differentiation from
8 the L2 layer was aborted in the *Slses* mutant. As a result, the *Slses* mutant exhibited wilted
9 anthers and produced no mature pollen grains (Fig. 2C, D). The phenotypes of the *Slses* mutant
10 were similar to those of the Arabidopsis *spl/nzz* mutant (Yang et al. 1999), indicating that *SISES*
11 is essential for anther primordia formation in tomato. We detected minimal mRNA expression
12 of *MS10³⁵* and *EMS1/EXS-like*, which are homologues of Arabidopsis *DYT1* and *EMS1/EXS*,
13 respectively, in the 1 mm bud of the *Slses* mutant (Fig. 7A, C). Given that both the tomato
14 *ms10³⁵* mutant and the Arabidopsis *dyl1* mutant exhibit normal PMC development and
15 abnormal tapetum development during pollen production in the anther (Zhang et al. 2006; Jeong
16 et al. 2014) and that *DYT1* acts downstream of *SPL/NZZ* in Arabidopsis (Zhang et al. 2006), it
17 is suggested that *SISES* also acts upstream of *MS10³⁵*. In addition, *MS10³⁵* mRNA expression
18 might be required for normal PMC development in tomato. Additionally, *EMS1/EXS* acts
19 upstream of *DYT1* and downstream of *SPL/NZZ*. The *ems1/exs* mutant produces anthers lacking

1 the tapetum but with extra PMCs in Arabidopsis (Zhao et al. 2002). This finding may suggest
2 that *EMS1/EXS-like* (*Solyc09g098420*) acts downstream of *SISES* and has similar functions as
3 its homologue in Arabidopsis.

4 The locule number and ovary size were significantly reduced in the *SlSES* mutant
5 compared with the WT (Fig. 1I, 4E), indicating that *SISES* is partly involved in determining
6 carpel number and ovary development. Although *SPL/NZZ* is essential for ovule primordia and
7 nucellus development, integuments develop normally in Arabidopsis (Schiefthaler et al. 1999;
8 Sieber et al. 2004a; Bencivenga et al. 2011). Interestingly, the *SlSES* mutant formed a pistil
9 (carpel) with ovules whose development arrested at the 2 to 3 mm bud stages, which was
10 associated with a failure in nucellus (MMC) and integument development (Fig. 3I, J), indicating
11 that *SISES* functions in nucellus development, similar to Arabidopsis and integument
12 development in tomato. *INO* plays a key role in the formation of the outer integuments in
13 Arabidopsis (Baker et al. 1997; Villanueva et al. 1999). *INO* expression is not affected in the
14 *spl/nzz* mutant compared with WT in Arabidopsis; therefore, the *spl/nzz* mutant forms
15 integuments normally (Balasubramanian and Schneitz 2002; Sieber et al. 2004a). In tomato, the
16 *SlSES* mutant did not exhibit any integument development (Fig. 3O), and *SlINO* expression in
17 the mutant is minimally detectable in the pistils of the 4.5 mm bud and anthesis stages (Fig.
18 7D), suggesting that the molecular regulation between *SPL/NZZ* and *INO* in Arabidopsis and
19 *SISES* and *SlINO* in tomato differs. *SISES* function is necessary for *SlINO* expression in tomato.

1 Arabidopsis forms ovules from two integuments (inner and outer integuments and bitegmic
2 ovules), whereas tomato forms ovules from a single integument (unitegmic ovules) (Wang and
3 Ren 2008; Skinner et al. 2016). Our results indicate that the characteristics of the single
4 integument in tomato are similar to that of the outer integument in Arabidopsis.

5 It has been demonstrated that exclusive and timely *WUS* expression in the nucellus is
6 critical for normal ovule development in Arabidopsis (Groß-Hardt et al. 2002). In the *Sl ses*
7 mutant, *SlWUS* mRNA expression was considerably downregulated compared with WT (Fig.
8 7E). In Arabidopsis, *WUS* expression in the nucellus was reduced in the *spl/nzz* mutants same
9 as our study (Sieber et al. 2004a), and this information suggests that *SlSES* expression is
10 required to induce *SlWUS* mRNA transcription in tomato. *ANT-like* and *BELI-like* mRNA
11 expression was downregulated in the pistils of the 4.5 mm bud in the *Sl ses* mutant compared
12 with the WT (Fig. 7G, I). These genes also play key roles in the development of ovules and
13 integuments in Arabidopsis (Elliott et al. 1996; Schneitz et al. 1997; Brambilla et al. 2007).
14 *ANT* is expressed within the developing integuments and funiculus. *BELI* is strongly expressed
15 within the developing integuments, and the *spl/nzz* mutant exhibits ectopic expression of *ANT*
16 and *BELI* within the ovule primordia in Arabidopsis (Balasubramanian and Schneitz 2000;
17 Bencivenga et al. 2012). *ANT* and *BELI* expression are required for the ovule integument
18 development in Arabidopsis (Schneitz et al. 1997). Ovule development, especially the ovule
19 integument (Fig. 3O), in the *Sl ses* mutant was arrested, suggesting reduced expression of *ANT*-

1 *like* and *BELI-like* genes, which was actually in accordance with our results. *PIN1* is an auxin
2 efflux facilitator, and the *pin1-5* mutant exhibits abnormal ovule development. Therefore,
3 *SPL/NZZ* expression in the ovule is required to induce *PIN1* transcription in Arabidopsis
4 (Bencivenga et al. 2012). Moreover, the class III homeodomain leucine zipper (HD-ZIPIII)
5 genes *CNA*, *PHB*, *PHV* and *REV* are expressed within the chalaza and restrict *WUS* and *INO*
6 expression to the nucellus and gynobasal side of the ovule in Arabidopsis (Kelley et al. 2009ab;
7 Yamada et al. 2016). *PIN1*, *CNA*, *PHB*, *PHV* and *REV* are required for precise spatiotemporal
8 expression for correct ovule development. In tomato, *PIN1-like*, *CNA-like*, *PHB/PHV-like* and
9 *SIREV* expression was reduced in the pistils of the 4.5 mm bud and/or anthesis flower, but the
10 reduction was comparatively small. Nevertheless, ovule development was arrested in the *Slses*
11 mutant (Fig. 7H, J-L). Given that the precise spatial expression of these genes is essential to
12 successfully form ovules, the control of the spatial expression and expression level of these
13 genes were collapsed in the *Slses* mutant.

14 *SPL/NZZ* expression is positively regulated by *AG* in Arabidopsis (Ito et al. 2004). On
15 the other hand, the highest mRNA expression point of *SISES* and *TAG1* was in the 1 mm bud
16 and the pistils of 4.5 mm bud, respectively (Fig. 6, 7D). This may represent that *TAG1* doesn't
17 regulate *SISES* expression or they need other factor to regulate the mRNA expression of *SISES*.
18 In any case, we could not judge there is such a relationship between *SISES* and *TAG1* in our
19 study. In order to clarify, we need to do further experiments.

1 In this study, we identified a key gene, *SISES*, that is responsible for both male and
2 female gametogenesis in tomato. *SISES* is a homologue of *SPL/NZZ* in Arabidopsis, and the
3 loss-of-function *SISES* mutant exhibits sexual sterility caused by a failure to form pollen and
4 ovules. Possible roles of *SISES* in ovule and anther development are proposed in Fig. 8 and are
5 compared with *SPL/NZZ* of Arabidopsis. In ovule development, *SPL/NZZ* positively regulates
6 *PIN1* and *WUS* expression and plays a key role in nucellus development in Arabidopsis (Sieber
7 et al. 2004a; Bencivenga et al. 2012). On the other hand, *SISES* is required to express *SIWUS*
8 and *SIINO*. Moreover, we demonstrate that *SISES* has a role in nucellus and integument
9 development via controlling *SIINO* expression in tomato (Fig. 8). In addition, in anther
10 development, *SPL/NZZ* positively regulates *BAMI/2* and *EMS1/EXS* expression and is essential
11 for archesporial cell development in Arabidopsis (Yang et al. 1999; Zhao et al, 2002; Hord et
12 al. 2006; Ma et al. 2012). *SISES* positively regulates *EMS1/EXS-like* expression and plays a key
13 role in archesporial cell development in tomato (Fig. 8). Isolation of the *SISES* gene may
14 positively impact gametogenesis research in tomato.

15

16 **Materials and Methods:**

17 *Plant materials and growth conditions*

18 The miniature tomato ‘Micro-Tom’ and the cultivated species ‘Ailsa Craig’ were used
19 in this study. The *Slses* mutant was obtained from gamma ray-irradiated Micro-Tom mutant

1 populations produced by the National BioResource Project (NBRP, <http://www.nbrp.jp/>) (Saito
2 et al. 2011). All lines were grown under a regimen of 16-h days and 8-h nights at a constant
3 temperature of 25°C under fluorescent lights at 100 to 150 $\mu\text{Em}^{-2}\text{s}^{-1}$.

4

5 *Histological and microscopic analysis*

6 Whole anthesis flowers and flower buds were collected and fixed in FAA (formalin :
7 acetic acid : 70% EtOH = 1 : 1 : 18), vacuum-infiltrated twice for 10 min each, and left overnight.
8 The fixed tissues were dehydrated with a *t*-butyl alcohol series from No. 1 to 6, which contained
9 *t*-butyl alcohol, ethanol and DW (Supplementary Table S4). Each step required a greater than
10 6-hour incubation at room temperature, and the No. 6 solution was used twice. The tissues in
11 the No. 6 solution were incubated at 60°C for 2 hours. Then, melted paraffin was added, and
12 the tissues were stored at 60°C overnight. The embedded tissues were sliced into 10- μm
13 sections using a microtome (MICROM HM325, Leica, <http://www2.leicabiosystems.com/>).
14 The sections were stained with 0.05% toluidine blue (pH 7.0). Paraffin was removed with
15 xylene, and the sections were mounted with entellan new (Merck Millipore). Then, sections
16 were observed under an optical microscope (Olympus BX53, [http://www.olympus-
18 lifescience.com/](http://www.olympus-
17 lifescience.com/)).

19 *Map-based cloning using DNA markers*

1 To conduct SolCAP analysis (<http://solcap.msu.edu/index.shtml>) for rough mapping,
2 an F₂ mapping population was constructed from a cross between *S. lycopersicum* ‘Ailsa Craig’
3 and the *Slses* heterozygous mutant in the ‘Micro-Tom’ background. Then, we used 28 F₂ plants
4 that exhibited the *Slses* mutation for this analysis. In total, 7,720 markers were used for this
5 experiment, and 2,098 markers showed polymorphism to distinguish between ‘Micro-Tom’ and
6 ‘Ailsa Craig’. Then, the region of the responsible gene was narrowed down.

7

8 *Genome re-sequencing and identification of responsible gene*

9 The genome sequence of the *Slses* mutant was obtained by Illumina HiSeq 2000. The
10 sequences of ‘Heinz 1706’ and ‘Micro-Tom’ Japan were used as reference genomes (Ariizumi
11 et al. 2014; Kobayashi et al. 2014). Linkage analysis was conducted with F₂ plants which
12 obtained crossed with the *ses* mutant and ‘Micro-Tom’, WT or ‘Ailsa Craig’. The marker (*SISES*
13 marker-1, Supplementary Table S2) was designed to amplify the 13-bp deletion region. The
14 amplicon sizes were different in the *ses* mutant and WT.

15

16 *Characterization of the identified gene*

17 The genome sequences and amino acid sequences were obtained from the Sol
18 Genomics Network (SGN) (<https://solgenomics.net/>). The estimation of the SISES protein
19 domains was conducted using the Conserved Domain Database (CDD)

1 (<http://www.ncbi.nlm.nih.gov/cdd>). The homologous genes from Arabidopsis and Tomato were
2 searched using The Arabidopsis Information Resource (TAIR) (<https://www.arabidopsis.org/>)
3 and SGN, respectively. We selected the homologous gene of which e-value is under '0.1'. The
4 phylogenetic tree was constructed with multiple sequence alignment using CLUSTALW
5 (<http://www.genome.jp/tools/clustalw/>). The amino acid homology of SISES and SPL/NZZ was
6 analyzed using the EMBOSS Needle (http://www.ebi.ac.uk/Tools/psa/emboss_needle/) in the
7 European Bioinformatics Institute (<http://www.ebi.ac.uk/>).

8

9 *Creation of transgenic plants*

10 The binary vector pIG121Hm was used for the creation of transgenic plants. The
11 complete *SISES* gene, which included 2563 bp of upstream sequence and 763 bp of downstream
12 sequence, was amplified with gene-specific primers (whole *SISES* gene primer, Supplementary
13 table S2) using the In-Fusion technique. The amplified fragments of the complete genomic
14 region were inserted into pIG121Hm digested by *HindIII* and *SalI* using the In-Fusion
15 technique (TAKARA) (Supplementary Fig. S3). Then, the construct was transformed into
16 *Agrobacterium tumefaciens* GV2260 by electroporation, and transformants were selected on
17 LB plates containing kanamycin (50 mg L⁻¹). The transformation construct was transformed
18 into 'Micro-Tom' WT and plants heterozygous or homozygous for the *Slses* mutation using an
19 *Agrobacterium*-mediated transformation method (Sun et al. 2006). The transgenic plants were

1 selected on MS plates containing kanamycin (100 mg L⁻¹). Ploidy and transgene copy number
2 transgene were assessed by flow cytometry and Southern blotting analysis, respectively, in the
3 T0 generation. Plants that were both diploid and single-copy transgenic were selected, and their
4 T1 or T2 generations that harbored the *Slses* homozygous mutation were used for further
5 analysis. The *Slses* homozygous mutants were selected using “*SISES* marker-2” and the
6 restriction enzyme *Bsl* I, we could distinguish among homozygous, heterozygous and WT by
7 electrophoresis band pattern (Supplementary Table S2; Supplementary Fig. S4). The presence
8 of transgene was confirmed by detecting “*NPT* II marker” using PCR (Supplementary Table
9 S2).

10

11 *Gene expression analysis*

12 mRNA expression levels were analyzed by quantitative RT-PCR (qRT-PCR) using
13 SYBR Premix Ex Taq II (TAKARA). The template cDNAs were synthesized from total RNA
14 extracted from 1 mm whole buds or pistils of unopened buds classified by bud length (3 mm,
15 4.5 mm, 6 mm, 1 day before anthesis) and anthesis flowers. Total RNA was extracted with an
16 RNeasy Mini Kit (Qiagen, <http://www.qiagen.com/>), and cDNA was synthesized with a
17 SuperScript VILO cDNA Synthesis Kit (Invitrogen, <http://www.lifetechnologies.com/>). SGN-
18 U316474 (*SAND*) mRNA was used as an internal control (Expósito-Rodríguez et al. 2008). The
19 expression level was calculated according to Pfaffl 2001. The primers were designed with

1 [Primer3Plus] (<http://primer3plus.com/cgi-bin/dev/primer3plus.cgi>) (Supplementary Table S5).

2

3 *Statistical analysis*

4 We used Student's *t* test ($P = 0.05$ and 0.01) to show the significance between two
5 values, WT and the *Slses* mutant. Then, Chi-squared (χ^2) test was performed to examine the
6 correctness of fit between the expected Mendelian ratios and segregation data for the *Slses*
7 mutation. We calculated χ^2 value on below conditions; degrees of freedom, 1; the expected
8 Mendelian ratios, WT : *Slses* = 3 : 1.

9

10 **Funding**

11 This work was supported by Science and Technology Research Promotion Program for
12 Agriculture, Forestry, Fisheries and Food Industry, Japan (grant no. 26013A) to HE. Seeds of
13 Micro-Tom (TOMJPF00001), Ailsa Craig (TOMJPF00004), and *Slses* were obtained from the
14 National BioResource Project, Ministry of Education, Culture, Sports, Science and Technology
15 (MEXT), Japan.

16

17 **Disclosures:**

18 The authors have no conflicts of interest to declare.

19

1 **Acknowledgements:**

2 We thank all of the members of our laboratory for helpful discussions throughout the work.

3

4 **References:**

5 Ariizumi, T., Kishimoto, S., Kakami, R., Maoka, T., Hirakawa, H., Suzuki, Y. et al. (2014)

6 Identification of the carotenoid modifying gene *PALE YELLOW PETAL 1* as an essential factor
7 in xanthophyll esterification and yellow flower pigmentation in tomato (*Solanum lycopersicum*).

8 *Plant J.* 79: 453-465.

9

10 Baker, S.C., Robinson-Beers, K., Villanueva, J.M., Gaiser, C. and Gasser, C.S. (1997)

11 Interactions among genes regulating ovule development in *Arabidopsis thaliana*. *Genetics* 145:
12 1109-1124.

13

14 Balasubramanian, S. and Schneitz, K. (2000) *NOZZLE* regulates proximal-distal pattern
15 formation, cell proliferation and early sporogenesis during ovule development in *Arabidopsis*
16 *thaliana*. *Development* 127: 4227-4238.

17

18 Balasubramanian, S. and Schneitz, K. (2002) *NOZZLE* links proximal-distal and adaxial-
19 abaxial pattern formation during ovule development in *Arabidopsis thaliana*. *Development* 129:

1 4291-4300.

2

3 Bencivenga, S., Colombo, L. and Masiero, S. (2011) Cross talk between the sporophyte and the
4 megagametophyte during ovule development. *Sex Plant Reprod.* 24: 113-121.

5

6 Bencivenga, S., Simonini, S., Benková, E. and Colombo, L. (2012) The Transcription Factors
7 BEL1 and SPL Are Required for Cytokinin and Auxin Signaling During Ovule Development
8 in *Arabidopsis*. *Plant Cell* 24: 2886-2897.

9

10 Brambilla, V., Battaglia, R., Colombo, M., Masiero, S., Bencivenga, S., Kater, M.M. et al.
11 (2007) Genetic and molecular interactions between BELL1 and MADS box factors support
12 ovule development in *Arabidopsis*. *Plant Cell* 19: 2544-2556.

13

14 Canales, C., Bhatt, A.M., Scott, R. and Dickinson, H. (2002) *EXS*, a Putative LRR Receptor
15 Kinase, Regulates Male Germline Cell Number and Tapetal Identity and Promotes Seed
16 Development in *Arabidopsis*. *Curr. Biol.* 12: 1718-1727.

17

18 Causier, B., Ashworth, M., Guo, W. and Davies, B. (2012) The TOPLESS Interactome: A
19 Framework for Gene Repression in *Arabidopsis*. *Plant Physiol.* 158: 423-438.

1

2 Chen, G.H., Sun, J.Y., Liu, M., Liu, J. and Yang, W.C. (2014) SPOROCTELESS Is a Novel
3 Embryophyte-Specific Transcription Repressor that Interacts with TPL and TCP Proteins in
4 *Arabidopsis*. *J. Genet. Genomics* 41: 617-625.

5

6 Elliott, R.C., Betzner, A.S., Huttner, E., Oakes, M.P., Tucker, W.Q., Gerentes, D. et al. (1996)
7 *AINTEGUMENTA*, an *APETALA2*-like gene of *Arabidopsis* with pleiotropic roles in ovule
8 development and floral organ growth. *Plant Cell* 8: 155-168.

9

10 Expósito-Rodríguez, M., Borges, A.A., Borges-Pérez, A. and Pérez, J.A. (2008) Selection of
11 internal control genes for quantitative real-time RT-PCR studies during tomato development
12 process. *BMC Plant Biol.* 8: 131.

13

14 Groß-Hardt, R., Lenhard, M. and Laux, T. (2002) *WUSCHEL* signaling functions in
15 interregional communication during *Arabidopsis* ovule development. *Genes Dev.* 16: 1129-
16 1138.

17

18 Hendelman, A., Stav, R., Zemach, H. and Arazi, T. (2013) The tomato NAC transcription factor
19 *SINAM2* is involved in flower-boundary morphogenesis. *J. Exp. Bot.* 64: 5497-5507.

1

2 Hord, C.L.H., Chen, C., DeYoung, B.J., Clark, S.E. and Ma, H. (2006) The BAM1/BAM2
3 receptor-like kinases are important regulators of *Arabidopsis* early anther development. *Plant*
4 *Cell* 18: 1667-1680.

5

6 Ito, T., Wellmer, F., Yu, H., Das, P., Ito, N., Alves-Ferreira, M. et al. (2004) The homeotic
7 protein AGAMOUS controls microsporogenesis by regulation of *SPOROCTELESS*. *Nature*
8 430: 356-360.

9

10 Jeong, H.J., Kang, J.H., Zhao, M., Kwon, J.K., Choi, H.S., Bae, J.H. et al. (2014) Tomato *Male*
11 *sterile 10³⁵* is essential for pollen development and meiosis in anthers. *J. Exp. Bot.* 65: 6693-
12 6709.

13

14 Kagale, S. and Rozwadowski, K. (2011) EAR motif-mediated transcriptional repression in
15 plants. *Epigenetics* 6: 141-146.

16

17 Kamiuchi, Y., Yamamoto, K., Furutani, M., Tasaka, M. and Aida, M. (2014) The *CUC1* and
18 *CUC2* genes promote carpel margin meristem formation during *Arabidopsis* gynoecium
19 development. *Front. Plant Sci.* 5: 1-9.

1

2 Kelley, D.R. and Gasser, C.S. (2009a) Ovule development: genetic trends and evolutionary
3 considerations. *Sex Plant Reprod.* 22: 229-234.

4

5 Kelley, D.R., Skinner, D.J. and Gasser, C.S. (2009b) Roles of polarity determinants in ovule
6 development. *Plant J.* 57: 1054-1064.

7

8 Kobayashi, M., Nagasaki, H., Garcia, V., Just, D., Bres, C., Mauxion, L.P. et al. (2014) Genome-
9 wide analysis of intraspecific DNA polymorphism in 'Micro-Tom', a model cultivar of Tomato
10 (*Solanum lycopersicum*). *Plant Cell Physiol.* 55: 445-454.

11

12 Long, J.A., Ohno, C., Smith, Z.R. and Meyerowitz, E.M. (2006) TOPLESS regulates apical
13 embryonic fate in *Arabidopsis*. *Science* 312: 1520-1523.

14

15 Ma, X., Feng, B. and Ma, H. (2012) AMS-dependent and independent regulation of anther transcriptome
16 and comparison with those affected by other *Arabidopsis* anther genes. *BMC Plant Biol.* 12: 23.

17

18 Mazzucato, A., Taddei, A.R. and Soressi, G.P. (1998) The *parthenocarpic fruit (pat)* mutant of
19 tomato (*Lycopersicon esculentum* Mill.) sets seedless fruits and has aberrant anther and ovule

1 development. *Development* 125: 107-114.

2

3 Minoia, S., Petrozza, A., D'Onofrio, P., Piron, F., Mosca, G., Sozio, G. et al. (2010) A new
4 mutant genetic resource for tomato crop improvement by TILLING technology. *BMC Res.*
5 *Notes* 3: 69

6

7 Pan, I.C., Li, C.W., Su, R.C., Cheng, C.P., Lin, C.S. and Chan, M.T. (2010) Ectopic expression
8 of an EAR motif deletion mutant of *SIERF3* enhances tolerance to salt stress and *Ralstonia*
9 *solanacearum* in tomato. *Planta* 232: 1075-1086.

10

11 Pfaffl, M.W. (2001) A new mathematical model for relative quantification in real-time RT-PCR.
12 *Nucleic Acids Res.* 29: e45.

13

14 Pinyopich, A., Ditta, G.S., Savidge, B., Liljegren, S.J., Baumann, E., Wisman, E. and Yanofsky,
15 M.F. (2003) Assessing the redundancy of MADS-box genes during carpel and ovule
16 development. *Nature* 424: 85-88.

17

18 Plackett, A.R.G., Thomas, S.G., Willson, Z.A. and Hedden, P. (2011) Gibberellin control of
19 stamen development: a fertile field. *Trends Plant Sci.* 16: 568-578.

1

2 Saito, T., Ariizumi, T., Okabe, Y., Asamizu, E., Hiwasa-Tanase, K., Fukuda, N. et al. (2011)

3 TOMATOMA: A Novel Tomato Mutant Database Distributing Micro-Tom Mutant Collections.

4 *Plant Cell Physiol.* 52: 283–296.

5

6 Sanders, P.M., Bui, A.Q., Weterings, K., McIntire, K.N., Hsu, Y.C., Lee, P.Y. et al. (1999)

7 Anther developmental defects in *Arabidopsis thaliana* male-sterile mutants. *Sex Plant Reprod.*

8 11: 297-322.

9

10 Schiefthaler, U., Balasubramanian, S., Sieber, P., Chevalier, D., Wisman, E. and Schneitz, K.

11 (1999) Molecular analysis of *NOZZLE*, a gene involved in pattern formation and early

12 sporogenesis during sex organ development in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA*

13 96: 11664-11669.

14

15 Schneitz, K., Hülskamp, M., Kopczak, S.D. and Pruitt, R.E. (1997) Dissection of sexual organ

16 ontogenesis: a genetic analysis of ovule development in *Arabidopsis thaliana*. *Development*

17 124: 1367-1376.

18

19 Scott, J.W. and Harbaugh, B.K. (1989) Micro-Tom - a miniature dwarf tomato. *Agricultural*

1 Experiment Station, Institute of Food and Agricultural Sciences. *University of Florida Circular*
2 S370: 1-6.

3

4 Shirasawa, K., Isobe, S., Hirakawa, H., Asamizu, E., Fukuoka, H., Just, D. et al. (2010) SNP
5 discovery and linkage map construction in cultivated tomato. *DNA Res.* 17: 381-391.

6

7 Sicard, A., Petit, J., Mouras, A., Chevalier, C. and Hernould, M. (2008a) Meristem activity
8 during flower and ovule development in tomato is controlled by the mini zinc finger gene
9 *INHIBITOR OF MERISTEM ACTIVITY*. *Plant J.* 55: 415-427.

10

11 Sicard, A., Hernould, M. and Chevalier, C. (2008b) The INHIBITOR OF MERISTEM
12 ACTIVITY (IMA) protein. *Plant Signal. Behav.* 3: 908-910.

13

14 Sieber, P., Gheyselinck, J., Gross-Hardt, R., Laux, T., Grossniklaus, U. and Schneitz, K. (2004a)
15 Pattern formation during early ovule development in *Arabidopsis thaliana*. *Dev. Biol.* 273: 321-
16 334.

17

18 Sieber, P., Petrascheck, M., Barberis, A. and Schneitz, K. (2004b) Organ Polarity in Arabidopsis.
19 NOZZLE Physically Interacts with Members of the YABBY Family. *Plant Physiol.* 135: 2172-

1 2185.

2

3 Sim, S.C., Durstewitz, G., Plieske, J., Wieseke, R., Ganal, M.W, Van Deynze, A. et al. (2012)
4 Development of a large SNP genotyping array and generation of high-density genetic maps in
5 tomato. *PLoS One* 7(7): e40563.

6

7 Skinner, D.J., Hill, T.A. and Gasser, C.S. (2004) Regulation of Ovule Development. *Plant Cell*
8 16: 32-46.

9

10 Skinner, D.L., Brown, R.H., Kuzoff, R.K. and Gasser, C.S. (2016) Conservation of the role of
11 *INNER NO OUTER* in development of unitegmic ovules of the Solanaceae despite a divergence
12 in protein function. *BMC Plant Biol.* 16: 143.

13

14 Sorensen, A.M., Kröber, S., Unte, U.S., Huijser, P., Dekker, K. and Saedler, H. (2003) The
15 *Arabidopsis* *ABORTED MICROSPORES (AMS)* gene encodes a MYC class transcription factor.
16 *Plant J.* 33: 413-413.

17

18 Sun, H.J., Uchii, S., Watanabe, S. and Ezura, H. (2006) A highly efficient transformation
19 protocol for Micro-Tom, a model cultivar for tomato functional genomics. *Plant Cell Physiol.*

1 47: 426–431.

2

3 Tao, Q., Guo, D., Wei, B., Zhang, F., Pang, C., Jiang, H. et al. (2013) The TIE1 transcriptional
4 repressor links TCP transcription factors with TOPLESS/TOPLESS-RELATED corepressors
5 and modulates leaf development in *Arabidopsis*. *Plant Cell* 25: 421-437.

6

7 The Tomato Genome Consortium. (2012) The tomato genome sequence provides insights into
8 fleshy fruit evolution. *Nature* 485: 635-641.

9

10 Villanueva, J.M., Broadhvest, J., Hauser, B.A., Meister, R.J. Schneitz, K. and Gasser, C.S.
11 (1999) *INNER NO OUTER* regulates abaxial - adaxial patterning in *Arabidopsis* ovules. *Genes*
12 *Dev.* 13: 3160-3169.

13

14 Wang, H., Jones, B., Li, Z., Frasse, P., Delalande, C., Regad, F. et al. (2005) The tomato *Aux/IAA*
15 transcription factor *IAA9* is involved in fruit development and leaf morphogenesis. *Plant Cell*
16 17: 2676-2692.

17

18 Wang, Z.F. and Ren, Y. (2008) Ovule morphogenesis in ranunculaceae and its systematic
19 significance. *Ann. Bot.* 101: 447-462.

1

2 Wei, B., Zhang, J., Pang, C., Yu, H., Guo, D., Jiang, H. et al. (2015) The molecular mechanism
3 of SPOROCTELESS/NOZZLE in controlling *Arabidopsis* ovule development. *Cell Res.* 25:
4 121-134.

5

6 Wilson, Z.A., Morroll, S.M., Dawson, J., Swarup, R. and Tighe, P.J. (2001) The *Arabidopsis*
7 MALE STERILITY1 (*MSI*) gene is a transcriptional regulator of male gametogenesis, with
8 homology to the PHD-finger family of transcription factors. *Plant J.* 28: 27-39.

9

10 Wilson, Z.A. and Zhang, D.B. (2009) From *Arabidopsis* to rice: Pathways in pollen
11 development. *J. Exp. Bot.* 60: 1479-1492.

12

13 Yamada, T., Sasaki, Y., Hashimoto, K., Nakajima, K. and Gasser, C.S. (2016) *CORONA*,
14 *PHABULOSA* and *PHAVOLUTA* collaborate with *BELL 1* to confine *WUSCHEL* expression to
15 the nucellus in *Arabidopsis* ovules. *Development* 143: 422-426.

16

17 Yang, S.L., Xie, L.F., Mao, H.Z., Pua, C.S., Yang, W.C., Jiang, L. et al. (2003) *TAPETUM*
18 *DETERMINANT1* Is Required for Cell Specialization in the *Arabidopsis* Anther. *Plant Cell* 15:
19 2792-2804.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

Yang, W.C., Ye, D., Xu, J. and Sundaresan, V. (1999) The *SPOROCTELESS* gene of *Arabidopsis* is required for initiation of sporogenesis and encodes a novel nuclear protein. *Genes Dev.* 13: 2108-2117.

Ye, Q., Zhu, W., Li, L., Zhang, S., Yin, Y. and Ma, H. (2010) Brassinosteroids control male fertility by regulating the expression of key genes involved in *Arabidopsis* anther and pollen development. *Proc. Natl. Acad. Sci. USA* 107: 6100-6105.

Zhang, W., Sun, Y., Timofejeva, L., Chen, C., Grossniklaus, U. and Ma, H. (2006) Regulation of *Arabidopsis* tapetum development and function by *DYSFUNCTIONAL TAPETUM1 (DYT1)* encoding a putative bHLH transcription factor. *Development* 133: 3085-3095.

Zhao, D.Z., Wang, G.F., Speal, B. and Ma, H. (2002) The *EXCESS MICROSPOROCTES1* gene encodes a putative leucine-rich repeat receptor protein kinase that controls somatic and reproductive cell fates in the *Arabidopsis* anther. *Genes Dev.* 16: 2021-2031.

1

2

3

4

5 **Tables:**

6

Table 1. Comparison of the numbers of flower organs between the *Sl ses* mutant and WT.

	Flower organ			
	Sepal	Petal	Anther	Pistil
WT	5.20 ± 0.09	5.05 ± 0.09	5.15 ± 0.08	1.00 ± 0.00
<i>Sl ses</i>	5.30 ± 0.11	5.25 ± 0.10	5.30 ± 0.11	1.00 ± 0.00

Values are means ± standard error. Statistical significance was analyzed using the Student's *t* test.

7

Table 2. Comparison of the phenotypes of F₁ and F₂ plants obtained by crossing the *Sl ses* mutant (heterozygous) and WT plants.

Cross	Generation	Phenotype		χ^2 value	P value
		WT	<i>Sl ses</i>		
<i>Sl ses</i> (Heterozygous) × WT	F ₁	17	0	-	-
	F ₂ (A)	118	40	0.008	0.927
	F ₂ (B)	149	0	-	-

Two patterns of F₂ population were observed: segregated (A) and non-segregated (B).

See supplementary Fig. S1.

The χ^2 value was calculated from the F₂ populations (A)

8

9

10

1 **Figure legends**

2 Figure 1.

3 The floral morphology of the *SlSES* mutant at the anthesis stage. (A, B) The flower, (C, D) the
4 anther and pollen grains, (E-H) the pistil. (A, C, E, G) WT, (B, D, F, H) *SlSES* mutant. Bar = 1
5 mm. (I) The diameter and length of the ovary. Statistical analysis was performed using the
6 Student's *t*-test. ** $P < 0.01$; * $P < 0.05$, $n = 10$, error bar = SE.

7

8 Figure 2.

9 Cross-sections of developing anthers in WT and the *SlSES* mutant. (A, C) Cross sections of
10 anthers at the anthesis stage, (B, D) enlarged view of a piece of anther at the anthesis stage, (E,
11 I) 1 to 2 mm bud, (F, J) 2 to 3 mm bud, (G, K) 4 to 5 mm bud, (H, L) 6 to 7 mm bud. (A, B, E-
12 H) WT, (C, D, I-L) the *SlSES* mutant. En, endothecium; Ep, epidermis; MC, meiotic cell; ML,
13 middle layer; Msp, microspore; P, pollen; PMC, pollen mother cell; S, style; T, tapetum; VB,
14 vascular bundle. Bar = 500 μm .

15

16 Figure 3.

17 Sections of developing pistils in WT and the *SlSES* mutant. (A, C) Cross-section, (B, D)
18 longitudinal section at the anthesis stage. (E, I) 1 to 2 mm bud, (F, J) 2 to 3 mm bud, (G, K) 4
19 to 5 mm bud, (H, L) 6 to 7 mm bud. Bar = 500 μm . (M) Ovule primordia of 2 to 3 mm bud, (N,

1 O) ovule primordia of 4 to 5 mm bud. Bar = 50 μ m. (A, B, E-H, M, N) WT, (C, D, I-L, O) the
2 *Slses* mutant. MMC, megaspore mother cell; O, ovule; OI, ovule integument; OP, ovule
3 primordia; P, placenta.

4

5 Figure 4.

6 Morphology of ripe red fruits in the WT and the *Slses* mutant. (A) Appearance of red ripe fruits,
7 (B) cross-section of the red fruits. Bar = 1 cm. (C) Fruit weight, (D) fruit size and (E) the number
8 of locules per fruit in the WT and the *Slses* mutant. n = 20, error bar = SE, statistical analysis
9 was performed using the Student's *t*-test, **P<0.01.

10

11 Figure 5.

12 Gene structure of *Solyc07g063670*. (A) The location and gene structure of *Solyc07g063670*.
13 The blue bar indicates chromosome 7, the gray box indicates the UTR, white boxes indicate
14 exons, and regions between the boxes indicate introns. (B) Genome and amino acid sequences
15 of WT and the *Slses* mutant at the mutation point. The upper black letters indicate genome
16 sequences, and the lower red letters indicate putative amino acids. Hyphens indicate the 13-bp
17 deletion. (C) The putative amino acid length and domain of the SISES protein. The putative
18 amino acid length and domain region were exhibited under and upper of SISES protein models
19 respectively. The domain is indicated with yellow and red boxes. * = stop codon. (D)

1 Phylogenetic tree of *SISES* and its homologous proteins in Arabidopsis. (E) The EAR motif of
2 *SISES* and *SPL/NZZ* at the C-terminal region. The red frame indicates the EAR motif.

3

4 Figure 6.

5 Expression analysis of *SISES* in the WT by qRT-PCR. *SISES* expression level in the WT in the
6 1 mm bud and in pistils from the 3 mm, 4.5 mm, 6 mm, 1 day before anthesis (DBA) and
7 anthesis stages. n = 3.

8

9 Figure 7.

10 Expression analysis of gametogenesis genes in the WT and the *Sises* mutant by qRT-PCR. (A)
11 *Male Sterile 10³⁵* (*MS10³⁵*), (B) *BARELY ANY MERISTEM1 and 2-like* (*BAM1/2-like*), (C)
12 *EXCESS MICROSPOROCTES1/EXTRA SPOROGENOUS CELLS-like* (*EMS1/EXS-like*), (D)
13 *Tomato AGAMOUS1* (*TAG1*), (E) *WUSCHEL* (*SIWUS*), (F) *INNER NO OUTER* (*SIINO*), (G)
14 *AINTEGUMENTA-like* (*ANT-like*), (H) *PIN-FORMED1-like* (*PIN1-like*), (I) *BELL1-like*
15 (*BEL1-like*), (J) *REVOLUTA* (*SIREV*), (K) *CORONA-like* (*CNA-like*), (L)
16 *PHABULOSA/PHAVOLUTA-like* (*PHB/PHV-like*), (M) *GOBLET* (*GOB*). mRNA expression
17 was analyzed at the 1 mm bud stage and in pistils from the 4.5 mm bud and anthesis stages. n
18 = 3, error bar = SE. Statistical analysis was performed using the Student's *t*-test. **P<0.01;
19 *P<0.05.

1

2 Figure 8.

3 A model for ovule (left) and anther (right) development regulated by *SISES* (*SPL/NZZ*) in
4 tomato and *Arabidopsis*. Solid arrows represent positive regulation, and dotted arrows indicate
5 gene function (red = tomato, blue = *Arabidopsis*). The gray box shows the model of integument
6 development in *Arabidopsis* (left) and tomato (right). I = integument, II = inner integument, OI
7 = outer integument, PMC = pollen mother cells, PPC = primary parietal cells, PSC = primary
8 sporogenous cells, SPC = secondary parietal cells.

9

10




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8