

# **Functional Analysis of MCAs and PIF4 in Plant Cold Signaling**

**January 2020**

**Na Renhu**

# **Functional Analysis of MCAs and PIF4 in Plant Cold Signaling**

**A Dissertation Submitted to  
the Graduate School of Life and Environmental Sciences,  
the University of Tsukuba  
in Partial Fulfillment of the Requirements  
for the Degree of Doctor of Philosophy in Science  
(Doctoral Program in Biological Sciences)**

**Na Renhu**

## TABLE OF CONTENT

Abbreviations .....	1
Chapter 1. General Introduction .....	1
1.1. Cold perception and cold signal molecules .....	1
1.2. Cold signaling pathways.....	1
1.3. Cold acclimation and light crosstalk .....	1
Chapter 2. Ca <sup>2+</sup> -Permeable Mechanosensitive Channels MCA1 and MCA2 Mediate Cold-Induced Cytosolic Ca <sup>2+</sup> Increase and Cold Tolerance in Arabidopsis.....	2
2.1. Introduction .....	2
2.2. Methods .....	6
2.2.1. Plant materials. ....	6
2.2.2. Monitoring of [Ca <sup>2+</sup> ] <sub>cyt</sub> changes following cold shock treatment.....	6
2.2.3. Plant freezing and chilling assay. ....	6
2.2.4. RNA preparation and quantitative RT-PCR.....	7
2.3. Results .....	8
2.3.1. MCA1 and MCA2 are involved in a cold-induced [Ca <sup>2+</sup> ] <sub>cyt</sub> increase. ....	8
2.3.2. Mutations in <i>MCA1</i> and <i>MCA2</i> result in cold sensitivity. ....	9
2.3.3. Down-regulation of cold-inducible genes is governed by a non- <i>CBF/DREB1</i> regulon.....	11
2.4. Discussion.....	12
Chapter 3. Cold-Induced Degradation of PIF4 is Mediated by Ubiquitin E3 Ligase HOS1 with ICE1 as a Scaffold Protein to Regulate Cold Signaling .....	26
3.1. Introduction .....	26
3.2. Method and material.....	26
3.2.1. Plant material and growth condition.....	26
3.2.2. BiFC assay.....	26
3.2.3. Plant freezing assay .....	26

3.2.4.	Quantitative RT-PCR analysis .....	26
3.2.5.	<i>In vitro</i> pull down assay.....	26
3.2.6.	Ubiquitination assay .....	26
3.2.7.	Yeast two hybrid assay .....	26
3.2.8.	Protein extraction and immunoblot analysis .....	26
3.3.	Results .....	26
3.3.1.	ICE1 interacts with PIF4 in nuclei. ....	26
3.3.2.	<i>PIF4</i> negatively regulates cold tolerance .....	26
3.3.3.	PIF4 is involved in <i>CBF</i> -dependent cold signaling.....	26
3.3.4.	PIF4 is degraded by low temperature .....	26
3.3.5.	PIF4 does not directly interact with HOS1.....	26
3.3.6.	PIF4 is ubiquitinated by HOS1, which is mediated by ICE1. ....	26
3.4.	Discussion.....	26
Chapter 4.	General Discussion .....	26
Summary.....		26
Acknowledgements .....		26
References .....		27

## Abbreviations

ABA, abscisic acid

RNA, ribonucleic acid

RNA-seq, RNA-sequencing

E-box, enhancer box

JA, jasmonic acid

GFP, green fluorescent protein

Col-0, Columbia-0

WT, wild type

MS, Murashige and Skoog medium

RT-PCR, real-time polymerase chain reaction

EGTA, ethylene glycol tetraacetic acid

GUS, beta-glucuronidases

ANOVA, analysis of variance

T-DNA, transfer deoxyribonucleic acid

SDS-PAGE, sodium dodecyl sulfate–polyacrylamide gel electrophoresis

CBB, Coomassie brilliant blue

DMSO, dimethyl sulfoxide

*E. coli*, *Escherichia coli*

## **Chapter 2. Ca<sup>2+</sup>-Permeable Mechanosensitive Channels MCA1 and MCA2 Mediate Cold-Induced Cytosolic Ca<sup>2+</sup> Increase and Cold Tolerance in Arabidopsis**

### **2.1. Introduction**

Calcium ions are used as secondary messengers in eukaryotic cells. The cytosolic Ca<sup>2+</sup> concentration, [Ca<sup>2+</sup>]<sub>cyt</sub>, fluctuates in response to a variety of stimuli, including mechanical stimulation, hormones, pathogens, light, and abiotic stresses such as low temperature (Sanders et al., 2002; White and Broadley, 2003; Dodd et al., 2010). The stimulus-specific spatiotemporal patterning of [Ca<sup>2+</sup>]<sub>cyt</sub> dynamics is called the Ca<sup>2+</sup> signature (Monshausen, 2012), and to create these signatures, Ca<sup>2+</sup> influx channels and Ca<sup>2+</sup> efflux transporters that permit transient increases in [Ca<sup>2+</sup>]<sub>cyt</sub> are required (McAinsh and Pittman, 2009).

How plant cells generate stimulus-specific Ca<sup>2+</sup> signals remains unknown. To identify the spatiotemporal patterning of [Ca<sup>2+</sup>]<sub>cyt</sub> dynamics, recombinant aequorin has been introduced as a reporter of [Ca<sup>2+</sup>]<sub>cyt</sub> changes in plant systems (Knight et al., 1991). In Arabidopsis plants expressing aequorin in the cytoplasm, low temperature triggers an immediate and transient rise in [Ca<sup>2+</sup>]<sub>cyt</sub> (Knight et al., 1991; Knight et al., 1996; Carpaneto et al., 2007). The final temperature and cooling rate are important for sensing low temperature in Arabidopsis (Knight, 2002). In mammals, many TRP (transient receptor potential) channels, which are a specific class of ion channels, function as intracellular Ca<sup>2+</sup> release channels (Gees et al., 2010). Some of these channels also function as thermosensors (Gees et al., 2010), and TRPA1 seems to act as a sensor for cold (Karashima et al., 2009; del Camino et al., 2010; Aubdool et al., 2014). Although no

proteins with high similarity to TRP channels have been identified in land-plant genomes, the genes for Cr-TRP proteins are encoded in the genomic sequence of the alga *Chlamydomonas reinhardtii* and show functional properties that are similar to those of mammalian TRP channels (Arias-Darraz et al., 2015).

Two Ca<sup>2+</sup>-permeable mechanosensitive channels, named MCA1 and MCA2 (*mid1*-complementing activity 1 and 2), have been identified in Arabidopsis (Nakagawa et al., 2007; Yamanaka et al., 2010; Nakano et al., 2011; Furuichi et al., 2012; Kamano et al., 2015). Both *MCA1* and *MCA2* complement deficiency of Ca<sup>2+</sup> uptake in yeast cells lacking a Ca<sup>2+</sup> channel composed of the Mid1 and Cch1 subunits (Nakagawa et al., 2007; Yamanaka et al., 2010). It should be noted that this complementation activity is detected under conditions that allow the Mid1/Cch1 channel to function as the sole Ca<sup>2+</sup> influx system in yeast cells, suggesting that MCA1 and MCA2 can directly mediate Ca<sup>2+</sup> influx in the cells lacking both Mid1 and Cch1. Electrophysiological studies have shown that both MCA1 and MCA2 produce stretch-activated currents when expressed in *Xenopus laevis* oocytes (Furuichi et al., 2012). These results with yeast cells and *Xenopus* oocytes suggest that MCA1 and MCA2 mediate Ca<sup>2+</sup> influx as mechanosensitive channels and are not accessory factors that facilitate Ca<sup>2+</sup> influx. Overexpression of *MCA1* enhances an increase in [Ca<sup>2+</sup>]<sub>cyt</sub> upon hypoosmotic shock (Nakagawa et al., 2007). The *mca2* mutant exhibits a defect in Ca<sup>2+</sup> uptake from the roots (Yamanaka et al., 2010). Structurally, MCA1 and MCA2 have 74% identity and 89% similarity in amino acid sequences (Nakagawa et al., 2007). Both have a single transmembrane segment and an EF-hand-like motif and coiled-coil motif in the N-terminal region, as well as a plac8 motif in the C-terminal region (Nakagawa et al., 2007; Kamano et al., 2015). MCA1-GFP and MCA2-GFP are localized to the plasma membrane (Nakagawa et al., 2007). MCA1 and

MCA2 form a homotetramer (Nakano et al., 2011; Shigematsu et al., 2014). Topological analysis has indicated that the EF-hand-like motif, the coiled-coil motif, and the plac8 motif are present in the cytoplasm (Kamano et al., 2015), suggesting that both channels recognize intracellular  $\text{Ca}^{2+}$ . The *MCA* genes are conserved in the plant kingdom (Kurusu et al., 2013), and an increase in  $[\text{Ca}^{2+}]_{\text{cyt}}$  as a result of hypo-osmotic shock is mediated by MCA proteins in rice and tobacco (Kurusu et al., 2012).

Application of the patch-clamp technique has demonstrated that  $\text{Ca}^{2+}$ -permeable channels are transiently activated by cold shock in *Arabidopsis* mesophyll cells (Carpaneto et al., 2007). In plants, extracellular freezing causes dehydration and mechanical stresses on the plasma membrane, and cold-acclimated plant plasma membranes become resistant to mechanical stress (Yamazaki et al., 2008). Expression of *CBF2* is induced not only by cold, but also by mechanical stress (Zarka et al., 2003). Therefore, it is assumed that mechanical stress may be one of the factors involved in cold acclimation.

Three CBF/DREB1 (C-repeat binding factor/DRE binding factor 1) transcription factors have been extensively studied. CBF/DREB1, belonging to the AP2/ERF (apetala/ethylene-responsive factor) superfamily, are important factors for cold acclimation in plants (Miura and Furumoto, 2013). *CBF/DREB1* genes are rapidly and transiently induced after cold treatment (Gilmour et al., 1998), and overexpression of *CBF/DREB1* constitutively enhances freezing tolerance (Stockinger et al., 1997; Liu et al., 1998). Under cold stress, CBF/DREB1 proteins bind to CRT/DRE *cis*-elements in the promoter of cold-regulated (*COR*) genes and induce transcription (Stockinger et al., 1997). However, gene expression analyses reveals that only 6.5% of the total *COR* genes are regulated by *CBF/DREB1* (Park et al., 2015). In addition to *CBF/DREB1* genes, 27

transcription factors that were up-regulated at an early stage after cold treatment were considered as first-wave transcription factors (Park et al., 2015). Use of the *cbf1/2/3* triple mutant showed that six first-wave transcription factors are partially regulated by *CBF/DREB1*, whereas the transcription factors *HSFC1*, *ZAT12*, and *CZF1*, which regulate cold-regulated genes (Vogel et al., 2005; Park et al., 2015), are not regulated by *CBF/DREB1* (Zhao et al., 2016). As acclimated *cbf1/2/3* triple mutants are more tolerant to freezing stress than non-acclimated ones (Jia et al., 2016), and the expression of a large number of cold-regulated genes is not affected by the *cbf1/2/3* triple mutation (Zhao et al., 2016), a *CBF/DREB1*-independent pathway may control cold tolerance. Overexpression of *HSFC1* enhances cold tolerance without an increase in expression of *CBF1*, *CBF2*, or *CBF3* (Park et al., 2015), suggesting that *HSFC1* is one of the important transcription factors controlling non-*CBF/DREB1* regulons and cold tolerance.

This research demonstrates that *MCA1* and *MCA2* are involved in a transient rise in  $[Ca^{2+}]_{\text{cyt}}$  upon cold shock. The cold-induced increase in  $[Ca^{2+}]_{\text{cyt}}$  was smaller in the *mca1* and *mca2* mutants than in the Col-0 wild type. The *mca1 mca2* double mutant exhibited increased sensitivity to chilling and freezing stresses. These results suggest that *MCA1* and *MCA2* are involved in cold-induced  $Ca^{2+}$  influx and that the reduced  $[Ca^{2+}]_{\text{cyt}}$  increase caused by the *mca1* and *mca2* mutations affects cold acclimation. As the *CBF/DREB1* genes and their regulon genes were not down-regulated in the *mca1 mca2* mutant, *MCA* may not be involved in the regulation of *CBF/DREB1*-dependent cold signaling.

## **2.2. Methods**

### **2.2.1. Plant materials.**

The Columbia-0 (Col-0) of *Arabidopsis* and its isogenic, transgenic lines *mca1*-null, *mca2*-null, and *mca1*-null *mca2*-null were previously described (Nakagawa et al., 2007; Yamanaka et al., 2010). The complementation lines *MCA1*<sub>pro::MCA1</sub> in *mca1*-null *mca2*-null and *MCA2*<sub>pro::MCA2</sub> in *mca1*-null *mca2*-null were also previously described (Nakagawa et al., 2007; Yamanaka et al., 2010).

### **2.2.2. Monitoring of [Ca<sup>2+</sup>]<sub>cyt</sub> changes following cold shock treatment.**

Apoaequorin-expressing seedlings grown at 22°C on MS medium supplemented with 0.8% agar and 1% sucrose under 16-h light conditions at 40-60 μM m<sup>-2</sup> s<sup>-1</sup> light intensity were used to monitor [Ca<sup>2+</sup>]<sub>cyt</sub> changes upon cold shock. A seedling was harvested 14 days after sowing and incubated overnight at 22°C in 2 ml of MS medium containing 2.5 μM coelenterazine in the dark to reconstitute aequorin. The seedling was transferred to fresh MS medium (400 μl) kept at 22°C in a tube (Microtech-Nition, #NU-063, Funabashi, Japan) and received an additional 500 μl of the same medium kept at 3, 10, or 22°C. Luminescence (L) from aequorin in the whole seedlings was measured using a luminometer (Microtech-Nition, Model NU-2500). At the end of each monitoring, 1 ml of 20% ethanol/2 M CaCl<sub>2</sub> solution was added to the medium (0.9 ml total) to measure the maximum luminescence (L<sub>max</sub>). The luminescence ratios (L/L<sub>max</sub>) are presented in Figures 2.1 and 2.2.

### **2.2.3. Plant freezing and chilling assay.**

Wild-type (ecotype Col-0), *mca1*, *mca2*, and *mca1 mca2* plants were grown at 24°C for

3 weeks in soil with fluorescent lighting (16 h/8 h light/dark photoperiod). These plants were then incubated at 4°C for 1 week for acclimation to low temperatures. For non-acclimation, 3-week-old plants were treated with the freezing temperature without incubation at 4°C. Whole-plant freezing assays were performed as previously described (Miura et al., 2011). Briefly, plants were incubated at 0°C for 1 h, and the temperature was lowered by 2°C h<sup>-1</sup> until it reached to the indicated temperature, and then held at the desired temperature for 1 h or 4 h for non-acclimated plants or cold-acclimated plants, respectively, in the incubator (IN602, Yamato Scientific Co., Ltd., Tokyo, Japan). After cold acclimation, the plants were incubated at 4°C overnight and transferred to 24°C. The survival ratio was determined 1 week after the freezing test.

For the chilling assay, 5-day-old plants were incubated at 4°C. After incubation for 1 month under constant illumination, the chlorophyll content of the plants was determined. Eighty percent acetone was added to leaves ground with liquid nitrogen. The mixture was shaken at 4°C. After centrifugation, the absorbances of the supernatant at 663 nm and 646 nm were measured. Total chlorophyll content was calculated as  $17.3 A_{646} + 7.18 A_{663}$  (Zakhleniuk et al., 2001).

Electrolyte leakage from fully developed rosettes of leaves of three-week-old plants was measured as previously described (Miura et al., 2007; Miura and Ohta, 2010). The sample was incubated in a refrigerated circular bath (TRL-11P, Thomas Kagaku Kikai, Co., Ltd., Japan). The conductivity was measured with a conductivity meter (CD-4302, Lutron Electronic Enterprise Co., Ltd., Taipei, Taiwan).

#### **2.2.4. RNA preparation and quantitative RT-PCR.**

Three-week-old wild-type (ecotype Col-0), *mca1*, *mca2*, and *mca1 mca2* plants were

subjected to cold treatment at 4°C for the indicated time. Isolation of total RNA, cDNA synthesis, and quantitative RT-PCR were performed as previously described (Miura et al., 2007). The primers used to detect *CBF/DREB1* and its regulon genes were also previously described (Miura et al., 2007). Other genes were detected with gene-specific primers for *At5g61820* (5'-GAGGCACCTGCGAGAAGCTTGAG-3' and 5'-GTAACCATCTTCCCGTTTCTGTC-3'), *At3g51660* (5'-GACCTCAAAACTTAGTGATGGTG-3' and 5'-TTAACTTGTTTGGTGATGCCTCC-3'), *At4g15490* (5'-CCTCCCATGGAAGGGACATTTGTAGA-3' and 5'-ACAAGCAATCGCAGGATGAGCCA-3'), *MCA1* (5'-AAGATTGCCACTGCAGCATCC-3' and 5'-ACGCCATTAGCTCATTACATGCTTC-3'), and *MCA2* (5'-AAGATCATTGCAACACCGTGGA-3' and 5'-GTGTCTTCAAGCAAAGACAAGGTTC-3').

## 2.3. Results

### 2.3.1. *MCA1* and *MCA2* are involved in a cold-induced $[Ca^{2+}]_{cyt}$ increase.

To monitor changes in  $[Ca^{2+}]_{cyt}$  (the cytosolic concentration of  $Ca^{2+}$ ), Arabidopsis seedlings expressing aequorin, a  $Ca^{2+}$  indicator (Nakagawa et al., 2007), that had been immersed in MS medium (400  $\mu$ l) at 22°C were exposed to low temperatures by the addition of MS medium (500  $\mu$ l) kept at 3, 10, or 22°C.  $[Ca^{2+}]_{cyt}$  in the wild type was significantly increased by a 3°C shock (Figure 2.1A green line and B), moderately by a 10°C shock (Figure 2.1A black line and C), and just a little by a 22°C shock (Figure 2.1A red line and D). On the other hand, the magnitude of the cold-induced  $[Ca^{2+}]_{cyt}$  increase was markedly lower in the *mca1*, *mca2*, and *mca1 mca2* mutants (Figure 2.1). Small increases seen in response to the 22°C shock in both the wild type and the mutants could

be a consequence of mechanical stress rather than cold stress, because MS medium (a fluid) was added to induce the response. These results suggest that MCA1 and MCA2 contribute to a  $[Ca^{2+}]_{cyt}$  increase upon cold shock.

Since MCA1 and MCA2 are present in the plasma membrane (Nakagawa et al., 2007; Yamanaka et al., 2010), the cold-induced  $[Ca^{2+}]_{cyt}$  increase could be brought about by  $Ca^{2+}$  influx. To examine this possibility, seedlings were preincubated for 30 min in MS medium including either a  $Ca^{2+}$  chelator, EGTA, or a plasma membrane ion channel blocker,  $La^{3+}$  or  $Gd^{3+}$ , and then monitored changes in  $[Ca^{2+}]_{cyt}$  upon cold shock. As expected, the cold-induced  $[Ca^{2+}]_{cyt}$  increase was inhibited by EGTA (Figure 2.2A, B),  $La^{3+}$  (Figure 2.2C, D), and  $Gd^{3+}$  (Figure 2.2E, F) in the wild type and in all the *mca* mutants, although the inhibition rates of the wild type were greater than those of the *mca* mutants. It should also be noted that significant  $[Ca^{2+}]_{cyt}$  increases remained in all the *mca* mutants, as well as in the wild type, suggesting that there is another cold-induced  $Ca^{2+}$  transport system(s) that is insensitive to the blockers used in the plasma membrane, or that is in the intracellular compartment.

Even though the *mca* mutants exhibited a reduced cold-induced  $[Ca^{2+}]_{cyt}$  increase, the mutants looked healthy when they grew under normal conditions (Figure 2.3A). To examine whether the *mca* mutation affects plant growth under normal conditions, fresh weight and chlorophyll contents were measured (Figure 2.3B, C). The *mca1*, *mca2*, and *mca1 mca2* plants had similar values, as did the wild type, suggesting that plant development in the aerial part is unaffected by *MCA1* and *MCA2*.

### **2.3.2. Mutations in *MCA1* and *MCA2* result in cold sensitivity.**

Since the *mca* mutants exhibited a reduced cold-induced  $[Ca^{2+}]_{cyt}$  increase, their cold

sensitivity was investigated. Three-week-old plants were incubated at 4°C for 1 week to acclimate to cold stress. These plants were then exposed to freezing temperatures (Figure 2.4A, B). Before this exposure, *mca1*, *mca2*, *mca1 mca2* and the wild-type plants looked healthy, as shown in Figure 2.3A. After this exposure, the survival of the *mca1* mutant was similar to that of the wild type. On the other hand, the *mca2* mutant exhibited a freezing-sensitive phenotype (Figure 2.4A, B). Furthermore, the *mca1 mca2* double mutant was more sensitive to freezing stresses than the *mca2* mutant (Figure 2.4A, B). Electrolyte leakage from the *mca1 mca2* double mutant was much higher (approximately 50%) than that of the *mca* single mutants and the wild type, even before it was subjected to freezing temperatures, and it increased as the freezing temperature was lowered (Figure 2.4C). At every freezing temperature employed (-3 to -9°C), the leakage was greatest in the double mutant. To confirm whether the freezing sensitivity of the *mca1 mca2* mutant was caused by the mutation in *MCA1* or *MCA2*, complement lines were produced. *MCA1pro::MCA1* or *MCA2pro::MCA2* was expressed in the *mca1 mca2* mutant (Figure 2.4D, E). Because the own promoter was used for expression of *MCA1* or *MCA2*, the expression level of *MCA1* in the complement lines, *MCA1pro::MCA1* in *mca1 mca2*, was similar to that of wild type and the *mca2* mutant (Figure 2.5). The expression level of *MCA2* in the complement lines, *MCA2pro::MCA2* in *mca1 mca2*, was slightly higher than that of WT and *mca1* (Figure 2.5). The sensitivity of *MCA2pro::MCA2*-expressing *mca1 mca2* mutant was recovered (Figure 2.4E). On the other hand, the survival ratio of *MCA1pro::MCA1*-expressing *mca1 mca2* was similar to that of *mca2* (Figure 2.4E). The wild-type, *mca1*, *mca2*, and *mca1 mca2* plants without cold acclimation were also treated with a freezing temperature for 1 h (Figure 2.4F). Before acclimation, the cold sensitivity of the *mca1 mca2* plants was slightly greater than that of wild-type, *mca1* and *mca2* plants

(Figure 2.4G). Furthermore, electrolyte leakage of the *mca1 mca2* mutant was a little higher than that of the wild type (Figure 2.4H). These results suggest that MCA mainly functions in the regulation of cold tolerance during cold acclimation.

To examine whether the mutant exhibits chilling sensitivity, wild-type, *mca1*, *mca2*, and *mca1 mca2* plants were incubated at 4°C for 1 month under continuous light conditions. The leaves of the *mca1 mca2* double mutant looked unhealthy (Figure 2.4I). Thus, to quantify chilling sensitivity, chlorophyll content was measured. The chlorophyll content in the *mca1 mca2* double mutant was approximately three-fourths that of the wild type (Figure 2.4J). No detectable difference was observed between the wild type and the *mca* single mutants. These results suggest that the double mutation in *MCA1* and *MCA2* results in hypersensitivity to cold stress in Arabidopsis plants.

### **2.3.3. Down-regulation of cold-inducible genes is governed by a non-*CBF/DREB1* regulon.**

The expression of the *CBF/DREB1* genes and their regulon genes, *COR15A*, *COR47*, and *RD29A*, was investigated. To perform this, three-week-old wild-type, *mca1*, *mca2*, and *mca1 mca2* plants were exposed to cold at 4°C for appropriate periods, and RNA prepared from the plants was subjected to a quantitative RT-PCR analysis. Interestingly, the expression of these genes was slightly up-regulated in the *mca1 mca2* double mutant, especially soon after the start of the cold treatment (Figure 2.6A). This increase could be the consequence of a compensatory response caused by a lack of the function of *MCA1* and *MCA2* and suggests that both proteins may control another cold signaling pathway. *CBF/DREB1* and its regulatory genes are only partly responsible for the acquisition of tolerance to freezing stress for cold acclimation (Park et al., 2015). Therefore, the

expression of cold-inducible genes that are governed by *HSFC1* but not by *CBF2* (Park et al., 2015), such as *At5g61820*, *At3g51660*, and *At4g15490*, which encode an unknown protein, a tautomerase/MIF superfamily protein, and a UDP-glycosyltransferase superfamily protein, respectively, was examined. Figure 2.6B shows that the expression of the three genes was significantly down-regulated in the *mca1 mca2* double mutant. Expression of MCA1 and MCA2 themselves was unchanged upon cold shock (Figure 2.6C). According to these results, it is plausible that MCA1 and MCA2 mediate cold tolerance by participating in a pathway other than the *CBF/DREB1* pathway.

#### 2.4. Discussion

The present study has demonstrated that MCA1 and MCA2 play a role in the transient rise in  $[Ca^{2+}]_{\text{cyt}}$  upon cold shock and are also involved in chilling and freezing tolerance. The *mca1 mca2* double mutant exhibited a lower cold-induced increase in  $[Ca^{2+}]_{\text{cyt}}$  than the wild type (Figure 2.1), as well as an increased sensitivity to cold stress (Figure 2.4). Furthermore, *CBF/DREB1* genes and their regulon genes were not down-regulated in the double mutant (Figure 2.6), suggesting that MCA may not regulate *CBF/DREB1*-dependent cold signaling.

Different stimuli produce different patterns of  $Ca^{2+}$  elevation and oscillations with different frequencies, and these are called  $Ca^{2+}$  signatures. As shown in Figure 2.1, low-temperature stress stimulates a transient increase in  $[Ca^{2+}]_{\text{cyt}}$  (Knight and Knight, 2012). MCA1 and MCA2 have been identified as plasma membrane proteins involved in  $Ca^{2+}$  influx in response to mechanical stimuli, such as touch, gravity, flexure, and turgor (Nakagawa et al., 2007; Yamanaka et al., 2010). In the *mca1* or *mca2* mutant, the magnitude of the cold-induced  $[Ca^{2+}]_{\text{cyt}}$  increase was lower, by approximately 40%, than

that in the wild type (Figure 2.1). These results suggest that MCA1 and MCA2 are partially involved in  $\text{Ca}^{2+}$  influx in response to cold shock. Application of a mechanosensitive  $\text{Ca}^{2+}$  channel blocker,  $\text{Gd}^{3+}$ , prevents the induction of cold-regulated genes (Sangwan et al., 2002). As two  $\text{Ca}^{2+}$  channel inhibitors,  $\text{La}^{3+}$  and  $\text{Gd}^{3+}$ , still reduced the cold-induced  $[\text{Ca}^{2+}]_{\text{cyt}}$  increase in the *mca* mutants, and the *mca* mutations were unable to block  $[\text{Ca}^{2+}]_{\text{cyt}}$  increases completely (Figure 2.2), other cold-activatable  $\text{Ca}^{2+}$  transport system(s) must exist in the plasma membrane and/or organellar membranes. Indeed, it is reported that the vacuole, the major intracellular  $\text{Ca}^{2+}$  store, is involved in a cold-induced  $\text{Ca}^{2+}$  release (Knight et al., 1996). In the present study, the bioluminescent intensity of aequorin for  $[\text{Ca}^{2+}]_{\text{cyt}}$  was not calibrated because of difficulties in the precise calibration, although there is a report describing a successful calibration specific for the isoform of aequorin and temperature that the authors used (Knight et al., 1996).

Plants employ several kinds of mechanisms to control  $\text{Ca}^{2+}$ -regulated gene expression (Galon et al., 2010). However, it is still unclear how cold-induced  $[\text{Ca}^{2+}]_{\text{cyt}}$  increases are recognized. One possible mechanism involves calmodulin-binding transcription factors (CAMTAs). CAMTAs possess calmodulin (CaM)-binding domains (Liu et al., 2015) and CAMTAs play a role in the regulation of gene expression in response to  $\text{Ca}^{2+}$  signals (Whalley and Knight, 2013). CAMTA3 is a positive regulator of *CBF2/DREB1C* expression and binds to the consensus sequence of a CGCG core motif, a *cis*-element for CAMTAs, in the promoter of *CBF2/DREB1C* (Doherty et al., 2009). The *camta2 camta3* double mutant is sensitive to freezing temperatures (Doherty et al., 2009). Microarray analyses demonstrated that the expression level of *HSFC1* (*At3g24520*) in the *camta1/2/3* mutant is lower than that in the wild type (Kim et al., 2013). CAMTA is one of possibilities how increased  $[\text{Ca}^{2+}]_{\text{cyt}}$  is recognized.


As shown in Figure 2.1, the *mca1 mca2* double mutation reduced about 40% of the  $[Ca^{2+}]_{\text{cyt}}$  increase. The transient increase in  $[Ca^{2+}]_{\text{cyt}}$  may be conducted by other  $Ca^{2+}$ -permeable channels that are responsible for 60% of the transient increase. One such channel could be AtGLR3.4, a member of the Arabidopsis homologs of ionotropic glutamate receptors, whose expression is up-regulated under cold stress (Meyerhoff et al., 2005). Another such channel could be the cyclic nucleotide-gated ion channel (CNGC) family. In rice, the expressions of 10 out of 16 *CNGC* genes are induced under cold stress (Nawaz et al., 2014).

The *mca1 mca2* double mutant exhibited increased sensitivity to chilling and freezing stresses, even though the single mutants did not exhibit a severe phenotype (Figure 2.4). Both MCA1 and MCA2 complement a  $Ca^{2+}$  uptake deficiency of yeast cells lacking a  $Ca^{2+}$  channel composed of Mid1 and Cch1 (Nakagawa et al., 2007; Yamanaka et al., 2010) and generate stretch-activated currents in *Xenopus* oocytes (Furuichi et al., 2012). Even though MCA1 and MCA2 have similar functions as  $Ca^{2+}$ -permeable mechanosensitive channels, their spatial expression patterns are not necessarily the same in whole plants (Yamanaka et al., 2010). *MCA1p::GUS* and *MCA2p::GUS* are expressed in vascular tissues of cotyledons, leaves and primary roots in common. On the other hand, *MCA1p::GUS* is expressed in the promeristem and adjacent elongation zone of the primary root, while *MCA2p::GUS* is not. *MCA2p::GUS* is expressed in mesophyll cells of cotyledons and leaves, but *MCA1p::GUS* is not. In addition, *MCA2p::GUS* is expressed more than *MCA1p::GUS* at the center of rosettes in a region corresponding to the shoot apical meristem. To survive freezing stress, the shoot apical meristem should be protected to recover plant growth (Pearce and McDonald, 1977). Based on the observation of differences in the spatial expression patterns of *MCA1p::GUS* and

*MCA2p::GUS* in whole plants, it is possible to speculate that the differences may allocate MCA1 and MCA2 a role in the acquisition of tolerance to cold stress. This allocation could explain why only the double mutant becomes hypersensitive to chilling and freezing stresses.


According to microarray analyses, several genes were up-regulated in *HSFC1*-overexpressed plants, but not in *CBF2*-overexpressed plants (Park et al., 2015). Some *HSFC1*-dependent and *CBF2*-independent cold-regulated genes, such as *At5g61820*, *At3g51660*, and *At4g15490*, encoding an unknown protein, a tautomerase/MIF superfamily protein, and the UDP-glycosyltransferase superfamily protein UGT84A3, respectively, were down-regulated in the *mca1 mca2* double mutant (Figure 2.6). It is possible that the MCA1/2-regulated  $\text{Ca}^{2+}$  signal is transduced to a *HSFC1*-dependent pathway to enhance cold tolerance. This possibility warrants further study.

In conclusion, two mechanosensitive  $\text{Ca}^{2+}$  channels, MCA1 and MCA2, are involved in a cold-induced transient  $[\text{Ca}^{2+}]_{\text{cyt}}$  increase in Arabidopsis, and in the regulation of cold tolerance through a pathway other than the *CBF/DREB1*-dependent pathway.




**Figure 2.1. Transient cold-induced increase in cytosolic  $Ca^{2+}$  is lower as a result of the *mca* mutation.**

(A) Relative luminescence of plants harboring aequorin was measured before and after the addition (indicated by the vertical arrow) of precooled solution (3°C or 10°C) or room temperature solution (22°C). The figures are of representative data. The peak luminescence after the addition of solution at 3°C (B,  $n \geq 10$ ), 10°C (C,  $n \geq 9$ ), and 22°C (D,  $n \geq 17$ ) is shown. Data represent the means  $\pm$  SD.  $n$  indicates the number of seedlings. \*,  $p < 0.05$ ; \*\*,  $p < 0.005$  versus the wild type. Significance was determined using unpaired Student's  $t$  tests.




**Figure 2.2. Effect of channel blockers and a Ca<sup>2+</sup> chelator on the cold-shock-induced [Ca<sup>2+</sup>]<sub>cyt</sub> increase.**

Thirty minutes before cold shock, 5 mM EGTA (A), 1 mM La<sup>3+</sup> (C), or 1 mM Gd<sup>3+</sup> (D) was added to the medium. Then, the relative luminescence of plant harboring aequorin was measured, as in Figure 2.1, before and after the application of precooled solution (3°C). The peak luminescence after the addition of solution at 3°C with or without 5 mM EGTA (B,  $n \geq 17$ ), 1 mM La<sup>3+</sup> (D,  $n \geq 9$ ), or 1 mM Gd<sup>3+</sup> (F,  $n \geq 9$ ) is shown. Data represent the means  $\pm$  SD. \*,  $p < 0.05$ ; \*\*,  $p < 0.005$  versus the wild type in each treatment. Significance was determined using unpaired Student's *t* tests.



**Figure 2.3. The growth of the *mca1 mca2* mutant is normal at room temperature.**


(A) Representative 3-week old plants of wild type, *mca1*, *mca2*, and *mca1 mca2* mutants (nine plants in each pot) are displayed. The bar indicates 1-cm length. Fresh weight (B) and chlorophyll content (C) of each plant were measured. Values represent the means  $\pm$  SD ( $n = 10$ ). No significant difference was observed with an unpaired Student's *t*-test.



**Figure 2.4. The *mca1 mca2* double mutant exhibited sensitivity to cold stress.**


(A) Freezing sensitivity of the *mca1 mca2* double mutant after cold acclimation. Three-week-old plants were incubated at 4°C for 1 week, then plants were used for freezing treatment. Photographs are representative plants 7 days after 4-h exposure to the indicated temperature. (B) Survival rates were determined for 9 plants after freezing treatment at the indicated temperature. The survival ratio was calculated from 9 plants per pots. Data represent the means  $\pm$  SD calculated from the data of 9 independent experiments. Differences between the values of each treatment were evaluated by one-way ANOVA followed by the Tukey-Kramer test. Difference of alphabet letters at each temperature indicates statistically significant difference ( $p < 0.05$ ). (C) Electrolyte leakage from cold-acclimated wild-type, *mca1*, *mca2*, and *mca1 mca2* plants after exposure to the indicated temperature (programmed to cool at a rate of 2°C h<sup>-1</sup>). Data represent the means  $\pm$  SD ( $n = 4$  leaves, each from a different plant). \*  $p < 0.05$  compared with the value of wild type at each temperature (one-way ANOVA followed by the Tukey-Kramer test). Data were representative experiments from 3 biological independent experiments. (D) Freezing sensitivity of the *mca1 mca2* double mutant harboring *MCA1pro::MCA1* (lines #2 and #11) or *MCA2pro::MCA2* (lines #2 and #5). Photographs are representative plants 7 days after 4-h exposure to -7°C. (E) The survival ratio at -7°C was calculated from 9 plants per pots. Data represent the means  $\pm$  SD calculated from the data of 9 independent experiments. Difference between values of each treatment were evaluated by one-way ANOVA followed by the Tukey-Kramer test. Difference of alphabet letters at each temperature indicates statistically significant difference ( $p < 0.05$ ). (F) Freezing sensitivity of the *mca1 mca2* double mutant without cold acclimation. Three-week-old plants were treated with a freezing temperature. Photographs are representative plants

after freezing treatment. Photographs are representative plants 7 days after 1-h exposure to the indicated temperature. (G) The survival ratio was calculated from 9 plants per pots. Difference between values of each treatment were evaluated by one-way ANOVA followed by the Tukey-Kramer test. Difference of alphabet letters at each temperature indicates statistically significant difference ( $p < 0.05$ ). (H) Electrolyte leakage from non-acclimated wild-type, *mca1*, *mca2*, and *mca1 mca2* plants after exposure to the indicated temperature. Data represent the means  $\pm$  SD ( $n = 3$  leaves, each from a different plant). \*  $p < 0.05$  compared with the value of wild types at each temperature (ANOVA followed by the Tukey-Kramer test). (I) Photographs are of representative wild type, *mca1*, *mca2*, and *mca1 mca2* mutants after incubation at 4°C for 1 month. Five-day-old plants were incubated at 4°C for 1 month. (J) The chlorophyll content of the plants was determined. Values represent the means  $\pm$  SD ( $n = 7$  plants per each genotype). Difference of alphabet letters indicates a significant difference ( $p < 0.05$ ) as determined by one-way ANOVA followed by the Tukey-Kramer test.



**Figure 2.5. The levels of the *MCA1* and *MCA2* transcripts in the mutant and complement lines.**

Total RNA was purified from 8 plant grown for 14 days. Transcript levels of the *MCA1* and *MCA2* genes were measured by quantitative real-time PCR as described in Methods in the main text. Values represent the means  $\pm$  SD ( $n = 3$ ). An asterisk indicates a significant difference from wild-type plants as determined by unpaired Student's *t*-tests.



**Figure 2.6.** The expression levels of several cold-regulated genes, which are not regulated by *CBF/DREB1*, were reduced in the *mca1 mca2* double mutant.

(A) Relative mRNA transcript levels of *CBF/DREB1* and its regulon genes, *COR15A*, *COR47*, and *RD29A*, in wild-type and *mca1 mca2* plants were determined by quantitative RT-PCR analyses. Three-week-old plants grown at 24°C were incubated at 4°C for the indicated time. Values represent the means  $\pm$  SD ( $n = 3$ ) from representative experiments from 3 biologically independent experiments. (B) Relative expression levels of cold-

regulated genes that are not *CBF/DREB1*-regulon genes<sup>30</sup> were determined by quantitative RT-PCR analyses. Values represent the means  $\pm$  SD ( $n = 3$ ) from representative experiments from 3 biologically independent experiments. (C) The expression levels of *MCA1* and *MCA2* in wild types and *mca1 mca2* double mutants were investigated by quantitative RT-PCR analyses. An asterisk indicates a significant difference from wild-type plants at each point ( $p < 0.05$ ) as determined by unpaired Student's *t*-tests.

## **Acknowledgements**

I would like to express my sincere gratitude to my academic advisor Dr. Kenji Miura for his valuable advice and continuing support. I also would like to show my gratitude to Dr. Tsuyoshi Furumoto, Dr. Hidetoshi Iida, Dr. Iwane Suzuki, Dr. Takuya Suzaki and Dr. Hiroaki Iwai for kind suggestions and comments. I thank all my lab members for providing me kind helps though my whole research.

## References

- Arias-Darraz, L., Cabezas, D., Colenso, C.K., Alegria-Arcos, M., Bravo-Moraga, F., Varas-Concha, I., Almonacid, D.E., Madrid, R., and Brauchi, S. (2015). A transient receptor potential ion channel in *Chlamydomonas* shares key features with sensory transduction-associated TRP channels in mammals. *The Plant cell* **27**, 177-188.
- Aubdool, A.A., Graepel, R., Kodji, X., Alawi, K.M., Bodkin, J.V., Srivastava, S., Gentry, C., Heads, R., Grant, A.D., Fernandes, E.S., Bevan, S., and Brain, S.D. (2014). TRPA1 is essential for the vascular response to environmental cold exposure. *Nature communications* **5**, 5732.
- Carpaneto, A., Ivashikina, N., Levchenko, V., Krol, E., Jeworutzki, E., Zhu, J.K., and Hedrich, R. (2007). Cold transiently activates calcium-permeable channels in *Arabidopsis* mesophyll cells. *Plant physiology* **143**, 487-494.
- del Camino, D., Murphy, S., Heiry, M., Barrett, L.B., Earley, T.J., Cook, C.A., Petrus, M.J., Zhao, M., D'Amours, M., Deering, N., Brenner, G.J., Costigan, M., Hayward, N.J., Chong, J.A., Fanger, C.M., Woolf, C.J., Patapoutian, A., and Moran, M.M. (2010). TRPA1 contributes to cold hypersensitivity. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **30**, 15165-15174.
- Dodd, A.N., Kudla, J., and Sanders, D. (2010). The language of calcium signaling. *Annual review of plant biology* **61**, 593-620.
- Doherty, C.J., Van Buskirk, H.A., Myers, S.J., and Thomashow, M.F. (2009). Roles for *Arabidopsis* CAMTA transcription factors in cold-regulated gene expression and freezing tolerance. *The Plant cell* **21**, 972-984.
- Furuichi, T., Iida, H., Sokabe, M., and Tatsumi, H. (2012). Expression of *Arabidopsis* MCA1 enhanced mechanosensitive channel activity in the *Xenopus laevis* oocyte plasma membrane. *Plant signaling & behavior* **7**, 1022-1026.
- Galon, Y., Finkler, A., and Fromm, H. (2010). Calcium-regulated transcription in plants. *Molecular plant* **3**, 653-669.
- Gees, M., Colsoul, B., and Nilius, B. (2010). The role of transient receptor potential cation channels in Ca<sup>2+</sup> signaling. *Cold Spring Harbor perspectives in biology* **2**, a003962.
- Gilmour, S.J., Zarka, D.G., Stockinger, E.J., Salazar, M.P., Houghton, J.M., and Thomashow, M.F. (1998). Low temperature regulation of the *Arabidopsis* CBF family of AP2 transcriptional activators as an early step in cold-induced COR gene expression. *The Plant journal : for cell and molecular biology* **16**, 433-442.
- Jia, Y., Ding, Y., Shi, Y., Zhang, X., Gong, Z., and Yang, S. (2016). The cbfs triple mutants reveal the essential functions of CBFs in cold acclimation and allow the definition of CBF regulons in *Arabidopsis*. *The New phytologist* **212**, 345-353.

- Kamano, S., Kume, S., Iida, K., Lei, K.J., Nakano, M., Nakayama, Y., and Iida, H.** (2015). Transmembrane Topologies of Ca<sup>2+</sup>-permeable Mechanosensitive Channels MCA1 and MCA2 in *Arabidopsis thaliana*. *The Journal of biological chemistry* **290**, 30901-30909.
- Karashima, Y., Talavera, K., Everaerts, W., Janssens, A., Kwan, K.Y., Vennekens, R., Nilius, B., and Voets, T.** (2009). TRPA1 acts as a cold sensor in vitro and in vivo. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 1273-1278.
- Kim, Y., Park, S., Gilmour, S.J., and Thomashow, M.F.** (2013). Roles of CAMTA transcription factors and salicylic acid in configuring the low-temperature transcriptome and freezing tolerance of *Arabidopsis*. *The Plant journal : for cell and molecular biology* **75**, 364-376.
- Knight, H., Trewavas, A.J., and Knight, M.R.** (1996). Cold calcium signaling in *Arabidopsis* involves two cellular pools and a change in calcium signature after acclimation. *The Plant cell* **8**, 489-503.
- Knight, M.R.** (2002). Signal transduction leading to low-temperature tolerance in *Arabidopsis thaliana*. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **357**, 871-875.
- Knight, M.R., and Knight, H.** (2012). Low-temperature perception leading to gene expression and cold tolerance in higher plants. *The New phytologist* **195**, 737-751.
- Knight, M.R., Campbell, A.K., Smith, S.M., and Trewavas, A.J.** (1991). Transgenic plant aequorin reports the effects of touch and cold-shock and elicitors on cytoplasmic calcium. *Nature* **352**, 524-526.
- Kurusu, T., Kuchitsu, K., Nakano, M., Nakayama, Y., and Iida, H.** (2013). Plant mechanosensing and Ca<sup>2+</sup> transport. *Trends in plant science* **18**, 227-233.
- Kurusu, T., Nishikawa, D., Yamazaki, Y., Gotoh, M., Nakano, M., Hamada, H., Yamanaka, T., Iida, K., Nakagawa, Y., Saji, H., Shinozaki, K., Iida, H., and Kuchitsu, K.** (2012). Plasma membrane protein OsMCA1 is involved in regulation of hypo-osmotic shock-induced Ca<sup>2+</sup> influx and modulates generation of reactive oxygen species in cultured rice cells. *BMC plant biology* **12**, 11.
- Liu, J., Whalley, H.J., and Knight, M.R.** (2015). Combining modelling and experimental approaches to explain how calcium signatures are decoded by calmodulin-binding transcription activators (CAMTAs) to produce specific gene expression responses. *The New phytologist* **208**, 174-187.
- Liu, Q., Kasuga, M., Sakuma, Y., Abe, H., Miura, S., Yamaguchi-Shinozaki, K., and Shinozaki, K.** (1998). Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*. *The Plant cell* **10**, 1391-1406.

- McAinsh, M.R., and Pittman, J.K.** (2009). Shaping the calcium signature. *The New phytologist* **181**, 275-294.
- Meyerhoff, O., Muller, K., Roelfsema, M.R., Latz, A., Lacombe, B., Hedrich, R., Dietrich, P., and Becker, D.** (2005). AtGLR3.4, a glutamate receptor channel-like gene is sensitive to touch and cold. *Planta* **222**, 418-427.
- Miura, K., and Ohta, M.** (2010). SIZ1, a small ubiquitin-related modifier ligase, controls cold signaling through regulation of salicylic acid accumulation. *J Plant Physiol* **167**, 555-560.
- Miura, K., and Furumoto, T.** (2013). Cold signaling and cold response in plants. *International journal of molecular sciences* **14**, 5312-5337.
- Miura, K., Ohta, M., Nakazawa, M., Ono, M., and Hasegawa, P.M.** (2011). ICE1 Ser403 is necessary for protein stabilization and regulation of cold signaling and tolerance. *The Plant journal : for cell and molecular biology* **67**, 269-279.
- Miura, K., Jin, J.B., Lee, J., Yoo, C.Y., Stirm, V., Miura, T., Ashworth, E.N., Bressan, R.A., Yun, D.J., and Hasegawa, P.M.** (2007). SIZ1-mediated sumoylation of ICE1 controls CBF3/DREB1A expression and freezing tolerance in Arabidopsis. *The Plant cell* **19**, 1403-1414.
- Monshausen, G.B.** (2012). Visualizing Ca(2+) signatures in plants. *Current opinion in plant biology* **15**, 677-682.
- Nakagawa, Y., Katagiri, T., Shinozaki, K., Qi, Z., Tatsumi, H., Furuichi, T., Kishigami, A., Sokabe, M., Kojima, I., Sato, S., Kato, T., Tabata, S., Iida, K., Terashima, A., Nakano, M., Ikeda, M., Yamanaka, T., and Iida, H.** (2007). Arabidopsis plasma membrane protein crucial for Ca<sup>2+</sup> influx and touch sensing in roots. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 3639-3644.
- Nakano, M., Iida, K., Nyunoya, H., and Iida, H.** (2011). Determination of structural regions important for Ca(2+) uptake activity in Arabidopsis MCA1 and MCA2 expressed in yeast. *Plant & cell physiology* **52**, 1915-1930.
- Nawaz, Z., Kakar, K.U., Saand, M.A., and Shu, Q.Y.** (2014). Cyclic nucleotide-gated ion channel gene family in rice, identification, characterization and experimental analysis of expression response to plant hormones, biotic and abiotic stresses. *BMC genomics* **15**, 853.
- Park, S., Lee, C.M., Doherty, C.J., Gilmour, S.J., Kim, Y., and Thomashow, M.F.** (2015). Regulation of the Arabidopsis CBF regulon by a complex low-temperature regulatory network. *The Plant journal : for cell and molecular biology* **82**, 193-207.
- Pearce, R.S., and McDonald, I.** (1977). Ultrastructural damage due to freezing followed by thawing in shoot meristem and leaf mesophyll cells of tall fescue (*Festuca arundinacea* Schreb.). *Planta* **134**, 159-168.
- Sanders, D., Pelloux, J., Brownlee, C., and Harper, J.F.** (2002). Calcium at the crossroads of

- signaling. *The Plant cell* **14 Suppl**, S401-417.
- Sangwan, V., Orvar, B.L., Beyerly, J., Hirt, H., and Dhindsa, R.S.** (2002). Opposite changes in membrane fluidity mimic cold and heat stress activation of distinct plant MAP kinase pathways. *The Plant journal : for cell and molecular biology* **31**, 629-638.
- Shigematsu, H., Iida, K., Nakano, M., Chaudhuri, P., Iida, H., and Nagayama, K.** (2014). Structural characterization of the mechanosensitive channel candidate MCA2 from *Arabidopsis thaliana*. *PloS one* **9**, e87724.
- Stockinger, E.J., Gilmour, S.J., and Thomashow, M.F.** (1997). *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 1035-1040.
- Vogel, J.T., Zarka, D.G., Van Buskirk, H.A., Fowler, S.G., and Thomashow, M.F.** (2005). Roles of the CBF2 and ZAT12 transcription factors in configuring the low temperature transcriptome of *Arabidopsis*. *The Plant journal : for cell and molecular biology* **41**, 195-211.
- Whalley, H.J., and Knight, M.R.** (2013). Calcium signatures are decoded by plants to give specific gene responses. *The New phytologist* **197**, 690-693.
- White, P.J., and Broadley, M.R.** (2003). Calcium in plants. *Annals of botany* **92**, 487-511.
- Yamanaka, T., Nakagawa, Y., Mori, K., Nakano, M., Imamura, T., Kataoka, H., Terashima, A., Iida, K., Kojima, I., Katagiri, T., Shinozaki, K., and Iida, H.** (2010). MCA1 and MCA2 that mediate Ca<sup>2+</sup> uptake have distinct and overlapping roles in *Arabidopsis*. *Plant physiology* **152**, 1284-1296.
- Yamazaki, T., Kawamura, Y., and Uemura, M.** (2008). Cryobehavior of the plasma membrane in protoplasts isolated from cold-acclimated *Arabidopsis* leaves is related to surface area regulation. *Plant & cell physiology* **49**, 944-957.
- Zakhleniuk, O.V., Raines, C.A., and Lloyd, J.C.** (2001). *pho3*: a phosphorus-deficient mutant of *Arabidopsis thaliana* (L.) Heynh. *Planta* **212**, 529-534.
- Zarka, D.G., Vogel, J.T., Cook, D., and Thomashow, M.F.** (2003). Cold induction of *Arabidopsis* CBF genes involves multiple ICE (inducer of CBF expression) promoter elements and a cold-regulatory circuit that is desensitized by low temperature. *Plant physiology* **133**, 910-918.
- Zhao, C., Zhang, Z., Xie, S., Si, T., Li, Y., and Zhu, J.K.** (2016). Mutational Evidence for the Critical Role of CBF Transcription Factors in Cold Acclimation in *Arabidopsis*. *Plant physiology* **171**, 2744-2759.