

DA
2214 (HG)
1999

Distribution of Cholinergic System Components in the Adult Newt Retina, and their Appearance and Maturation during Retinal Development and Regeneration

Eun Woo CHEON

June, 1999

Submitted in partial fulfillment of the requirements for the degree
of Doctor of Philosophy in Science, in the Doctoral Program in
Biological Sciences, University of Tsukuba.



00301771

Contents

	page
Title	
Contents -----	i
Abbreviations -----	v
I Abstract -----	1
II General introduction -----	4
1 Development and cellular organization of the retina	
2 Regeneration of the retina	
3 Neurochemical differentiation of the retina	
4 Aims of the present study	
Chapter I	
Localization of the cholinergic system in the normal adult newt retina -----	10
1 Introduction -----	11
2 Materials and Methods -----	13
2.1 Preparation of the normal adult retina	
2.2 Immunocytochemistry	
2.3 α -bungarotoxin binding histochemistry	
2.4 Acetylcholinesterase histochemistry	
3 Results -----	17
3.1 Cellular organization of the normal newt retina	
3.2 Choline acetyltransferase immunoreactivity	

3.3	Acetylcholinesterase activity	
3.4	Acetylcholine receptor immunoreactivity	
3.4.1	Muscarinic acetylcholine receptors (mAChRs)	
3.4.2	Nicotinic acetylcholine receptors (nAChRs)	
4	Discussion	22
4.1	Localization of cholinergic neurons	
4.2	Acetylcholine receptors (AChRs)	
4.2.1	Localization of mAChRs	
4.2.1	Localization of nAChRs	
Chapter II		
Appearance and maturation of the cholinergic system during retinal development		31
1	Introduction	32
2	Materials and Methods	34
2.1	Preparation of developing retina	
2.2	Immunocytochemistry and histochemistry	
3	Results	36
3.1	Definition of stages of developing retinas	
3.2	Choline acetyltransferase	
3.3	Acetylcholinesterase	
3.4	Acetylcholine receptors	
3.4.1	Muscarinic acetylcholine receptors (mAChRs)	
3.4.2	Nicotinic acetylcholine receptors (nAChRs)	
4	Discussion	41
4.1	The onset of AChE activity at an earlier embryonic stage than that of ChAT immunoreactivity	

- 4.2 The onset of mAChRs at an earlier embryonic stage than that of ChAT
- 4.3 The later appearance of nAChRs than that of mAChRs

Chapter III	
Appearance and maturation of the cholinergic system during retinal regeneration -----	46
1 Introduction -----	47
2 Materials and Methods -----	50
2.1 Preparation of regenerating retina	
2.2 Immunocytochemistry and histochemistry	
3 Results -----	52
3.1 Definition of stages of regenerating retina	
3.2 Choline acetyltransferase	
3.3 Acetylcholinesterase	
3.4 Acetylcholine receptors	
3.4.1 Muscarinic acetylcholine receptors (mAChRs)	
3.4.2 Nicotinic acetylcholine receptors (nAChRs)	
4 Discussion -----	57
4.1 The onset of AChE activity in regenerating retina is earlier than that of ChAT immunoreactivity	
4.2 The onset of mAChRs in regenerating retina is earlier than that of ChAT immunoreactivity	
4.3 The onset of nAChRs in regenerating retina is later than that of mAChRs	

III General discussion -----	61
1 Development versus regeneration	
2 Possible non-cholinergic functions of AChE	
3 Future perspective	
IV Acknowledgements -----	67
V References -----	68
VI Table -----	90
VII Figures and legends -----	92

Abbreviations

CNS : central nervous system
PCL : pigment cell layer
RCL : receptor cell layer
ONL : outer nuclear layer
OPL : outer plexiform layer
INL : inner nuclear layer
IPL : inner plexiform layer
GCL : ganglion cell layer
R : photoreceptor cell
H : horizontal cell
B : bipolar cell
A : amacrine cell
G : ganglion cell
RPE : retinal pigment epithelium
ACh : acetylcholine
ChAT : choline acetyltransferase
AChE : acetylcholinesterase
iso-OMPA : tetraisopropylpyrophosphoramide
ERG : electroretinogram
 α -BTX : α -bungarotoxin
AChRs : acetylcholine receptor
mAChR : muscarinic acetylcholine receptor
nAChR : nicotinic acetylcholine receptor
PBS : phosphate buffer saline
PB : phosphate buffer
SD : standard deviation