

DB  
2007  
2004  
MG

**Neogene echinoid fossils of the northern part of  
Ibaraki Prefecture, northeastern Kanto, Japan**

**A Dissertation Submitted to  
the Graduate School of Life and Environmental Sciences,  
the University of Tsukuba  
in Partial Fulfillment of the Requirements  
for the Degree of Doctor of Philosophy in Science**

**Yoshibumi KIKUCHI**

寄贈  
菊池芳文氏

05009576

## Contents

• Abstract	iii
• Introduction	1
• Objectives and methods of the study	4
<b>Part A. Geological setting of echinoid fossil-bearing strata in the northern part of Ibaraki Prefecture</b>	
• I. Daigo area	9
• II. Yamagata area	31
• III. Central Sheared Zone in southern part of Suifu area	39
• IV. Omiya area	47
• V. Naka area	59
• VI. Kanasago to Hitachiota area	66
• VII. Southern part of Kanasago area	78
• VIII. Eastern part of Hitachiota area	85
• IX. Ishinazaka area	93
• X. Kuji area	100
• XI. Hitachi area	105
• XII Tokai area	117
• XIII. Hitachinaka area	126

**Part B. Paleontological study of the Neogene echinoid fossils  
from the northern part of Ibaraki Prefecture**

• Systematic classification . . . . .	135
• Localities of echinoid fossils . . . . .	221
• Distribution of time and space and ecology of echinoid fossils from the northern part of Ibaraki Prefecture . . . . .	226
• Assemblages and paleoenvironments of echinoid fossils . . . . .	249
• Faunal change of echinoid fossils . . . . .	271
• Origin and migration of the Neogene echinoid fossils . . . . .	281
• Results on paleontological study of echinoid fossils . . . . .	285
• Results on stratigraphical study . . . . .	291
• Acknowledgements . . . . .	295
• References cited . . . . .	297
• Plates 1-22	