

### References cited

- Agassiz, A., 1879; Preliminary report on the echini of the exploring expedition H.M.S. "Challenger" Sir C. Wyville Thomson Chief of civilian staff. *Proc. Amer. Acad., Arts. Sci.*, vol. 14 (n.s., vol. 6), (1878-1879), p. 182-261.
- , 1881; Repot on the Echinoidea dredged by H.M.S. Challenger during the years 1873-1876. *Rep. Challenger-Exped. Zool*, vol. 3, no. 9, p. 1-321, pls.1-45.
- and Clark, H. L., 1907; Preliminary reports on the echini collected in 1906, from May to December, among the Aleutian Islands, in Bering Sea, and along the coasts of Kamtchatka, Sakhalin, Korea and Japan, by the U. S. Fish Commission Steamer "Albatross" in Charge of Lieut. Commander, L. M. Garrett, U.S.N. Commanding. *Bull. Mus. Com. Zool.*, vol. 51, no. 5, 109-139.
- Akiba, F., 1988; *Diatom fossil assemblage included in the molluscan fossils from the Pliocene Kume Formation of Ibaraki Prefecture, Japan*. Abstracts of the 1988 Annual Meeting of the Palaeontological Society of Japan, p. 63. (in Japanese, title translated)

- Akutsu, J., 1952; Geomorphology and geology of Hitachi Omiya and its environment, Ibaraki Prefecture. *Bull. Utsunomiya Univ.*, no. 2, p. 1-19. (*in Japanese*)
- Amano, k. and Takahashi, H., 1986; *Tanakura Sheared Zone area*, In. Omori, M., Hayama, Y. and Horiguchi, M., eds., *Geology of Japan 3, Kanto district*, p. 132-134. Kyoritsu Printed Company, Tokyo. (*in Japanese, title translated*)
- , Koshiya, S., Takahashi, H., Noda, H. and Yagishita, K., 1989; *Tectonics and sedimentation in Tanakura Shear Zone*. p. 55-86, pl. 1. Excursion Guidebook at the 96th Annual Meeting of Geological Society of Japan. (*in Japanese*)
- and Takahashi, H., 1984; *Geology and Tertiary System in the area of Tanakura SHERA Zone*. p. 6-9, Rep. Spec. Res. Project, Ibaraki Univ., Mito. (*in Japanese, title translated*)
- Aoki, S., 1933; *Echinodermata*. In, *Geology and Paleontology*, p. 1-60, Iwanami Lectures, Iwanami-shoten, Tokyo. (*in Japanese*)
- Baranova, Z. I., 1955; New species and subspecies of echinodermata from the Bering Sea. *Travaux Inst. Zool., Acad. Sci., SSSR*, vol. 18, p. 334-342. (*in Russian, title translated*)
- Castex, L., 1930; Révision des échinides du Nummulitique du Department des Landes. *Act. Soc., Linn. Bordeaux*, vol. 82,

p. 5-72, pls. 1-4.

Checchia-Rispoli, G., 1950; Su la echinidi ecoceni della Migiurtina.

*Bull. Ufficio Geol. Italia*, vol. 70, p. 21-39, pls. 1-2.

Clark, A. H., 1942; Geology and biology of North Atlantic deep-sea

Cores between New Fundland and Ireland; Part 6,

Echinodermata. *U. S. G. S. Prof. Pap*, vol. 196-D, p. 111-

117, pl. 22.

————— and Rowe, F. W. E., 1971; *Monograph of shallow-water*

*Indo-west Pacific echinoderms*. P.1-238, pls. 1-31. Trustees

of the British Museum (Natural History), London.

Clark, H. L., 1912; Hawaiian and other Pacific Echini; the Pedinidae,

Phymosomatidae, Stomopneustidae, Echinidae,

Temnopleuridae, Strongylocentrotidae and Echinometridae.

*Mem. Mus. Comp. Zool.*, vol. 34, p. 205-383, pls. 90-121.

—————, 1914; *Ditto.*; the Clypeasterina; the Clypeasteridae,

Arachnoididae, Laganidae, Fibulariidae and Scutellidae.

*Ibid.*, vol. 46, no. 1, p. 1-80, pls. 122-143.

—————, 1917; *Ditto.*; the Echinoidae, Nucleolitidae,

Urechinidae, Echinocorythidae, Calymnidae, Pourtalesiidae,

Palaeostomatidae, Aeropsidae, palaeopneustidae and

Spatangidae. *Ibid.*, vol. 46, no. 2, p. 81-284, pls. 144

- , 1925; *Catalogue of the Recent sea-urchins (Echinoidea) in the collection of the British Museum*, p. 1-250, pls. 1-12. British Museum Natural History, London.
- Cooke, C. W., 1957; *Geology of Saipan, Mariana Islands, Part 3; Paleontology chapter J., Echinoids. U. S. G. S. Prof. Pap., 280J*, p. 361-364, pl. 119.
- Cotteau, G. H., 1885-1889; *Paléontologie Française ou description des fossils de la France. Terrain tertiaire, Échinides éocènes*, vol. 1, p. 1-669, pls. 1-200.
- Dörlein, L., 1885; *Seeigel von Japan und Liu-Kiu-Inseln. Arch. Naturg.*, vol. 51, Band 1, p. 73-112.
- , 1903; *Bericht über die von Herren Prof. Semon bei Amboina und Thursday Island gesammelten Echinoidea: in Semon, Forschungsreisen. Jena. Denkscher. Gesell.*, Band 8, p. 683-726, pls. 58-65.
- Fell, H. B., 1966; *Diadematacea. In, Moor, R. C. ed., Treaties on invertebrate paleontology, Part U, Echinodermata 3(1)*, p. U340-U366. Geol. Soc. Amer. and Univ. Kansas Press, Kansas.
- Fisher, A. G., 1966; *Spatangoids. In, Moor, R. C. ed., Treaties on invertebrate paleontology, Part U, Echinodermata 3(2)*, p.

- U543-U640. Geol. Soc. Amer. and Univ. Kansas Press, Kansas.
- Fujiyama, I., 1982; *Cenozoic echinoid*. In, Fujiyama, I., Hamada, T. and Yamagiwa, N. eds., *Encyclopedia of Japanese fossils*, p. 380-385, pls. 190-192, Hokuryukan, Tokyo. (*in Japanese, title translated*)
- Ghiold, J. B. and Hoffman, A., 1989; Biogeography of spatangoid echinoids. *N. Jb. Geol. Paläont. Abh.*, vol. 178, no. 1, p. 59-83.
- Hashimoto, W. and Shibata, M., 1960; A find of *Palaeopneustes* from Totsuka-ku, Yokohama City. *Trans. Proc. Palaeont. Soc. Japan*, N. S., no. 40, p. 337-338, pl. 38.
- Hatai, K, 1936; Neogene Brachiopoda from Japan. *Jour. Geol. Geogr.*, vol. 12, nos. 3-4, p. 283-324, pls. 34-35.
- , 1940; Cenozoic Brachiopoda from Japan. *Sci. Rep., Tohoku Imp. Univ., 2nd Ser.*, vol. 20, p. 222-253.
- Hayashi, T. and Miura, Y., 1973; The Cenozoic sediments in the southern part of Okazaki City, Central Japan. *Rep. Aichi Educ. Univ., (Nat. Sci.)*, vol. 22, no. 2, p. 133-150, pls. 1-2. (*in Japanese with English abstract*)
- Henderson, R. A., 1975; Cenozoic spatangoid echinoids from New Zealand. *N. Z. Geol. Surv., Paleont. Bull.*, no. 46, p. 1

-192, pls. 1-18.

Horikoshi, M., Ohta, S., Okiyama, M., Shigei, M., Imajima, M., Takeda, T., Gamo, S., Noda, H., Irimura, S., Nakamura, K., Hiruta, S., Kito, K., Hoshino, T. and Okimura, O., 1983; *Preliminary catalogue of benthic organisms collected at each station during various cruises of R/Vs Tansei-Marui and Hakuho-Marui, Ocean Research Institute, University of Tokyo (1966-1982)*. In Horikoshi, M., Ohta, S., Shirayama, Y. and Tsuchida, E. eds., p. 1-160, Ocean Res. Inst., Univ. Tokyo.

Ishii, Y., Uno, T., Oki, N., Omori, S., Kasai, K., Kurosawa, M., Komori K. and Nagashima, T., 1974; *Rocks and fossils in Ibaraki*. In Prof. Oyama, T. Mem.Vol., p. 69-141, pls. 1-31. Earth Sci. Educ. Ibaraki Univ., Mito. (*in Japanese, title translated*)

Kamei, T. and Kamiya, H., 1981; *On the fossil teeth of Stegolophodon pseudolatidens (Yabe) from the Miocene Bed of the Abukuma mountains. Mem. Fac. Sci., Kyoto Univ., (Geol. & Mineral), vol. 47, no. 2, p. 165-176, pls. 1-2.*

Kamiya, H., 1969; *Tertiary System along the Kuji River, Ibaraki Prefecture, especially on its facies change. Jour. Geol. Soc,*

Japan, vol. 75, no. 3, p. 157-170. (*in Japanese with English abstract*)

Kato, T., 1914; On elephant fossil from the Kuji-cho, Ibaraki Prefecture, Japan. *Jour. Geol. Soc. Japan*, vol. 21, no. 251, p. 345-349, pl. 18. (*in Japanese, title translated*)

Kier, P. M. and Lawson, M. H., 1978; Index of Living and Fossil echinoids 1924-1970. *Smith. Contr. Paleobiol.*, no. 34, p. 1-182.

Kikuchi, Y., 2000; *Cenozoic echinoid fossils from the Okinawa-jima, Okinawa Prefecture, Southwestern Japan*. p. 11-45, pls. 1-8. Report of the Science Research Funds 1997-1999 by the Japanese Ministry of Education. (*in Japanese with English abstract*)

————— and Nikaido, A., 1985; The first occurrence of abyssal echinoid *Pourtalesia* from the Middle Miocene Tatsukuroiso Mudstone Member in Ibaraki Prefecture, northeastern Honshu, Japan. *Ann. Rept., Inst. Geosci., Univ. Tsukuba*, no. 11, p. 32-34.

————— and —————, 1987; Gregarious occurrence of echinoid *Linthia tokunagai* from the Pliocene Kume Formation in Ibaraki Prefecture, northeastern Honshu, Japan. *Ibid.*, no. 13, p. 87-91.

————— and —————, 1996; Miocene holothurian fossils from the middle Miocene Nawashiroda Formation in Ibaraki Prefecture, northern Kanto, Japan. *Prof.H.Igo Commem. Vol.*, p. 155-162.

—————, ————— and Noda, H., 1992; Occurrence of marine accretional lapilli from the Pliocene Hanareyama Formation in the northern part of Ibaraki Prefecture, Japan. *Ann. Rept., Inst. Geosci., Univ. Tsukuba*, no. 18, p. 48-52.

—————, ————— and Sugaya, M., 1991; Hard substratum boring shells from the Pliocene Kume Formation in Ibaraki Prefecture, northern Kanto, Japan. *Ibid.*, no. 17, p. 48-51.

Kinoshita, K., 1935; *Geological map of the 1/75,000 Sukegawa*. Geol. Surv. Japan. (in Japanese)

Kochibe, T., 1892; *Geology of Jyohoku (Jyohoku Chishitsu Hen)*. *Rika Kai-shi, Univ. Tokyo*, no. 4, p. 1-153, pls. 1-7. (in Japanese title translated)

Koda, Y., Yanagisawa, Y., Hasegawa., Y., Otsuka H. and Aizawa, M., 2003; A middle Miocene mandible of *Stegolophodon* (Proboscidea, Mammalia) discovered in Katsura Village, Ibaraki Prefecture, eastern Japan. *Earth Science (Chikyu Kagaku)*, vol. 57, nos. 1/2, p. 49-59. (in Japanese with English



Abstract).

Koehler, R., 1914; An account of the Echinoidea. *Echinoderma of Indian Museum, part. VIII, Echinoidea*, no. 1, p. 1-258, pls.1-20.

Koehler, R., 1922; *Ditto. Ibid, Part IV, Echinoidea*, no. 2, p. 1-167, pls. 1-15.

Koizumi, I., 1973; The stratigraphic ranges of marine planktonic diatoms and diatom biostratigraphy in Japan. *Mem. Geol. Soc. Japan*, no. 8, p. 35-44. (in Japanese with English abstract)

——— and Terunuma, Y., 1985; Geologic age, based on diatoms, of Odontoceti (Physeteridae, Mammalia) from the Taga Formation at Nukata-nango, Ibaraki Prefecture. *Jour. Geol. Soc. Japan*. vol.91, no. 11, p. 805-809. (in Japanese)

Lambert, J. and Jeannet, A., 1935; Contribution à l' étude des Échinides tertiaires des Iles de la Sonde. *Mém. Soc. Paléont. Suisse*, vol. 56, p. 1-72, pl. 1-4.

——— and Thiély, P., 1925; *Essai de nomenclature raisonnée des Échinides*. fasc. 8-9, p. 513-607, pls. 12-13, 15.

Liao, Y., 1987; The Echinoderms of the Xisha Islands Guandong Province, China. III, Echinoidea. *Stud. Mar. Sinica*, vol. 12, p. 107-127, pls. 1-5. (in Chinese, with English abstract)

- Lovén, S., 1883; On *Pourtalesia*, a genus of Echinoidea. *Kongl. Svenska Vetenskaps-Akade. Handlingar*, vol. 19, no. 7, p. 1-96, pls. 1-21.
- Maruyama, T., 1984; Miocene diatom biostratigraphy of on shore sequence on the Pacific side on northeast Japan with reference to DSDP H438A (Part 2). *Tohoku Univ., Sci. Rep., 2nd Ser. (Geol.)*, vol. 55, no. 1, p. 77-140, pls. 11-15.
- Masuda, K., 1953; A new species of *Patinopecten* from Ibaraki Prefecture. *Short Paper, IGPS*, no. 5, p. 41-50, pls. 5-6.
- Matsuura, N., Fujii, S., Sekido, S. and Kawai, A., 1984; Data of fossils in Kaga, Ishikawa Prefecture, central Japan. *Rep., Ishikawa Prefectural Education Center*, no. 23, p. 1-54, pls. 3-7. (in Japanese, title translated)
- , and Hotsuta, O., 1986; Data of fossils in Noto, Ishikawa Prefecture, central Japan. *Ibid.*, no. 27, p. 1-30, pls. 2-4. (in Japanese title translated)
- Mazzetti, A. G., 1983; Catalogo degli Echinidi del Mar Rosso e descrizione di specie nuove. *Atti d. Soc. d. Natural. d. Modena, ser. 3*, vol. 12, p. 238-242.
- , 1894; Gli Echinidi del Mar. Rosso. *Mem. Real. Accad. di Sci. Lettr, Art. Modena, ser. 2*, vol. 10, p. 211-228.

- Meijere, J. C. H. De., 1904; Die Echinoidea der Siboga-Expedition.  
*Siboga-Exped.Monogr.*, vol. 43, p. 1-252, pls. 1-23.
- Minato, M., 1949; Fossil echinoderms in Hokkaido. *Cenozoic Res.*, no. 1, p. 14-17. (*in Japanese, title translated*)
- , 1950; On some Paleogene fossils in Hokkaido. *Jour. Geol. Soc. Japan*, vol. 56, no. 655, p. 157-159.
- Mironov, A. N., 1980; Two modes of formation of deep-sea echinoid fauna. *Oceanol.*, vol.20, no, 4. p. 462-465.
- Mizuno, Y., 1992; Fossil echinoids from the Miocene Morozaki Group. *Bull. Mizunami Foss. Mus.*, no. 19, p. 337-346, pls. 52-53. (*in Japanese, with English abstract*)
- , 1993; Echinoids fossil from the Morozaki Group. In, *Fossils from the Morozaki Group*. P. 145-156, pls. 1-4. Tokai Fossil Society Nagoya. (*in Japanese with English abstract*)
- , Hachiya, K. and Yamaoka, M., 1989; Some fossil echinoids from the Miocene Morozaki Group in the Chita Peninsula, central Japan. *Kaseki-no-Tomo*, no. 35, p. 1-19, pls. 1-4. (*in Japanese*)
- Morishita, A., 1953a; Fossil species of Palaeopneustidae from Japan. *Trans. proc. Palaeont. Soc. Japan, N.S.*, no. 9, p. 27-29, pl. 1.

- , 1953b; Neogene echinoids from Gifu Prefecture, Japan.  
*Ibid.*, no. 10, p. 61-64, pl. 6.
- , 1953c; On some Neogene echinoids from Nagano Prefecture,  
Japan. *Mem. Coll. Sci., Univ. Kyoto, Ser. B*, vol. 20, no.4,  
art. 1, p. 217-226, pl. 1.
- , 1954; Tertiary echinoids from the environs of Ise-Bay.  
*Ibid.*, vol. 21, no. 2, art. 8, p. 223-230, pl. 7.
- , 1956; On some fossil echinoids from Kyushu, Japan.  
*Ibid.*, vol. 23, no. 2, art. 3, p. 193-202, pls. 1-4.
- , 1967; *Fossil echinoids from Anan-Cho*. In, *Fossils from  
Anan-Cho*. p. 105-110, pls. 17-19. Bord of Educt. Anan-Cho,  
Nagano Prefecture. (*in Japanese, title translated*)
- , 1969; On some echinoids found in Suruga Bay. *Publ. Seto  
Mar. Biol. Lab.*, Vol. 16, no. 6, p. 363-379, pls. 21-25.
- , 1974; Miocene echinoids from the Mizunami district.  
*Bull. Mizunami Foss. Mus.*, no. 1, p. 205-214, pls. 64-66.  
(*in Japanese, title translated*)
- , 1983; Fossil species of *Palaeopneustes* from Japan.  
*Ibid.*, no. 10, p. 103-106, pls. 28-31.
- and Itoigawa, J., 1986; *Illustrated paleoecology*. p.  
1-173. Asakura-Shoten, Tokyo (*in Japanese*)

- Mortensen, Th., 1904; The Danish Expedition to Siam, 1899-1900: II, Echinoidea; Part 1. *Kongl. Dansk. Vidensk. Selsk. Skr., ser. 7*, vol. 7, no. 1, p. 1-124, pls. 1-7.
- , 1907; Echinoidea; Part 2. *Danish Inglof-Expedition*, vol. 4, no. 2, p. 1-200, pls. 1-19.
- , 1943; A monograph of the Echinoidea: Vol. 3, pt. 1, Aulodonta, with Additions to Vol. 2 (*Lepidocentroida and Stirodonta*), p. 1-370, pls. 1-77. Köbenhavn and London.
- , 1948a; Ditto: Vol. 4, pt. 1; *Holectypoida, Cassiduloida*, p. 1-363, pls. 1-14. Köbenhavn and London.
- , 1948b; Ditto: Vol. 4, pt. 2; *Clypeastroidea, Arachnoididae, Fibulariidae, Laganidae and Scutellidae*, p. 1-471, pls. 1-72. Köbenhavn and London.
- , 1948c; Report on the Echinoidea collected by the United States Fisheries Steamer "Albatross" during the Philippine Expedition, 1907-1910: Part 3, The Echinoneidae, Echinolampadidae, Clypeastridae, Arachnoididae, Laganidae, Fibulariidae, Urechinidae, Echinocorythidae, Palaeostomatidae, Micrasteridae, Palaeopneustidae, Hemiasteridae and Spatangidae. *U. S. Nat. Mus., Bull.* vol. 14, pt. 3, p. 93-140.

—————, 1950; *A monograph of the Echinoidea: Vol. 5, pt. 1; Spatangoida I; Protosternata, Meridosternata, Amphisternata I; Palaeopneustidae, Palaeostomatidae, Aeropsidae, Toxasteridae, Micrasteridae and Hemiasteridae*, p. 1-422, pls. 1-25. Köbenhavn and London.

—————, 1951; *Ditto: Vol. 5, pt. 2; Spatangoida II; Amphisternata II; Spatagidae, Loveniidae, Pericosmidae, Shizasteridae and Brissidae*, p. 1-593, pls. 1-64. Köbenhavn and London.

Nagao, T., 1928; *Palaeogene fossils of the Island of Kyushu, Japan: Part 2. Sci. Rept., Tohoku Imp. Unive., 2 nd Ser. (Geol.)*, Vol. 12, no. 1, p. 11-140, pls. 1-7.

Nakashima, T., 1992MS; *Geology of the Yamagata district, southern part of the Tanakura Sheared Zone in Ibaraki Prefecture. The Graduation Thesis of the Collage of Natural Science, the University of Tsukuba*, p. 1-64, pls. 1-15. (*in Japanese with English abstract*)

Nathorst, A. G., 1883; *Contribution à la flore fossile du Japon. Kungl. Svenska Vet. Akad. Handle.*, vol. 20, p. 3-92, pls. 1-12.

Nikaido, A. and Kikuchi, Y., 1983; *Some vertebrae of shark from the Miocene Nawashiroda Formation, Ibaraki Prefecture*,

- northeast Japan. *Jour. Geol. Soc. Japan*, vol. 89, no. 5, p. 299-301. (in Japanese)
- Nishimura, S., 1974; *Completion of the Sea of Japan: The approach from Biogeography*. p. 1-236. Tsukiji Syokan, Tokyo. (in Japanese, title translated)
- , 1981; *Sea and life of the earth*. p. 1-284. Kaimei Sha, Tokyo. (in Japanese, title translated)
- Nishio, T., 1961; On some Pliocene echinoids from Chiba Prefecture. *Bull. Tokyo Gakugei Univ.*, no. 12, p. 129-131, pl. 1.
- Nisiyama, S., 1936; On the occurrence of *Temnotrema rubrum* in the Pleistocene of Tsurumi, Kwanto Region. *Jour. Geol. Soc. Japan*, vol. 43, no. 509, p. 122-128, pl. 1.
- , 1966; The echinoid fauna from Japan and adjacent regions, Part 1. *Palaeont. Soc. Japan, Spec. Pap.*, no. 11, p. 1-277, pls. 1-18.
- , 1968; The echinoid fauna from Japan and adjacent regions, Part 2. *Ibid.*, no. 13, p. 1-491, pls. 19-30.
- Noda, H., 1973; Cenozoic Arcidae of Japan. *Sci. Rep., Tohoku Univ.*, 2nd Ser., Geol., Spec. Vol., no. 6, p. 205-215.
- , 1975; Turriculid gastropoda of Japan. *Ibid.*, vol. 45, no. 2, p. 51-82, pls. 9-12.

————— and Amano, K., 1977; Geological significance of *Anadara amacula elongata* from the Pliocene Kume Formation, Ibaraki Prefecture, Japan. *Ann. Rep., Inst. Geosci., Univ. Tsukuba*, no. 10, p. 37-41.

————— and Kikuchi, Y., 1980; *Phanerolepida expansilabrum* (Kuroda) (Gastropoda) from the Miocene Nantaisan Volcanic Breccia, Ibaraki Prefecture, Northeastern Japan. *Venus (Japan Jour. Malac.)*, vol. 39, no. 1, p. 69-73, pl. 1. (in Japanese with English abstract)

————— and —————, 1995; Miocene *Aturia* (Cephalopoda, Mollusca) from Ibaraki, Northern Kanto, Japan. *Ann. Rep., Inst. Geosci., Univ. Tsukuba*, no. 21, p. 27-31.

—————, ————— and Nikaido, A., 1989; Paleobiographical significance of the first occurrence of the extinct Miocene genus *Acilana* (Mollusca; Bivalvia) from the Pliocene Kume Formation in Ibaraki Prefecture, northern Kanto, Japan. *Prof H. Matsuo Mem. Vol.*, p. 53-64, pl. 1. (in Japanese with English abstract)

—————, ————— and —————, 1993; Molluscan fossils from the Pliocene Kume Formation in Ibaraki Prefecture, Northeastern Kanto, Japan. *Sci. Rep., Inst. Geosci., Univ. Tsukuba, Sec.*



B, vol. 14, p. 115-204.

———, ——— and ———, 1994; Middle Miocene molluscan fauna from the Tamagawa Formation in Ibaraki Prefecture, northern Kanto, Japan — Arcid-Potamid Fauna in the Tanakura Tectonic Zone —. *Ibid.*, vol. 15, p. 81-102.

———, Watanabe, R. and Kikuchi, Y., 1995; Pliocene marine molluscan fauna from the Hitachi Formation in the northeastern part of Ibaraki Prefecture, Japan. *Ibid.*, vol. 16, p. 39-93.

Ogasawara, K., 1993; Neogene paleogeography and marine climate of Japanese Islands based on shallow-marine molluscs. *Palaeogeogr. Palaeoclimat. Palaeoecol.*, vol. 108, no. 1994, p. 335-351.

———, 2001; Notes on paleoceanographic background for change of the Japanese Cenozoic molluscan faunas. *Sebutsu-Kagaku*, vol. 53, no. 3, p. 185-191. (in Japanese)

——— and Masuda, K., 1989; Paleobathymetric indexes of the Neogene molluscs in Tohoku District and their implications. *Mem. Geol. Soc. Japan*, no. 32, p. 217-227.

——— and Nagasawa, K., 1992; Tropical molluscan associations in the Middle Miocene marginal sea of the Japanese Islands:

an example of molluscs from the Oyama Formation, Tsuruoka City, northeast Honshu, Japan. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, vol. 167, p. 1224-1246.

Ohyama, T. and Sakurai, K., 1966; On the Argonautinae from the Sakachi Formation, Neogene Tertiary in Ibaraki. *Bull. Fac. Lib. Arts. Sci., Ibaraki Univ., Nat. Sci.*, no. 17, p. 29-37, pl. 1. (in Japanese with English abstract)

Okamoto, K., Katsuhara, M., Ueno, Y. and Sumiyoshi, O., 1990; Molluscan assemblages from the Miocene Bihoku Group in the Kaisekidani Area, Myauchi-cho, Shobara City, Southwest Japan — Study of the Bihoku Group III —. *Bull. Mizunami Foss. Mus.*, no. 17, p. 35-49, pls. 9-11. (in Japanese with English abstract)

Omori, M., 1948; structure of the Tertiary deposits along the drainage of the Kuji-gawa River. *Jour. Geol. Soc. Japan*, vol. 54, no. 638, p. 151. (in Japanese)

———, 1958; On the geological history of the Tertiary System in the southern part of the Abukuma mountainland with special reference to the geological meaning of the Tanakura Sheared Zone. *Sci. Rep., Tokyo Kyoiku Daigaku. Sec. C*, no. 6, p. 75-83. (in Japanese with English abstract)

————— and Suzuki, K., 1950; Stratigraphical study of the Taga Series in south margin of the Abukuma Plateau (Study on Cenozoic History of southwestern margin of Abukuma Plateau, Part 1). *Jour. Geol. Soc. Japan*, vol. 56, no. 658, p. 369-378. (in Japanese with English abstract)

—————, Horikoshi, K., Suzuki, K. and Fujita, Y., 1953; On the Tanakura Sheared Zone in the SW margin of the Abukuma Mountainland (Study of Cenozoic history of southwestern margin of Abukuma Plateau, Part 3). *Jour. Geol. Soc. Japan*, vol. 59, no. 693, p. 217-223. (in Japanese with English abstract)

Omori, N. and Owada, T., 1985; Late Cenozoic gravel bed found in the southernmost Abukuma Plateau containing gravels with burrows of boring shells. *Jour. Geol. Soc. Japan*. vol. 91, no. 7, p. 477-479. (in Japanese)

Otsuki, K., 1975; Geology of the Tanakura Shear Zone and adjacent area. *Contr. Inst. Geol. Paleont., Tohoku Univ.*, no. 76, 1-71. (in Japanese with English abstract)

Oyama, T., 1960a; Miocene fossil plants from the Kamikanazawa, Daigo-machi, Ibaraki Prefecture, Japan. *Tohoku Univ. Sci. Rept., Spec. Vol.*, 4, p. 488-490. (in Japanese, with English

abstract)

———, 1960b; On the conclusion of the Oarai Flora from the Upper Cretaceous Oarai Formation in Ibaraki Prefecture, Japan, Part 1. *Bull. Fac. Lib. Arts, Sci., Ibaraki Univ., Nat. Sci.*, no. 11, p. 75-105.

———, 1961; *Ditto*, (Part 2). *Ibid.*, no. 12, no. 61-102.

Ozaki, H., 1958; Stratigraphical and paleontological studies on the Neogene and Pleistocene formations of the Tyosi district. *Bull. Natn. Sci. Mus.*, vol. 4, no. 1, p. 1-182, pls. 1-24.

———, and Saito, T., 1954; Stratigraphical studies of the Tertiary System in the environs of Ota-machi, Kuji-gun, Ibaraki Prefecture (Geology of Ibaraki Prefecture, Part 2). *Bull. Fac. Lib. Arts, Ibaraki Univ., Nat. Sci.*, no. 4, p. 87-94. (in Japanese with English abstract)

——— and ———, 1955; The Cretaceous System along the coast of Nakaminato City, Ibaraki Prefecture (Geology of Ibaraki Prefecture, Part 3). *ibid.*, no. 5, p. 37-49, pl. 1. (in Japanese with English abstract)

Rowe, F. W. E. and Gates, J., 1995; Echinodermata. In Wells, A. ed., *Zoological Catalogue of Australia*. vol. 33, p. 1-510.

Saba, M., Tomida, Y. and Kimoto, T., 1982; Echinoderm fauna of Ise

Bay, and the northern and the middle parts of Kumano-nada.  
*Bull. Mie. Pref. Mus., Nat. Sci.*, no. 4, p. 1-82, pls. 1-34.  
(in Japanese with English abstract)

Saito, T., 1952; On the stratigraphical order of the vicinity of Ota-machi, Ibaraki Prefecture, part 1. *Bull. Fac. Lib. Arts, Sci., Ibaraki Univ., Nat. Sci.*, no. 2, p. 129-148. (in Japanese with English abstract)

———, 1956; The Tertiary System in the environs of Nakaminato City (Geology of Ibaraki Prefecture, Part 4). *Ibid.*, no. 6, p. 39-51. (in Japanese with English abstract)

———, 1959; Notes on some Cretaceous fossils from the Nakaminato City, Ibaraki Prefecture, Japan, Part 1. *Ibid.*, no. 8, p. 83-94.

———, 1959; *Ditto.*, *Ibid.*, no. 9, p. 79-85. (in Japanese with English abstract)

———, 1961; The Upper Cretaceous System of Ibaraki and Fukushima Prefectures, Japan. Part. 1. *Ibid.*, no. 12, p. 103-144.

———, Takahashi, H. and Amano, K., 1992; *Geological map of the Neogene along Tanakura Fault*. 1 sheet. *Fac. Sci., Ibaraki Univ., Mito.* (in Japanese)

Sakamoto, T., Tanaka, K., Soya, T., Noma, T. and Matsuno, K., 1972;

- Geology of the Nakaminato district.* Geol. Surv. Japan, Quadrangle Ser., 1/50,000, Tokyo 8, p. 1-94, and 1map.
- Shigei, M., 1973; A check list of echinoids found in Sagami Bay with brief note on each species. *Jour. Fac. Sci., Univ. Tokyo, Sec. 4*, no. 13, p. 1-33.
- , 1974; *Echinoids*. In, Uchida, T. ed., *Systematic zoology*. vol. 8, no. 2, p. 208-332. Nakayama Book Co., Tokyo. (in Japanese title translated)
- , 1975; A new species of the heart-urchin, an extant species of *Brissopatagus* (Echinoidea; Spatangoida), from Sagami Bay. *Jour. Fac. Sci., Univ. Tokyo, Ser. 4*, vol. 13, p. 333-339.
- , 1981; A study on the echinoid fauna of the East China Sea and the coastal water of southern Korea, Kyushu, Ryukyu and Taiwan. *Publ. Seto Mar. Biol. Lab.*, vol. 26, nos. 1-3, p. 192-241.
- , 1986; *The sea urchins of Sagami Bay*. p. 1-204, (in English), p. 1-173, (in Japanese), pls. 1-126. Maruzen Book Co., Tokyo.
- Shikama, T. and Omori, M., 1952; Notes on an occurrence of *Dicrocerus* in the Daigo Group of the Ibaraki Prefecture, Japan. *Proc. Japan Acad.*, vol. 28, no. 10, p. 567-572.
- and Kase, T., 1976; Molluscan fauna of the Miocene Morozaki

Group in the southern part of Chita Peninsula, Aich Prefecture,  
Japan. *Sci. Rep., Yokohama Natn. Univ., Sec. 2*, no. 23, p.  
1-25, pls. 1-2.

Smith, A., 1984; *Echinoid paleobiology*. p. 1-190. George Allen &  
Union, London.

Sudo, I., Takahashi, M. and Yanagisawa Y., 2002; Diatom  
biostratigraphy of the Miocene Tsuchishio Formation in the  
Hiki Hills area (Aketo Section), Saitama Prefecture, central  
Japan. *Jour. Geol. Soc. Japan*, vol. 109, no. 1, p. 48-62.  
(in Japanese with English abstract)

Suzuki, K., 1954a; A study of the sedimentation on the Neogene System  
in the southern part of the Joban Coal Field. *Misc. Rep.,  
Res. Inst. Nat. Resour.*, no. 35, p. 14-27, (in Japanese with  
English abstract)

———, 1954b; Stratigraphical study of the Neogene System in the  
neighbourhood of Abukuma Plateau, Part 1. *Stud. Geol. Miner.,  
Inst. Tokyo Univ. Educ.*, no. 3, p. 75-96. (in Japanese with  
English abstract)

———, and Omori, M., 1953; Stratigraphic study of Tertiary in  
Ohta-machi, Ibaraki Prefecture (Historical study of Cenozoic  
in southwestern part of the Abukuma Mountains, Part 2). *Jour.*

*Geol. Soc. Japan*, vol. 59, no. 589, p. 34-46. (in Japanese with French résumé, title translated)

Takada, T., 1999MS; *Biostratigraphical study by Tertiary diatom - for the explanation of teaching materials -*. Master Thesis of Program, Master of Arts in Education, the University of Tsukuba, p. 1-87. (in Japanese)

Takahashi, H., 1984; *Miocene faunal and floral facies in the area of the Tanakura Sheared Zone*. Rep. Tokutei-Kenkyu, Ibaraki Univ., p. 10-15. (in Japanese, title translated)

————— and Amano, K., 1984; *Miocene transgression in and around the Tanakura Shear Zone*. *Bull. Coll. General Educ., Ibaraki Univ.*, no. 16, p. 149-162.

Takahashi, H., 1986; *Characteristics of the molluscan assemblage in the Pliocene Hitachi-ota area, Ibaraki Prefecture, central Japan*. *Monogr. Mizunami Foss. Mus.*, no. 6, p. 91-103, pls. 12-13. (in Japanese with English abstract)

—————, 2001; *Arcid-Potamid assemblage from Miocene formation in the Tanakura Shear Zone*. *Biological Science (Tokyo)*, vol. 53, no. 3, p.168-177. (in Japanese with English abstract)

Takahashi, M. and Ikeda, Y., 1984; *Tertiary Volcanic rocks of the*



- Tanakura *Sheared Zone and adjacent regions*. Rep. Tokutei-Kenkyu, Ibaraki Univ., p. 16-17. (*in Japanese, title translated*)
- Tokunaga, S., 1900; Japanese echini. *Zool. Mag. Tokyo*, vol.12, no. 145, p. 379-405. (*in Japanese, title translated*)
- , 1903; On the fossil echinoids of Japan. *Jour. Coll. Sci., Imp. Univ. Tokyo*, vol. 17, art. 12, p. 1-27, pls. 1-4.
- , 1906a; Japanese echini: Plate. *Zool. Mag. Tokyo*, vol. 18, pls. 8-3. (*in Japanese*)
- , 1908; Japanese echini. *Ibid.*, vol. 20, no. 235, pls. 20-12. (*in Japanese, title translated*)
- , 1927a; A fossil asteroid found in the Tertiary strata of Hitachi Prov. *Annot. Zool. Japon*, vol. 12, no. 1, p. 323-326.
- , 1927b; Geology of Joban coal-field. *Mem. fac. Sci., Eng. Waseda Univ.*, no. 5, p. 1-136, pls. 1-10. (*in Japanese*)
- Tomida, S. and Itoigawa, J., 1989; Occurrence of *Harutungia* (Gastropoda; Janthinidae) from the Pliocene Hatsuzaki Group in Hitachi City, Ibaraki Prefecture. *Bull. Mizunami Foss. Mus.*, no. 16, p. 125-129, pl. 23.
- Utinomi, H., 1954; A check list of echinoids found in the Kii region. *Publ. Seto Mar. Biol. Lab.*, vol. 3, no. 3, p. 339-358.

- , 1960; Echinoids from Hokkaido and the neighbouring subarctic water. *Ibid.*, vol. 8, no. 2, p. 337-349, pls. 37-40.
- Watanabe, R., 1993MS; *Taphonomical study of boring fossils — Microboring fossils from Hatsuzaki, Hitachi City, Ibaraki Prefecture, northeastern Japan —*. Master Thesis of Program in Geoscience, University of Tsukuba, p. 1-47, pls. 1-11. (in Japanese with English abstract)
- Yabe, H., 1950; Controversies relation to the Kuji Proboscidean molars. *Proc. Japan Acad.*, vol. 26, no. 8, p. 29-35.
- Yamada, H., 1888; *Geological map of the 1/200,000 Mito*. Geol. Surv. Japan. (in Japanese)
- Yamana, I., 1978; Catalog of fossils (33) — Fossils of Tottori Group and Tari Formation —. *Miscel. Rep., Tottori Pref. Mus.*, no. 19, p. 4-36, pls. 1-6. (in Japanese, title translated)
- Yanagisawa, Y., 1990; Age of fossil Argonautidae (Cephalopoda) from Hokuriku Province (central Honshu, Japan) based on diatom biostratigraphy. *Bull. Geol. Surv. Japan*, vol. 41, no. 3, p. 115-127, pl. 1. (in Japanese with English abstract).
- and Akiba, F., 1998; Refined Neogene diatom biostratigraphy for the northwest Pacific around Japan, with an introduction of code numbers for selected diatom bio-

horizons. *Jour. Geol. Soc. Japan*, vol. 104, no. 6, p. 395  
-414.

———, Nakamura, K., Suzuki, Y., Sawamura, K., Yoshida, F.,  
Tanaka, Y., Honda, Y. and Tanahashi, M., 1989; Tertiary  
biostratigraphy and subsurface geology of the Futaba  
district, Joban Coalfield, northeast Japan. *Bull. Geol.  
Surv. Japan*, vol. 40, no. 8, p. 405-467, pls. 1-12. (*in  
Japanese with English abstract*)

Yoshioka, T., Takizawa, F., Takahashi, M., Miyazaki, K., Banno, Y.,  
Yanagisawa, Y., Kubo, K., Seki, Y., Komazawa, M. and  
Hiroshima, T., 2001; *Geological map of Japan 1:200,000,  
Mito, 2nd Edition*). 1 sheet. Geological Survey of Japan.  
(*in Japanese with English abstract*)

Yoshiwara, S., 1899; On some new fossil echinoids of Japan. *Jour.  
Geol. Soc. Japan*, vol. 6, no. 65, p. 1-4, pl. 2.