

Acknowledgements

The present author wishes to express his hearty gratitudes to Professor Hiroshi Noda of the University of Tsukuba and Dr. Hisao Nakagawa, former Professor of Tohoku University, for their criticism and advice on this work and gave him great encouragements.

The author is grateful to Professor Kenshiro Ogasawara of the University of Tsukuba, Dr. Kunihiro Ishizaki, Professor Emeritus of Tohoku University and Professor Kenshiro Otsuki of Tohoku University for their valuable suggestions to complete the present paper.

In preparation of this paper, the author had a great deal of assistance from Dr. Takeshi Matsui, Professor Emeritus of Nippon University, Professor Yo Uesugi of Tsuru University, Professor Motoyoshi Oda of Kumamoto University, Dr. Sadako Takeuti of Saito Ho-on-kai Museum of Natural History, Mr. Shizuo Oda of the Division of Social Education of Tokyo Metropolis, and Mr. Mamoru Hosono of the Tokyo Shizenshi-kiko.

The author thanks Mr. Koichi Minagawa, an independent enterpriser, for his thermal demagnetization measurement of ferromagnetic minerals in the Sanshokuaisu Pumice (SIP), the Kyoto Fission Track & Co., Ltd. for the fission track age measurements of zircons in the Biotite bearing Ash Zone (Bi) and Hachioji Biotite Pumice (HBP).

The present author would like to appreciate to Messrs. Hiroyuki Ishikawa of the Historical Research Center of Saitama Prefecture, Itsuo Masuda of the Archaeological Center of Saitama Prefecture, Kozo Yoshida, Osamu Sakamoto, Takahisa Nishida, Ki'ichi Obata and Takashi Nirei, all staffs of the Saitama Prefectural Museum of Natural History for their valuable supports to this paper.