

## References

- (1) Adrian, K., Bernhard, M. K., Breiting, H., Ogilvie, A. Expression of purinergic receptors (ionotropic P2X1-7 and metabotropic P2Y1-11) during myeloid differentiation of HL60 cells. *Biochim. Biophys. Acta.*, 1492, 127-138, 2000.
- (2) Asahi, K., Honma, Y., Hazeki, K., Sassa, T., Kubohara, Y., Sakurai, A., Takahashi, N. Cotylenin A, a plant-growth regulator, induces the differentiation in murine and human myeloid leukemia cells. *Biochem. Biophys. Res. Commun.*, 238, 758-763, 1997.
- (3) Asahi, K., Sakurai, A., Takahashi, N., Kubohara, Y., Okamoto, K., Tanaka, Y. DIF, morphogen of *Dictyostelium discoideum*, induces the erythroid differentiation in murine and human leukemia cells. *Biochem. Biophys. Res. Commun.*, 208, 1036-1039, 1995.
- (4) Barcizewski, J., Mielcarek, M., Stobiecki, M., Siboska, G., Clark, B. F. Identification of 6-furfuryladenine (kinetin) in human urine. *Biochem. Biophys. Res. Commun.*, 279(1), 69-73, 2000.
- (5) Barciszewski, J., Siboska, G.E., Pedersen, B.O., Clark, B.F.C., Rattan, S.I.S. A mechanism for the in vivo formation of N<sup>6</sup>-furfuryladenine, kinetin, as a secondary oxidative damage product of DNA. *FEBS Lett.*, 414, 457-460, 1997.
- (6) Becker, T., Gerke, V., Kube, E., Weber, K. S100P, a novel Ca (2+)-binding protein from human placenta. cDNA cloning, recombinant protein expression and Ca<sup>2+</sup> binding properties. *Eur. J. Biochem.*, 207(2), 541-547, 1992.
- (7) Bierregaard, M. D., Jurlander, J., Klausen, P., Borregaard, N., Cowland, J. B. The in vitro profile of transcription factors during neutrophil

- differentiation in human bone marrow. *Blood*, 101(11), 4322-4332, 2003.
- (8) Bindokas, V.P., Jordan, J., Lee, C.C., Miller, R.J. Superoxide production in rat hippocampal neurons: selective imaging with hydroethidine. *J. Neurosci.*, 16, 1324-1336, 1996.
- (9) Brault, M., Caiveau, O., Pendron, J., Maldiney, R., Sotta, B., Miginiac, E. Deterction of membrane-bound cytokinin-binding proteins in *Arabidopsis thaliana* cells. *Eur. J. Biochem.* 260, 512-519, 1999.
- (10) Brinster, R.L. The effect of cells transferred into the mouse blastocyst on subsequent development. *J. Exp. Med.*, 140, 1049-1056, 1974.
- (11) Bruce, M.I., Zwar, J.A., Kefford, N. P. Chemical structure and plant kinin activity - the activity of urea and thiourea derivatives. *Life Sci.*, 4, 461-466, 1965.
- (12) Brumm, A.F., Potter, V.R., Siekevitz, P. Nucleotide metabolism V. Incorporation of P<sup>32</sup> into acid-soluble nucleotides of rat liver *in vivo*. *J. Biol. Chem.*, 22, 13-722, 1956.
- (13) Buck, M., Griffiths, E. Regulation of aromatic amino acid transport by tRNA: role of 2-methylthio-N<sup>6</sup>-( $\Delta^2$ -isopentenyl)-adenosine. *Nucleic Acid Res.*, 9, 401-414, 1981.
- (14) Budd, R. C. Activation-induced cell death. *Curr. Opin. Immunol.*, 13 (3), 356-362, 2001.
- (15) Buell, G., Michel, A. D., Lewis, C., Collo, G., Humphrey, P. P. A., Surprenant, A. P2X1 receptor activation in HL60 cells. *Blood*, 87, 2659-2664, 1996.
- (16) Camici, M., Turriani, M., Tozzi, M.G., Turchi, G., Cos, J., Alemany, C., Miralles, A., Noe, V., Ciudad, C. Purine enzyme profile in human colon-carcinoma cell lines and differential sensitivity to deoxycoformycin and 2'-deoxyadenosine in combination. *Int. J. Cancer*, 62, 176-183, 1995.

- (17) Castaigne, S., Chomienne, C., Daniel, M.T., Ballerini, P., Berger, R., Fenaux, P., Degos, L. All-*trans* retinoic acid as a differentiation therapy for acute promyelocytic leukemia. I. Clinical results, *Blood*, 76, 1704-1709, 1990.
- (18) Chomeczynski, P. A reagent for the single-step simultaneous isolation of RNA, DNA and proteins from cell and tissue samples, *Biotechniques*, 15:532-534, 1993.
- (19) Collins, S.J., Bonder, A., Ting, R., Gallo, R.C. Induction of morphological and functional differentiation of human promyelocytic leukemia cells (HL-60) by compounds which induce differentiation of murine leukemia cells. *Int. J. Cancer.*, 25: 213-218, 1980.
- (20) Collins, S.J., Gallo, R.C., Gallagher, R. E. Continuous growth and differentiation of human myeloid leukemia cells in suspension culture. *Nature*, 270, 347-349, 1977.
- (21) Crans, H, N., Sakamoto, K. M. Transcription factors and translocations in lymphoid and myeloid leukemia. *Leukemia*, 15(3), 313-331, 2001.
- (22) Devery, J. M., King, N. J., Geczy, C. L. Acute inflammatory activity of the S100 protein CP-10: activation of neutrophils in vivo and in vitro. *J. Immunol.*, 152, 1888-1897, 1994.
- (23) Donato, R. Functional roles of S100 proteins, calcium-binding proteins of the EF-hand type. *Biochim. Biophys. Acta.*, 1450, 191-231, 1999.
- (24) Donato, R. S100: a multigenic family of calcium-modulated proteins of the EF-hand type with intracellular and extracellular functional roles. *Int. J. Biochem. Cell. Biol.*, 33 (7), 637-668, 2001.
- (25) Emoto, Y., Kobayashi, R., Akatsuka, H., Hidaka, H. Purification and characterization of a new member of the S100 protein family from human placenta. *Biochem. Biophys. Res. Commun.*, 182, 1246-1253, 1992.
- (26) Fajkus, J., Fulneckova, J., Hulanova, M., Berkova, K., Riha, K., Matyasek,

- R. Plant cells express telomerase activity upon transfer to callus culture, without extensively changing telomere lengths. *Mol. Gen. Genet.*, 260, 470-474, 1998.
- (27) Fibach, E., Peled, T., Treves, A., Kornberg, A., Rachmilewitz, E.A., Modulation of the maturation of human leukemic promyelocytes (HL-60) to granulocytes or macrophages. *Leuk. Res.*, 6 (6), 781-790, 1982.
- (28) French, B. T., Patrick, D. E., Grever, M. R., Trewyn, R. W. Queuine, a tRNA anticodon wobble base, maintains the proliferative and pluripotent potential of HL-60 cells in the presence of the differentiating agent 6-thioguanine. *Proc. Natl. Acad. Sci. USA*, 88, 370-374, 1991.
- (29) Froidi, G., Gallo, E., Ragazzi, E., Caparrotta, L. 6-Benzylaminopurine; a plant-derived cytokinin inducing positive inotropism by P2-purinoreceptors. *Planta Medica*, 65, 245-249, 1999.
- (30) Gillissen, B., Burkle, L., Andre, B., Kuhn, C., Rentsch, D., Brandl, B., Frommer, W.B. A new family of high-affinity transporters for adenine, cytosine, and purine derivatives in *Arabidopsis*. *Plant Cell*, 12, 291-300, 2000.
- (31) Gootwine, E., Webb, C.G., Sachs, L. Participation of myeloid leukaemic cells injected into embryos in haematopoietic differentiation in adult mice. *Nature*, 299, 63-65, 1982.
- (32) Grutter, M. G. Caspase: key players in programmed cell death. *Curr. Opin. Struct. Biol.*, 10 (6), 649-655, 2000.
- (33) Gusella, J.F., Weil, S.C., Tsiftoglou, A.S., Volloch, V., Neumann, J. R., Keys, C., Houseman, D. E. Hemin dose not cause commitment of murine erythroleukemia (MEL) cells to terminal differentiation. *Blood*, 56, 481-487, 1980.
- (34) Heithoff, D. M., Sinsheimer, R. L., Low, D. A., Mahan, M. J. An essential role for DNA adenine methylation in bacterial virulence. *Science*,

284, 967-970, 1999.

- (35) Honma, Y. A novel therapeutic strategy against monocytic leukemia with deoxyadenosine analogs and adenosine deaminase inhibitors. *Leuk. Lymphoma*, 42(5), 953-962, 2001.
- (36) Honma, Y. Cotylenin A—a plant growth regulator as a differentiation inducing agent against myeloid leukemia. *Leuk. Lymphoma*, 43(6), 1169-1178, 2002.
- (37) Honma, Y., Ishii, Y., Sassa, T., Asahi, K. Treatment of human promyelocytic leukemia in the SCID mouse model with cotylenin A, an inducer of myelomonocytic differentiation of leukemia cells. *Leukemia Research*, 27, 1019-1025, 2003.
- (38) Honma, Y., Ishii, Y., Yamamoto-Yamaguchi, Y., Sassa, T., Asahi, K. Cotylenin A, a differentiation-inducing agent, and IFN-alpha cooperatively induce apoptosis and have an antitumor effect on human non-small cell lung carcinoma cells in nude mice. *Cancer Research*, 63, 3659-66, 2003.
- (39) Honma, Y., Takenaga, K., Kasukabe, T., Hozumi, M. Induction of differentiation of cultured human promyelocytic leukemia cells by retinoids. *Biochem. Biophys. Res. Commun.*, 95, 507-512, 1980.
- (40) Horgan, R. Cytokinins, in *Advances Plant Physiology*, Wilkins, M. B., Ed., Pitman Medical, London, 1984.
- (41) Hozumi, M. Fundamentals of chemotherapy of myeloid leukemia by induction of leukemia cell differentiation. *Adv. Cancer Res.*, 38, 121-169, 1983.
- (42) Hsiao, G., Shen, M. Y., Lin, K. H., Chou, C. Y., Tzu, N. H., Lin, C. H., Chou, D. S., Chen, T. F., Sheu, J. R. Inhibitory activity of kinetin of free radical formation of activated platelets in vitro and on thrombus formation in vivo. *Eur. J. Pharmacol.*, 465 (3), 281-7, 2003.

- (43) Huan, M., Ye, Y., Chen, S., Chai, J., Lu, J., Lin, Z., Gu, L., Wang, Z. Use of all-*trans* retinoic acid in the treatment of acute promyelocytic leukemia. *Blood*, 72, 567-572, 1988.
- (44) Hungness, E. S., Luo, G. J., Pritts, T. A., Sun, X., Robb, B. W., Hershko, D., Hasselgren, P. O. Transcription factors C/EBP-beta and delta regulate IL-6 production in IL-1 beta-stimulated human enterocytes. *J. Cell. Physiol.*, 192(1), 64-70, 2002.
- (45) Hutt, J. A., O'Rourke, J. P., DeWille, J. Signal transducer and activator of transcription 3 activates CCAAT enhancer-binding protein delta gene transcription in G0 growth-arrested mouse mammary epithelial cells and involuting mouse mammary gland. *J. Biol. Chem.*, 275(37), 29123-29131, 2000.
- (46) Inoue, H., Hirose, M. Superoxide dismutase mimetic activity of cytokinin-copper (II) complexes. *Biochem. Biophys. Res. Commun.*, 137, 372-377, 1986.
- (47) Inoue, T., Higuchi, M., Hashimoto, Y., Seki, M., Kobayashi, M. Kato, T., Tabata, S., Shinozaki, K., Kakimoto, T. Identification of CRE1 as a cytokinin receptor from *Arabidopsis*. *Nature*, 409, 1060-1063, 2001.
- (48) Ishii, Y., Kiyota, H., Sakai, S., Honma, Y. Induction of differentiation of human myeloid leukemia cells by jasmonates, plant hormones. (submitting)
- (49) Ishii, Y., Sakai, S., Honma, Y. Cytokinin-induced differentiation of human myeloid leukemia HL-60 cells is associated with the formation of nucleotides, but not with incorporation into DNA or RNA. *Biochimica et Biophysica Acta*, 1643(1-3), 11-24.
- (50) Ishii, Y., Sakai, S., Honma, Y. Pyridinyl imidazole inhibitor SB203580 activates p44/42 mitogen-activated protein kinase and induces the differentiation of human myeloid leukemia cells. *Leukemia Research*, 25, 813-20, 2001.

- (51) Jouanneau, J., Teyssendier de la Serve, B., Peaud-Lenoel, C. Frequency of N<sup>6</sup>-benzyladenine incorporation into major RNA reactions of tobacco cell suspensions grown at stimulatory or cytostatic cytokinin concentration. *Plant Physiol.*, 74, 663-668, 1984.
- (52) Kanatani, H., Nakanishi, K., Kuroiwa, T. Isolation and identification on meiosis inducing substance in star fish *Asteria amurensis*. *Nature*, 221, 273-274, 1969.
- (53) Kasukabe, T., Honma, Y., Hozumi, M. Characterization of lysozyme synthesized by differentiated mouse myeloid leukemia cells. *Biochimica et Biophysica Acta*, 586, 615-623, 1979.
- (54) Keeshan, K., Santilli, G., Corradini, F., Perrotti, D., Calabretta, B. Transcription activation function of C/EBP alpha is required for induction of granulocytic differentiation. *Blood*, 102 (4), 1267-1275, 2003.
- (55) Kerkhoff, C., Eue, I., Sorg, C. The regulatory role of MRP8 (S100A8) and MRP14 (S100A9) in the transendothelial migration of human leukocytes. *Pathobiology*, 67(5-6), 230-232, 1999.
- (56) Kretz, K. A., Katze, J. R., Trewyn, R. W. Guanine analog-induced differentiation of human promyelocytic leukemia cells and changes in queuine modification of tRNA. *Mol. Cell. Biol.*, 7, 3613-3619, 1987.
- (57) Kroemer, G., Zamzami, N., Susin, S.A. Mitochondrial control of apoptosis. *Immunol. Today*, 18, 44- 51, 1997.
- (58) Kurosaki, F., Takahashi, S., Shudo, K., Okamoto, T., Isogai, Y., Structural and biological link between urea- and purine-cytokinins. *Chem. Pharm. Bull.*, 29, 3751-3753, 1981.
- (59) Leonard, N. J., Fujii, T. The synthesis of compounds possessing kinetin activity. The use of a blocking group at the 9-position of adenine for the biosynthesis of 1-substituted adenines. *Proc. Natl. Acad. Sci. USA*, 51, 73-75, 1964.

- (60) Leonard, N. J., Henderson, T. R. Purine ring rearrangements to the development of cytokinin activity. Mechanism of the rearrangement of 3-benzyladenine to N<sup>6</sup>-benzyladenine. *J. Am. Chem. Soc.*, 93, 4990-4999, 1975.
- (61) Letham, D. S. Zeatin, a factor inducing cell division from *Zea mays*. *Life Sci.*, 8, 596, 1963.
- (62) Li, P.F., Dietz, R., von Harsdorf, R. p53 regulates mitochondrial membrane potential through reactive oxygen species and induces cytochrome c-independent apoptosis blocked by Bcl-2. *EMBO J.*, 18, 6027-6036, 1999.
- (63) MacDougald, O. A., Lane, M. D. Transcriptional regulation of gene expression during adipocyte differentiation. *Annu. Rev. Biochem.*, 64, 345-373, 1995.
- (64) Makishima, M., Honma, Y., Hozumi, M., Sampi, K., Hattori, M., Ishikawa, I., Ogura, H., Motoyoshi, K. Effects of novel uracil analogs on proliferation and differentiation of human myeloid leukemia cells. *Exp. Hematol.*, 20, 879-885, 1992.
- (65) Makishima, M., Kanatani, Y., Yamamoto-Yamaguchi, Y., Honma, Y. Enhancement of activity of 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> for growth inhibition and differentiation induction of human myelomonocytic leukemia cells by tretinoin tocopheril, an  $\alpha$ -tocopherol ester of all-*trans* retinoic acid. *Blood*, 87, 3384-3394, 1996.
- (66) Mancini, M., Sedghinasab, M., Knowlton, K., Tam, A., Hockenbery, D., Anderson, B. O. Flow cytometric measurement of mitochondrial mass and function: a novel method for assessing chemoresistance. *Ann. Surg. Oncol.*, 5(3), 287-295, 1998.
- (67) Manitz, M. P., Horst, B., Seeliger, S., Strey, A., Skryabin, B. V., Gunzer, M., Frings, W., Schonlau, F., Roth, J., Sorg, C., Nacken, W. Loss of



- S100A9 (MRP14) results in reduced interleukin-8-induced CD11b surface expression, a polarized microfilament system, and diminished responsiveness to chemoattractants in vitro. *Mol. Cell. Biol.*, 23(3), 1034-1043, 2003.
- (68) Maytin, E. V., Habener, J. F. Transcription factor C/EBP alpha, C/EBP beta, and CHOP(Gadd153) expressed during the differentiation program of keratinocytes in vitro and in vivo. *J. Invest. Dermatol.*, 110(3), 238-246, 1998.
- (69) Miller, C. O., Skoog, F., Okamura, F. S., von Salta, M. H., Strong, F. M. Isolation, structure and synthesis of kinetin, a substance promoting cell division. *J.Am.Chem.Soc.*, 78, 1345, 1956.
- (70) Mintz, B., Illumensee, L. Normal genetically mosaic mice produced from malignant teratocarcinoma cell. *Proc. Natl. Acad. Sci. U.S.A.*, 72, 3585-3589, 1975.
- (71) Miranda, M. B., McGuire, T. F., Johnson, D. E. Importance of MEK-1/-2 signaling in monocytic and granulocytic differentiation of myeloid cell lines. *Leukemia*, 16(4), 683-692, 2002.
- (72) Miyaura, C., Abe, E., Kuribayashi, T., Tanaka, H., Konno, K., Nishi, Y., Suda, T.  $1\alpha,25$ -Dihydroxyvitamin D<sub>3</sub> induces differentiation of human myeloid leukemia cells. *Biochem. Biophys. Res. Commun.*, 102, 937-943, 1981.
- (73) Mok, M.C. Chemistry of Adenine Cytokinins. In *Cytokinins - Chemistry, Activity and Function* (D.W.S. Mok and M.C. Mok, Eds.), pp. 15-34. CRC Press., Boca Raton, FL., 1994.
- (74) Mok, M. C. Cytokinins and plant development -an overview. In *Cytokinins - Chemistry, Activity and Function* (D.W.S. Mok and M.C. Mok, Eds.), pp. 155-166. CRC Press., Boca Raton, FL., 1994.
- (75) Mok, M.C., Martin, R.C., Mok, D.S. Cytokinins: biosynthesis,

- metabolism and perception. *In Vitro Cell. Dev. Biol.-Plant*, 36, 102-107, 2000.
- (76) Mok, M. C., Mok, D. W. S., Armstrong, D. J., Shudo, K., Isogai, Y., Okamoto, T. Cytokinin activity of N-phenyl-N'-(1,2,3-thiadiazol-5-yl) urea (thidiazuron). *Phytochemistry*, 21, 1509-1511, 1982.
- (77) Momparler, R.L., Bouchard, J., Samson, J. Induction of differentiation and inhibition of DNA methylation in HL-60 myeloid leukemic cells by 5-AZA-2'-deoxycytidine. *Leuk. Res.*, 9, 1361-1366, 1985.
- (78) Murai, N., Taller, B.J., Armstrong, D. J., Skoog, F. Kinetin incorporated into tobacco callus ribosomal RNA and transfer RNA preparations. *Plant Physiol.* 60, 197-20, 1977.
- (79) Ngernprasirtsiri, J., Akazawa, T. Modulation of DNA methylation and gene expression in cultured sycamore cells treated by hypomethylating base analog. *Eur. J. Biochem.*, 194, 513-520, 1990.
- (80) Niitsu, N., Hayashi, Y., Sugita, K., Honma, Y. Sensitization by 5-aza-2'-deoxycytidine of leukaemia cells with *MLL* abnormalities to induction of differentiation by all-*trans* retinoic acid and 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub>. *Br. J. Haematol.*, 112, 315-326, 2001.
- (81) Nishida, E., Gotoh, Y. The MAP Kinase cascade is essential for diverse signal transduction pathway. *Trends Biochem. Sci.*, 18,128-131, 1993.
- (82) Nishio, M., Zushi, S., Ishii, T., Furuya, T., Syono, K. Mass fragmentgraphic determination of indole-3-acetic acid in callus tissues of *Panax ginseng* and *Nicotiana tabacum*. *Chem. Pharm. Bull*, 24, 2038-2042, 1976.
- (83) Ohno R, Naoe T, Hirano M, Kobayashi M, Hirai H, Tubaki K, Oh, H. Treatment of myelodysplastic syndromes with all-*trans* retinoic acid. Leukemia Study Group of the Ministry of Health and Welfare. *Blood*, 81, 1152-1154, 1993.

- (84) Oka, H., Asahi, K., Morishima, H., Sanada, M., Shiratori, K., Iimura, Y., Sakurai, T., Uzawa, J., Iwadare, S., Takahashi, N. Differanisole A, a new differentiation inducing substance. *J. Antibiot.*, 38, 1100-1102, 1985.
- (85) Okamoto, T., Shudo, K., Takahashi, S., Kawachi, E., Isogai, Y. 4-Pyridylureas are surprisingly potent cytokinins. The structure-activity relationship. *Chem. Pharm. Bull.*, 29, 3748-3750, 1981.
- (86) Olsen, A., Siboska, G.E., Clark, B.F.C., Rattan, S.I.S. N<sup>6</sup>-Furfuryladenine, kinetin, protects against Fenton reaction-mediated oxidative damage to DNA. *Biochem. Biophys. Res. Commun.*, 265, 499-502, 1999.
- (87) Olson, M., Kornbluth, S. Mitochondria in apoptosis and human disease. *Curr. Mol. Med.*, 1(1), 91-122, 2001.
- (88) Pabst, T., Mueller, B. U., Zhang, P., Radomska, H. S., Narravula, S., Schnittger, S., Behre, G., Hiddemann, W., Tenen, D. G. Dominant-negative mutations of CEBPA, encoding CCAAT/enhancer binding protein-alpha (C/EBPalpha), in acute myeloid leukemia. *Nat. Genet.*, 27(3), 263-270, 2001.
- (89) Passay, R. J., Xu, K., Hume, D. A., Geczy, C. L. S100A8: emerging function and regulation. *J. Leukoc. Biol.*, 66 (4), 549-556, 1999.
- (90) Pinto, V. Attadia, A. Fusco, F. Ferrara, O. A., Spada, P., Di Fiore, P. 5-Aza-2'-deoxycytidine induces terminal differentiation of leukemic blasts from patients with acute myeloid leukemias. *Blood*, 64, 922-929, 1984.
- (91) Podesta, A.H., Mullins, J., Pierce, G.B., Wells, R.S. The neural stage mouse embryo in control of neuroblastoma. *Proc. Natl. Acad. Sci. U.S.A.*, 81, 7608-7611. 1984.
- (92) Radomska, H. S., Huettner, C. S., Zhang, P., Cheng, T., Scadden, D. T., Tenen, D. G. CCAAT/Enhancer binding protein alpha is a regulatory switch sufficient for induction of granulocytic development from bipotential myeloid progenitors. *Mol. Cell. Biol.*, 18 (7), 4301-4314, 1998.

- (93) Ramji, D. P., Foka, P. CCAAT/enhancer-binding proteins: structure, function and regulation. *Biochem J.*, 365 (Pt 3), 561-575, 2002.
- (94) Rattan, SI., Clark, BF., Kinetin delays the onset of ageing characteristics in human fibroblasts. *Biochem. Biophys. Res. Commun.*, 201(2) 665-72, 1994.
- (95) Reutelingsperger, C. P., van Heerde, W. L., Annexin V, the regulator of phosphatidylserine-catalyzed inflammation and coagulation during apoptosis. *Cell Mol Life Sci.*, 53 (6), 527-532, 1997.
- (96) Rosel, F. A simple and rapid method for detection of apoptosis in human cells. *Nucleic Acids Research*, 20 (19), 5243, 1992.
- (97) Rovera, G., Santoli, D., and Damsky, C. Human myelocytic leukemia cells in culture differentiate into macrophage-like cells when treated with a phorbol diester. *Proc. Natl. Acad. Sci. USA*, 76: 2779-2783, 1979.
- (98) Ryckman, C., Vandal, K., Rouleau, P., Talbot, M., Tessier, P. A. Proinflammatory activities of S100: proteins S100A8, S100A9, and S100A8/A9 induce neutrophil chemotaxis and adhesion. *J. Immunol.*, 170 (6), 3233-3242, 2003.
- (99) Sassa, T. Promotion of the enlargement of cotyledons by cotylenin A and B. *Agr. Biol. Chem.*, 34, 1588-1589, 1970.
- (100) Schulze-Gahmen, U., Brandsen, J., Jones, H. D., Morgan, D. O., Meijer, L., Versely, J., Kim, S.-H. Multiple modes of ligand recognition: crystal structures of cyclin-dependent protein kinase 2 in complex with ATP and two inhibitors, olomoucine and isopentenyladenine. *Proteins*, 22, 378-391, 1995.
- (101) Shantz, E. M. and Steward, F. C. The identification of compound A from coconut milk as 1,3-diphenylurea. *J. Am. Chem. Soc.*, 77, 6351-6353, 1955.

- (102) Sharma, S. P., Kaur, J., Rattan, S. I. Increased longevity of kinetin-fed *Zaprionus* fruitflies is accompanied by their reduced fecundity and enhanced catalase activity. *Biochem. Mol. Bio. Int.*, 41(5), 869-75, 1997.
- (103) Sharma, S. P., Kaur, P., Rattan, S. I. Plant growth hormone kinetin delays ageing, prolongs the lifespan and slow down development of the fruitfly *Zaprionus paravittiger*. *Biochem. Biophys. Res. Commun.*, 216 (3), 1067-71, 1995.
- (104) Sherman, M. L., Sharfman, T. D., Colman, M. S., Kufe, D. W. Tiazofurin induction of mouse erythroleukemia cell hemoglobin production in the absence of commitment or changes in protooncogene expression. *Blood*, 73, 431-434, 1989.
- (105) Shivdasani, R. A., Orkin, S. H. The transcriptional control of hematopoiesis. *Blood*, 87(10), 4025-4039. 1996.
- (106) Skoog, F., Armstrong, D. J. Cytokinins. *Science*, 148, 532-533, 1970.
- (107) Smith, L. T., Hohaus, S., Gonzalez, D. A., Dziennis, S. E., Tenen, D. G. PU.1 (Spi-1) and C/EBP alpha regulate the granulocyte colony-stimulating factor receptor promoter in myeloid cells. *Blood*, 88 (4), 1234-1247, 1996.
- (108) Smith, M. L., Fornace, A. J. Mammalian DNA damage-inducible genes associated with growth arrest and apoptosis. *Mutat. Res.*, 340, 109-124, 1996.
- (109) Strong, F. M. Kinetin and kinins. *In Topics in microbial chemistry*. John Wiley and Sons, Inc., pp.98-157, 1959.
- (110) Suzuki, K., Hino, M., Hato, F., Tatsumi, N., Kitagawa, S. Cytokine-specific activation of distinct mitogen-activated protein kinase subtype cascades in human neutrophils stimulated by granulocyte colony-stimulating factor, granulocyte-macrophage colony-stimulating factor, and tumor necrosis factor-alpha. *Blood*, 93 (1), 341-349, 1999.
- (111) Tanimoto, S., Harada, H. Effect of cytokinin and anticytokinin on the

- initial stage of adventitious bud differentiation in the epidermis of *Torenia* stem segments. *Plant Cell Physiol.*, 23, 1371-137, 1982.
- (112) Teysseudier de la Serve, B., Jouanneau, J., Peaud-Lenoel, C. Incorporation of N<sup>6</sup>-benzyladenine into messenger poly(A)-RNA of tobacco cells incubated at stimulatory or cytostatic cytokinin concentration. *Plant Physiol.*, 74, 669-674, 1984.
- (113) Trumpp-Kallmeyer, S., Rubin, J.R., Humblet, C., Hamby, J.M., Showalter, H.D. Development of a binding model to protein tyrosine kinase for substituted pyrido[2,3-*d*]pyridine inhibitors. *J. Med. Chem.*, 41, 1752-1763, 1998.
- (114) Turner, J. G., Ellis, C., Devoto, A. The jasmonate signal pathway. *Plant Cell*, (Supplement 2002), S153-S164, 2002.
- (115) van der Woude, M., Braaten, B., Low, D. Epigenetic phase variation of the pap operon in *Escherichia coli*. *Trends Microbiol*, 4, 5-9, 1996.
- (116) Vogl, T., Propper, C., Hartmann, M., Strey, A., Strupat, K., van den Bos, C., Sorg, C., Roth, J. S100A12 is expressed exclusively by granulocytes and acts independently from MRP8 and MRP14. *J. Biol. Chem.*, 274(36), 25291-25296, 1999.
- (117) Warner, G. J., Rusconi, C.P., White, I. E., Faust, J. R. Identification and sequencing of two isopentenyladenosine-modified transfer RNAs from Chinese hamster ovary cells. *Nucleic Acid Res.*, 26, 5533-5535, 1998.
- (118) Weber, H. Fatty acid-derived signals in plants. *TRENDS Plant Sci.*, 7, 217-224, 2002.
- (119) Wiesner, J. Die Elementarstruktur und das Wachstum der lebenden Substanz, A.Holder, Vienna, 1892.
- (120) Yamaki, T., Shimojo, E., Ichimura, H., Nishinari, N., Namekawa, K., Yoshizawa, K. Possible role of indoleacetic acid in animal cell division and cancer growth. *J.UOEH*, 1(4), 471-485, 1979.

- (121) Yamamoto-Yamaguchi, Y., Makishima, M., Kanatani, Y., Kasukabe, T., Honma, Y. Reversible differentiation of human monoblastic leukemia U937 cells by ML-9, an inhibitor of myosin light chain kinase. *Experimental Hematology*, 24, 682-689, 1996.
- (122) Yamamoto-Yamaguchi, Y., Okabe-Kado, J., Kasukabe, T., Honma, Y. Induction of differentiation of human myeloid leukemia cells by immunosuppressant macrolides (rapamycin and FK506) and calcium/calmodulin-dependent kinase inhibitors. *Exp. Hematol.*, 29, 582-588, 2001b.
- (123) Yamamoto-Yamaguchi, Y., Yamada, K., Ishii, Y., Asahi, K., Tomoyasu, S., Honma, Y. Induction of the monocytic differentiation of myeloid leukemia cells by cotylenin A, a plant growth regulator. *Br. J. Haematol.*, 112, 697-705, 2001a.
- (124) Yen, A., Roberson, M.S., Varvayanis, S., Lee, A.T. Retinoic acid induced mitogen-activated protein (MAP)/extracellular signal-regulated kinase (ERK) kinase-dependent MAP kinase activation needed to elicit HL-60 cell differentiation and growth arrest, *Cancer Res.*, 58, 3163-3172. 1998.
- (125) Yukawa, K., Tanaka, T., Tsujii, S., Akira, S. Expression of CCAAT/Enhancer-binding proteins beta and delta and their activities are intensified by cAMP signaling as well as Ca<sup>2+</sup>/calmodulin kinases activation in hippocampal neurons. *J. Biol. Chem.*, 273(47), 31345-31351, 1998.
- (126) Zamzami, N., Marchetti, P., Castedo, M., Zanin, C., Vayssiere, J.L., Petit, P.X., Kroemer, G. Reduction in mitochondrial potential constitutes an early irreversible step of programmed lymphocyte death in vivo. *J. Exp. Med.*, 181, 1661-1672, 1995.
- (127) Zhang, D.E., Zhang, P., Wang, N. D., Hetherington, C. J., Darlington, G. J., Tenen, D. G. Absence of granulocyte colony-stimulating factor signaling

and neutrophil development in CCAAT enhancer binding protein  $\alpha$ -deficient mice. Proc. Natl. Acad. Sci. USA, 94, 569-574, 1997.