

References

Abeles, F.B. (1973) Ethylene in plant biology. Academic Press, New York.

Adams, D.O. and Yang, S.F. (1979) Ethylene biosynthesis: identification of 1-aminocyclopropane-1-carboxylic acid as an intermediate in the conversion of methionine to ethylene. *Proc. Natl. Acad. Sci. USA* 76:170-174.

Atsmon, K. and Galun, E. (1960) A morphogenetic study of staminate, pistillate and hermaphrodite flowers in *Cucumis sativus* (L.). *Phytomorphology* 10: 110-115.

Atsmon, D. and Tabbak, C. (1979) Comparative effects of gibberellin, silver nitrate and aminoethoxyvinyl glycine on sexual tendency and ethylene evolution in the cucumber plant (*Cucumis sativus* L.). *Plant Cell Physiol.* 20: 1547-1555.

Beyer, E.J. (1976) Silver ion: a potent antiethylene agent in cucumber and tomato. *HortScience* 11:195-196.

Boller, T., Herner, R.C. and Kende, H. (1979) Assay for and enzymatic formation of an ethylene precursor, 1-aminocyclopropane-1-carboxylic acid. *Planta* 145: 293-303.

Darwin, C. (1876) Effects of Cross and Self-Fertilization in the Vegetable Kingdom. John Murray, London.

Dauphin-Guerin, B., Teller, G. and Durand, B. (1980) Different endogenous cytokinins between male and female *Mercurialis annua* L. *Planta* 148: 124-129.

Dellaporta, S.L. and Calderon-Urrea, A. (1993) Sex determination in flowering plants. *Plant Cell* 5: 1241-1251.

Destefano-Beltran, L.J.C., Caeneghem, W.V., Gielen, J., Richard, L., Montegu, M.V. and Straeten, D.V.D. (1995) Characterization of three members of the ACC synthase gene family in *Solanum tuberosum* L. *Mol. Gen. Genet.* 246: 496-508.

Dong, J.G., Kim, W.T., Yip, W.K., Thompson, G.A., Li, L., Bennett, A.B. and Yang, S.F. (1991) Cloning of a cDNA encoding 1-aminocyclopropane-1-carboxylate synthase and expression of its mRNA in ripening apple fruit. *Planta* 185: 38-45.

Durand, R. and Durand, B. (1984) Sexual differentiation in higher plants. *Physiol. Plant.* 60: 267-274.

Durand, B. and Durand, R. (1991) Sex determination and reproductive organ differentiation in *Mercurialis*. *Plant Sci.* 80: 49-65.

Emerson, R.A. and Emerson, S.H. (1922) Genetic interrelations of two andromonoecious types of maize, *dwarf* and *anther ear*. *Genetics* 7: 203-236.

Frankel, R. and Galun, E. (1977) *Pollination Mechanisms, Reproduction and Plant Breeding*. Springer-Verlag, Berlin.

Galun, E. (1959) Effect of gibberelic acid and naphthaleneacetic acid in sex expression and some morphological characters in the cucumber plants. *Phyton* 13: 1-8.

Galun, E. (1961) Study of the inheritance of sex expression in the cucumber, the interactions of major genes with modifying genetic and non-genetic factors. *Genetica* 32: 134-163.

Galun, E., Jung, Y. and Lang, A. (1962) Culture and sex modification of male cucumber buds in vitro. *Nature* 194: 596-598.

Galun, E., Jung, Y. and Lang, A. (1963) Morphogenesis of floral buds of cucumber cultured in vitro. *Develop. Biol.* 6: 370-387.

Hemphill, D.D., Baker, L.R. and Sell, H.M. (1972) Different sex phenotypes of *Cucumis sativus* L. and *C. melo* L. and their endogenous gibberellin activity. *Euphytica* 21: 285-291.

Heslop-Harrison, J. (1957) The experimental modification of sex expression in plants. *Biol. Rev.* 32:38-90.

Huang, P.L., Parks, J.E., Rottmann, W.H. and Theologis, A. (1991) Two genes encoding 1-aminocyclopropane-1-carboxylate synthase in zucchini (*Cucurbita pepo*) are clustered and similar but differentially regulate. *Proc. Natl. Acad. Sci. USA* 88: 7021-7025.

Iwahori, S., Lyons, J.M. and Smith, O.E. (1970) Sex expression in cucumber plants as affected by 2-chloroethylphosphonic acid, ethylene, and growth regulators. *Plant Physiol.* 46: 412-415.

Kende, H. (1993) Ethylene biosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 44: 283-307.

Kubicki, B. (1969a) Investigations on sex determination in cucumber (*Cucumis sativus* L.). *Genet. Pol.* 10: 69-85.

Kubicki, B. (1969b) Investigations on sex determination in cucumber (*Cucumis sativus* L.). VII. Andromonoecism and hermaphroditism. *Genet. Pol.* 10: 101-120.

Kubicki, B. (1969c) Investigations on sex determination in cucumber (*Cucumis sativus* L.). VI. Androecism. *Genet. Pol.* 10: 87-99.

Lazarte, J.E. and Palser, B.F. (1979) Morphology, vascular anatomy and embryology of pistillate and staminate flowers of *Asparagus officinalis*. *Am. J. Bot.* 66:753-764.

Liang, X., Abel, S., Keller, J.A., Shen, N.F. and Theologis, A. (1992) The 1-aminocyclopropane-1-carboxylate synthase gene family of *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* 89: 11046-11050.

Louis, J.-P. (1989) Genes for the regulation of sex differentiation and male fertility in *Mercurialis annua* L. *J. Hered.* 80: 104-111.

Louis, J.-P., Augur, C. and Teller, G. (1990) Cytokinins and differentiation processes in *Mercurialis annua*. *Plant Physiol.* 94: 1535-1541.

Malepszy, S. and Niemirowicz-Szczytt, K. (1991) Sex determination in cucumber (*Cucumis sativus*) as a model system for molecular biology. *Plant Science* 80: 39-47.

McMurray, A.L. and Miller, C.H. (1968) Cucumber sex expression modified by 2-chloroethanephosphonic acid. *Science* 162: 1397-1398.

Mitchell, W.D. and Wittwer, S.H. (1962) Chemical regulation of sex expression and vegetative growth in *Cucumis sativus*. *Science* 136: 880-881.

Mohan Ram, H.Y. and Nath, R. (1964) The morphology and embryology of *Cannabis sativa* Linn. *Phytomorphology* 14: 414-429.

Nakagawa, N., Mori, H., Yamazaki, K. and Imaseki, H. (1991) Cloning of a complementary DNA for auxin-induced 1-aminocyclopropane-1-carboxylate synthase and differential expression of the gene by auxin and wounding. *Plant Cell Physiol.* 32: 1153-1163.

Nakajima, N., Mori, H., Yamazaki, K. and Imaseki, H. (1990) Molecular cloning and sequence of a complementary DNA encoding 1-aminocyclopropane-1-carboxylate synthase induced by tissue wounding. *Plant Cell Physiol.* 31: 1021-1029.

Nakatsuka, A., Shimoi, S., Kubo, Y. and Inaba, A. (1997) Expression and internal feedback regulation of ACC synthase and ACC oxidase genes in ripening tomato fruit. *Plant Cell Physiol.* 38: 1103-1110.

Okamoto, H., Tanaka, Y. and Sakai, S. (1995) Molecular cloning and analysis of the cDNA for an auxin-regulated calmodulin gene. *Plant Cell Physiol.* 36: 1531-1539.

Olson, D.C., White, J.A., Edlman, L., Harkins, R.N. and Kende, H. (1991) Differential expression of two genes for 1-aminocyclopropane-1-carboxylate synthase in tomato fruits. *Proc. Natl. Acad. Sci. USA* 88: 5340-5344.

O'Neill, S.D., Nedeau, J.A., Zhang, X., Bui, A.Q. and Halevy, A.H. (1993) Interorgan regulation of ethylene biosynthetic genes by pollination. *Plant Cell* 5: 419-432.

Park, K.Y., Drory, A. and Woodson, W.R. (1992) Molecular cloning of an 1-aminocyclopropane-1-carboxylate synthase from senescing carnation flower petals. *Plant Mol. Biol.* 18: 377-386.

Pierce, L.K. and Wehner, T.C. (1990) Review of genes and linkage groups in cucumber. *HortScience* 25: 605-615.

Prescott, A. and Martin, C. (1987) A rapid method for the quantitative assessment of levels of specific mRNAs in plants. *Plant Mol. Biol. Rep.* 4: 219-224.

Rood, S.B., Paris, R.P. and Major, D.J. (1980) Changes of endogenous gibberellin-like substances with sex reversal of the apical inflorescence of corn. *Plant Physiol.* 66: 793-796.

Rottmann, W.H., Peter, G.F., Oeller, P.W., Keller, J.A., Shen, N.F., Nagy, B.P., Taylor, L.P., Campbell, A.D. and Theologis, A. (1991) 1-Aminocyclopropane-1-carboxylate synthase in tomato is encoded by a multigene family whose transcription is induced during fruit and floral senescence. *J. Mol. Biol.* 222: 937-961.

Rudich, J., Baker, L.R., Scott, J.W. and Sell, H.M. (1976) Phenotypic stability and ethylene evolution in androecious cucumber. *J. Amer. Soc. Hort. Sci.* 101: 48-51.

Rudich, J., Halevy, A.H. and Kedar, N. (1969) Increase in femaleness of three cucurbits by treatment with ethrel, an ethylene-releasing compound. *Planta* 86: 69-76.

Rudich, J., Halevy, A.H. and Kedar, N. (1972) Ethylene evolution from cucumber plants as related to sex expression. *Plant Physiol.* 49: 998-999.

Sanger, F.S., Nicklen, S. and Coulson, A.R. (1977) DNA sequencing with chain-terminating inhibitor. *Proc. Natl. Acad. Sci. USA* 74: 5463-5467.

Shannon, S. and De La Guardia, M.D. (1969) Sex expression and the

production of ethylene induced by auxin in cucumber (*Cucumis sativus* L.). *Nature* 223: 186.

Shifriss, O. (1961) Sex control in cucumbers. *J. Hered.* 52: 5-12.

Shifriss, O., George, W.L. and Quinones, J.A. (1964) Gynodioecism in cucumber. *Genetics* 49: 285-291.

Trebitsh, T., Staib, J.E. and O'Neill, S.D., (1997) Identification of a 1-aminocyclopropane-1-carboxylic acid synthase gene linked to the female (*F*) locus that enhances female sex expression in cucumber. *Plant Physiol.* 113: 987-995.

Trebitsh, T., Rudich, J. and Riov, J. (1987) Auxin, biosynthesis of ethylene and sex expression in cucumber (*Cucumis sativus*). *Plant Growth Regul.* 5: 105-113.

Wittwer, S.H. and Bukovac, M.J. (1958) The effect of gibberellins on economic crops. *Econ. Bot.* 12: 213-255.

Woodson, W.R., Park, K.Y., Drory, A., Larsen, P.B. and Wang, H. (1992) Expression of ethylene biosynthetic pathway transcripts in senescing carnation flowers. *Plant Physiol.* 99: 526-532.

Yang, S.F. and Hoffman, N.E. (1984) Ethylene biosynthesis and its regulation in higher plants. *Annu. Rev. Plant Physiol.* 35: 155-189.

Ye, D., Oliveira, M., Veuskens, J., Wu, Y., Installe, P., Hinnisdaels, S.,

Truong, A.T., Brown, S., Mouras, A. and Negrutiu, I. (1991) Sex determination in the dioecious *Melandrium*. The X/Y chromosome system allows complementary cloning strategies. *Plant Sci.* 80: 93-106.

Yin, T. and Quinn, J.A. (1995) Tests of a mechanistic model of one hormone regulating both sexes in *Cucumis sativus* (Cucurbitaceae). *Am. J. Bot.* 82: 1537-1546.

Zarembinski, T.I. and Theologis, A. (1993) Anaerobiosis and plant growth hormone induced two gene encoding 1-aminocyclopropane-1-carboxylate synthase in rice (*Oryza sativus* L.). *Mol. Biol. Cell* 4:363-373.

Acknowledgements

I wish to express my sincere appreciation to Professor S. Sakai of the University of Tsukuba for his invaluable suggestions and encouragement throughout the course of my research.

My sincere thanks are also due to Emeritus Professor T. Fujii (University of Tsukuba), Professor H. Kamada (University of Tsukuba) and Associate Professor S. Satoh (University of Tsukuba) for their valuable suggestions and criticisms. Special thanks must be accorded to all members of my laboratory for their invaluable discussions and encouragement.

I am grateful to Dr. S. Matsuura and Dr. H. Mizusawa (Tohoku Seed Co.) for kindly providing seeds of isogenic cucumber lines and for their valuable informations and suggestions.

I am also grateful to Professor I. Inoue (University of Tsukuba) and Professor J. Hayashi (University of Tsukuba) for their valuable suggestions and constructive criticisms for the preparation of the manuscript.