

LA
25/8
2000

(176)

Mechanism of Mixture Suppression in Newt Olfactory Receptor Neurons

Hiroshi YAMADA

A dissertation submitted to the Doctoral
Program in Biological Sciences, the
University of Tsukuba in partial fulfillment of
the requirements for the degree of Doctor of
Philosophy (Science)

寄	贈
山	平成
田	年
博	月
氏	日

January, 2001

01301565

ABSTRACT	1
INTRODUCTION	4
1 Excitatory transduction cascade	5
2 Inhibitory responses and mixture suppression	6
METHODS	9
1 Animal and cell preparation	10
2 Whole cell patch clamp	10
3 Perfusion system	11
4 EOG recordings	12
5 Solutions	13
RESULTS	14
1 Targets of odor inhibitory action	15
1.1 Odor inhibition of IBMX-induced depolarization	15
1.2 Odorants inhibit IBMX-induced conductance, but not cell's basal K ⁺ conductance	16
1.3 Odorants inhibit cAMP-induced current	17
1.4 Odorants inhibit 8-Br-cGMP induced current	18
2 Property of odor inhibitory effect	18
2.1 Dose-dependence of odor inhibition	18
2.2 Interaction between odor excitatory and inhibitory effects	19
2.3 Non-cell-specific suppression by odorants	19
3 Mutual suppression caused by odor inhibitory effect	20
3.1 Odorants inhibit odor-induced depolarization	20
3.2 Mixture suppression is dependent on individual odor excitation	21
3.3 Mutual suppression observed in EOG	22
DISCUSSION	25
1 Targets of odor inhibitory action	26

1.1	Inhibitory effects on cAMP cascade	26
1.2	Inhibitory effects on other conductances	27
2	Dose-range of odor inhibitory effect	28
3	Inhibitory responses that caused by odor suppression of cAMP-induced current	29
4	Odor representation in ORNs	30
4.1	Mixture suppression	30
4.2	Putative representation of mixture odorant in ORNs	31
	ACKNOWLEDGEMENTS	33
	REFERENCES	35
	FIGURES AND LEGENDS	43