

ACKNOWLEDGEMENTS

I wish to express my special gratitude to my Academic Advisor, Dr. Kei Nakatani who introduce me to work on this research field. I also earnestly thank Dr. Takehiko Saito, Dr. Masahiro Yamada, and Dr. Shiraiwa Yoshihiro for their critical reading of this paper and for their valuable comments.

Thanks are also to my colleagues, Miss. Masami Yokoyama and Miss Rieko Niikura for many valuable discussion and encouragement."

I would like to express my heartfelt gratitude to my parents, Seikai and Kyohko. I sincerely thank them for the silent, constant, and truly gratifying support.

REFERENCES

- Ache, B.W., and Zhainazarov, A. (1995) Dual second-messenger pathways in olfactory transduction. *Curr. Opin. Neurobiol.*, 5, 461-466.
- Bakalyar, H.A., and Reed, R.R. (1990) Identification of a specialized adenylyl cyclase that may mediate odorant detection. *Science* 250, 1403-1406.
- Bell, G.A., Laing, D.G., and Panhuber, H. (1987) Odour mixture suppression: evidence for a peripheral mechanism in human and rat. *Brain Res.* 426, 8-18.
- Berglund, B., and Olsson, M.J. (1993a) Odor-intensity interaction in binary mixtures. *J. Exp. Psychol. Hum. Percept.* 19, 302-314.
- Berglund, B., and Olsson, M.J. (1993b) Odor-intensity interaction in binary and ternary mixtures. *Percept. Psychophys.* 53, 475-482.
- Breer, H., Boekhoff, I., and Tareilus, E. (1990) Rapid kinetics of second messenger formation in olfactory transduction. *Nature* 345, 65-68.
- Brooks, S.P., and Storey, K.B. (1992) Bound and determined: a computer program for making buffers of defined ion concentrations. *Anal. Biochem.* 201, 119-126.
- Buck, L., and Axel, R. (1991) A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* 65, 175-187.
- Buck, L. (1996) Information coding in the vertebrate olfactory system. *Annu. Rev. Neurosci.* 19, 517-544.
- Cain, W.S. (1975) Odor intensity: mixtures and masking. *Chem. Senses Flavor* 1, 339-352.
- Collins, A., Somlyo, A.V., and Hilgemann, D.W. (1992) The giant cardiac membrane patch method: stimulation of outward Na^+ - Ca^{2+} exchange current by MgATP. *J. Physiol. (Lond.)* 454, 27-57.
- Daniel, P.C., and Derby, C.D. (1991) Mixture suppression in behavior: the antennular flick response in the spiny lobster towards binary odorant

- mixtures. *Physiol. Behav.* 49, 591-601
- Derby, C.D., Girardot, M.N., and Daniel, P.C. (1991) Responses of olfactory receptor cells of spiny lobsters to binary mixtures. I. Intensity mixture interactions. *J. Neurophysiol.* 66, 112-130.
- Duchamp, A., Revial, M.F., Holley, A. and MacLeod, P. (1974) Odor discrimination by frog olfactory receptors. *Chem. Senses and Flavor* 1, 213-233.
- Duchamp-Viret, P., Duchamp, A., and Sicard, G. (1990) Olfactory discrimination over a wide concentration range. Comparison of receptor cell and bulb neuron abilities. *Brain Res.* 517, 256-262.
- Duchamp-Viret, P., Chaput, M.A., and Duchamp, A. (1999) Odor response properties of rat olfactory receptor neurons. *Science* 284, 2171-2174.
- Fadool, D.A., and Ache, B.W. (1992) Plasma membrane inositol 1,4,5-trisphosphate-activated channels mediate signal transduction in lobster olfactory receptor neurons. *Neuron* 9, 907-918.
- Firestein, S., Darrow, B., and Shepherd, G.M. (1991) Activation of the sensory current in salamander olfactory receptor neurons depends on a G protein-mediated cAMP second messenger system. *Neuron*, 6, 825-835.
- Firestein, S., Picco, C., and Menini, (1993) A. The relation between stimulus and response in olfactory receptor cells of the tiger salamander. *J. Physiol. (Lond.)* 468, 1-10.
- Firestein, S., and Werblin, F. (1989) Odor-induced membrane currents in vertebrate-olfactory receptor neurons. *Science* 244, 79-82.
- Frings, S., and Lindemann, B. (1991) Current recording from sensory cilia of olfactory receptor cells in situ. I. The neuronal response to cyclic nucleotides. *J. Gen. Physiol.* 97, 1-16.
- Getchell, T.V. (1986) Functional properties of vertebrate olfactory receptor

- neurons. *Physiol. Rev.* 66, 772-818.
- Getchell, T.V., and Shepherd, G.M. (1978a) Responses of olfactory receptor cells to step pulses of odour at different concentrations in the salamander. *J. Physiol. (Lond.)* 282, 521-540.
- Getchell, T.V., and Shepherd, G.M. (1978b) Adaptive properties of olfactory receptors analysed with odour pulses of varying durations. *J. Physiol. (Lond.)* 282, 541-560.
- Hamill, O.P., Marty, A., Neher, E., Sakmann, B., and Sigworth, F.J. (1981) Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflugers Arch.* 391, 85-100.
- Hildebrand, J.G., and Shepherd, G.M. (1997) Mechanisms of olfactory discrimination: converging evidence for common principles across phyla. *Annu. Rev. Neurosci.* 20, 595-631.
- Hodgkin, A.L., McNaughton, P.A., Nunn, B.J., and Yau, K.W. (1984) Effect of ions on retinal rods from *Bufo marinus*. *J. Physiol. (Lond.)* 350, 649-680.
- Jinks, A., and Laing, D.G. (1999a) A limit in the processing of components in odour mixtures. *Perception*, 28, 395-404.
- Jinks, A., and Laing, D.G. (1999b) Temporal processing reveals a mechanism for limiting the capacity of humans to analyze odor mixtures. *Brain Res. Cogn. Brain Res.* 8, 311-325
- Kang, J., and Caprio, J. (1995) In vivo responses of single olfactory receptor neurons in the channel catfish *Ictalurus punctatus*. *J. Neurophys.* 73, 172-177, 1995.
- Kleene, S.J. (1992) Basal conductance of frog olfactory cilia. *Pflugers Arch.* 421, 374-80.
- Kleene, S.J. (1993) Origin of the chloride current in olfactory transduction. *Neuron* 11, 123-132.

- Kleene, S.J., and Gesteland, R.C. (1991a) Transmembrane currents in frog olfactory cilia. *J. Membr. Biol.* 120, 75-81.
- Kleene, S.J., and Gesteland, R.C. (1991b) Calcium-activated chloride conductance in frog olfactory cilia. *J. Neurosci.* 11, 3624-3629.
- Kleene, S.J., Gesteland, R.C., and Bryant, S.H. (1994) An electrophysiological survey of frog olfactory cilia. *J Exp. Biol.* 195, 307-328.
- Kramer, R.H., and Siegelbaum, S.A. (1992) Intracellular Ca^{2+} regulates the sensitivity of cyclic nucleotide-gated channels in olfactory receptor neurons. *Neuron* 9, 897-906.
- Kurahashi, T. (1989) Activation by odorants of cation-selective conductance in the olfactory receptor cell isolated from the newt. *J. Physiol. (Lond.)* 419, 177-192.
- Kurahashi, T. (1990) The response induced by intracellular cyclic AMP in isolated olfactory receptor cells of the newt. *J. Physiol. (Lond.)* 430, 355-371.
- Kurahashi, T., Lowe, G., and Gold, G.H. (1994) Suppression of odorant responses by odorants in olfactory receptor cells. *Science* 265, 118-120.
- Kurahashi, T., and Yau, K.W. (1993) Co-existence of cationic and chloride components in odorant-induced current of vertebrate olfactory receptor cells. *Nature* 363, 71-74.
- Laing D.G., Cain W.S., McBride, R.L., and Ache, B.W. (ed.) (1989a) *Perception of Complex Smells and Tastes.* Academic Press, Sydney.
- Laing D.G., Panhuber, H., and Slotnick, B.M. (1989b) Odor masking in the rat. *Physiol. Behav.* 45, 689-694.
- Laing, D.G., Panhuber, H., Willcox, M.E., and Pittman, E.A. (1984) Quality and intensity of binary odor mixtures. *Physiol. Behav.* 33, 309-319.

- Laska, M., and Hudson, R. (1993) Discriminating parts from the whole: determinants of odor mixture perception in squirrel monkeys, *Saimiri sciureus*. J. Comp. Physiol. A 173, 249-256.
- Lowe, G., and Gold, G.H. (1991) The spatial distributions of odorant sensitivity and odorant-induced currents in salamander olfactory receptor cells. J. Physiol. (Lond.) 442, 147-168.
- Lowe, G., and Gold, G.H. (1993) Nonlinear amplification by calcium-dependent chloride channels in olfactory receptor cells. Nature 366, 283-286.
- Lowe, G., Nakamura, T., and Gold, G.H. (1989) Adenylate cyclase mediates olfactory transduction for a wide variety of odorants. Proc. Natl. Acad. Sci. USA 86, 5641-5645.
- Michel, W.C., and Ache, B.W. (1992) Cyclic nucleotides mediate an odor-evoked potassium conductance in lobster olfactory receptor cells. J. Neurosci. 12,3979-3984.
- Michel, W.C., and Ache, B.W. (1994) Odor-evoked inhibition in primary olfactory receptor neurons. Chem. Senses 19, 11-24
- Michel W.C., McClintock, T.S., and Ache, B.W. (1991) Inhibition of lobster olfactory receptor cells by an odor-activated potassium conductance. J. Neurophysiol. 65, 446-453.
- Morales, B., Labarca, P., and Bacigalupo, J. (1995) A ciliary K⁺ conductance sensitive to charibdotxin underlies inhibitory responses in toad olfactory receptor neurons. FEBS Lett. 359, 41-44.
- Morales, B., Madrid, R., and Bacigalupo, J. (1997) Calcium mediates the activation of the inhibitory current induced by odorants in toad olfactory receptor neurons. FEBS Lett. 402, 259-264.
- Morales, B., Ugarte, G., Labarca, P., and Bacigalupo, J. (1994) Inhibitory K⁺

- current activated by odorants in toad olfactory neurons. Proc. R. Soc. Lond. B 257, 235-242.
- Mori, K., and Yoshihara, Y. (1995) Molecular recognition and olfactory processing in the mammalian olfactory system. Prog. Neurobiol. 45, 585-619.
- Nakamura, T., and Gold, G.H. (1987) A cyclic nucleotide-gated conductance in olfactory receptor cilia. Nature 325, 442-444.
- Nawy, S. (1999) The metabotropic receptor mGluR6 may signal through G(o), but not phosphodiesterase, in retinal bipolar cells. J. Neurosci. 19, 2938-2944.
- Ottoson, D. (1956) Analysis of the electrical activity of the olfactory epithelium. Acta Physiol. Scand. 35 (Suppl. 122) 1-83.
- Raming, K., Krieger, J., Strotmann, J., Boekhoff, I., Kubick, S., Baumstark, C., and Breer, H. (1993) Cloning and expression of odorant receptors. Nature 361, 353-356
- Reuter, D., Zierold, K., Schroder, W.H., and Frings, S. A (1998) depolarizing chloride current contributes to chemoelectrical transduction in olfactory sensory neurons in situ. J. Neurosci. 18, 6623-6630.
- Sicard, G., and Holley, A. (1984) Receptor cell responses to odorants: similarities and differences among odorants. Brain Res. 292, 283-296.
- Sklar, P.B., Anholt, R.R., and Snyder, S.H. (1986) The odorant-sensitive adenylate cyclase of olfactory receptor cells. Differential stimulation by distinct classes of odorants. J. Biol. Chem. 261, 15538-15543.
- Trotier, D. A. (1986) patch-clamp analysis of membrane currents in salamander olfactory receptor cells. Pflugers Arch. 407, 589-595.
- Vogler, C., and Schild, D. (1999) Inhibitory and excitatory responses of olfactory receptor neurons of *xenopus laevis* tadpoles to stimulation with

amino acids. *J. Exp. Biol.* 202, 997-1003.

Yamada, H., and Nakatani, K. Odor-induced hyperpolarization and suppression of cAMP-activated current in newt olfactory receptor neurons. *Chem. Senses* in press.

Zimmerman, A.L., Yamanaka, G., Eckstein, F., Baylor, D.A., and Stryer, L. (1985) Interaction of hydrolysis-resistant analogs of cyclic GMP with the phosphodiesterase and light-sensitive channel of retinal rod outer segments. *Proc. Natl. Acad. Sci. U S A* 82, 8813-8817.