

REFERENCES

- Abe, A., Koide, H., Kohno, T., and Watabe, K. (1995) A *Bacillus subtilis* spore coat polypeptide gene, *cotS*. *Microbiology* 141: 1433-1442.
- Akita, M., Sasaki, S., Matsuyama, S., and Mizushima, S. (1990) SecA interacts with secretory proteins by recognizing the positive charge at the amino terminus of the signal peptide in *Escherichia coli*. *J. Biol. Chem.* 265: 8164-8169.
- Althoff, S., Selinger, D., and Wise, J. A. (1994) Molecular evolution of SRP cycle components: functional implications. *Nucleic Acids Res.* 22: 1933-1947.
- Altman, E., Kumamoto, C. A., and Emr, S. D. (1991) Heat-shock proteins can substitute for SecB function during protein export in *Escherichia coli*. *EMBO J.* 10: 239-245.
- Bauer, T., Little, S., Stover, A. G., and Driks, A. (1999) Functional regions of the *Bacillus subtilis* spore coat morphogenetic protein CotE. *J. Bacteriol.* 181: 7043-7051.
- Bernstein, H. D., Poritz, M. A., Strub, K., Hoben, P. J., Brenner, S., and Walter, P. (1989) Model for signal sequence recognition from amino-acid sequence of 54K subunit of signal recognition particle. *Nature* 340: 482-486.
- Bochkareva, E. S., Lissin, N. M., and Girshovich, A. S. (1988) Transient association of newly synthesized unfolded proteins with the heat-shock GroEL protein. *Nature* 336: 254-257.
- Bourne, H. R., Sanders, D. A., and McCormick, F. (1990) The GTPase superfamily: a conserved

switch for diverse cell function. *Nature* 348: 125-132.

Brown, S. (1987) Mutations in the gene for EF-G reduce the requirement for 4.5S RNA in the growth of *E. coli*. *Cell* 49: 825-33.

Bunai, K., Yamada, K., Hayashi, K., Nakamura, K., and Yamane, K. (1999) Enhancing effect of *Bacillus subtilis* Ffh, a homologue of the SRP54 subunit of the mammalian signal recognition particle, on the binding of SecA to precursors of secretory proteins *in vitro*. *J. Biochem.* 125: 151-159.

Collier, D. N., Bankaitis, V. A., Weiss, J. B., and Bassford, P. J., Jr. (1988) The antifolding activity of SecB promotes the export of the *E. coli* maltose-binding protein. *Cell* 53: 273-283.

Connolly, T., and Gilmore, R. (1986) Formation of a functional ribosome-membrane junction during protein translocation requires the participation of a GTP-binding protein. *J. Cell Biol.* 103: 2253-2261.

Connolly, T., and Gilmore, R. (1989) The signal recognition particle receptor mediates the GTP-dependent displacement of SRP from the signal sequence of the nascent polypeptide. *Cell* 57: 599-610.

Connolly, T., Rapiejko, P. J., and Gilmore, R. (1991) Requirement of GTP hydrolysis for dissociation of the signal recognition particle from its receptor. *Science* 252: 1171-1173.

Cutting, S., Driks, A., Schmidt, R., Kunkel, B., and Losick, R. (1991) Forespore-specific transcription of a gene in the signal transduction pathway that governs pro-sigma K processing in *Bacillus subtilis*. *Genes Dev.* 5:456-466.

de Gier, J. W., Mansournia, P., Valent, Q. A., Phillips, G. J., Luirink, J., and von Heijne, G. (1996) Assembly of a cytoplasmic membrane protein in *Escherichia coli* is dependent on the signal recognition particle. FEBS Lett. 399: 307-309.

Dobberstein, B. (1994) On the beaten pathway. Nature 367: 599-600.

Donovan, W. P., Zheng, L., Sandman, K., and Losick, R. (1987) Genes encoding spore coat polypeptides from *Bacillus subtilis*. J. Mol. Biol. 196: 1-10.

Driks, A. (1999). *Bacillus subtilis* Spore coat. Microbiol. Mol. Biol. Rev. 63: 1-20

Driks, A., Roels, S., Beall, B., Moran, Jr., C.P., and Losick, R. (1994) Subcellular localization of proteins involved in the assembly of the spore coat of *Bacillus subtilis*. Genes Dev. 8: 234-244.

Economou, A. (1998) Bacterial pre-protein translocase: mechanism and conformational dynamics of a processive enzyme. Mol. Microbiol. 27: 511-518.

Fekkes, P., and Driessens, A. J. M. (1999) Protein targeting to the bacterial cytoplasmic membrane. Microbiol. Mol. Biol. Rev. 63: 161-173.

Gilmore, R., Walter, P., and Blobel, G. (1982) Protein translocation across the endoplasmic reticulum. II. Isolation and characterization of the signal recognition particle receptor. J. Cell Biol. 95: 470-477.

Görlich, D., Hartmann, E., Prehn, S., and Rapoport, T. A. (1992a) A protein of the endoplasmic reticulum involved early in polypeptide translocation. Nature 357: 47-52.

Görlich, D., Prehn, S., Hartmann, E., Kalies, K.-U., and Rapoport, T. A. (1992b) A mammalian homolog of SEC61p and SECYp is associated with ribosomes and nascent polypeptides during translocation. *Cell* 71: 489-503.

Hartl, F.-U., Lecker, S., Schiebel, E., Hendrick, J. P., and Wickner, W. (1990) The binding cascade of SecB to SecA to SecY/E mediates preprotein targeting to the *E. coli* plasma membrane. *Cell* 63: 269-279.

Hartl, F.-U., and Wiedmann, M. (1993) A signal recognition particle in *Escherichia coli*? *Curr. Biol.* 3: 86-89.

Haldenwang, W. G. (1995) The sigma factors of *Bacillus subtilis*. *Microbiol. Rev.* 59: 1-30.

Hayat, M. A. (1972) Basic electron microscopy techniques, p. 80. Van Nostrand Reinhold Co., New York.

Henriques, A. O., Beall, B. W., and Moran, C. P. Jr. (1997) CotM of *Bacillus subtilis*, a member of the α -crystallin family of stress proteins, is induced during development and participates in spore outer coat formation. *J. Bacteriol.* 179: 1887-1897.

Henner, D. J. (1990) Inducible expression of regulatory gene in *Bacillus subtilis*. *Methods Enzymol.* 185: 223-228.

High, S., and Doberstein, B. (1991) The signal sequence interacts with the methionine-rich domain of the 54-kD protein of signal recognition particle. *J. Cell Biol.* 113: 229-233.

Honda, K., Nakamura, K., Nishiguchi, M., and Yamane, K. (1993) Cloning and characterization of a *Bacillus subtilis* gene encoding a homologue of the 54-kilodalton subunit of mammalian signal recognition particle and *Escherichia coli* Ffh. *J. Bacteriol.* 175: 4885-4894.

Hofmeister, A. (1988) Activation of the proprotein transcription factor prosigma E is associated with its progression through three patterns of subcellular localization during sporulation in *Bacillus subtilis*. *J. Bacteriol.* 180: 2426-2433.

Hofmeister, A., Londono-Vallejo, A., Harry, E., Stragier, P., and Losick, R. (1995) Extracellular signal protein triggering the proteolytic activation of a developmental transcription factor in *Bacillus subtilis*. *J. Bacteriol.* 180:2426-2433.

Igo, M. M., and Losick, R. (1986) Regulation of a promoter that is utilized by minor forms of RNA polymerase holoenzyme in *Bacillus subtilis*. *J. Mol. Biol.* 191: 615-624.

Jeong, S. M., Yoshikawa, H., and Takahashi, H. (1993) Isolation and characterization of the *secE* homologue gene of *Bacillus subtilis*. *Mol. Microbiol.* 10: 133-142.

Joly, J. C., and Wickner, W. (1993) The SecA and SecY subunits of translocase are the nearest neighbors of a translocating preprotein, shielding it from phospholipids. *EMBO J.* 12: 255-263.

Kakeshita, H., Takamatsu, H., Amikura, R., Nakamura, K., Watabe, K., and Yamane, K. (2000a) Effect of Depletion of FtsY, a Homologue of the α Subunit of Mammalian Signal Recognition Particle Receptor, on Spore Morphology and Protein Composition of Spore Coat Layer in *Bacillus subtilis*. *FEMS letter*. In press.

Kakeshita, H., Oguro, A., Amikura, R., Nakamura, K., and Yamane, K. (2000b) Expression of

the *ftsY* gene, encoding a homologue of the α subunit of mammalian signal recognition particle receptor, is controlled by different promoters in vegetative and sporulating cells of *Bacillus subtilis*. *Microbiol.* 146: 2595-2603.

Karnovsky, M. J. (1965) A formaldehyde-glutaraldehyde fixative of high osmolarity for use in electron microscopy. *J. Cell Biol.* 27: 137A-138A.

Kumamoto, C. A., and Beckwith, J. (1985) Evidence for specificity at an early step in protein export in *Escherichia coli*. *J. Bacteriol.* 163: 267-274.

Kumamoto, C. A. (1991) Molecular chaperones and protein translocation across the *Escherichia coli* inner membrane. *Mol. Microbiol.* 5: 19-22.

Kumamoto, C. A., and Francetic, O. (1993) Highly selective binding of nascent polypeptides by an *Escherichia coli* chaperone protein in vivo. *J. Bacteriol.* 175: 2184-2188.

Kunst, F., Ogasawara, N., Moszer, I., and others. (1997) The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. *Nature* 390: 249-256.

Lill, R., Cunningham, K., Brundage, L. A., Ito, K., Oliver, D., and Wickner, W. (1989) SecA protein hydrolyzes ATP and is an essential component of the protein translocation ATPase of *Escherichia coli*. *EMBO J.* 8: 961-966.

Losick, R. and Stragier, P. (1992) Crisscross regulation of cell-type-specific gene expression during development in *B. subtilis*. *Nature* 355: 601-604

Luirink, J., and Dobberstein, B. (1994) Mammalian and *Escherichia coli* signal recognition

particles. Mol. Microbiol. 11: 9-13.

Luirink, J., ten Hagen-Jongman, C. M., van der Weijden, C. C., Oudega, B., High, S., Dobberstein, B., and Kusters, R. (1994) An alternative protein targeting pathway in *Escherichia coli*: studies on the role of FtsY. EMBO J. 13: 2289-2296

Lütcke, H. (1995) Signal recognition particle (SRP), a ubiquitous initiator of protein translocation. Eur. J. Biochem. 228: 531-550

Lütcke, H., High, S., Wood, H., Giner, A., Tollervey, D., and Dobberstein, B. (1992) The methionine-rich domain of the 54 kDa subunit of signal recognition particle is sufficient for the interaction with signal sequences. EMBO J. 11: 1543-1551.

Matsudaira, P. (1987) Sequence from picomole quantities of proteins electroblotted onto polyvinylidene difluoride membranes. J. Mol. Chem. 262: 10035-10038.

Meyer, D. I., Krause, E., and Dobberstein, B. (1982) Secretory protein translocation across membranes-the role of the 'docking protein' Nature 297: 647-650.

Micka, B., Groch, N., Heinemann, U., and Marahiel, M. A. (1991) Molecular cloning, nucleotide sequence, and characterization of the *Bacillus subtilis* gene encoding the DNA-binding protein HBsu. J. Bacteriol. 173: 3191-3198.

Miller, J. D., Bernstein, H. D., and Walter, P. (1994) Interaction of *E. coli* Ffh/4.5S ribonucleoprotein and FtsY mimics that of mammalian signal recognition particle and its receptor. Nature 367: 657-659.

Miyakawa, Y., and Komano, T. (1981) Study on the cell cycle of *Bacillus subtilis* using temperature-sensitive mutants I. Isolation and genetic analysis of the mutants defective in septum formation. *Mol. Gen. Genet.* 181: 207-214.

Moriya, S., Tsujikawa, E., Hassan, A. K., Asai, K., Kodama, T., and Ogasawara, N. (1998) A *Bacillus subtilis* gene-encoding protein homologous to eukaryotic SMC motor protein is necessary for chromosome partition. *Mol. Microbiol.* 29: 179-187

Moriyama, R., Fukuoka, H., Miyata, S., Kudoh, S., Hattori, A., Kozuka, S., Yasuda, Y., Tochikubo, K., and Makino, S. (1999) Expression of a germination-specific amidase, SleB, of bacilli in the forespore compartment of sporulating cells and its localization on the exterior side of the cortex in dormant spores. *J. Bacteriol.* 181:2373-2378.

Nakamura, K., Imai, Y., Nakamura, A., and Yamane, K. (1992) Small cytoplasmic RNA of *Bacillus subtilis*: functional relationship with human signal recognition particle 7S RNA and *Escherichia coli* 4.5S RNA. *J. Bacteriol.* 174: 2185-2192.

Nakamura, K., Nakamura, A., Takamatsu, H., Yoshikawa, H., and Yamane, K. (1990a) Cloning and characterization of a *Bacillus subtilis* gene homologous to *E. coli secY*. *J. Biochem.* 107: 603-607.

Nakamura, K., Nishiguchi, M., Honda, H., and Yamane, K. (1994) The *bacillus subtilis* SRP54 homologue, Ffh, has an intrinsic GTPase activity and forms a ribonucleoprotein complex with small cytoplasmic RNA in vivo. *Biochem. Biophys. Res. Commun.* 199:1394-1399.

Nakamura, K., Takamatsu, H., Akiyama, Y., Ito, K., and Yamane, K. (1990b) Complementation of the protein transport defect of an *Escherichia coli secY* mutant (*secY24*) by *Bacillus subtilis*

secY homologue. FEBS lett. 273: 75-78.

Nakamura, K., Yahagi, S., Yamazaki, T., and Yamane, K. (1999) *Bacillus subtilis* histone-like protein, Hbsu is an integral component of a SRP-like particle that can bind the Alu domain of small cytoplasmic RNA. J. Biol. Chem. 274: 13569-13579.

Nakane, A., Takamatsu, H., Oguro, A., Sadaie, Y., Nakamura, K., and Yamane, K. (1995) Acquisition of azide-resistance by elevated SecA ATPase activity confers azide-resistance upon cell growth and protein translocation in *Bacillus subtilis*. Microbiol. 141: 113-121.

Nicholson, W.L., and Setlow, P. (1990) Sporulation, germination and outgrowth. In: Molecular microbiological method for *Bacillus*. (Harwood, C.R., and Cutting, S.M., Eds), pp. 391-450. John Wiley and Sons, Ltd., Chichester, United Kingdom.

Nishiguchi, M., Honda, H., Amikura, R., Nakamura, K., and Yamane, K. (1994) Structural requirements of *Bacillus subtilis* small cytoplasmic RNA for cell growth, sporulation, and extracellular enzyme production. J. Bacteriol. 176: 157-165.

Nishiyama, K., Hanada, M., and Tokuda, H. (1994) Disruption of the gene encoding p12 (SecG) reveals the direct involvement and important function of SecG in the protein translocation of *Escherichia coli* at low temperature. EMBO J. 13: 3272-3277.

Oguro, A., Kakeshita, H., Honda, K., Takamatsu, H., Nakamura, K., and Yamane, K. (1995). *srB*: a *Bacillus subtilis* gene encoding a homologue of the α -subunit of the mammalian signal recognition particle receptor. DNA Research 2: 95-100.

Oguro, A., Kakeshita, H., Takamatsu, H., Nakamura, K., and Yamane, K. (1996). The effect of

Srb, a homologue of the mammalian SRP receptor α -subunit, on *Bacillus subtilis* growth and protein translocation. *Gene* 172: 17-24.

Oguro, A., Kakeshita, H., Nakamura, K., Yamane, K., Wang, W., and Bechhofer, D.H. (1998) *Bacillus subtilis* RNase III cleaves both 5'- and 3'-sites of the small cytoplasmic RNA precursor. *J. Biol. Chem.* 1998 273:19542-195427.

Phillips, G. J., and Silhavy, T. J. (1992) The *E. coli ffh* gene is necessary for viability and efficient protein export. *Nature* 359: 744-746.

Pohlschröder, M., Prinz, W. A., Hartmann, E., and Beckwith, J. (1997) Protein translocation in the three domains of life: variations on a theme. *Cell* 91:563-566.

Popham, D. L., Illades-Aguiar, B., and Setlow, P. (1995) The *Bacillus subtilis dacB* gene, encoding penicillin-binding protein 5*, is part of a three-gene operon required for proper spore cortex synthesis and spore core dehydration. *J. Bacteriol.* 177: 4721-4729.

Popham, D. L., Helin, J., Costello, C. E., and Setlow, P. (1996) Analysis of the peptidoglycan structure of *Bacillus subtilis* endospores. *J. Bacteriol.* 178: 6451-6458.

Poritz, M. A., Bernstein, H. D., Strub, K., Zopf, D., Wilhelm, H., and Walter, P. (1990) An *E. coli* ribonucleoprotein containing 4.5S RNA resembles mammalian signal recognition particle. *Science* 250: 1111-1117.

Pugsley, A. (1993) The complete general secretory pathway in gram-negative bacteria. *Microbiological Reviews* 57: 50-108.

Rapoport, T. A. (1992) Transport of proteins across the endoplasmic reticulum membrane. Science 258: 931-936.

Ribes, V., Romosch, K., Giner, A., Dobberstein, B., and Tollervey, D. (1990) *E. coli* 4.5S RNA is part of a ribonucleoprotein particle that has properties related to the signal recognition particle. Cell 63: 591-600.

Riezman, H. (1997) The ins and outs of protein translocation. Science 278:1728-1729.

Roels, S., Driks, A., and Losick, R. (1992) Characterization of *spoIVA*, a sporulation gene involved in coat morphogenesis in *Bacillus subtilis*. J. Bacteriol. 174: 575-585.

Roels, S., and Losick, R. (1995) Adjacent and divergently oriented operons under the control of the sporulation regulatory protein GerE in *Bacillus subtilis*. J. Bacteriol. 177: 6263-6275.

Römisch, K., Webb, J., Herz, J., Prehn, S., Frank, R., Vingron, M., and Dobberstein, B. (1989) Homology of 54K protein of signal-recognition particle, docking protein and two *E. coli* proteins with putative GTP-binding domains. Nature 340: 478-482.

Römisch, K., Webb, J., Lingerbach, K., Gausepohl, H., and Dobberstein, B. (1990) The 54-kD protein of signal recognition particle contains a methionine- rich RNA binding domain. J. Cell Biol. 111: 1793-1802.

Ryter, A. (1965) Etude morphologique de la sporulation de *bacillus subtilis*. Ann. Inst. Pasteur. (Paris) 108: 40-60.

Sadaie, Y., and Kada, T. (1985) *Bacillus subtilis* gene involved in cell division, sporulation, and

exoenzyme secretion. *J. Bacteriol.* 163: 648-653.

Sadaie, Y., Takamatsu, H., Nakamura, K., and Yamane, K. (1991) Sequencing reveals similarity of the wild-type *div+* gene of *Bacillus subtilis* to the *Escherichia coli secA* gene. *Gene* 98: 101-105.

Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989) Molecular cloning: a laboratory manual, 2nd ed. Cold Harbor Laboratory Press, Cold Spring Harbor, N.Y.

Sanger, F., Nicklen, S., and Coulson, A. R. (1977) DNA sequencing with chain terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74: 5463-5467.

Schaeffer, P., Millet, J., and Aubert, J. P. (1965) Catabolite repression of bacterial sporulation. *Proc. Natl. Acad. Sci. USA* 54: 704-711.

Schatz, G., and Dobberstein, B. (1996) Common principles of protein translocation across membranes. *Science* 271: 1519-1526.

Schiebel, E., Driessen, A. J. M., Hartl, F.-U., and Wickner, W. (1991) DmH⁺ and ATP function at different steps of the catalytic cycle of preprotein translocase. *Cell* 64: 927-939.

Seluanov, A., and Bibi, E. (1997) FtsY, the prokaryotic signal recognition particle receptor homologue, is essential for biogenesis of membrane proteins. *J. Biol. Chem.* 272: 2053-2055.

Serrano, M., Zilhao, E., Ricca, E., Ozin, A. J., Moran Jr., C. P., and Henriques, A. O. (1999) A *Bacillus subtilis* secreted protein with a role in endospore coat assembly and function. *J. Bacteriol.* 181: 3632-3643.

Siegel, V., and Walter, P. (1988) Binding sites of the 19-kDa and 68/72-kDa signal recognition particle (SRP) protein on SRP RNA as determined by protein-RNA "footprinting". Proc. Natl. Acad. Sci. USA 85: 1801-1805.

Stevens, C.M., Daniel, R., Illing, N., and Errington, J. (1992) Characterization of a sporulation gene, *spoIVa*, involved in spore coat morphogenesis in *Bacillus subtilis*. J. Bacteriol. 174: 586-594.

Stöver, A. G., and Driks, A. (1999) Secretion, localization, and antibacterial activity of TasA, a *Bacillus subtilis* spore-associated protein. J. Bacteriol. 181: 1664-1672

Stragier, P., and Losick, R. (1996) Molecular genetics of sporulation in *Bacillus subtilis*. Annu. Rev. Genet. 30: 297-341

Struck, J. C. R., Hartmann, R. K., Toschka, H. Y., and Erdmann, V. A. (1989) Transcription and processing of *Bacillus subtilis* small cytoplasmic RNA. Mol. Gen. Genet. 215: 478-482.

Struck, J. C. R., Vogel, D. W., Ulbrich, N., and Erdmann, V. A. (1988) The *Bacillus subtilis* scRNA is related to the 4.5S RNA from *Escherichia coli*. Nucleic Acids Res. 16: 2719.

Takamatsu, H., Imamura, A., Kodama, T., Asai, A., Ogasawara, N and Watabe, K. (2000) The *yabG* gene of *Bacillus subtilis* encodes a sporulation specific protease which is involved in processing of several spore coat proteins. FEMS Microbiol. Lett 192: 33-38.

Takamatsu, H., Fuma, S., Nakamura, K., Sadaie, Y., Shinkai, A., Matsuyama, S., Mizushima, S., and Yamane, K. (1992) In vivo and in vitro characterization of the *secA* gene product of *Bacillus*

subtilis. J. Bacteriol. 174: 4308-4316.

Takamatsu, H., Kodama, T., Nakayama, T. and Watabe, K. (1999) Characterization of the *yrbA* gene of *Bacillus subtilis* involved in the resistance and germination of spores. J. Bacteriol. 181: 4986-4994.

Takamatsu, H., Nakane, A., Sadaie, Y., Nakamura, K., and Yamane, K. (1994) The rapid degradation of mutant SecA protein in the *B. subtilis secA341* (ts) mutant causes a protein translocation defect in the cell. Biosci. Biotech. and Biochem. 58: 1845-1850.

Triglia, T., Peterson, M. G., and Kemp, D. J. (1988) A procedure for in vitro amplification of DNA segments that lie outside the boundaries of known sequence. Nucleic Acids Res. 16 : 8186.

Ulbrandt, N. D., Newitt, J. A., and Bernstein, H. D. (1997) The *E. coli* signal recognition particle is required for the insertion of a subset of inner membrane proteins. Cell 88: 187-196.

von Heijne, G. (1990a) Protein targeting signals. Curr. Opin. Cell Biol. 2: 604-608.

von Heijne, G. (1990b) The signal peptide. J. Membr. Biol. 115:195-201.

von Heijne, G. (1998) Life and death of a signal peptide. Nature 396: 111-113.

Walter, P., and Blobel., G. (1982) Signal recognition particle contains a 7S RNA essential for protein translocation across the endoplasmic reticulum. Nature 299: 691-698.

Walter, P., Ibrahimi, I., and Blobel, G. (1981) Translation of proteins across the endoplasmic reticulum. I. Signal recognition protein (SRP) binds to in-vitro-assembled polysomes

synthesizing secretory protein. *J. Cell Biol.* 91: 545-550.

Walter, P., and Johnson, A. E. (1994) Signal sequence recognition and protein targeting to the endoplasmic reticulum membrane. *Annu. Rev. Cell Biol.* 10: 87-119.

Wild, J., Altoman, E., Yura, T., and Gross, C. A. (1992) DnaK and DnaJ heat shock proteins participate in protein export in *Escherichia coli*. *Genes Dev.* 6: 1165-1172.

Wiedmann, M., Kurzchalia, T. V., Bielka, H., and Rapoport, T. A. (1987) Direct probing of the interaction between the signal sequence of nascent preprolactin and the signal recognition particle by specific cross-linking. *J. Cell Biol.* 104: 201-208.

Wolin, S. L. (1994) From the elephant to *E. coli*: SRP-dependent protein targeting. *Cell* 77: 787-790.

Yamazaki, T., Yahagi, S., Nakamura, K., and Yamane, K. (1999) Depletion of *Bacillus subtilis* histone-like protein, HBsu, causes defective protein translocation and induces upregulation of small cytoplasmic RNA. *Biochem. Biophys. Res. Commun.* 258:211-214.

Zhang, J., Ichikawa, H., Halberg, R., Kroos, L., and Aronson, A. I. (1994) Regulation of the transcription of a cluster of *Bacillus subtilis* spore coat genes. *J. Mol. Biol.* 240: 405-415.

Zhang, J., Fitz-James, P. C., and Aronson, A. I. (1993) Cloning and characterization of a cluster of genes encoding polypeptides present in the insoluble fraction of the spore coat of *Bacillus subtilis*. *J. Bacteriol.* 175: 3757-3766.

Zheng, L., Donovan, W. P., Fitz-James, P. C., and Losick, R. (1988) Gene encoding a

morphogenic protein required in the assembly of the outer coat of the *Bacillus subtilis* endospore.
Genes Dev. 2: 1047-1054.

Zheng, L., Halberg, R., Roels, S., Ichikawa, H., Kroos, L., and Losick, R. (1992) Sporulation regulatory protein GerE from *Bacillus subtilis* binds to and can activate or repress transcription from promoters for mother-cell-specific genes. J. Mol. Biol. 226: 1037-1050.

Zopf, D., Bernstein, H. D., Johnson, A. E., and Walter, P. (1990) The methionine-rich domain of the 54kd protein subunit of the signal recognition particle contains an RNA binding site and can be crosslinked to a signal sequence. EMBO J. 9: 4511-4517.

Zopf, D., Bernstein, H. D., and Walter, P. (1993) GTPase domain of the 54-kD subunit of the mammalian signal recognition particle is required for protein translocation but not for signal sequence binding. J. Cell Biol. 120: 1113-1121.