

ACKNOWLEDGEMENTS

I would like to express my sincere thanks to Prof. Kazuhisa Nakayama, Graduate School of Pharmaceutical Sciences, Kyoto University, for his valuable guidance and suggestion through this work. I would also like to thank Prof. Yoshimasa Tanaka, Institute of Biological Sciences, University of Tsukuba, for his encouragement.

I would like to express my gratitude to Prof. Hiroshi Ohno, Prof. Nobuhiro Nakamura, Dr. Hye-Won Shin, Dr. Shinichiro Yoshimura and Mr. Yusuke Yamada for their helpful comments and discussion, and Dr. Hiroyuki Takatsu, Mr. Michinari Sakurai, Miss Kaori Yoshino, Mr. Yohei Kato and Miss Hiromi Kobayashi for their technical advice.

I would like to thank Mr. Hidehisa Shimizu for his encouragement through this work.

I am also grateful for the great kind supports of the members of the laboratory; Dr. Kazuo Kasai, Dr. Kyoko Toda, Dr. Chisa Shinotsuka, Mr. Kazumasa Kawamoto, Miss Mutsumi Futatsumori, Mr. Yusaku Yoshida, Mr. Masafumi Kitamura, Mr. Hironori Taka and Miss Hiroko Mitsuhashi.

Finally, I would like to thank my family and all my friends very much.