

REFERENCES

1. Schmid, S. L. and Damke, H. (1995) Coated vesicles: a diversity of form and function. *FASEB J.* **9**, 1445-1453
2. Schekman, R. and Orci, L. (1996) Coat proteins and vesicle budding. *Science* **271**, 1526-1533
3. Lewin, D. A. and Mellman, I. (1998) Sorting out adaptors. *Biochim. Biophys. Acta* **1401**, 129-145
4. Hirst, J. and Robinson, M.S. (1998) Clathrin and adaptors. *Biochim. Biophys. Acta* **1404**, 173-193
5. Kirchhausen, T. (1999) Clathrin. *Annu. Rev. Cell Dev. Biol.* **15**, 705-732
6. Bonifacino, J. S. and Dell'Angelica, E.C. (1999) Molecular basis of the recognition of tyrosine-based sorting signals. *J. Cell Biol.* **145**, 923-926
7. Le Borgne, R., Griffiths, G. and Hoflack, B. (1996) Mannose 6-phosphate receptors and ADP-ribosylation factors. *J. Biol. Chem.* **271**, 2162-2170
8. Futter, C.E., Gibson, A., Allechin, E.H., Maxwell, S., Ruddock, L.J., Odorizzi, G., Domingo, D., Trowbridge, I.S. and Hopkins, C.R. (1998) In polarized MDCK cells basolateral vesicles arise from clathrin- γ -adaptin-coated domains on endosomal tubules. *J. Cell Biol.* **141**, 611-623
9. Mallard, F., Antony, C., Tenza, D., Salamero, J., Goud, B. and Johannes, L. (1998) Direct pathway from early/recycling endosomes to the Golgi apparatus revealed through the study of shiga toxin B-fragment transport. *J. Cell Biol.* **143**, 973-990
10. Sorkina, T., Bild, A., Tebar, F. and Sorkin, A. (1999) Clathrin, adaptors and eps15 in endosomes containing activated epidermal growth factor receptors. *J. Cell Sci.* **112**, 317-327
11. Dittié, A. S., Hajibagheri, N. and Tooze, S.A. (1996) The AP-1 adaptor complex binds to immature secretory granules from PC12 cells, and is regulated by ADP-ribosylation factor. *J. Cell Biol.* **132**, 523-536
12. Klumperman, J., Kuliawat, R., Griffith, J.M., Geuze, H.J. and Arvan, P. (1998) Mannose 6-phosphate receptors are sorted from immature secretory granules via

- adaptor protein AP-1, clathrin and syntaxin6-positive vesicles. *J. Cell Biol.* **141**, 359-371
13. Odorizzi, G., Cowles, C.R. and Emr, S.D. (1998) The AP-3 complex: a coat of many colours. *Trends Cell Biol.* **8**, 282-288
 14. Simmen, T., Honing, S., Icking, A., Tikkanen, R. and Hunziker, W. (2002) AP-4 binds basolateral signals and participates in basolateral sorting in epithelial MDCK cells *Nat. Cell Biol.* **4**, 154-159
 15. Takatsu, H., Sakurai, M., Shin, H.-W., Murakami, K. and Nakayama, K. (1998) Identification and characterization of novel clathrin adaptor-related proteins. *J. Biol. Chem.* **273**, 24693-24700
 16. Lewin, D.A., Sheff, D., Ooi, C.E., Whitney, A., Yamamoto, E., Chicione, L.M., Webster, P., Bonifacino, J. S. and Mellman, I. (1998) Cloning expression, and localization of a novel γ -adaptin-like molecules. *FEBS Lett.* **435**, 263-268
 17. Heuser, J.E. and Keen, J. (1988) Deep-etch visualization of proteins involved in clathrin assembly. *J. Cell Biol.* **107**, 877-886
 18. Ohno, H., Stewart, J., Fournier, M.-C., Bosshart, H., Rhee, I., Miyatake, S., Saito, T., Gallusser, A., Kirchhausen, T. and Bonifacino, J.S. (1995) Interaction of tyrosine-based sorting signals with clathrin-associated proteins. *Science* **269**, 1872-1875
 19. Ohno, H., Fournier, M.-C., Poy, G. and Bonifacino, J.S. (1996) Structural determinations of interaction of tyrosine-based sorting signals with the adaptor medium chains. *J. Biol. Chem.* **271**, 29009-29015
 20. Marsh, M. and McMahon, H.T. (1999) The structural era of endocytosis. *Science* **285**, 215-220
 21. Gallusser, A. and Kirchhausen, T. (1993) The β 1 and β 2 subunits of the AP complexes are the clathrin coat assembly components. *EMBO J.* **12**, 5237-5244
 22. Shih, W., Gallusser, A. and Kirchhausen, T. (1995) A clathrin-binding site in the hinge of the beta2 chain of mammalian AP-2 complexes. *J. Biol. Chem.* **270**, 31083-31090
 23. Dell'Angelica, E.C., Klumperman, J., Stoorvogel, W. and Bonifacino, J.S. (1998) Association of the AP-3 adaptor complex with clathrin. *Science* **280**, 431-434

24. Slepnev, V.I. and De Camili, P. (2000) Accessory factors in clathrin-dependent synaptic vesicle endocytosis. *Nat. Rev. Neurosci.* **1**, 161-172
25. Traub, L. (2003) Sorting it out: AP-2 and alternate clathrin adaptors in endocytic cargo selection. *J. Cell Biol.* **163**, 203-208
26. Page, L.J., Sowerby, P.J., Lui, W.W.Y., and Robinson, M.S. (1999) γ -synergin: and EH domain-containing protein that interacts with γ -adaptin. *J. Cell Biol.* **146**, 993-1004
27. Hille-Rehfeld, A. (1995) Mannose 6-phosphate receptors in sorting and transport of lysosomal enzymes. *Biochim. Biophys. Acta* **1241**, 177-194
28. Klumperman, J., Hille, A., Veenendaal, T., Oorschot, V., Stoorevogel, W., von Figura, K. and Geuze, H.J. (1993) Differences in the endosomal distributions of the two mannose 6-phosphate receptors. *J. Cell Biol.* **121**, 997-1010
29. Huang, F., Nesterov, A., Carter, R.E. and Sorokin, A. (2001) Trafficking of yellow-fluorescent-protein-tagged μ 1 subunit of clathrin adaptor AP-1 complex in living cells. *Traffic* **2**, 345-357
30. Waguri, S., Dewitte, F., Le Borgne, R., Rouille, Y., Uchiyama, Y., Dubremetz, J.F. and Hoflack, B. (2003) Visualization of TGN to endosome trafficking through fluorescently labeled MPR and AP-1 in living cells. *Mol. Biol. Cell* **14**, 142-155
31. Nakagawa, T., Setou, M., Seog, D., Ogasawara, K., Dohmae, N., Takio, K. and Hirokawa, N. (2000) A novel motor, KIF13A, transports mannose-6-phosphate receptor to plasma membrane through direct interaction with AP-1 complex. *Cell* **103**, 569-581
32. Meyer, C., Zizioli, D., Lausmann, S., Eskelinen, E.L., Hamann, J., Saftig, P., von Figura, K. and Schu, P. (2000) μ 1A-adaptin-deficient mice: lethality, loss of AP-1 binding and rerouting of mannose 6-phosphate receptors. *EMBO J.* **19**, 2193-2203
33. Poussu, A., Lohi, O. and Lehto, V.-P. (2000) Vear, a novel Golgi-associated protein with VHS and γ -adaptin "ear" domain. *J. Biol. Chem.* **275**, 7176-7183
34. Boman, A.L., Zhang, C.J., Zhu, X. and Kahn, R.A. (2000) A family of ADP-ribosylation factor effectors that can alter membrane transport through the trans-Golgi. *Mol. Biol. Cell* **4**, 1241-1255
35. Hirst, J. Lui, W.W., Bright, N.A., Totty, N.A., Seaman, M.N.J. and Robinson, M.S.

- (2000) A family of proteins with γ -adaptin and VHS domains that facilitate trafficking between the trans-Golgi network and the vacuole/lysosomes. *J. Cell Biol.* **149**, 67-79
36. Dell'Angelica, E., Puertollano, R., Mullins, C., Aguilar, R.C., Vargas, J.D., Hartnell, L.M. and Bonifacino, J.S. (2000) GGAs: a family of ADP ribosylation factor-binding proteins related to adaptors and associated with the Golgi complex. *J. Cell Biol.* **149**, 81-93
37. Takatsu, H., Yoshino, K. and Nakayama, K. (2000) Adaptor γ ear homology domain conserved in γ -adaptin and GGA proteins that interact with γ -synergin. *Biochem. Biophys. Res. Comm.* **271**,719-725
38. Puertollano, R., Randazzo, P.A., Presley, J.F., Hartnell, L.M. and Bonifacino, J.S. (2001) The GGAs promote ARF-dependent recruitment of clathrin to the TGN. *Cell* **105**, 93-102
39. Puertollano, R., Aguilar, R.C., Gorshkova, I., Crouch, R.J. and Bonifacino, J.S. (2001) Sorting of mannose 6-phosphate receptors mediated by the GGAs. *Science* **292**, 1712-1716
40. Nielsen, M.S., Madsen, P., Christensen, E.I., Nykjær, A., Gliemann, J., Kasper, D., Pohlmann, R. and Petersen, C. M. (2001) The sortilin cytoplasmic tail conveys Golgi-endosome transport and binds the VHS domain of the GGA2 sorting proteins. *EMBO J.* **20**, 2180-2190
41. Zhu, Y., Doray, B., Poussu, A., Lehto, V.-P. and Kornfeld, S. (2001) Binding of GGA2 to the lysosomal enzyme sorting motif of the mannose 6-phosphate receptor. *Science* **292**, 1716-1718
42. Takatsu, H., Katoh, Y., Shiba, Y. and Nakayama, K. (2001) Golgi-localizing, γ -adaptin ear homology domain, ADP-ribosylation factor-binding (GGA) proteins interact with acidic dileucine sequences within the cytoplasmic domains of sorting receptors through their Vps27p/Hrs/STAM (VHS) domains. *J. Biol. Chem.* **276**, 28541-28545
43. Takatsu, H., Yoshino, K., Toda, K. and Nakayama, K. (2002) GGA proteins associate with Golgi membranes through interaction between their GGAH domains and ADP-ribosylation factors. *Biochem. J.* **365**, 369-378

44. Hinners, I. and Tooze, S. A. (2003) Changing directions: clathrin-mediated transport between the Golgi and endosomes. *J. Cell Sci.* **116**, 763-771
45. Wakasugi, M., Waguri, A., Kametaka, S., Tomiyama, Y., Kanamori, S., Shiba, Y., Nakayama, K. and Uchiyama, Y. (2003) Predominant expression of the short form of GGA3 in human cell lines and tissues. *Biochem. Biophys. Res. Comm.* **306**, 687-692
46. Gonzalez, L.Jr. and Scheller, R.H. (1999) Regulation of membrane trafficking: structural insights from a Rab/effector complex. *Cell* **96**, 755-758
47. Pfeffer, S.R. (1999) Transport-vesicle targeting: tethers before SNAREs. *Nat. Cell Biol.* **1**, E17-E22
48. Olkkonen, V.M. and Stenmark, H. (1997) Role of Rab GTPases in membrane traffic. *Int. Rev. Cytol.* **176**, 1-85
49. Zerial, M. and McBride, H. (2001) Rab proteins as membrane organizers. *Nat. Rev. Mol. Cell Biol.* **2**, 107-119
50. Stenmark, H., Vitale, G., Ullrich, O. and Zerial, M. (1995) Rabaptin-5 is a direct effector of the small GTPase Rab5 in endocytic membrane fusion. *Cell* **83**, 423-432
51. Simonsen, A., Lippé, R., Christoforidis, S., Gaullier, J.-M., Brech, A., Callaghan, J., Toh, B.-H., Murphy, C., Zerial, M. and Stenmark, H. (1998) EEA1 links PI(3)K function to Rab5 regulation of endosome fusion. *Nature* **394**, 494-498
52. Christoforidis, S., McBride, H. M., Burgoyne, R. D. and Zerial, M. (1999) The Rab5 effector EEA1 is a core component of endosome docking. *Nature* **397**, 621-625
53. Christoforidis, S., Miaczynska, M., Ashman, K., Wilm, M., Zhao, L., Yip, S.-C., Waterfield, M. D., Backer, J. M. and Zerial, M. (1999) Phosphatidylinositol-3-OH kinases are Rab5 effectors. *Nat. Cell Biol.* **1**, 249-252
54. McBride, H. M., Rybin, V., Murphy, C., Giner, A., Teasdale, R. and Zerial, M. (1999) Oligomeric complexes link Rab5 effectors with NSF and drive membrane fusion via interactions between EEA1 and syntaxin13. *Cell* **98**, 377-386
55. Simonsen, A., Gaullier, J.-M., D'Arrigo, A. and Stenmark, H. (1999) The Rab5 effector EEA1 interacts directly with syntaxin6. *J. Biol. Chem.* **274**, 28857-28860
56. Vitale, G., Rybin, V., Christoforidis, S., Thornqvist, P.-Ö., McCaffrey, M., Stenmark, H.

- and Zerial, M. (1998) The Rab-binding domains mediate the interaction of Rabaptin-5 with GTP-bound Rab4 and Rab5. *EMBO J.* **17**, 1941-1951
57. Nagelkerken, B., van Anken, E., van Raak, M., Gerez, L., Mohrmann, K., van Uden, N., Holthuisen, J., Pelkmans, L. and van der Sluijs, P. (2000) Rabaptin-4, a novel effector of the small GTPase rab4a, is recruited to perinuclear recycling vesicles. *Biochem. J.* **346**, 593-601
58. Mohrmann, K. and van der Sluijs, P. (1999) Regulation of membrane transport through the endocytic pathway by rabGTPases. *Mol. Membr. Biol.* **16**, 81-87
59. Chavrier, P., van der Sluijs, P., Mishal, Z., Nagelkerken, B. and Gorvel, J.-P. (1997) Early endosome membrane dynamics characterized by flow cytometry. *Cytometry* **29**, 41-49
60. Sönnichsen, B., De Renzis, S., Nielsen, E., Rietdorf, J. and Zerial, M. (2000) Distinct membrane domains on endosomes in the recycling pathway visualized by multicolor imaging of Rab4, Rab5, and Rab11. *J. Cell Biol.* **149**, 901-913
61. Shin, H.-W., Shinotsuka, C., Torii, S., Murakami, K. and Nakayama, K. (1997) Identification and subcellular localization of a novel mammalian dynamin-related protein homologous to yeast Vps1p and Dnm1p. *J. Biochem.* **122**, 525-530
62. Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A. and Struhl, K. (1987) *Current Protocols in Molecular Biology*, pp. 13.6.2-13.6.5, John Wiley and Sons, Inc., New York
63. Neuhaus, E.M., Horstmann, H. and Almers, W. (1998) Ethane-freezing/methanol-fixation of cell monolayers: a procedure for improved preservation microscope. *J. Struct. Biol.* **121**, 326-342
64. Subramaniam, N.V., Loh, E., Horstmann, H., Habermann, A., Xu, Y., Coe, J., Griffiths, G. and Hong, W. (2000) Preferential association of syntaxin with the early endosome. *J. Cell Sci.* **113**, 997-1008
65. Klausner, R.D., Donaldson, J.G. and Lippincott-Schwartz, J. (1992) Brefeldin A: insights into the control of membrane traffic and organelle structure. *J. Cell Biol.* **116**, 1071-1080

66. Jackson, C.L. (2000) Brefeldin A revealing the fundamental principles governing membrane dynamics and protein transport. *Subcell. Biochem.* **34**, 233-272
67. Stamnes, M.A. and Rothman, J.E. (1993) The binding of AP-1 clathrin adaptor particle to Golgi membrane requires ADP-ribosylation factor, a small GTP-binding protein. *Cell* **73**, 999-1005
68. Zhu, Y., Traub, L.M. and Kornfeld, S. (1998) High affinity binding of AP-1 adaptor complex to trans-golgi network membranes devoid of mannose 6-phosphate receptors. *Mol. Biol. Cell* **9**, 1323-1337
69. Donaldson, J.G., Lippincott-Schwartz, J., Bloom, G.S., Kreis, T.E. and Klausner, R.D. (1990) Dissociation of a 110-kD peripheral membrane protein from the Golgi apparatus I an early event in brefeldin A action *J. Cell Biol.* **111**, 2295-2306
70. Mukherjee, S., Ghosh, R.N., and Maxfield, F.R. (1997) Endocytosis. *Physiol. Rev.* **77**, 759-803
71. Rubino, M., Miaczynska, M., Lippé, R. and Zerial, M. (2000) Selective membrane recruitment of EEA1 suggests role in directional transport of clathrin-coated vesicles to early endosomes. *J. Biol. Chem.* **275**, 3745-3748
72. Prekeris, R., Klumperman, J., Chen, Y.A. and Scheller, R.H. (1998) Syntaxin13 mediates cycling of plasma membrane proteins via tubulovesicular recycling endosomes. *J. Cell Biol.* **143**, 957-971
73. Bock, J.B., Klumperman, J., Davanger, S. and Scheller, R.H. (1997) Syntaxin6 functions in trans-Golgi network vesicle trafficking. *Mol. Biol. Cell* **8**, 1261-1271
74. McLauchan, H., Newell, J., Morrice, N., Osborne, A., West, M. and Smythe, E. (1997) A novel role for Rab5-GDI in ligand sequestration into clathrin-coated pits. *Curr. Biol.* **8**, 34-45
75. Gillingham, A.K., Koumanov, F., Pryor, P.R., Reaves, B.J. and Holman, G.D. (1999) Association of AP1 adaptor complexes with GLUT4 vesicles. *J. Cell Sci.* **112**, 4793-4800
76. Cormont, M., Tanti, J.-F., Zahraoui, A., Van Obberghen, E., Tavitian, A. and Le Marchand-Brustel, Y. (1993) Insulin and okadaic acid induce Rrab4 redistribution. *J.*

77. Cao, X. and Barlowe, C. (2000) Asymmetric requirements for a Rab GTPase and SNARE proteins in fusion of COPII vesicles with acceptor membranes. *J. Cell Biol.* **149**, 55-65
78. Brennwald, P. (2000) Reversal of fortune. Do Rab GTPases act on the target membrane? *J. Cell Biol.* **149**, 1-3
79. Horiuchi, H., Lippé, R., McBride, H.M., Rubino, M., Woodman, P., Stenmark, H., Rybin, V., Wilm, M., Ashman, K., Mann, M. and Zerial, M. (1997) A novel Rab5 GDP/GTP exchange factor complexed to Rabaptin-5 links nucleotide exchange to effector requirement and function. *Cell* **90**, 1149-1159
80. Umeda, A., Meyerholz, A. and Ungewickell, E. (2000) Identification of the universal cofactor (auxilin2) in clathrin coat dissociation. *Eur. J. Cell. Biol.* **79**, 336-342
81. Fernandez-Chaco, R., Achiriloaie, M., Janz, R., Albanesii, J.P. and Südhof, T.C. (2000) SCAMP1 function in endocytosis. *J. Biol. Chem.* **275**, 12752-12756
82. Costaguta, G., Stefan, C.J., Bensen, E.S., Emr, S.D. and Payne, G.S. (2001) Yeast Gga proteins function with clathrin in Golgi to endosome transport. *Mol. Biol. Cell* **12**, 1885-1896
83. Misra, S., Puertollano, R., Kato, Y., Bonifacino, J.S. and Hurley, J. H. (2002) Structural basis for acidic-cluster dileucine sorting-signal recognition by VHS domains. *Nature* **415**, 933-937.
84. Shiba, T., Takatsu, H., Nogi, T., Matsugaki, N., Kawasaki, M., Igarashi, N., Suzuki, M., Kato, R., Earnest, T., Nakayama, K. and Wakatsuki, S. (2002) Structural basis for recognition of acidic-cluster dileucine sequence by GGA1. *Nature* **415**, 937-941
85. Leslie, A.G.W. *Crystallographic Computing*, (Oxford University Press, Oxford, 1990).
86. Collaborative Computational Project, Number 4. (1994) *Acta Crystallogr. D* **50**, 760-763
87. Terwilliger, T.C. and Berendzen, J. (1999) Evaluation of macromolecular electron-density map quality using the correlation of local r.m.s. density. *Acta Crystallogr. D* **55**, 849-861
88. Terwilliger, T.C. (2000) Sigma2R, a reciprocal-space measure of the quality of

- mactomolecular electron-density maps. *Acta Crystallogr. D* **56**, 965-972
89. Perrakis, A., Morris, R. and Lamzin, V.S. (1999) Automated protein model building combined with iterative structure refinement. *Nat. Struct. Biol.* **6**, 458-446
 90. Jones, T.A., Zou, J.Y., Cowan, S.W. and Kjeldgaard, M. (1991) Improved methods for building protein models in electron density maps and the location of errors in these model. *Acta Crystallogr. A* **47**, 110-119
 91. Laskowski, R.A., MacArthur, M.W., Moss, D.S. and Thornton, J.M. (1993) PROCHECK: a program to check the stereochemical quality of protein structures. *J. Appl. Cryst.* **26**, 283-291
 92. Kraulis, P.J. (1991) MOLSCRIPT: a program to produce both detailed and schematic plots of protein structures. *J. Appl. Cryst.* **24**, 946-950
 93. Nicholls, A., Sharp, K.A. and Honig, B. (1991) Protein folding and association: insights from the interfacial and thermodynamic properties of hydrocarbons. *Proteins* **11**, 281-296
 94. Owen, D.J., Vallis, Y., Noble, M.E., Hunter, J.B., Dafforn, T.R., Evans, P.R. and McMahon, H.T. (1999) A structural explanation for the binding of multiple ligands by the alpha-adaptin appendage domain. *Cell* **97**, 805-815
 95. Traub, L.M., Downs, M.A., Westrich, J.L. and Fremont, D.H. (1999) Crystal structure of the alpha appendage of AP-2 reveals a recruitment platform for clathrin-coat assembly. *Proc. Natl. Acad. Sci. USA* **96**, 8907-8912
 96. Owen, D.J., Vallis, Y., Pearse, B.M., McMahon, H.T. and Evans, P.R. (2000) The structure and function of the β 2-adaptin appendage domain. *EMBO J.* **19**, 4216-4227
 97. Burk, O., Worpenberg, S., Haenig, B. and Klempnauer, K. H. (1997) tom-1, novel v-myb target gene expressed in AMV- and E26-transformed myelomonocytic cells. *EMBO J.* **16**, 1371-1380
 98. Hicke, L. (2001) Protein relulation by monoubiquitin. *Nat. Rev. Mol. Cell Biol.* **2**, 195-201
 99. Katzmann, D.J., Odorizzi, G. and Emr, S. D. (2002) Receptor downregulation and multivesicular-body sorting. *Nat. Rev. Mol. Cell Biol.* **3**, 893-905

100. Di Fiore, P. P., Polo, S. and Hofmann, K. (2003) When ubiquitin meets ubiquitin receptors: a signaling connection. *Nat. Rev. Mol. Cell Biol.* **4**, 491-497
101. Schnell, J.D. and Hicke, L. (2003) Non-traditional functions of ubiquitin and ubiquitin-binding proteins. *J. Biol. Chem.* **278**, 35857-35860
102. Treler, M., Staszewski, L.M. and Bohmann, D. (1994) Ubiquitin-dependent c-jun degradation in vivo is mediated by the delta domain. *Cell* **78**, 787-798
103. Fujimuro, M., Sawada, H. and Yokosawa, H. (1994) Production and characterization of monoclonal antibodies specific to multi-ubiquitin chains of polyubiquitinated proteins. *FEBS Lett.* **349**, 173-180
104. Takada, K., Nasu, H., Tsukada, Y., Ohkawa, K., Fujimuro, M., Sawada, H. and Yokosawa, H. (1995) Immunoassay for the quantification of intracellular multi-ubiquitin chains. *Eur. J. Biochem.* **233**, 42-47
105. Collins, B.M., Watson, P.J. and Owen, D.J. (2003) The structure of the GGA1-GAT domain reveals the molecular basis for ARF binding and membrane association of GGAs. *Dev. Cell* **4**, 321-332
106. Suer, S., Misra, S., Saidi, L.F. and Hurley, J.H. (2003) Structure of the GAT domain of human GGA1: a syntaxin amino-terminal domain fold in an endosomal trafficking adaptor. *Proc. Natl. Acad. Sci. USA* **100**, 4451-4456
107. Shiba, T., Kawasaki, M., Takatsu, H., Nogi, T., Matsugaki, N., Igarashi, N., Suzuki, M., Kato, R., Nakayama, K. and Wakatsuki, S. (2003) Molecular mechanism of membrane recruitment of GGA by ARF in lysosomal protein transport. *Nat. Struct. Biol.* **10**, 386-393
108. Zhu, G., Zhai, P., He, X., Terzyan, S., Zhang, R., Joachimiak, A., Tang, J. and Zhang, X.C. (2003) Crystal structure of the human GGA1 GAT domain. *Biochemistry* **42**, 6392-639
109. Shih, S.C., Katzmann, D.J., Schnell, J.D., Sutanto, M., Emr, S. D. and Hicke, L. (2002) Epsins and Vps27p/Hrs contain ubiquitin-binding domains that function in receptor endocytosis. *Nat. Cell Biol.* **4**, 389-393
110. Donaldson, K.M., Yin, H., Gekakis, N., Supek, F. and Joazeiro, C. A. P. (2002)

- Ubiquitin signals protein trafficking via interaction with a novel ubiquitin binding domain in the membrane fusion regulator, Vps9p. *Curr. Biol.* **13**, 258-262
111. Beal, R., Deveraux, Q., Rechsteiner, M. and Pickart, C. (1996) Surface hydrophobic residues of multiubiquitin chains essential for proteolytic targeting. *Proc. Natl. Acad. Sci. USA* **93**, 861-866
112. Nakatsu, F., Sakuma, M., Matsuo, Y., Arase, H., Yamasaki, S., Nakamura, N., Saito, T. and Ohno, H. (2000) A di-leucine signal in the ubiquitin moiety. Possible involvement in ubiquitination-mediated endocytosis. *J. Biol. Chem.* **275**, 26213-26219
113. Polo, S., Sigismund, S., Faretta, M., Guidi, M., Capua, M. R., Bossi, G., Chen, H., De Camilli, P. and Di Fiore, P. P. (2002) A single motif responsible for ubiquitin recognition and monoubiquitination in endocytic proteins. *Nature* **416**, 451-455
114. Shih, S.C., Prag, G., Francis, S.A., Sutanto, M.A., Hurley, J.H. and Hicke, L. (2003) A ubiquitin-binding motif required for intramolecular monoubiquitylation, the CUE domain. *EMBO J.* **22**, 1273-1281
115. Oldham, C.E., Mohney, R.P., Miller, S.L.H., Hanes, R.N. and O'Bryan, J.P. (2002) The ubiquitin-interacting motifs target the endocytic adaptor protein epsin for ubiquitination. *Curr. Biol.* **12**, 1112-1116
116. Klapisz, E., Sorokina, I., Lemeer, S., Pijnenburg, M., Verkleij, A.J. and van Bergen en Henegouwen, P. M. P. (2002) A ubiquitin-interacting motif (UIM) is essential for Eps15 and Eps15R ubiquitination. *J. Biol. Chem.* **277**, 30746-30753
117. Davies, B.A., Topp, J.D., Sfeir, A.J., Katzmann, D.J., Carney, D.S., Tall, G.G., Friedberg, A.S., Deng, L., Chen, Z. and Horazdovsky, B.F. (2003) Vps9p CUE domain ubiquitin binding is required for efficient endocytic protein traffic. *J. Biol. Chem.* **278**, 19826-19833
118. Mattera, R., Arighi, C.N., Lodge, R., Zerial, M. and Bonifacio, J.S. (2003) Divalent interaction of the GGAs with the Rabaptin-5-Rabex-5 complex. *EMBO J.* **22**, 78-88
119. Kang, R.S., Daniels, C.M., Francis, S.A., Shih, S.C., Salerno, W.J., Hicke, L. and Radhakrishnan, I. (2003) Solution structure of a CUE-ubiquitin complex reveals a conserved mode of ubiquitin binding. *Cell* **113**, 621-630

120. Prag, G., Misra, S., Jones, E.A., Ghirlando, R., Davies, B.A., Horazdovsky, B.F. and Hurley, J.H. (2003) Mechanism of ubiquitin recognition by the CUE domain of Vps9p. *Cell* **113**, 609-620
121. Wang, B., Alam, S.L., Meyer, H.H., Payne, M., Stemmler, T.L., Davis, D.R. and Sundquist, W.I. (2003) Structure and ubiquitin interactions of the conserved zinc finger domain of Npl4. *J. Biol. Chem.* **278**, 20225-20234
122. Swanson, K.A., Kang, R.S., Stemenova, S.D., Hicke, L. and Radhakrishnan, I. (2003) Solution structure of Vps27 UIM-ubiquitin complex important for endosomal sorting and receptor downregulation. *EMBO J.* **22**, 4597-4606
123. Horak, J. (2003) The role of ubiquitin in down-regulation and intracellular sorting of membrane proteins: insights from yeast. *Biochim. Biophys. Acta.* **1614**, 139-155
124. Katzmann, D.J., Babst, M. and Emr, S. (2001) Ubiquitin-dependent sorting into the multivesicular body pathway requires the function of a conserved endosomal protein sorting complex. *Cell* **106**, 145-155
125. Reggiori, F. and Pelham, H.R.B. (2001) A transmembrane ubiquitin ligase required to sort membrane proteins into multivesicular bodies. *EMBO J.* **20**, 5176-5186
126. Helliwell, S., Losko, S. and Kaiser, C.A. (2001) Components of a ubiquitin ligase complex specify polyubiquitination and intracellular trafficking of the general amino acid permease. *J. Cell Biol.* **153**, 649-662
127. Soetens, O., De Craene, J.O. and Andre, B. (2001) Ubiquitin is required for sorting to the vacuole of the yeast general amino acid permease, Gap1. *J. Biol. Chem.* **276**, 43949-43957
128. Beck, T., Schmidt, A. and Hall, M.N. (1999) Starvation induces vacuolar targeting and degradation of the tryptophan permease in yeast. *J. Cell Biol.* **146**, 1227-1237
129. Umebayashi, K. and Nakano, A. (2003) Ergosterol is required for targeting of tryptophan permease to the yeast plasma membrane. *J. Cell Biol.* **161**, 1117-1131
130. Giorgino, F., de Robertis, O., Laviola, L., Montrone, C., Perrini, S., McCowen, K.C. and Smith, R.J. (2000) The sentrin-conjugating enzyme mUbc9 interacts with GLUT4 and GLUT1 glucose transporters and regulates transporter levels in skeletal muscle cells.

Proc Natl. Acad. Sci. USA **97**, 1125-1130.

131. Lalioti, V.S., Vergarajauregui, S., Pulido, D. and Sandoval IV. (2002) The insulin-sensitive glucose transporter, GLUT4, interacts physically with Daxx. Two proteins with capacity to bind Ubc9 and conjugated to SUMO1. *J Biol. Chem.* **277**, 19783-19791.
132. Bogan, J.S., Hendon, N., McKee, A.E., Tsao, T.S. and Lodish, H.F. (2003) Functional cloning of TUG as a regulator of GLUT4 glucose transporter trafficking. *Nature* **425**,727-733.
133. Ortolan, T.G., Tongaonkar, P., Lambertson, D., Chen, L., Schaubert, C. and Madura, K. (2000) The DNA repair protein rad23 is a negative regulator of multi-ubiquitin chain assembly. *Nat. Cell Biol.* **2**, 601-608
134. Meyer, H.H., Wang, Y. and Warren, G. (2002) Direct binding of ubiquitin conjugates by the mammalian p97 adaptor complexes, p47 and Ufd1-Npl4. *EMBO J.* **21**, 5645-5652
135. Cohen, M., Stutz, F. and Dargemont, C. (2003) Deubiquitination, a new player in Golgi to endoplasmic reticulum retrograde transport. *J. Biol Chem.* **278**, 51989-51992.
136. Cohen, M., Stutz, F., Belgareh, N., Haguenaer-Tsapis, R. and Dargemont, C. (2003) Ubp3 requires a cofactor, Bre5, to specifically de-ubiquitinate the COPII protein, Sec23. *Nat. Cell Biol.* **5**, 661-667
137. Shenoy, S.K., McDonald, P.H., Kohout, T.A. and Lefkowitz, R.J. (2001) Regulation of receptor fate by ubiquitination of activated beta 2-adrenergic receptor and beta-arrestin. *Science* **294**, 1307-1313
138. Shenoy, S.K. and Lefkowitz, R.J. (2003) Trafficking patterns of beta-arrestin and G protein-coupled receptors determined by the kinetics of beta-arrestin deubiquitination. *J. Biol. Chem.* **278**,14498-14506
139. Duncan, M.C. and Payne, G.S. (2003) ENTH/ANTH domains expand to the Golgi. *Trends Cell Biol.* **13**, 211-215
140. Doray, B., Ghosh, P., Griffith, J., Geuze, H.J. and Kornfeld, S. (2002) Cooperation of GGAs and AP-1 in packaging MPRs at the trans-Golgi network. *Science* **297**,1700-

141. Hirst, J., Lindsay, M.R. and Robinson, M.S. (2001) GGAs: roles of the different domains and comparison with AP-1 and clathrin. *Mol. Biol. Cell* **12**, 3573-3588
142. Black, M.W. and Pelham, H.R. (2000) A Selective Transport Route from Golgi to Late Endosomes that Requires the Yeast GGA Proteins. *J. Cell Biol.* **151**, 587-600
143. Ha, S.A., Torabinejad, J., DeWald, D.B., Wenk, M.R., Lucast, L., De Camilli, P., Newitt, R.A., Aebersold, R. and Nothwehr, S.F. (2003) The synaptojanin-like protein Inp53/Sjl3 functions with clathrin in a yeast TGN-to-endosome pathway distinct from the GGA protein-dependent pathway. *Mol. Biol. Cell* **14**, 1319-1333.
144. Miller, G.J., Mattera, R., Bonifacino, J.S. and Hurley, J.H. (2003) Recognition of accessory protein motifs by the gamma-adaptin ear domain of GGA3. *Nat. Struct. Biol.* **8**, 599-606
145. Collins, B.M., Praefcke, G.J., Robinson, M.S. and Owen, D.J. (2003) Structural basis for binding of accessory proteins by the appendage domain of GGAs. *Nat. Struct. Biol.* **8**, 607-613