

Acknowledgments

I would like to express my most sincere thanks to Dr. Ryuichi Ikeda, Professor of University of Tsukuba, for his valuable suggestions filled with his deep insight into science. Under his guidance, I could learn experimental as well as theoretical knowledge on NMR, NQR, solid state physics and the other related fields.

I am very grateful to Dr. Hiroyuki Ishida in Okayama University and Dr. Sadamu Takeda in Hokkaido University for helpful discussions and suggestions. I also thank to Dr. Hideta Ishihara in Saga University for measurements of ^{35}Cl NQR frequency.

I am much obliged to Dr. Hiroshi Kitagawa and Dr. Shin'ichi Ishimaru in University of Tsukuba for their numerous help and valuable advice.

I am very grateful to Dr. Riki Seki, Dr. Hitoshi Shoji in University of Tsukuba, Dr. Isoroku Nagasawa in Fukuoka University of Education, and Dr. Noriko Yamamuro in Tokyo Denki University, for their appropriate advice and help.

Thanks are given to Mr. Hiroshi Miyoshi and Mr. Satoshi Takahashi for their assistance in the experimental work. I am thankful to my seniors, juniors and friends for their kindness.

I greatly thank my grand mother, parents and sister for their affection and patient encouragement.

It remains for me to thank Ms. Miho Yamauchi for her tender heart.